

CENTRAL APPROXIMATION PROCESSES

To Prof. Dr. Hellmuth Kneser, on his 75th birthday, 16 April, 1973

BERND DRESELER AND WALTER SCHEMPF

(Received January 19, 1973)

1. Introduction. The first investigations which are concerned with the saturation behaviour of approximation processes of the convolution type on general locally compact *abelian* topological groups are due to H. Buchwalter [7]. His brief sketch was followed by the recent papers [17, 18], [42] which contain a detailed and extended treatment of the locally compact abelian case by means of the quasimeasure concept. Inspired by the results of S. Pawelke [35], the study of saturation theory on compact (*not necessarily abelian*) groups was initiated by the junior author [16]. Some of the results of his thesis were announced in [15]. The methods employed in these papers are based on the Peter-Weyl decomposition theorem and remain to a certain extent in the *algebraic* domain.

On the other hand the constructive approximation theory on Lie groups has made some progress in the past years. In this connection we shall refer the reader to the papers by P. L. Butzer-H. Johnen [9], H. Johnen [25-27], R. A. Mayer [29-33] and D. L. Ragozin [37-40]. It is the primary objective of the present paper to apply the general methods as developed in [16] to the study of *the saturation behaviour of central approximation processes on compact Lie groups G* . In particular, the approximations by the heat-diffusion semigroup and by the Poisson semigroup which both are, as is well known, closely connected with hypoelliptic partial differential equations, will be treated (Sections 4 and 5). Since it is shown in [15, 16] that the techniques to deduce our saturation theorem for central approximation processes on compact Lie groups (Theorems 2 and 3 in Section 3 *infra*) apply also for zonal approximation processes on compact homogeneous spaces associated with Riemannian symmetric pairs, we shall deal in Section 6 with the interplay between zonal and central approximation processes on the compact Euclidean unit sphere S_3 . Finally, Section 7 is devoted to an investigation of the approximation processes of the de La Vallée Poussin type and of the Fejér type on the special unitary group $SU(2)$.

Acknowledgment. We record our gratitude to Prof. Dr. H. Ehlich, Ruhr-Universität Bochum, for his support of our joint work.

2. Notation, Terminology and Preliminaries. Throughout this paper G will denote a *real compact connected Lie group* with neutral element $1 \in G$ and Haar measure ν which is assumed to be normalized by $\nu(G) = 1$. Let $\mathcal{E}(G) = \bigcap_{m \geq 0} \mathcal{E}_c^m(G)$ be the complex vector space of all infinitely differentiable complex-valued functions on G . We shall provide $\mathcal{E}(G)$ with the topology of uniform convergence of the functions and of all their derivatives. Then the strong topological dual $\mathcal{E}'(G)$ of the Fréchet space $\mathcal{E}(G)$ consists of all *complex distributions* on G . The space $\mathcal{E}'(G)$ is a complex locally convex topological algebra with respect to the (jointly continuous) convolution product

$$(R, S) \rightsquigarrow R * S$$

of distributions. Moreover, the Dirac measure ε_1 forms the neutral element of $\mathcal{E}'(G)$ and the mapping

$$S \rightsquigarrow S^*$$

which is the transpose of the continuous linear mapping

$$\mathcal{E}(G) \ni f \rightsquigarrow (f^*: G \ni x \rightsquigarrow \bar{f}(x^{-1}) \in \mathbb{C}) \in \mathcal{E}(G),$$

is an involution of $\mathcal{E}'(G)$. The involutive topological \mathbb{C} -algebra $\mathcal{E}'(G)$ is said to be *the group algebra* of G .

In the sequel, $\mathcal{M}(G) = \mathcal{E}'^0(G)$ will denote the involutive Banach algebra of all complex Radon measures on G under convolution and $\mathcal{N}(G)$ will stand for the complex involutive Banach algebras $L^p(G) = L^p(G; \nu)$, $p \in \llbracket 1, +\infty \llbracket$, and $\mathcal{E}^0(G)$ simultaneously. Thus the injections of the sequence

$$\mathcal{N}(G) \rightarrow \mathcal{M}(G) \rightarrow \mathcal{E}'(G)$$

are monomorphisms with respect to the category of involutive topological \mathbb{C} -algebras. It will be convenient to denote the canonical norms of $\mathcal{N}(G)$ and $\mathcal{M}(G)$ by the same symbol $\|\cdot\|$.

It is well known (J. Dieudonné [14]) that $L^2(G)$ is a complex (complete) Hilbert algebra with respect to the standard scalar product. In view of the spectral theorem of Ambrose-Gurevič let the external Hilbert sum

$$(1) \quad L^2(G) = \bigoplus_{\lambda \in A(G)} \alpha_\lambda$$

be *the spectral decomposition* of $L^2(G)$. Using the picturesque terminology of J. Dieudonné [13] for some ring theoretic concepts, *the feet* $\{\alpha_\lambda \mid \lambda \in A(G)\}$

are minimal self-adjoint two-sided ideals of $L^2(G)$ and also of $\mathcal{M}(G)$. Moreover, each foot $\alpha_\lambda (\lambda \in A(G))$ is a lame (complete) Hilbert subalgebra of $L^2(G)$ which has a (self-adjoint) *identity element* u_λ and the (finite) dimension

$$\dim_C \alpha_\lambda = u_\lambda(1) = n_\lambda^2 \quad (\lambda \in A(G)) .$$

Here the natural number

$$n_\lambda = \|u_\lambda\| = \text{long}_{L^2(G)}(\alpha_\lambda)$$

denotes the *longitude* of the foot α_λ . In the following α_{λ_0} will always denote the *trivial* foot of dimension $n_{\lambda_0}^2 = 1$ which consists of the constant complex-valued functions on G . It should be observed that *the socle* of the ring $L^2(G)$, i.e. the (algebraic) direct sum

$$\mathfrak{S}_{L^2(G)} = \coprod_{\lambda \in A(G)} \alpha_\lambda ,$$

which is a vector subspace of $\mathcal{E}(G)$, is an everywhere norm dense two-sided ideal of $\mathcal{A}(G)$. Moreover, according to the Peter-Weyl theorem, there exists a bijection η of the set $\{\alpha_\lambda \mid \lambda \in A(G)\}$ onto the dual object \hat{G} of G such that *the group character*

$$\chi_\lambda = \frac{1}{n_\lambda} u_\lambda$$

associated with the foot α_λ and the equivalence class $\eta(\alpha_\lambda) \in \hat{G}$ of continuous irreducible unitary representations of G are interrelated according to the equation

$$\text{Tr} (\eta(\alpha_\lambda)) = \bar{\chi}_\lambda$$

for any parameter $\lambda \in A(G)$.

Choose an orientation on G and equip G with a (fixed) Riemannian structure such that the associated metric g is invariant by both right and left translations. Since G may be identified with the Riemannian homogeneous manifold $G \times G$ modulo the diagonal of $G \times G$ such a choice is indeed possible. Let $\Delta = \Delta_g$ be the Laplace-Beltrami operator on G with respect to the Riemannian metric g . Then

$$\Delta: \mathcal{E}(G) \ni f \rightsquigarrow \text{div}_g (\text{grad}_g f) = * d * df$$

where $*$ denotes the Hodge star with respect to the oriented Riemannian structure of G . It is a well known consequence of Schur's lemma that (1) represents a discrete spectral decomposition of $L^2(G)$ with respect to the self-adjoint positive-definite elliptic differential operator $-\Delta$. For the corresponding eigenvalues $\zeta_\lambda (\lambda \in A(G))$ we have

$$\zeta_{\lambda_0} = 0, \quad \zeta_{\lambda} > 0 \quad \text{for } \lambda \in \Lambda(G) \setminus \{\lambda_0\}$$

and the countable family

$$\text{Spec}(G, \mathfrak{g}) = \{\zeta_{\lambda} \mid \lambda \in \Lambda(G)\}$$

is said to be *the (discrete) spectrum* of the compact connected oriented Riemannian manifold (G, \mathfrak{g}) . In particular, the space of all harmonic functions $G \rightarrow \mathbb{C}$ coincides with the trivial foot α_{λ_0} .

3. Approximation Processes on $\mathcal{A}(G)$. Let T be a non-empty directed set of parameters. A family $(I_t)_{t \in T}$ of continuous \mathbb{C} -linear mappings $I_t: \mathcal{A}(G) \rightarrow \mathcal{A}(G)$ is said to form an *approximation process* on the complex Banach space $\mathcal{A}(G)$ if it converges pointwise with respect to the section filter of the directed set T towards the identity automorphism of $\mathcal{A}(G)$, i.e. if

$$\lim_{t \in T} \|I_t(f) - f\| = 0$$

holds for each function $f \in \mathcal{A}(G)$.

As usual $\mathbb{R}_+^* =]0, +\infty[= \mathbb{R}_+ \setminus \{0\}$ will denote the open positive real half-line.

DEFINITION 1. The approximation process $(I_t)_{t \in T}$ on $\mathcal{A}(G)$ is said to have *the saturation structure* $(\varphi; \mathcal{A}(G); V)$ iff the following conditions are satisfied:

(I) There exists a mapping $\varphi: T \rightarrow \mathbb{R}_+^*$ fulfilling $\lim_{t \in T} \varphi(t) = 0$ such that $f \in \mathcal{A}(G)$ and

$$(2) \quad \|I_t(f) - f\| = o(\varphi(t)) \quad (t \in T)$$

imply

$$f \in \alpha_{\lambda_0}.$$

(II) There exists a vector subspace V of $\mathcal{A}(G)$ such that $f \in \mathcal{A}(G)$ and the condition

$$(3) \quad \|I_t(f) - f\| = O(\varphi(t)) \quad (t \in T)$$

imply $f \in V$.

(III) Conversely, if $f \in V$ then f satisfies the condition (3).

In this case, V is called *the Favard space* of the saturation structure $(\varphi; \mathcal{A}(G); V)$.

After having reviewed these general concepts, let us now turn to an important special class of approximation processes (of the convolution type) on $\mathcal{A}(G)$. We shall agree to retain the preceding notations. In addition, denote by $\mathcal{ZM}(G)$ *the center* of the complex involutive Banach

algebra $\mathcal{M}(G)$ so that $\mathcal{Z}\mathcal{M}(G)$ is a closed involutive subalgebra of $\mathcal{M}(G)$. Observe that $\mathcal{A}(G)$ is in particular a Banach $\mathcal{Z}\mathcal{M}(G)$ -module.

DEFINITION 2. A family $(\mu_t)_{t \in T}$ of complex Radon measures which belong to the convolution algebra $\mathcal{Z}\mathcal{M}(G)$ is said to form a *central approximate unit for $\mathcal{A}(G)$* if the relations

$$\mu_t(G) = 1 \text{ for any } t \in T$$

and

$$\lim_{t \in T} \|\mu_t * f - f\| = 0$$

hold for any function $f \in \mathcal{A}(G)$.

Define for any parameter $t \in T$ the continuous linear mapping

$$I_t: \mathcal{A}(G) \ni f \mapsto \mu_t * f \in \mathcal{A}(G).$$

Then $(I_t)_{t \in T}$ is said to be a *central approximation process which is generated on $\mathcal{A}(G)$ by the approximate unit $(\mu_t)_{t \in T}$* .

REMARK 1. Since $\mathcal{A}(G)$ is a Banach $L^1(G)$ -module, any central approximate unit for $L^1(G)$ is also a central approximate unit for $\mathcal{A}(G)$. See E. Hewitt-K. A. Ross [22], Chapter VIII.

Before presenting a general saturation theorem for central approximation processes, we shall make a few observations and introduce some further concepts. Since each foot a_λ is a two-sided ideal of the complex Banach algebra $\mathcal{M}(G)$ it follows that $\mathcal{Z}\mathcal{M}(G) * u_\lambda$ is a vector subspace of the center $\mathcal{Z}a_\lambda$ for each $\lambda \in \Lambda(G)$. On the other hand, since a_λ is a *simple* complex algebra, its center $\mathcal{Z}a_\lambda$ is isomorphic with the field \mathbb{C} . Denoting by X the Gelfand functor, i.e. the contravariant functor from the category of complex commutative Banach algebras with identity to the category of compact topological spaces which assigns to each object of the first category its space of characters, we obtain the following

THEOREM 1. For any $\lambda \in \Lambda(G)$ define

$$c_\lambda: \mathcal{Z}\mathcal{M}(G) \ni \mu \mapsto \frac{1}{n_\lambda^2} \int_G \bar{u}_\lambda(x) d\mu(x) \in \mathbb{C}.$$

Then we have $c_\lambda \in X(\mathcal{Z}\mathcal{M}(G))$ and the identity

$$\mu * u_\lambda = c_\lambda(\mu) \cdot u_\lambda$$

obtains for any measure $\mu \in \mathcal{Z}\mathcal{M}(G)$.

REMARK 2. Y. Kawada [28] has shown that the identity

$$X(\mathcal{Z}L^1(G)) = \{c_\lambda \mid \mathcal{Z}L^1(G)\} \mid \lambda \in \Lambda(G)\}$$

obtains. See also E. Hewitt-K. A. Ross [22], Chapter VII.

DEFINITION 3. A central approximate unit $(\mu_t)_{t \in T}$ for $\mathcal{A}(G)$ is said to be a (central) *pro-saturation measure of type* $(\varphi; \psi)$ with respect to $\mathcal{A}(G)$ iff the following requirements are satisfied:

There exists a function $\varphi: T \rightarrow \mathbf{R}_+^*$ satisfying $\lim_{t \in T} \varphi(t) = 0$ and a function $\psi: A(G) \rightarrow \mathbf{C}$ where

$$\psi(\lambda) \neq 0 \quad \text{for } \lambda \in A(G) \setminus \{\lambda_0\}$$

in such a way that

$$\lim_{t \in T} \frac{c_\lambda(\mu_t) - 1}{\varphi(t)} = \psi(\lambda)$$

holds for all $\lambda \in A(G)$.

REMARK 3. Let K be a closed subgroup of G , with normalized Haar measure ν_K , such that (G, K) forms a Riemannian *symmetric pair*. See, for instance, R. R. Coifman-G. Weiss [12]. Then (G, K) is a Gelfand pair, i.e. the convolution algebra $\mathcal{M}(K \backslash G / K)$ of all complex Radon measures on G which are bi-invariant with respect to K , is a commutative involutive Banach algebra with the identity element ν_K . Denote by

$$L^2(G/K) = \bigoplus_{\lambda \in A_K(G)} \mathfrak{h}_\lambda(G/K)$$

the spectral decomposition of the complex Hilbert space $L^2(G/K)$ which is deduced from (1) and by $(\omega_\lambda)_{\lambda \in A_K(G)}$ the family of associated *zonal spherical functions*. Then the mapping

$$c_\lambda: \mathcal{M}(K \backslash G / K) \ni \mu \mapsto \frac{1}{n_\lambda} \int_G \bar{\omega}_\lambda(x) d\mu(x) \in \mathbf{C}$$

is an element of the space $X(\mathcal{M}(K \backslash G / K))$ for any $\lambda \in A_K(G)$ and we have

$$\mu * \omega_\lambda = c_\lambda(\mu) \cdot \omega_\lambda$$

for all measures $\mu \in \mathcal{M}(K \backslash G / K)$.

Thus, by a generalization of Definition 3, the concept of *zonal pro-saturation measure* and, similarly, of *zonal approximation process* can be considered for symmetric pairs. Details of a saturation theory which is generalized in this direction may be found in [16] and [19]. Clearly, if the subgroup K is the diagonal of $G \times G$, the notion of central pro-saturation measure for G and of zonal pro-saturation measure for $(G \times G)/K$ coalesce. We shall return to this question in Example (ii) of Section 6 infra.

We shall assign to $\mathcal{A}(G)$ a complex Banach space $\mathcal{B}(G)$ according

to the following rule:

$$\mathcal{B}(G) = \begin{cases} \mathcal{E}^0(G) & \text{if } \mathcal{A}(G) = L^1(G); \\ L^1(G) & \text{if } \mathcal{A}(G) = \mathcal{E}^0(G); \\ L^{p'}(G) & \text{if } \mathcal{A}(G) = L^p(G) \text{ and } p \in]1, +\infty[. \end{cases}$$

As usual, p' denotes the dual exponent of p . Under these circumstances, we shall refer to $(\mathcal{A}(G), \mathcal{B}(G))$ as an *adapted pair*. Bearing in mind this terminology, we establish the following saturation theorem.

THEOREM 2. *Let $(I_t)_{t \in T}$ be an approximation process which is generated on $\mathcal{A}(G)$ by the central pro-saturation measure $(\mu_t)_{t \in T}$ of type $(\varphi; \psi)$ with respect to $\mathcal{A}(G)$. Suppose that $(\mathcal{A}(G), \mathcal{B}(G))$ is an adapted pair and define*

$$\mathcal{V}_\psi(\mathcal{A}(G)) = \{f \in \mathcal{A}(G) \mid \psi(\lambda) \cdot u_\lambda * f = u_\lambda * \rho, \rho \in \mathcal{B}'(G), \lambda \in A(G)\}.$$

Then $(I_t)_{t \in T}$ has the saturation structure $(\varphi; \mathcal{A}(G); V)$ where the Favard space V satisfies the condition

$$V \subset \mathcal{V}_\psi(\mathcal{A}(G)).$$

PROOF. (I) Suppose that the function $f \in \mathcal{A}(G)$ satisfies the condition (2), i.e. that

$$\lim_{t \in T} \frac{\|\mu_t * f - f\|}{\varphi(t)} = 0$$

holds. Then we obtain

$$\lim_{t \in T} \frac{c_\lambda(\mu_t) - 1}{\varphi(t)} \cdot u_\lambda * f = 0,$$

hence

$$\psi(\lambda) \cdot u_\lambda * f = 0$$

for any $\lambda \in A(G)$. It follows $u_\lambda * f = 0$ for $\lambda \in A(G) \setminus \{\lambda_0\}$. Let $g = f - u_{\lambda_0} * f$. Then $g \in L^1(G)$ and $u_\lambda * g = 0$ for each index $\lambda \in A(G)$. A familiar limiting argument yields $g = 0$. Hence $f = u_{\lambda_0} * f \in \alpha_{\lambda_0}$.

(II) Suppose that $f \in \mathcal{A}(G)$ satisfies the condition (3). For brevity, define the function

$$Q_t(f) = \frac{1}{\varphi(t)} (\mu_t * f - f) \in \mathcal{A}(G)$$

for any $t \in T$. The task will be to show that the existence of a section S of the directed set T and the existence of the upper bound

$$\sup_{t \in S} \|Q_t(f)\| < +\infty$$

imply the existence of an element $\rho \in \mathcal{B}'(G)$ which verifies the identity

$$(4) \quad \psi(\lambda) \cdot u_\lambda * f = u_\lambda * \rho$$

for each $\lambda \in A(G)$. We provide $\mathcal{B}'(G)$ with the strong dual topology $\beta(\mathcal{B}'(G), \mathcal{B}(G))$. Since $(\mathcal{A}(G), \mathcal{B}(G))$ is an adapted pair, there exists a natural isometric isomorphism of the complex involutive Banach algebra $\mathcal{A}(G)$ onto a closed subalgebra of $\mathcal{B}'(G)$. Let $\mathcal{A}(G)$ be identified with its image under this isomorphism. Owing to the Alaoglu theorem, there exists a cofinal subset T_0 of S and an element $\rho \in \mathcal{B}'(G)$ such that

$$\lim_{t \in T_0} Q_t(f) = \rho$$

holds with respect to the weak dual topology $\sigma(\mathcal{B}'(G), \mathcal{B}(G))$. In particular, for any $\lambda \in A(G)$, the identity

$$(5) \quad \lim_{t \in T_0} u_\lambda * Q_t(f) = u_\lambda * \rho$$

holds with respect to the topology of pointwise convergence. On the other hand, the equality

$$(6) \quad \lim_{t \in T_0} u_\lambda * Q_t(f) = \psi(\lambda) \cdot u_\lambda * f$$

obtains pointwise for any $\lambda \in A(G)$. Combining (5) and (6) we obtain the desired identity (4).

This completes the proof of the saturation theorem.—

REMARK 4. Theorems 4, 5 and Theorems 7, 8 *infra* will show that, in general, the “set theoretic estimate” $\mathcal{V}_\psi(\mathcal{A}(G))$ for the Favard space V can *not* be improved.

In order to establish a further characterization of the space $\mathcal{V}_\psi(\mathcal{A}(G))$, we shall recall the concept of relative completion in the sense of E. Gagliardo [21], N. Aronszajn-E. Gagliardo [1] and H. Berens [4].

Let $(X; \|\cdot\|_X)$, $(Y; \|\cdot\|_Y)$ be complex Banach spaces. Suppose that the inclusion $X \subset Y$ holds and that the canonical injection $X \hookrightarrow Y$ is continuous and has norm ≤ 1 . The completion $[X, Y]_0$ of X relative to Y is the complex vector space of all limits with respect to the norm topology of Y of those sequences $(x_n)_{n \geq 1}$ in X which are bounded with respect to $\|\cdot\|_X$. The natural norm

$$[X, Y]_0 \ni y \rightsquigarrow \inf_{(x_n)_{n \geq 1}} \{ \sup_{n \geq 1} \|x_n\|_X \mid \lim_{n \geq 1} \|y - x_n\|_Y = 0 \} = \|y\|_0$$

turns $[X, Y]_0$ into a complex Banach space. It is to be emphasized that

the identity $[X, Y]_0 = X$ holds in the case when the space X is reflexive.

Let $(\mathcal{A}(G), \mathcal{B}(G))$ be an adapted pair and $\psi: A(G) \rightarrow C$ be any mapping. The graph norm

$$\mathcal{V}_\psi(\mathcal{A}(G)) \ni f \mapsto \|f\|_{\mathcal{V}} = \|f\| + \|\rho\|$$

where $\rho \in \mathcal{B}'(G)$ is the unique element associated with f according to the equations

$$\psi(\lambda) \cdot u_\lambda * f = u_\lambda * \rho \quad (\lambda \in A(G)),$$

turns $\mathcal{V}_\psi(\mathcal{A}(G))$ into a complex Banach space. In a similar way, define the complex vector space

$$\mathcal{W}_\psi(\mathcal{A}(G)) = \{f \in \mathcal{A}(G) \mid \psi(\lambda) \cdot u_\lambda * f = u_\lambda * g, g \in \mathcal{A}(G), \lambda \in A(G)\}.$$

Then $\mathcal{W}_\psi(\mathcal{A}(G))$ is a Banach space when equipped with the graph norm

$$\mathcal{W}_\psi(\mathcal{A}(G)) \ni f \mapsto \|f\|_{\mathcal{W}} = \|f\| + \|g\|.$$

Now $\mathcal{V}_\psi(\mathcal{A}(G))$ admits the following characterization:

THEOREM 3. *Retain the above notations and assumptions - then the identity*

$$\mathcal{V}_\psi(\mathcal{A}(G)) = [\mathcal{W}_\psi(\mathcal{A}(G)), \mathcal{A}(G)]_0$$

obtains.

PROOF. Let us suppose that $\mathcal{A}(G) = L^1(G)$ holds and let $f \in \mathcal{V}_\psi(L^1(G))$ be fixed. In view of a result due to E. Hewitt-K. A. Ross [22], Chapter VII, it is possible to construct a central approximate unit $(h_n \nu)_{n \geq 1}$ for $L^1(G)$ such that $h_n \in \mathcal{E}_{L^2(G)}$ and $\|h_n \nu\| = 1$ for any integer $n \geq 1$. Let $\rho \in \mathcal{M}(G)$ be the Radon measure associated with f such that $\psi(\lambda) u_\lambda * f = u_\lambda * \rho$ holds for any $\lambda \in A(G)$. This being so, it then follows that

$$h_n * f \in \mathcal{W}_\psi(L^1(G)) \quad (n \geq 1);$$

$$\sup_{n \geq 1} \|h_n * f\|_{\mathcal{W}} \leq \|f\| + \|\rho\|;$$

$$\lim_{n \rightarrow \infty} \|h_n * f - f\| = 0.$$

Thus $f \in [\mathcal{W}(L^1(G)), L^1(G)]_0$.

Conversely, let $f \in [\mathcal{W}(L^1(G)), L^1(G)]_0$ be given. Choose sequences $(f_n)_{n \geq 1}$ and $(g_n)_{n \geq 1}$ in the space $L^1(G)$ such that

$$\lim_{n \rightarrow \infty} \|f_n - f\| = 0;$$

$$\psi(\lambda) \cdot u_\lambda * f_n = u_\lambda * g_n \quad (n \geq 1, \lambda \in A(G));$$

$$\sup_{n \geq 1} \|f_n\|_{\mathcal{W}} = \sup_{n \geq 1} (\|f_n\| + \|g_n\|) < +\infty.$$

By virtue of the Alaoglu theorem, the subsets of $\mathcal{M}(G)$ which are bounded with respect to the strong dual topology are relatively sequentially compact with respect to the vague topology. This fact implies the existence of a subsequence $(g_{n_k}\nu)_{k \geq 1}$ of the sequence $(g_n\nu)_{n \geq 1}$ and of a Radon measure $\rho \in \mathcal{M}(G)$ such that

$$\lim_{k \rightarrow \infty} g_{n_k}\nu = \rho$$

holds with respect to the vague topology of $\mathcal{M}(G)$. Consequently, the equations

$$\lim_{k \rightarrow \infty} u_\lambda * g_{n_k} = u_\lambda * \rho \quad (\lambda \in A(G))$$

and

$$\lim_{k \rightarrow \infty} \psi(\lambda)u_\lambda * f_{n_k} = u_\lambda * \rho \quad (\lambda \in A(G))$$

hold with respect to the topology of pointwise convergence. In particular

$$\psi(\lambda)u_\lambda * f = u_\lambda * \rho$$

obtains for any $\lambda \in A(G)$, i.e. $f \in \mathcal{V}_\psi(L^1(G))$ and $\|f\|_0 = \|f\|_\psi$.

Summing up, we have proved the equality $\mathcal{V}_\psi(\mathcal{A}(G)) = [\mathcal{W}_\psi(\mathcal{A}(G)), \mathcal{A}(G)]_0$ in the case $\mathcal{A}(G) = L^1(G)$. If $\mathcal{A}(G) = \mathcal{E}^0(G)$, the proof follows in a similar fashion. Finally, for $\mathcal{A}(G) = L^p(G)$ and $p \in]1, +\infty[$, the statement of the theorem is immediate from the reflexivity of the spaces $L^p(G)$.—

In the special case when $G = T$ is the one-dimensional torus group, a proof of Theorem 3 based on the convergence theorem for Fejér means may be found in the monograph by P. L. Butzer-R. J. Nessel [10], Chapter 10.

4. The Heat-Diffusion Approximation Process. Let us agree to retain the notations and conventions introduced in the preceding sections; in particular \mathbf{R}_+^* denotes the open positive real half-line, equipped with the induced Lebesgue measure dt .

We first define by means of $\text{Spec}(G, g)$ the family of functions

$$(7) \quad g_\lambda: \mathbf{R}_+^* \ni t \mapsto \exp(-\zeta_\lambda t) \quad (\lambda \in A(G)).$$

Let now the value $t \in \mathbf{R}_+^*$ be fixed and consider, following E. M. Stein [44, 45], the continuous endomorphism $\exp(\Delta t)$ of the complex Banach space $L^2(G)$. Since the positivity of the resolvent of Δ on \mathbf{R}_+^* implies that the restricted operator

$$\exp(\Delta t): \mathcal{E}^0(G) \rightarrow \mathcal{E}^0(G)$$

is a *positive* linear mapping, there exists exactly one (positive) Radon measure $\mu_t \in \mathcal{M}(G)$ such that $\mu_t(G) = 1$ and

$$\exp(\Delta t): \mathcal{E}^0(G) \ni f \mapsto f * \mu_t \in \mathcal{E}^0(G)$$

holds. By an extension to $L^2(G)$, we infer the identities

$$u_\lambda * \exp(\Delta t)f = \exp(\Delta t)(u_\lambda * f) \quad (\lambda \in \Lambda(G))$$

for any function $f \in L^2(G)$. The spectral decomposition (1) then shows that

$$\exp(\Delta t)f = \sum_{\lambda \in \Lambda(G)} q_\lambda(t)u_\lambda * f$$

holds in the complete Hilbert algebra $L^2(G)$. Consequently, the kernel function

$$K = \sum_{\lambda \in \Lambda(G)} q_\lambda \otimes u_\lambda$$

is an element of the complex Fréchet space $L^2_{loc}(\mathbf{R}_+^* \times G)$.

As usual, let $\mathcal{D}(\mathbf{R}_+^* \times G)$ be the complex vector space of all infinitely differentiable complex-valued functions on the manifold $\mathbf{R}_+^* \times G$ whose support is compact. Provide the space $\mathcal{D}(\mathbf{R}_+^* \times G)$ with the canonical $\mathcal{L}\mathcal{F}$ topology and let $\mathcal{D}'(\mathbf{R}_+^* \times G)$ be its strong topological dual which consists of all complex distributions on $\mathbf{R}_+^* \times G$. Then we have

$$\mathcal{D}'(\mathbf{R}_+^* \times G) = \mathcal{D}'(\mathbf{R}_+^*) \hat{\otimes} \mathcal{E}'(G)$$

and we may identify the Fréchet space $L^2_{loc}(\mathbf{R}_+^* \times G)$ with its image under the continuous injection

$$h \mapsto h \cdot (dt \otimes \nu).$$

Let the heat-diffusion operator $\partial/\partial t - \Delta$ be operating from the space $\mathcal{D}'(\mathbf{R}_+^* \times G)$ into itself as a continuous linear mapping. We observe that for an arbitrary *finite* subset A_0 of $\Lambda(G)$ the function $K_0 = \sum_{\lambda \in A_0} q_\lambda \otimes u_\lambda$ which belongs to $\mathcal{E}(\mathbf{R}_+^* \times G)$ satisfies the heat-diffusion equation $(\partial/\partial t - \Delta)K_0 = 0$. We obtain therefore the identity

$$\left(\frac{\partial}{\partial t} - \Delta\right)K = 0 \quad \text{in } \mathcal{D}'(\mathbf{R}_+^* \times G).$$

On the other hand it is a well known fact that the parabolic differential operator $\partial/\partial t - \Delta$ is *hypoelliptic*: see, for instance, F. Trèves [46], Chapter 7. It follows that the singular support of K in $\mathbf{R}_+^* \times G$ is empty, i.e.

$$K \in \mathcal{E}(\mathbf{R}_+^* \times G) = \mathcal{E}(\mathbf{R}_+^*) \hat{\otimes} \mathcal{E}(G).$$

Therefore we have

$$\mu_t = K(t, \cdot)\nu \quad (t \in \mathbf{R}_+^*),$$

i.e. the Radon measure μ_t admits the central function $K(t, \cdot)$ as its Lebesgue-Nikodym density with respect to the base ν .

From the fact that $(\exp(\Delta t))_{t \in \mathbf{R}_+^*}$ forms a strongly continuous one-parameter Markovian semigroup in $\mathcal{A}(G)$ which admits Δ as its infinitesimal generator (see too N. Dunford-J. T. Schwartz [20], Chapter XIV), we infer that the generating family $(\mu_t)_{t \in \mathbf{R}_+^*}$ is a central approximate unit for $\mathcal{A}(G)$. Moreover we have the following result.

THEOREM 4. *The heat-diffusion approximation process $(\exp(\Delta t))_{t \in \mathbf{R}_+^*}$ on $\mathcal{A}(G)$ is central. It has the saturation structure $(\varphi; \mathcal{A}(G); \mathcal{V}_\psi(\mathcal{A}(G)))$ where the mappings φ, ψ are defined in the following way:*

$$\varphi: \mathbf{R}_+^* \ni t \mapsto t; \quad \psi: \Lambda(G) \ni \lambda \mapsto -\zeta_\lambda.$$

PROOF. For the character $c_\lambda \in X(\mathcal{Z}\mathcal{M}(G))$ defined in Theorem 1 we obtain in the present case obviously

$$c_\lambda(\mu_t) = q_\lambda(t)$$

for any pair $(\lambda, t) \in \Lambda(G) \times \mathbf{R}_+^*$. It follows that $(\mu_t)_{t \in \mathbf{R}_+^*}$ forms a central pro-saturation measure of type $(\varphi; \psi)$ where the functions φ and ψ are defined as indicated above. Let V denote the Favard space of $(\exp(\Delta t))_{t \in \mathbf{R}_+^*}$ with respect to $\mathcal{A}(G)$. By virtue of Theorem 2 which ensures the validity of conditions (I) and (II) in Definition 1, and in view of Theorem 3 it suffices to prove that the inclusion

$$[\mathcal{W}_\psi(\mathcal{A}(G)), \mathcal{A}(G)]_0 \subset V$$

holds.

(III) Let any function $f \in [\mathcal{W}_\psi(\mathcal{A}(G)), \mathcal{A}(G)]_0$ be given. Choose sequences $(f_n)_{n \geq 1}$ and $(g_n)_{n \geq 1}$ in $\mathcal{A}(G)$ such that

$$\lim_{n \rightarrow \infty} \|f_n - f\| = 0;$$

$$\psi(\lambda) \cdot u_\lambda * f_n = u_\lambda * g_n \quad (n \geq 1, \lambda \in \Lambda(G));$$

$$\sup_{n \geq 1} \|f_n\|_{\mathcal{W}} = \sup_{n \geq 1} (\|f_n\| + \|g_n\|) = M < +\infty.$$

A direct computation shows that the identity

$$\exp(\Delta t)f_n - f_n = \int_0^t \exp(\Delta \tau)g_n d\tau$$

obtains for any $n \geq 1$ and $t \in \mathbf{R}_+^*$. It follows

$$\|\exp(\Delta t)f - f\| \leq M \cdot t$$

for $t \in \mathbf{R}_+^*$. Thus we have

$$\|\exp(\Delta t)f - f\| = O(\varphi(t)) \quad (t \rightarrow 0+),$$

i.e. $f \in V$ and the proof is complete.—

Let $\text{Dom}(\Delta)$ denote the domain of the infinitesimal generator Δ of the heat-diffusion semigroup $(\exp(\Delta t))_{t \in \mathbb{R}_+}$ in $\mathcal{A}(G)$. Taking into account a general theorem due to H. Berens [4], we obtain furthermore the following result.

COROLLARY. *The approximation process $(\exp(\Delta t))_{t \in \mathbb{R}_+^*}$ on $\mathcal{A}(G)$ has the relative completion*

$$V = [\text{Dom}(\Delta), \mathcal{A}(G)]_0$$

as its Favard space.

5. The Poisson Approximation Process. We continue to follow the treatises of E. M. Stein [44, 45] in replacing the family $(q_\lambda)_{\lambda \in \Lambda(G)}$ in (7) by the functions

$$p_\lambda: \mathbb{R}_+^* \ni t \mapsto \exp(-\sqrt{\zeta_\lambda} t) \quad (\lambda \in \Lambda(G)).$$

Then the Poisson kernel

$$P = \sum_{\lambda \in \Lambda(G)} p_\lambda \otimes u_\lambda$$

associated with the compact Lie group G belongs to the space $L_{loc}^2(\mathbb{R}_+^* \times G)$. Denote by

$$\underline{\Delta} = \frac{\partial^2}{\partial t^2} + \Delta$$

the Laplace-Beltrami operator of the Riemannian manifold $\mathbb{R}_+^* \times G$ and assume that $\underline{\Delta}$ acts on the space $\mathcal{D}'(\mathbb{R}_+^* \times G)$ of distributions. Then P is a harmonic distribution on $\mathbb{R}_+^* \times G$, i.e.

$$\underline{\Delta}P = 0 \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times G).$$

Therefore an application of Weyl's regularity lemma shows that P is a harmonic function on $\mathbb{R}_+^* \times G$. In particular, we have $P \in \mathcal{E}(\mathbb{R}_+^* \times G)$. Define the Radon measures

$$\pi_t = P(t, \cdot) \nu \quad (t \in \mathbb{R}_+^*)$$

on G . Then the family $(\pi_t)_{t \in \mathbb{R}_+^*}$ forms a central approximate unit for $\mathcal{A}(G)$. The associated strongly continuous one-parameter Markovian semigroup $(\exp(-\sqrt{-\Delta} t))_{t \in \mathbb{R}_+}$ has $-\sqrt{-\Delta}$ as its infinitesimal generator.

THEOREM 5. *The Poisson approximation process $(\exp(-\sqrt{-\Delta} t))_{t \in \mathbb{R}_+^*}$ on $\mathcal{A}(G)$ is central. Define the mappings*

$$\varphi: \mathbb{R}_+^* \ni t \mapsto t; \quad \psi: \Lambda(G) \ni \lambda \mapsto -\sqrt{\zeta_\lambda}.$$

Then $(\exp(-\sqrt{-\Delta} t))_{t \in \mathbb{R}_+^*}$ has the saturation structure $(\varphi; \mathcal{A}(G); \mathcal{V}_\psi(\mathcal{A}(G)))$

and the identity

$$\mathcal{V}_\psi(\mathcal{A}(G)) = [\text{Dom}(-\sqrt{-\Delta}), \mathcal{A}(G)]_0$$

obtains for the associated Favard space.

The proof is based on Theorems 2 and 3 *supra* and follows by an application of the method outlined in Section 4. We omit the details.

REMARK 5. Let $\alpha \in]0, 1[$. Needless to say, there are analogous results concerning the saturation behaviour of the central approximation processes $(\exp(-(-\Delta)^\alpha t))_{t \in \mathbb{R}_+^*}$ on $\mathcal{A}(G)$. A discussion of this case, of course, depends upon the concepts of *fractional* differentiation and integration. See, for instance, P. L. Butzer-H. Berens [8], Chapter II, and the recent paper by H. Bavinck [3]. We shall return to these problems within our general framework in a forthcoming paper.

6. Applications I. (i) Let G be an n -dimensional *abelian* compact connected Lie group. Then G is isomorphic with the Euclidean torus $T^n = \mathbb{R}^n/\mathbb{Z}^n = (\mathbb{R}/\mathbb{Z})^n$ of dimension n . We shall identify the integral lattice \mathbb{Z}^n with its dual one, i.e. with the lattice of all vectors $x \in \mathbb{R}^n$ such that the scalar product $(\lambda | x)$ belongs to \mathbb{Z} for all $\lambda \in \mathbb{Z}^n$. Then we have

$$A(T^n) = \hat{T}^n = \mathbb{Z}^n.$$

If $\mathbb{R}^n \ni x \mapsto \hat{x} \in T^n$ denotes the canonical epimorphism, we obtain for any wave number $\lambda \in \mathbb{Z}^n$ in the present case

$$n_\lambda = 1;$$

$$u_\lambda = \chi_\lambda: T^n \ni \hat{x} \mapsto \exp(2\pi i(\lambda | x)) \in \mathbb{C};$$

$$c_\lambda: \mathcal{M}(T^n) \ni \mu \mapsto \mathcal{F}\mu(\lambda) \in \mathbb{C};$$

where $\mathcal{F}: \mathcal{M}(T^n) \rightarrow \mathcal{E}^0(\mathbb{Z}^n)$ stands for the classical Fourier transformation.

Let the manifold \mathbb{R}^n be endowed with its canonical orientation and with its canonical Riemannian structure such that the corresponding metric g_0 induces the Euclidean norm $|\cdot|$. Denote by g_0/\mathbb{Z}^n the unique flat Riemannian metric on T^n such that (\mathbb{R}^n, g_0) is the universal Riemannian covering manifold of $(T^n, g_0/\mathbb{Z}^n)$. Then we obtain

$$\text{Spec}(T^n, g_0/\mathbb{Z}^n) = \{4\pi^2|\lambda|^2 \mid \lambda \in \mathbb{Z}^n\}$$

and the heat-diffusion kernel K admits in the present case the following expansion in a *theta series*:

$$K: (\mathbb{R}_+^* \times T^n) \ni (t, \hat{x}) \mapsto \sum_{\lambda \in \mathbb{Z}^n} \exp(2\pi i((\lambda | x) + 2\pi i t(\lambda | \lambda))) = \Theta_0(\tau_t, x).$$

Here $\theta_0(\tau_t, \cdot)$ denotes the classical theta function of characteristic 0 and of modulus

$$\tau_t = 4\pi it \cdot 1_n \quad (t \in \mathbf{R}^*)$$

where 1_n stands for the (n, n) identity matrix. Therefore τ_t belongs to the Siegel upper-half space of degree n : see, for instance, J. Igusa [24], Chapter I.

An application of Theorems 4 and 5 in the present situation furnishes results due to R. J. Nessel-A. Pawelke [34]. If we assume $n = 1$, the Cauchy problem which is solved by the heat-diffusion semigroup $(\exp(\Delta t))_{t \in \mathbf{R}_+}$ in $\mathcal{A}(T)$ is known as the “Fourier torus problem” (A. Sommerfeld [43], Kapitel III). Detailed treatments of this problem and of the corresponding Dirichlet problem from the saturation theoretic point of view may be found in the monographs by P. L. Butzer-H. Berens [8] and P. L. Butzer-R. J. Nessel [10]. Also see [17] for a study of saturation on abelian quotient groups.

(ii) Now suppose that the compact connected Lie group G is diffeomorphic with the standard sphere S_{n-1} of an Euclidean space $\mathbf{R}^n (n \geq 1)$. Since S_0 is not connected and S_1 is diffeomorphic with the torus T as treated in Example (i) supra, we may suppose $n \geq 3$. Then, according to a well-known result due to H. Samelson [41], *only the choice $n = 4$ is possible*. Indeed, the Lie group $SU(2)$, the group of unit quaternions, is diffeomorphic with the sphere S_3 and $SU(2)$ acts as a subgroup of the special orthogonal group $SO(4)$ in a natural way transitively by rotations on S_3 . As usual, let

$$\Lambda(SU(2)) = \frac{1}{2}N,$$

be the family of weights, where $N = \mathbf{Z}_+$.

Let the compact manifold S_3 be oriented to the exterior and let it be endowed with the Riemannian structure induced by the canonical structure of \mathbf{R}^4 . Denote by g the Riemannian metric of S_3 . A direct computation (cf. M. Berger-P. Gauduchon-E. Mazet [6]) shows that

$$n_\lambda = 2\lambda + 1 \quad \left(\lambda \in \frac{1}{2}N \right);$$

$$\text{Spec}(SU(2), g) = \left\{ 4\lambda(\lambda + 1) \mid \lambda \in \frac{1}{2}N \right\}.$$

In order to obtain explicitly the units u_λ and the characters χ_λ ($\lambda \in (1/2)N$) of the spectral decomposition of $L^2(SU(2))$, fix the North pole

$1 = (0, 0, 0, 1) \in R^4$ of S_3 . Then we may consider $SO(3)$ as the stabilizer of 1 in the rotation group $SO(4)$ and identify S_3 with the compact homogeneous manifold $SO(4)/SO(3)$. Since $(SO(4), SO(3))$ is a Riemannian *symmetric pair*, we obtain for the unit u_λ the identity

$$u_\lambda = Z_1^{2\lambda} \quad \left(\lambda \in \frac{1}{2}N \right)$$

where $Z_1^{2\lambda}$ denotes the zonal surface spherical harmonic of degree 2λ with pole 1 on S_3 . See, for instance, R. R. Coifman-G. Weiss [11, 12]. Thus, for any $\lambda \in (1/2)N$, the character χ_λ takes the following form:

$$\chi_\lambda: S_3 \ni (x_j)_{1 \leq j \leq 4} \rightsquigarrow \frac{1}{2\lambda + 1} Z_1^{2\lambda}(x_4) = \prod_{2\lambda}^{(\frac{1}{2}, \frac{1}{2})}(x_4).$$

Here $\prod_{2\lambda}^{(1/2, 1/2)}(X) \in R[X]$ denotes the ultraspherical polynomial (Gegenbauer polynomial) in the indeterminate X of degree 2λ with exponent $1/2$ i.e. the Čebyšev polynomial of the second kind

$$\prod_{2\lambda}^{(\frac{1}{2}, \frac{1}{2})}(X) = \sum_{0 \leq m \leq [2\lambda]} (-1)^m \frac{(2\lambda - m)!}{m!(2\lambda - 2m)!} (2X)^{2(\lambda - m)}.$$

The heat-diffusion kernel K and the Poisson kernel P admit therefore the following expansions in Laplace series:

$$K: R_+^* \times S_3 \ni (t, (x_j)_{1 \leq j \leq 4}) \rightsquigarrow \sum_{\lambda \in (1/2)N} \exp(-4\lambda(\lambda + 1)t) \cdot (2\lambda + 1) \prod_{2\lambda}^{(\frac{1}{2}, \frac{1}{2})}(x_4);$$

$$P: R_+^* \times S_3 \ni (t, (x_j)_{1 \leq j \leq 4}) \rightsquigarrow \sum_{\lambda \in (1/2)N} \exp(-2\sqrt{\lambda(\lambda + 1)}t) \cdot (2\lambda + 1) \prod_{2\lambda}^{(\frac{1}{2}, \frac{1}{2})}(x_4).$$

Equip the sphere S_3 with the normalized Lebesgue surface measure which may be identified with the Haar measure ν of $SU(2)$. Then an application of Theorems 4 and 5 to the special unitary group $SU(2)$ yields results which are concerned with the saturation behaviour of the approximation of functions $f \in \mathcal{A}(S_3)$ performed by Laplace expansions in Čebyšev polynomials of the second kind.

At this juncture, bearing in mind the notion of zonal approximation process [19] as outlined in Remark 3 supra and the fact that the commutative complex Banach algebras $\mathcal{M}(SO(3) \setminus SO(4)/SO(3))$ and $\mathcal{M}(SU(2))$ may be identified, we shall summarize the preceding considerations in the following

THEOREM 6. *On the compact unit sphere S_3 in R^4 , the zonal approximation processes with respect to the compact homogeneous manifold $SO(4)/SO(3)$ and the central approximation processes with respect to the special unitary group $SU(2)$ coalesce.*

For further results concerning saturation theory on spheres we refer to S. Pawelke [35] and H. Berens-P. L. Butzer-S. Pawelke [5]. Also see S. Pawelke [36]. Details for saturation on general compact homogeneous spaces may be found in [15, 16].

7. Applications II. In this section we shall continue to study central approximation processes on the Lie group $SU(2)$. In order to adopt the conventions of R. A. Mayer [29, 30], let us now parametrize the spectral decomposition of $L^2(SU(2))$ by the set

$$A(SU(2)) = N^* .$$

Henceforth we will write

$$k = 2\lambda + 1 \quad \left(\lambda \in \frac{1}{2} N \right)$$

for the weights of $SU(2)$. Let us identify $SU(2)$ and S_3 as in Section 6. Then we have

$$u_k : S_3 \ni (x_j)_{1 \leq j \leq 4} \rightsquigarrow Z_1^{k-1}(x_4) \quad (k \in N^*) .$$

$$\chi_k : S_3 \ni (x_j)_{1 \leq j \leq 4} \rightsquigarrow \prod_{k-1}^{(\frac{1}{2}, \frac{1}{2})}(x_4)$$

(iii) In the algebra $\mathcal{A}(SU(2))$ define the sequence $(\mu_n)_{n \geq 0}$ by letting

$$\mu_n = \sum_{1 \leq k \leq n+1} \left[\binom{2n+2}{n+k+1} / \binom{2n+2}{n+2} \right] u_k \nu \quad (n \in N) .$$

Obviously $\mu_n(SU(2)) = 1$. Since $\mu_n \geq 0$ (R. A. Mayer [30]) it follows

$$\|\mu_n\| = 1 \quad (n \in N) .$$

Consequently, the continuous linear mappings

$$V_n : L^1(SU(2)) \ni f \rightsquigarrow \mu_n * f \in L^1(SU(2)) \quad (n \in N)$$

are contractions. Therefore $(V_n)_{n \geq 0}$ forms an equicontinuous family which converges pointwise on the socle $\mathcal{S}_{L^2(SU(2))}$ towards the identity automorphism of $L^1(SU(2))$. Since $\mathcal{S}_{L^2(SU(2))}$ is everywhere norm dense in $L^1(SU(2))$, it follows from Remark 1 supra that $(\mu_n)_{n \geq 1}$ is a central approximate unit for $\mathcal{A}(SU(2))$ which generates the approximation process $(V_n)_{n \geq 1}$ of the de La Vallée Poussin type on $\mathcal{A}(SU(2))$. In the present case we obtain

$$c_k(\mu_n) = \begin{cases} \binom{2n+2}{n+k+1} / \binom{2n+2}{n+2} & \text{if } 1 \leq k \leq n+1 ; \\ 0 & \text{if } n+1 < k ; \end{cases}$$

whence

$$\lim_{n \rightarrow \infty} \frac{c_k(\mu_n) - 1}{1/n} = 1 - k^2$$

for any $k \in N^*$. Therefore $(\mu_n)_{n \geq 1}$ forms a central pro-saturation measure of type $(n \rightsquigarrow 1/n, k \rightsquigarrow 1 - k^2)$ with respect to $\mathcal{A}(\text{SU}(2))$.

THEOREM 7. *The approximation process $(V_n)_{n \geq 1}$ on $\mathcal{A}(\text{SU}(2))$ of the de La Vallée Poussin type is central and has the saturation structure $(\varphi; \mathcal{A}(\text{SU}(2)); \mathcal{W}_\psi(\text{SU}(2)))$ where*

$$\varphi: N^* \ni n \rightsquigarrow \frac{1}{n}; \quad \psi: N^* \ni k \rightsquigarrow 1 - k^2.$$

PROOF. Let V denote the Favard space of $(V_n)_{n \geq 1}$ with respect to $\mathcal{A}(\text{SU}(2))$. In view of Theorems 2 and 3 it suffices to prove that the inclusion

$$[\mathcal{W}_\psi(\text{SU}(2)), \mathcal{A}(\text{SU}(2))]_0 \subset V$$

holds.

(III) Let the function $f \in [\mathcal{W}_\psi(\text{SU}(2)), \mathcal{A}(\text{SU}(2))]_0$ be given. There are sequences $(f_n)_{n \geq 1}$ and $(g_n)_{n \geq 1}$ in the space $\mathcal{A}(\text{SU}(2))$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|f_n - f\| &= 0; \\ \psi(k)u_k * f_n &= u_k * g_n \quad (n \geq 1, k \geq 1); \\ \sup_{n \geq 1} \|f_n\|_{\mathcal{W}} &= \sup_{n \geq 1} (\|f_n\| + \|g_n\|) = M < +\infty. \end{aligned}$$

Since the identity

$$\psi(k)c_k(\mu_j) = -j(j + 2)(c_k(\mu_j) - c_k(\mu_{j-1}))$$

holds for any pair $(k, j) \in N^* \times N^*$, we obtain for all integers $r \geq m \geq 1$:

$$V_m(f_n) - V_r(f_n) = \sum_{m+1 \leq j \leq r} \frac{1}{j(j + 2)} V_j(g_n) \quad (n \in N^*).$$

Thus the estimate

$$\|V_m(f_n) - f_n\| \leq M \cdot \sum_{m+1 \leq j} \frac{1}{j(j + 2)} \leq \varphi(m) \cdot M \quad (m \in N^*)$$

obtains. Hence $f \in V$ and the proof is complete.

(iv) Finally, let

$$\mu_{n-1} = \left(\sum_{1 \leq k \leq n} k^2 \right)^{-1} \left(\sum_{1 \leq k \leq n} k\chi_k \right)^2 \quad (n \in N^*).$$

As in the preceding section it can be shown that the sequence $(\mu_{n-1})_{n \geq 1}$

forms a central approximate unit for $\mathcal{A}(\text{SU}(2))$. Since the identity

$$\mu_{n-1} = \sum_{1 \leq k \leq 2n-1} \left(1 - \frac{6(2n+1)^2[(1/2)k^2] - k^2(k^2 - 1)}{8kn(n+1)(2n+1)} \right) u_k \nu \quad (n \in N^*)$$

obtains (R. A. Mayer [30]), it follows

$$\lim_{n \rightarrow \infty} \frac{c_k(\mu_{n-1}) - 1}{1/(n-1)} = -\frac{3}{2} \frac{[(1/2)k^2]}{k}$$

for $k \in N^*$.

THEOREM 8. *The central approximation process $(F_{n-1})_{n \geq 1}$ on $\mathcal{A}(\text{SU}(2))$ of the Fejér type which is generated by the approximate unit $(\mu_{n-1})_{n \geq 1}$ for $\mathcal{A}(\text{SU}(2))$ has the saturation structure $(\varphi; \mathcal{A}(\text{SU}(2)))$; $V_\psi(\mathcal{A}(\text{SU}(2)))$ where*

$$\varphi: N^* \ni n \rightsquigarrow \frac{1}{n}, \quad \psi: N^* \ni k \rightsquigarrow -\frac{3}{2} \frac{[(1/2)k^2]}{k}.$$

Since the proof of Theorem 8 is similar to the preceding one, we may omit the details. For further results concerning summation processes on the special unitary group $\text{SU}(2)$ we refer to R. A. Mayer [31-33].

8. Conclusion. Let us briefly review Sections 4, 5 and 6. As it was pointed out by E. M. Stein [45], his construction of the heat-diffusion semigroup $(\exp(\Delta t))_{t \in R_+}$ and of the Poisson semigroup $(\exp(-\sqrt{-\Delta} t))_{t \in R_+}$ which is based on the celebrated paper by G. A. Hunt [23], can be extended to *non-compact* connected Lie groups. In this connection also see R. Azencott [2].

On the other hand, it is a well known fact that the expansions in series of special functions require in general the study of *non-compact* Lie groups. For these reasons we shall return to saturation theory on Lie groups in a forthcoming paper. In particular we shall draw upon the theory of nuclear topological vector spaces.

REFERENCES

[1] N. ARONSZAIN AND E. GAGLIARDO, Interpolation spaces and interpolation methods, Ann. Mat. Pura Appl. 68 (1965), 51-118.
 [2] R. AZENCOTT, Espaces de Poisson des groupes localement compacts, Lecture Notes in Mathematics, Vol. 148. Berlin-Heidelberg-New York: Springer 1970.
 [3] H. BAVINCK, A special class of Jacobi series and some applications, J. Math. Anal. Appl. 37 (1972), 767-797.
 [4] H. BERENS, Interpolationsmethoden zur Behandlung von Approximationsprozessen auf Banachräumen, Lecture Notes in Mathematics, Vol. 64. Berlin-Heidelberg-New York: Springer 1968.

- [5] H. BERENS, P. L. BUTZER AND S. PAWELKE, Limitierungs-verfahren von Reihen mehrdimensionaler Kugelfunktionen und deren Saturationsverhalten, *Publ. Res. Inst. Math. Sci. Ser. A.* 4 (1968), 201-268.
- [6] M. BERGER, P. GAUDUCHON AND E. MAZET, *Le spectre d'une variété riemannienne*, Lecture Notes in Mathematics, Vol. 194. Berlin-Heidelberg-New York: Springer 1971.
- [7] H. BUCHWALTER, Saturation sur un groupe abélien localement compact, *C. R. Acad. Sci. Paris* 250 (1960), 808-810.
- [8] P. L. BUTZER AND H. BERENS, *Semi-groups of operators and approximation*, Berlin-Heidelberg-New York: Springer 1967.
- [9] P. L. BUTZER AND H. JOHNNEN, Lipschitz spaces on compact manifolds, *J. Functional Analysis* 7 (1971), 242-266.
- [10] P. L. BUTZER AND R. J. NESSEL, *Fourier analysis and approximation. Vol. 1: One-dimensional theory*, Basel-Stuttgart: Birkhäuser 1971.
- [11] R. R. COIFMAN AND G. WEISS, Representations of compact groups and spherical harmonics, *Enseignement Math.* 14 (1968), 121-173.
- [12] R. R. COIFMAN AND G. WEISS, *Analyse harmonique non-commutative sur certains espaces homogènes*, Lecture Notes in Mathematics. Vol. 242, Berlin-Heidelberg-New York: Springer 1971.
- [13] J. DIEUDONNÉ, Sur le socle d'un anneau et les anneaux simples infinis, *Bull. Soc. Math. France* 70 (1942), 46-75.
- [14] J. DIEUDONNÉ, Representaciones de grupos compactos y funciones esfericas, *Cursos y seminarios de matemática*, Fasc. 14. Universidad de Buenos Aires 1964.
- [15] B. DRESELER, Saturationstheorie auf kompakten topologischen Gruppen. In: *Proceedings of the Conference on Approximation Theory at Poznań 1972*. Edited by Z. Ciesielski, Dordrecht: D. Reidel Publishing Company 1974.
- [16] B. DRESELER, Saturation auf kompakten topologischen Gruppen und homogenen Räumen. *Dissertation. Universität Mannheim* 1972.
- [17] B. DRESELER AND W. SCHEMPP, Saturation on locally compact abelian groups. *Manuscripta Math.* 7 (1972), 141-174.
- [18] B. DRESELER AND W. SCHEMPP, Saturation on locally compact abelian groups: An extended theorem. *Manuscripta Math.* 8 (1973), 271-286.
- [19] B. DRESELER AND W. SCHEMPP, Zonal approximation processes. *Math. Z.* 133 (1973), 81-92.
- [20] N. DUNFORD AND J. T. SCHWARTZ, *Linear operators. Part II: Spectral theory*. New York-London: Interscience Publishers 1963.
- [21] E. GAGLIARDO, A unified structure in various families of function spaces. Compactness and closure theorems. In: *Proceedings of the International Symposium on Linear Spaces*, pp. 237-241. Jerusalem: Jerusalem Academic Press 1961.
- [22] E. HEWITT AND K. A. ROSS, *Abstract harmonic analysis. Vol. II: Structure and analysis for compact groups*. Berlin-Heidelberg-New York: Springer 1970.
- [23] G. A. HUNT, Semi-groups of measures on Lie groups. *Trans. Amer. Math. Soc.* 81 (1956), 264-293.
- [24] J. IGUSA, *Theta functions*. Berlin-Heidelberg-New York: Springer 1972.
- [25] H. JOHNNEN, Stetigkeitsmoduli und Approximationstheorie auf kompakten Lie-Gruppen. *Dissertation. Technische Hochschule Aachen* 1970.
- [26] H. JOHNNEN, Darstellungen von Liegruppen und Approximationsprozesse auf Banachräumen. *J. Reine Angew. Math.* 254 (1972), 160-187.
- [27] H. JOHNNEN, Best approximation on the unitary group. In: *Colloquia Mathematica Societatis János Bolyai*, 5. Hilbert space operators, pp. 295-303. Amsterdam-London: North-Holland Publishing Comp. 1972.
- [28] Y. KAWADA, Über den Dualitätssatz der Charaktere nichtkommutativer Gruppen. *Proc.*

- Phys.-Math. Soc. Japan 24 (1942), 97-109.
- [29] R. A. MAYER, Localization and summability for Fourier series on compact groups. Ph. D. Thesis. Columbia University 1964.
 - [30] R. A. MAYER, Summation of Fourier series on compact groups. Amer. J. Math. 89 (1967), 661-692.
 - [31] R. A. MAYER, Fourier series of differentiable functions on $SU(2)$. Duke Math. J. 34 (1967), 549-554.
 - [32] R. A. MAYER, Localization for Fourier series on $SU(2)$. Trans. Amer. Math. Soc. 130 (1968), 414-424.
 - [33] R. A. MAYER, On the Fourier series of certain characteristic functions on $SU(2)$. Duke Math. J. 38 (1971), 63-75.
 - [34] R. J. NESSEL AND A. PAWELKE, Über Favardklassen von Summationsprozessen mehrdimensionaler Fourierreihen. Compositio Math. 19 (1968), 196-212.
 - [35] S. PAWELKE, Saturation und Approximation bei Reihen mehrdimensionaler Kugelfunktionen. Dissertation. Technische Hochschule Aachen 1969.
 - [36] S. PAWELKE, Über die Approximationsordnung bei Kugelfunktionen und algebraischen Polynomen. Tôhoku Math. J. 24 (1972), 473-486.
 - [37] D. L. RAGOZIN, Approximation theory on compact manifolds and Lie groups with applications to harmonic analysis. Ph. D. Thesis. Harvard University 1967.
 - [38] D. L. RAGOZIN, Approximation Theory on $SU(2)$. J. Approximation Theory 1 (1968), 464-475.
 - [39] D. L. RAGOZIN, Constructive polynomial approximation on spheres and projective spaces. Trans. Amer. Math. Soc. 162 (1971), 157-170.
 - [40] D. L. RAGOZIN, Uniform convergence of spherical harmonic expansions. Math. Ann. 195 (1972), 87-94.
 - [41] H. SAMELSON, Über die Sphären die als Gruppenräume auftreten. Comment. Math. Helv. 13 (1940), 144-155.
 - [42] W. SCHEMPP, Zur Theorie der saturierten Approximationsverfahren auf lokalkompakten abelschen Gruppen. In: Proceedings of the Conference on Approximation Theory at Poznań 1972. Edited by Z. Ciesielski, Dordrecht: D. Reidel Publishing Company 1974.
 - [43] A. SOMMERFELD, Vorlesungen über theoretische Physik. Band VI: Partielle Differentialgleichungen der Physik. Wiesbaden: Dieterich'sche Verlagsbuchhandlung 1947.
 - [44] E. M. STEIN, Variations on the Littlewood-Paley theme. In: Lectures in Modern Analysis and Applications III, pp. 1-17. Lecture Notes in Mathematics, Vol. 170. Berlin-Heidelberg-New York: Springer 1970.
 - [45] E. M. STEIN, Topics in harmonic analysis related to the Littlewood-Paley theory. Annals of Mathematics Studies, Number 63. Princeton, New Jersey: Princeton University Press 1970.
 - [46] F. TRÉVES, Linear partial differential equations with constant coefficients. New York-London-Paris: Gordon and Breach 1966.

DR. BERND DRESELER
 PROF. DR. WALTER SCHEMPP
 LEHRSTUHL FÜR MATHEMATIK I
 GESAMTHOCHSCHULE SIEGEN
 D-5930 HÜTTENTAL-WEIDENAU, HÖLDERLINSTRASSE 3
 FEDERAL REPUBLIC OF GERMANY

