

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/47487>

Please be advised that this information was generated on 2022-08-25 and may be subject to change.

## Cerebral Venous Sinus Thrombosis: Prevention of Recurrent Thromboembolism B.F.L. van Nuenen, M. Munneke and B.R. Bloem

*Stroke*. 2005;36:1822-1823

doi: 10.1161/01.STR.0000176582.17934.82

*Stroke* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2005 American Heart Association, Inc. All rights reserved.

Print ISSN: 0039-2499. Online ISSN: 1524-4628

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://stroke.ahajournals.org/content/36/9/1822>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Stroke* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Stroke* is online at:  
<http://stroke.ahajournals.org/subscriptions/>

# Letters to the Editor

Stroke welcomes Letters to the Editor and will publish them, if suitable, as space permits. They should not exceed 750 words (including references) and may be subject to editing or abridgment. Please submit letters in duplicate, typed double-spaced. Include a fax number for the corresponding author and a completed copyright transfer agreement form (published in every issue).

## Protein Z Levels, Protein Z G79A Polymorphism, and Prothrombotic Conditions

To the Editor:

We read with great interest the article on protein Z gene polymorphisms, protein Z concentrations, and ischemic stroke published by Staton et al.<sup>1</sup> The authors examine protein Z concentrations and 2 common polymorphisms of the protein Z gene in patients with a first-ever ischemic stroke and conclude that high levels of protein Z are prothrombotic and that the 79AA genotype of the G79A polymorphism is protective for the ischemic stroke.

The conclusions drawn by Staton et al seem questionable because of several flaws.

First, the hypothesis that high levels of protein Z are prothrombotic is, to date, not supported by the pathophysiology, because the role of protein Z has been demonstrated to be the inhibition of activated factor X through the formation of a complex with protein Z–dependent protease inhibitor. Consequently, low levels of protein Z are expected to be prothrombotic. In line with this theory, the majority of the studies in this field have reported low levels of protein Z in association with prothrombotic conditions. Actually, only 2 studies have reported that high levels of protein Z are prothrombotic, but methodological biases are detectable. In the former, the investigators modified the commercial assay and used a normal plasma pool.<sup>2</sup> In the latter, the control group included subjects with a previous vascular event.<sup>3</sup>

Second, Staton et al reported lower levels of protein Z plasma in controls with respect to all the other case-control studies present in the literature, including reports with a greater number of subjects analyzed. This might be attributable to the presence of subjects with a previous vascular event (not better clarified) included in the control group. Actually, patients with vascular diseases have been reported to have low levels of protein Z.<sup>4</sup>

Third, the significant result of the 79A allele as a protective factor for the ischemic stroke has been performed by pooling data from a previous study. Actually, a pooled analysis in case-control studies, especially for the evaluation of genetic polymorphisms, is rather a questionable approach. To perform this kind of analysis, the 2 populations should be homogenous and similar for the main parameters investigated. Conversely, the Staton et al do not explain how they tested the homogeneity of the 2 populations, particularly with regard to genetic background and prevalence of the traditional cardiovascular risk factors. Too many differences (first of all, age) between such distant and different populations are detectable to correctly perform this analysis.

In fact, the analysis performed in their single population demonstrated no significant relationship between both the polymorphisms and the ischemic stroke; the only significant result derives from pooling data from the study by Lichy et al<sup>5</sup> and is also determined by an overall low number of homozygotes for the AA genotype (n=32), thereby not reaching a sufficient statistical power.

In consideration of all these limitations, we do not believe that the conclusions stated by Staton et al, ie, high levels of protein Z prothrombotic and 79A allele protective for the ischemic stroke, can be drawn from this study. Because low protein Z plasma levels have been reported in several studies to be associated with different prothrombotic conditions, it can be hypothesized that protein Z is either simply a marker of the disease or influenced

by the acute phase, but it is at least questionable, to date, to postulate that high levels of protein Z are associated with prothrombotic conditions and that 79AA genotype is protective for the ischemic stroke.

Francesco Sofi, MD  
Francesca Cesari, BS  
Gian Franco Gensini, MD  
Rosanna Abbate, MD  
Sandra Fedi, BS

Department of Medical and Surgical Critical Care  
Thrombosis Centre, University of Florence and  
Department of Heart and Vessels  
Azienda Ospedaliero-Universitaria Careggi  
Florence, Italy

1. Staton J, Sayer M, Hankey GJ, Cole V, Thom J, Eikelboom JW. Protein Z gene polymorphisms, protein Z concentrations, and ischemic stroke. *Stroke*. 2005;36:1123–1127.
2. Kobelt K, Biasiutti FD, Mattle HP, Lammle B, Wuillemin WA. Protein Z in ischaemic stroke. *Br J Haematol*. 2001;114:169–173.
3. McQuillan AM, Eikelboom JW, Hankey GJ, Baker R, Thom J, Staton J, Yi Q, Cole V. Protein Z in ischemic stroke and its etiologic subtypes. *Stroke*. 2003;34:2415–2419.
4. Fedi S, Sofi F, Brogi D, Tellini I, Cesari F, Sestini I, Gazzini A, Comeglio M, Abbate R, Gensini GF. Low protein Z plasma levels are independently associated with acute coronary syndromes. *Thromb Haemost*. 2003;90:1173–1178.
5. Lichy S, Kropp S, Dong-Si T, Genius J, Dolan T, Hampe T, Stoll F, Renner K, Grond-Ginsbach C, Grau A. A common polymorphism of the protein Z gene is associated with protein Z plasma levels and with risk of cerebral ischemia in the young. *Stroke*. 2004;35:40–45.

### Response:

Dr Sofi and colleagues question our conclusion that the consistency of the association between protein Z genotype, elevated blood concentrations of protein Z, and ischemic stroke strengthens the evidence that increased blood concentrations of protein Z concentrations are causally associated with the risk of ischemic stroke.<sup>1</sup>

First, they propose that our conclusion is not supported by pathophysiology, citing methodological bias as an explanation for previous reports of elevated blood concentrations protein Z in stroke patients. However, they provide no explanation of how the use of a normal plasma pool may have biased the results of the study by Kobelt et al.<sup>2</sup> In the study by McQuillan et al,<sup>3</sup> removal of control subjects with a previous vascular event from the analysis does not change the results or conclusions.

Second, as indicated by Sofi et al, blood concentrations of protein Z in our study were lower than previously reported in several other studies, including their own.<sup>4</sup> The reason for this is unclear, although could potentially be related to prolonged storage of blood samples before measuring protein Z levels.<sup>3</sup> However, irrespective of the cause, our within-study comparisons of protein Z concentrations in stroke cases and controls remain valid because the duration of storage of samples for cases and controls was similar and all laboratory samples were collected, processed, stored, and analyzed in the same manner.

Third, Sofi and colleagues question our approach of pooling results from case control studies to clarify the association between protein Z polymorphisms and ischemic stroke. However, this is a well-established technique that is commonly used for aggregating previous research when individual studies have

insufficient power to detect an association.<sup>5</sup> Taken together with the data by Lichy et al,<sup>6</sup> we believe that our study results are consistent with the conclusion that elevated blood concentrations of protein Z are prothrombotic.

**John W. Eikelboom**  
Department of Medicine  
McMaster University  
Hamilton, Canada

**Janelle Staton**  
Department of Hematology  
Royal Perth Hospital  
Perth, Australia

**Graeme J. Hankey**  
School of Medicine and Pharmacology  
University of Western Australia  
Perth, Australia

1. Staton J, Sayer M, Hankey GJ, Cole V, Thom J, Eikelboom JW. Protein Z gene polymorphisms, protein Z concentrations, and ischemic stroke. *Stroke*. 2005;36:1123–1127.
2. Kobelt K, Biasutti FD, Mattle HP, Lammler B, Willemin WA. Protein Z in ischemic stroke. *Br J Haematol*. 2001;114:169–173.
3. McQuillan AM, Eikelboom JW, Hankey GJ, Baker R, Thom J, Staton J, Ti Q, Cole V. Protein Z in ischemic stroke and its etiologic subtypes. *Stroke*. 2003;34:2415–2419.
4. Fedi S, Sofi F, Brogi D, Tellini I, Cesari F, Sestini I, Gazzini A, Comeglio M, Abbate R, Gensini GF. Low protein Z plasma levels are independently associated with acute coronary syndromes. *Thromb Haemost*. 2003;90:1173–1178.
5. Attia J, Phakkinian A, D'Este C. Meta analysis of molecular association studies: methodological lessons for genetic epidemiology. *J Clin Epidemiol*. 2003;56:297–303.
6. Lichy C, Kropp S, Dong-Si T, Genius J, Dolan T, Hampe T, Stoll F, Renner K, Grond-Ginsbach C, Grau A. A common polymorphism of the protein Z gene is associated with protein Z plasma levels and with risk of cerebral ischemia in the young. *Stroke*. 2004;35:40–45.

## Cerebral Venous Sinus Thrombosis: Prevention of Recurrent Thromboembolism

To the Editor:

Patients who have recovered from cerebral venous sinus thrombosis (CVST) are at risk for sustaining a recurrence of CVST or other thrombotic events (Table). For example, in the recent International Study on Cerebral Vein and Dural Sinus Thrombosis (ISCVT) that was performed in 624 patients with cerebral vein or dural sinus thrombosis, the cumulative risk for recurrent CVST or other thrombotic events after CVST was 6.5%.<sup>1</sup> Recurrent thrombosis is potentially fatal. Therefore, an important clinical question is whether recurrence of CVST or other thrombotic events can be prevented in this group.

Currently, the standard care for patients with a first episode of CVST is anticoagulation therapy, which is usually prescribed for 3

to 6 months in the absence of an identifiable cause.<sup>2</sup> However, controlled data about the benefit and optimal duration of oral anticoagulation in patients with CVST are missing.<sup>3</sup> More prolonged treatment may be required, because the cumulative risk for recurrent CVST or other thrombotic events after 3 to 6 months of oral anticoagulation therapy ranges between 5.5 and 26% (calculated over a mean follow-up period of 18 to 77.8 months).<sup>2,4,5</sup> At the time of these recurrences, which mostly occurred within 1 year, none of the patients received anticoagulation therapy. In the recent ISCVT study, 58.8% of the patients with a recurrence of thrombotic event were not undergoing anticoagulation therapy.<sup>1</sup> The significance of these observations is underscored by the fact that recurrence of cerebral venous thrombosis after stopping anticoagulation therapy can occasionally run a fatal course.<sup>6</sup>

The need for more chronic treatment is further supported by recent observations on the effects of prolonged anticoagulant therapy in patients with idiopathic venous thromboembolism (either pulmonary or deep venous thrombosis).<sup>7</sup> This study evaluated the efficacy of long-term, low-intensity warfarin therapy (international normalized ratio, 1.5 to 2.0) in preventing recurrent thrombotic events among patients with idiopathic venous thromboembolism who had completed at least 3 months of therapy with conventional-dose warfarin. Compared with placebo, long-term and low-intensity warfarin afforded a risk reduction for recurrent thromboembolism of 64% over a mean follow-up period of 2.1 years.

The practical implication was that clinical practice should presumably consist of this prolonged, low-intensity anticoagulant regime, at least for patients with a prior event of pulmonary or deep venous thrombosis.<sup>7</sup> In this particular study, patients with CVST were not specifically studied. However, it seems justified to extrapolate the findings on pulmonary embolism or deep venous thrombosis to CVST, because the underlying risk factors associated with these conditions are identical.<sup>8</sup> Thus, applying this potential risk reduction of 64% to the ISCVT study, long-term anticoagulation therapy could have prevented recurrent thromboembolism in 15 of the 24 patients where recurrence occurred in the absence of anticoagulants. In other words, prolonged anticoagulation in the ISCVT study could have resulted in a cumulative risk for recurrent CVST and other thrombotic events of 4.1%, instead of 6.5%.

We, therefore, suggest that, to reduce recurrence of thrombotic events after CVST, it may be prudent to treat these patients with long-term anticoagulation therapy. Future research should determine the optimal duration and intensity of oral anticoagulation therapy that is necessary to optimally reduce the risk for recurrence of thrombotic events.

**B.F.L. van Nuenen, MD**  
**M. Munneke, PhD**  
**B.R. Bloem, MD**

Department of Neurology  
Radboud University Nijmegen Medical Centre  
The Netherlands

### Cumulative Risk for Recurrence of Thrombotic Events After a First Episode of CVST (Overview of Published Studies)

Study	No. of Patients	Mean Follow-Up (Months)	Duration of OAC Treatment (Months)	Cumulative Recurrence of Thromboembolism (%)
ISCVT study <sup>1</sup>	624	18	Mean of 7.7	6.5
Preter et al <sup>4</sup>	77	78	3 to 4	26.0
Breteau et al <sup>2</sup>	55	36	6 (56.4% of patients) 6–36 (12.7% of patients) >36 (30.9% of patients)	5.5*
VENOPORT study <sup>7</sup>	Not reported	22	Not reported	5.0

\*Two patients with protein S deficiency. OAC indicates oral anticoagulation.

1. Ferro JM, Canhao P, Stam J, Bousser MG, Barinagarrementeria F; ISCVT Investigators. Prognosis of cerebral vein and dural sinus thrombosis: results of the International Study on Cerebral Vein and Dural Sinus Thrombosis (ISCVT). *Stroke*. 2004;35:664–670.
2. Breteau G, Mounier-Vehier F, Godefroy O, Gauvrit JY, Mackowiak-Cordoliani MA, Girot M, Bertheloot D, Henon H, Lucas C, Leclerc X, Fourrier F, Pruvo JP, Leys D. Cerebral venous thrombosis 3-year clinical outcome in 55 consecutive patients. *J Neurol*. 2003;250:29–35.
3. Masuhr F, Mehraein S, Einhaupl K. Cerebral venous and sinus thrombosis. *J Neurol*. 2004;251:11–23.
4. Preter M, Tzourio C, Ameri A, Bousser MG. Long-term prognosis in cerebral venous thrombosis. Follow-up of 77 patients. *Stroke*. 1996;27:243–246.
5. Ferro JM, Lopes MG, Rosas MJ, Ferro MA, Fontes J. Long-term prognosis of cerebral vein and dural sinus thrombosis. Results of the VENOPORT study. *Cerebrovasc Dis*. 2002;13:272–278.
6. van den Berg JS, Boerman RH, vd Stolpe A, Kremer HP. Cerebral venous thrombosis: recurrence with fatal course. *J Neurol*. 1999;246:144–146.
7. Ridker PM, Goldhaber SZ, Danielson E, Rosenberg Y, Eby CS, Deitcher SR, Cushman M, Moll S, Kessler CM, Elliott CG, Paulson R, Wong T, Bauer KA, Schwartz BA, Miletich JP, Bounameaux H, Glynn RJ; PREVENT Investigators. Long-term, low-intensity warfarin therapy for the prevention of recurrent venous thromboembolism. *N Engl J Med*. 2003;348:1425–1434.
8. de Bruijn. *Cerebral Versus Sinus Thrombosis. Clinical and Epidemiological Studies*. Amsterdam: Thesis Publishers; 1998.

**Kazushi Tsuda, MD, FAHA**  
 Division of Cardiology  
 Department of Medicine  
 Wakayama Medical University  
 Wakayama, Japan

1. Shen H, Chen GJ, Harvey BK, Bickford PC, Wang Y. Inosine reduces ischemic brain injury in rats. *Stroke*. 2005;36:654–659.
2. Tsuda K, Tsuda S, Goldstein M, Masuyama Y. Inhibitory effects of verapamil on [<sup>3</sup>H]-acetylcholine release in the central nervous system of Sprague-Dawley rats. *Clin Exp Pharmacol Physiol*. 1994;21:527–531.
3. Tsuda K, Tsuda S, Goldstein M, Nishio I, Masuyama Y. Glutamatergic regulation of [<sup>3</sup>H]-acetylcholine release in striatal slices of normotensive and spontaneously hypertensive rats. *Neurochem Int*. 1996;29:231–237.
4. Otori Y, Kusaka S, Kawasaki A, Morimura H, Miki A, Tano Y. Protective effect of nilvadipine against glutamate neurotoxicity in purified retinal ganglion cells. *Brain Res*. 2003;961:213–219.

### Response:

We thank Dr Tsuda for expressing interest in our article.<sup>1</sup> As indicated by Dr Tsuda, increasing evidence has suggested that calcium channels are involved in glutamate-mediated neurodegeneration. In our study, we reported that inosine suppresses glutamate-evoked neuronal excitation in cerebral cortex. It is possible that inosine may have indirect actions on calcium channels through the inhibition of glutamate response during cerebral ischemia. Currently, there are no reports suggesting that calcium-channel blockers have synergistic effects on inosine-mediated response in the central nervous system. Our preliminary studies showed that an N-type calcium channel blocker had a trend toward increasing inosine-mediated protection against ischemic brain injury in adult Sprague-Dawley rats. Animals were injected intracerebroventricularly, before middle cerebral artery occlusion (MCAo), with  $\omega$ -conotoxin GVIA (0.1  $\mu$ g/10  $\mu$ L, n=6) or  $\omega$ -conotoxin GVIA+inosine (25 nmol, n=7). Compared with our previous data,<sup>1</sup>  $\omega$ -conotoxin GVIA alone did not significantly reduce the size of infarction. Coadministration of  $\omega$ -conotoxin GVIA and inosine significantly reduced volume of infarction compared with the stroke animals pretreated with vehicle ( $52.1 \pm 28.9$  mm<sup>3</sup> versus  $187.1 \pm 9.0$  mm<sup>3</sup>;  $P < 0.05$ ; 1-way ANOVA). Although the mean of infarction volume is further reduced by  $\omega$ -conotoxin GVIA in the animals pretreated with inosine, the difference in protection between  $\omega$ -conotoxin GVIA+inosine ( $52.1 \pm 28.9$  mm<sup>3</sup>) and inosine alone ( $118.5 \pm 18.4$  mm<sup>3</sup>) is not statistically significant. We observed that the majority of animals that received  $\omega$ -conotoxin GVIA with or without inosine developed seizures after MCAo. Other studies have shown nifedipine or felodipine suppressed hypoxia-evoked inosine release in rat cerebral cortex.<sup>2</sup> The combined effects of inosine and calcium channel blockers in cerebral ischemia are complicated by such observations. Further studies are needed to definitively address the interaction of inosine and calcium channels.

**Hui Shen, MD**  
**Guann-Juh Chen, MD**  
**Brandon Harvey, PhD**  
**Yun Wang, MD, PhD**

Neural Protection and Regeneration Section  
 National Institute on Drug Abuse  
 Baltimore, Md

1. Shen H, Chen G-J, Harvey BK, Bickford PL, Wang Y. Inosine reduces ischemic brain injury in rats. *Stroke*. 2005;36:654–659.
2. Phillis JW, O'Regan MH, Walter GA. Effects of nifedipine and felodipine on adenosine and inosine release from the hypoxic rat cerebral cortex. *J Cereb Blood Flow Metab*. 1988;8:179–185.

## Inosine, Calcium Channels, and Neuroprotection Against Ischemic Brain Injury

To the Editor:

We read with great interest the article by Shen et al,<sup>1</sup> dealing with neuroprotective effects of inosine against ischemic brain injury. The results of their study demonstrated that intracerebroventricular administration of inosine before middle cerebral artery occlusion in rats resulted in a higher level of locomotor activity and less cerebral infarction. In addition, they indicated that coadministration of selective A3 receptor antagonist MRS1191 significantly attenuated inosine-mediated protection. In the electrophysiological study, it was shown that inosine antagonized glutamate-induced excitation in cerebral cortical neurons. The authors proposed that inosine may inhibit glutamate postsynaptic responses and reduce cerebral infarction via the activation of A3 receptors.

Several studies have shown the mechanisms for glutamate-induced excitation of neural cells. In a study we presented earlier, changes in acetylcholine (ACh) release evoked by glutamate was investigated in rat central nervous system.<sup>2,3</sup> In an in vitro study, we showed that glutamate increased the release of ACh from rat striatum, which was inhibited by the *N*-methyl-D-aspartate type of glutamate receptor antagonist MK-801.<sup>3</sup> Furthermore, it was demonstrated that the calcium-channel blocker verapamil significantly reduced glutamate-evoked ACh release and that the inhibitory effect was more pronounced in the presence of magnesium.<sup>2</sup> It might be possible that the calcium-channel blocker could inhibit the *N*-methyl-D-aspartate receptor-dependent calcium current in neural cells. It was also shown that the calcium-channel blocker nilvadipine blocked glutamate-evoked intracellular calcium increase with a concomitant reduction in glutamate-induced apoptosis in purified retinal ganglion cells,<sup>4</sup> suggesting that the calcium-channel blocker might inhibit glutamate-induced apoptotic cell death by blocking calcium influx. Therefore, we would like to know whether inosine and calcium-channel blockers might have a synergistic effect in neural protection in the present study by Shen et al. Although the authors demonstrated that inosine did not alter basal glutamate release, nor did it reduce ischemia-evoked glutamate overflow in the central nervous system, it can be speculated that inosine might interfere with glutamate receptor-mediated calcium conductance in neural cells and might have therapeutic potential as a neuroprotective agent against stroke-induced damage.



### Comment on the *Phosphodiesterase 4D* Replication Study by Bevan et al

To the Editor:

We observe with interest the results of the study by Bevan et al,<sup>1</sup> where they attempted to replicate our findings associating the *PDE4D* gene with stroke. We appreciate the effort by the authors and their presentation of their data in some detail. The latter allowed us to compare their results with ours, including some unpublished results. In our original publication,<sup>2</sup> the main findings are summarized in Figure 4, where haplotypes are grouped into 3 categories: the at-risk haplotypes, the “wild-type,” and the protective haplotype. Although the definition of the at-risk haplotypes involved multiple markers, the protective haplotype, in our data, is characterized by one single-nucleotide polymorphism, specifically allele A of SNP45. This makes it particularly suitable for a replication in a different population. In Table I of Bevan et al, allele A has frequency of 15.4% in 621 stroke patients and 18.7% in 848 controls. We calculated that this corresponds to 191 copies of allele A and 1051 Gs in stroke patients, and 317 As and 1379 Gs in controls. Applying Fisher exact test gives a 2-sided probability value of 0.02 and a 1-sided probability value of 0.01. For the same allele counts, estimated risk of A relative to G is 0.79, or equivalently risk of G relative to A is 1.26. This agrees well with our estimated relative risk of 1.33 for allele G for all stroke patients (in Table 1 of Gretarsdottir et al<sup>2</sup>). Hence, even though Bevan et al did not consider this result as significant, which could be attributable to the use of

somewhat different statistical tests and adjustment for multiple comparisons, we are nonetheless encouraged by these results, especially because we have obtained similar results for SNP45 in a replication study of ours (Gretarsdottir et al, unpublished data, 2005). In summary, we appreciate the difficulties of getting definitive results for a replication study when the effect of the variant may only be modest and proper care has to be taken to adjust for multiple testing of variants and phenotypes. And we agree with the authors that further investigation is needed to develop a better understanding of the role of *PDE4D* in stroke.

**Solveig Gretarsdottir, PhD**  
**Jeffrey Gulcher, MD, PhD**  
**Gudmar Thorleifsson, PhD**  
**Augustine Kong, PhD**  
**Kari Stefansson, MD, DrMed**  
*DeCODE Genetics*  
*Reykjavik, Iceland*

1. Bevan S, Porteous L, Sitzer M, Markus HS. Phosphodiesterase 4D gene, ischemic stroke, and asymptomatic carotid atherosclerosis. *Stroke*. 2005; 36:949–953.
2. Gretarsdottir S, Thorleifsson G, Reynisdottir ST, Manolescu A, Jonsdottir S, Jonsdottir T, Gudmundsdottir T, Bjarnadottir S, Einarsson OB, Gudjonsdottir HM, Hawkins M, Gudmundsson G, Gudmundsdottir H, Andrason H, Gudmundsdottir AS, Sigurdardottir M, Chou TT, Nahmias J, Goss S, Sveinbjörnsdottir S, Valdimarsson E, Jakobsson F, Agnarsson U, Gudnason V, Kong A, Thorgeirsson G, Fingerle J, Gurney M, Frigge ML, Kong A, Stefansson K, Gulcher JR. The gene encoding phosphodiesterase 4D confers risk of ischemic stroke. *Nat Genet*. 2003;35:131–138.