

Certain Class of Analytic Functions Defined by Ruscheweyh Derivative with Varying Arguments

RABHA MOHAMED EL-ASHWAH

Department of Mathematics, Faculty of Science, Damietta University, New Damietta 34517, Egypt

e-mail : r_elashwah@yahoo.com

MOHAMED KAMAL AOUF

Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 33516, Egypt

e-mail : mkaouf127@yahoo.com

AHMED HASSAN AND ALAA HASSAN*

Department of Mathematics, Faculty of Science, Zagazig University, Zagazig 44519, Egypt

e-mail : aam_hassan@yahoo.com, alaahassan1986@yahoo.com

ABSTRACT. In this paper we derive some results for certain new class of analytic functions defined by using Ruscheweyh derivative with varying arguments.

1. Introduction

Let \mathcal{A} denote the class of functions of the form:

$$(1.1) \quad f(z) = z + \sum_{k=2}^{\infty} a_k z^k,$$

which are analytic and univalent in the open unit disc $U = \{z \in \mathbb{C} : |z| < 1\}$. Given

* Corresponding Author.

Received September 17, 2012; accepted April 22, 2013.

2010 Mathematics Subject Classification: 30C45.

Keywords and Phrases: analytic functions, univalent, Hadamard product, Ruscheweyh derivative, extreme points.

two functions $f, g \in \mathcal{A}$, where $f(z)$ is given by (1.1) and $g(z)$ is given by

$$(1.2) \quad g(z) = z + \sum_{k=2}^{\infty} b_k z^k,$$

the Hadamard product $(f * g)(z)$ is defined by

$$(1.3) \quad (f * g)(z) = z + \sum_{k=2}^{\infty} a_k b_k z^k = (g * f)(z).$$

By using the Hadamard product, Ruscheweyh [7] defined

$$(1.4) \quad D^\gamma f(z) = \frac{z}{(1-z)^{\gamma+1}} * f(z) \quad (\gamma \geq -1).$$

Also Ruscheweyh [7] observed that

$$(1.5) \quad D^n f(z) = \frac{z(z^{n-1}f(z))^{(n)}}{n!} \quad (n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}; \mathbb{N} = \{1, 2, \dots\}).$$

The symbol $D^n f(z)$ ($n \in \mathbb{N}_0$) was called the n -th order Ruscheweyh derivative of $f(z)$ by Al-Amiri [1]. We note that

$$D^0 f(z) = f(z) \text{ and } D^1 f(z) = z f'(z).$$

It is easy to see that

$$(1.6) \quad D^n f(z) = z + \sum_{k=2}^{\infty} \delta(n, k) a_k z^k,$$

where

$$(1.7) \quad \delta(n, k) = \binom{n+k-1}{n}.$$

In [4] Attiya and Aouf defined the class $Q(n, \lambda, A, B)$ as follows:

Definition 1. ([4]) Let $Q(n, \lambda, A, B)$ denote the subclass of \mathcal{A} consisting of functions $f(z)$ of the form (1.1) such that

$$(1.8) \quad (1-\lambda)(D^n f(z))' + \lambda(D^{n+1} f(z))' \prec \frac{1+Az}{1+Bz}$$

$$(\lambda \geq 0; -1 \leq A < B \leq 1; 0 < B \leq 1; n \in \mathbb{N}_0; z \in U).$$

Specializing the parameters λ, A, B and n , we can obtain different classes studied by various authors:

- (i) $Q(0, \lambda, 2\alpha - 1, 1) = R(\lambda, \alpha)$ ($0 \leq \alpha < 1, \lambda \geq 0$) (see Altintas [3])
- (ii) $Q(0, 0, 2\alpha - 1, 1) = T^{**}(\alpha)$ ($0 \leq \alpha < 1$) (see Sarangi and Uralegaddi [8] and Al-Amiri [2])
- (iii) $Q(n, 0, 2\alpha - 1, 1) = Q_n(\alpha)$ ($0 \leq \alpha < 1, n \in \mathbb{N}_0$) (see Uralegaddi and Sarangi [11])
- (iv) $Q(0, 0, (2\alpha - 1)\beta, \beta) = P^*(\alpha, \beta)$ ($0 \leq \alpha < 1, 0 < \beta \leq 1$) (see Gupta and Jain[5])
- (v) $Q(0, 0, ((1 + \mu)\alpha - 1)\beta, \mu\beta) = P^*(\alpha, \beta, \mu)$ ($0 \leq \alpha < 1, 0 < \beta \leq 1, 0 < \mu \leq 1$) (see Owa and Aouf [6]).

Also we note that:

$$(i) \quad Q(0, \lambda, A, B) = R(\lambda, A, B) \tag{1.9}$$

$$= \left\{ f(z) \in \mathcal{A} : f'(z) + \lambda z f''(z) \prec \frac{1 + Az}{1 + Bz} \quad (\lambda \geq 0; -1 \leq A < B \leq 1; 0 < B \leq 1; z \in U) \right\}$$

$$(ii) \quad Q(n, 0, A, B) = Q_n(A, B) = \tag{1.10}$$

$$\left\{ f(z) \in \mathcal{A} : (D^n f(z))' \prec \frac{1 + Az}{1 + Bz} \quad (\lambda \geq 0; -1 \leq A < B \leq 1; 0 < B \leq 1; n \in \mathbb{N}_0; z \in U) \right\}$$

Silverman [9] defined the class of univalent functions $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$ for which $arg(a_k)$ prescribed in such a way that $f(z)$ is univalent if and only if $f(z)$ is starlike as follows:

Definition 2.([9]) A function $f(z)$ of the form (1.1) is said to be in the class $V(\theta_k)$ if $f \in \mathcal{A}$ and $arg(a_k) = \theta_k$ for all $k \geq 2$. If further more there exist a real number δ such that $\theta_k + (k - 1)\delta \equiv \pi \pmod{2\pi}$ ($k \geq 2$), then $f(z)$ is said to be in the class $V(\theta_k, \delta)$. The union of $V(\theta_k, \delta)$ taken over all possible sequences $\{\theta_k\}$ and all possible real numbers δ is denoted by V .

Let $VQ(n, \lambda, A, B)$ denote the subclass of V consisting of functions $f(z) \in Q(n, \lambda, A, B)$.

We note that $VQ(0, 0, 2\alpha - 1, 1) = C_\alpha$ ($0 \leq \alpha < 1$) = $\{f \in V : \mathbf{Re}\{f'(z)\} > \alpha\}$, studied by Srivastava and Owa [10].

Also we note that by specializing the parameters λ, A, B and n we can obtain different classes with varying arguments:

- (i) $VQ(0, \lambda, 2\alpha - 1, 1) = VR(\lambda, \alpha)$ ($0 \leq \alpha < 1, \lambda \geq 0$)
- (ii) $VQ(0, 0, 2\alpha - 1, 1) = VT^{**}(\alpha)$ ($0 \leq \alpha < 1$)
- (iii) $VQ(n, 0, 2\alpha - 1, 1) = VQ_n(\alpha)$ ($0 \leq \alpha < 1, n \in \mathbb{N}_0$)

- (iv) $VQ(0, 0, (2\alpha - 1)\beta, \beta) = VP^*(\alpha, \beta)$ ($0 \leq \alpha < 1, 0 < \beta \leq 1$)
 (v) $VQ(0, 0, ((1 + \mu)\alpha - 1)\beta, \mu\beta) = VP^*(\alpha, \beta, \mu)$ ($0 \leq \alpha < 1, 0 < \beta \leq 1, 0 < \mu \leq 1$)
 (vi) $VQ(0, \lambda, A, B) = VR(\lambda, A, B)$ ($\lambda \geq 0, -1 \leq A < B \leq 1, 0 < B \leq 1$)
 (vii) $VQ(n, 0, A, B) = VQ_n(A, B)$ ($-1 \leq A < B \leq 1, 0 < B \leq 1, n \in \mathbb{N}_0$)

In this paper we obtain coefficient bounds for functions in the class $VQ(n, \lambda, A, B)$, further we obtain distortion bounds and the extreme points for functions in this class.

2. Coefficient Estimates

Unless otherwise mentioned, we assume in the reminder of this paper that, $\lambda \geq 0, -1 \leq A < B \leq 1, 0 < B \leq 1, n \in \mathbb{N}_0, \delta(n, k)$ and C_k are given by (1.7) and (2.2) respectively and $z \in U$.

Theorem 1. *Let the function $f(z)$ defined by (1.1) be in V . Then $f(z) \in VQ(n, \lambda, A, B)$, if and only if*

$$(2.1) \quad \sum_{k=2}^{\infty} k\delta(n, k)C_k |a_k| \leq (B - A)(n + 1),$$

where

$$(2.2) \quad C_k = (1 + B)[n + 1 + \lambda(k - 1)].$$

Proof. Suppose that $f(z) \in VQ(n, \lambda, A, B)$. Then

$$(2.3) \quad h(z) = (1 - \lambda)(D^n f(z))' + \lambda(D^{n+1} f(z))' = \frac{1 + Aw(z)}{1 + Bw(z)},$$

where

$$w \in H = \{w \text{ analytic, } w(0) = 0 \text{ and } |w(z)| < 1, z \in U\}.$$

Thus we get

$$w(z) = \frac{1 - h(z)}{Bh(z) - A}.$$

Therefore

$$h(z) = 1 + \sum_{k=2}^{\infty} \frac{k[n + 1 + \lambda(k - 1)]\delta(n, k)}{n + 1} a_k z^{k-1},$$

and $|w(z)| < 1$ implies

$$(2.4) \quad \left| \frac{\sum_{k=2}^{\infty} \frac{k[n + 1 + \lambda(k - 1)]\delta(n, k)}{n + 1} a_k z^{k-1}}{(B - A) + B \sum_{k=2}^{\infty} \frac{k[n + 1 + \lambda(k - 1)]\delta(n, k)}{n + 1} a_k z^{k-1}} \right| < 1.$$

Since $f(z) \in V$, $f(z)$ lies in the class $V(\theta_k, \delta)$ for some sequence $\{\theta_k\}$ and a real number δ such that

$$\theta_k + (k - 1)\delta \equiv \pi \pmod{2\pi} \quad (k \geq 2).$$

Set $z = re^{i\delta}$ in (2.4), we get

$$(2.5) \quad \left| \frac{\sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| r^{k-1}}{(B-A) - B \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| r^{k-1}} \right| < 1.$$

Since $\text{Re}\{w(z)\} < |w(z)| < 1$, we have

$$(2.6) \quad \text{Re} \left\{ \frac{\sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| r^{k-1}}{(B-A) - B \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| r^{k-1}} \right\} < 1.$$

Hence

$$(2.7) \quad \sum_{k=2}^{\infty} kC_k\delta(n,k) |a_k| r^{k-1} \leq (B-A)(n+1).$$

Letting $r \rightarrow 1$ in (2.7), we get (2.1).

Conversely, $f(z) \in V$ and satisfies (2.1). Since $r^{k-1} < 1$. So we have

$$\begin{aligned} \left| \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| z^{k-1} \right| &\leq \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| r^{k-1} \\ &\leq (B-A) - B \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} |a_k| r^{k-1} \\ &\leq \left| (B-A) - B \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} a_k z^{k-1} \right| \\ &\leq \left| (B-A) + B \sum_{k=2}^{\infty} \frac{k[n+1+\lambda(k-1)]\delta(n,k)}{n+1} a_k z^{k-1} \right| \end{aligned}$$

which gives (2.4) and hence follows that

$$(1 - \lambda) (D^n f(z))' + \lambda (D^{n+1} f(z))' = \frac{1 + Aw(z)}{1 + Bw(z)}$$

that is $f(z) \in VQ(n, \lambda, A, B)$. This completes the proof of Theorem 1.

Corollary 1. Let the function $f(z)$ defined by (1.1) be in the class $VQ(n, \lambda, A, B)$. Then

$$|a_k| \leq \frac{(B-A)(n+1)}{kC_k\delta(n,k)} \quad (k \geq 2).$$

The result (2.1) is sharp for the function $f(z)$ defined by

$$(2.8) \quad f(z) = z + \frac{(B-A)(n+1)}{kC_k\delta(n,k)} e^{i\theta_k} z^k \quad (k \geq 2).$$

3. Distortion Theorems

Theorem 2. Let the function $f(z)$ defined by (1.1) be in the class $VQ(n, \lambda, A, B)$. Then

$$(3.1) \quad |z| - \frac{B-A}{C_2} |z|^2 \leq |f(z)| \leq |z| + \frac{B-A}{C_2} |z|^2.$$

The result is sharp.

Proof. We employ the same technique as used by Silverman [9]. In view of Theorem 1, since

$$(3.2) \quad \Phi(k) = C_k\delta(n, k),$$

is an increasing function of k ($k \geq 2$), we have

$$\Phi(2) \sum_{k=2}^{\infty} |a_k| \leq \sum_{k=2}^{\infty} \Phi(k) |a_k| \leq (B-A)(n+1),$$

that is

$$(3.3) \quad \sum_{k=2}^{\infty} |a_k| \leq \frac{(B-A)(n+1)}{\Phi(2)} = \frac{(B-A)}{2C_2}.$$

Thus we have

$$|f(z)| \leq |z| + \sum_{k=2}^{\infty} |a_k| |z|^k \leq |z| + |z|^2 \sum_{k=2}^{\infty} |a_k|,$$

Thus

$$|f(z)| \leq |z| + \frac{(B-A)}{2C_2} |z|^2.$$

Similarly, we get

$$|f(z)| \geq |z| - \sum_{k=2}^{\infty} |a_k| |z|^k \geq |z| - |z|^2 \sum_{k=2}^{\infty} |a_k|.$$

Thus

$$|f(z)| \geq |z| - \frac{(B - A)}{2C_2} |z|^2.$$

This completes the proof of Theorem 2. Finally the result is sharp for the function

$$(3.4) \quad f(z) = z + \frac{(B - A)}{2C_2} e^{i\theta_2} z^2,$$

at $z = \pm |z| e^{-i\theta_2}$.

Corollary 2. Under the hypotheses of Theorem 2, $f(z)$ is included in a disc with center at the origin and radius r_1 given by

$$(3.5) \quad r_1 = 1 + \frac{(B - A)}{2C_2}.$$

Theorem 3. Let the function $f(z)$ defined by (1.1) be in the class $VQ(n, \lambda, A, B)$. Then

$$(3.6) \quad 1 - \frac{(B - A)}{C_2} |z| \leq |f'(z)| \leq 1 + \frac{(B - A)}{C_2} |z|.$$

The result is sharp.

Proof. Similarly $\frac{\Phi(k)}{k}$ is an increasing function of $k(k \geq 2)$, where $\Phi(k)$ is defined by (3.2). In view of Theorem 1, we have

$$\frac{\Phi(2)}{2} \sum_{k=2}^{\infty} k |a_k| \leq \sum_{k=2}^{\infty} \Phi(k) |a_k| \leq (B - A)(n + 1),$$

that is

$$\sum_{k=2}^{\infty} k |a_k| \leq \frac{(B - A)}{\Phi(2)} = \frac{(B - A)}{C_2}.$$

Thus we have

$$(3.7) \quad |f'(z)| \leq 1 + |z| \sum_{k=2}^{\infty} k |a_k| \leq 1 + \frac{(B - A)}{C_2} |z|.$$

Similarly

$$(3.8) \quad |f'(z)| \geq 1 - |z| \sum_{k=2}^{\infty} k |a_k| \geq 1 - \frac{(B - A)}{C_2} |z|.$$

Finally, we can see that the assertions of Theorem 3 are sharp for the function $f(z)$ defined by (3.4). This completes the proof of Theorem 3.

Corollary 3. Under the hypotheses of Theorem 3, $f'(z)$ is included in a disc with center at the origin and radius r_2 given by

$$(3.9) \quad r_2 = 1 + \frac{(B-A)}{C_2}.$$

4. Extreme Points

Theorem 4. Let the function $f(z)$ defined by (1.1) be in the class $VQ(n, \lambda, A, B)$, with $\arg a_k = \theta_k$, where $\theta_k + (k-1)\delta \equiv \pi \pmod{2\pi}$ ($k \geq 2$). Define

$$f_1(z) = z$$

and

$$f_k(z) = z + \frac{(B-A)(n+1)}{kC_k\delta(n,k)} e^{i\theta_k} z^k \quad (k \geq 2; z \in U).$$

Then $f(z) \in VQ(n, \lambda, A, B)$ if and only if $f(z)$ can be expressed in the form $f(z) = \sum_{k=1}^{\infty} \mu_k f_k(z)$, where $\mu_k \geq 0$ and $\sum_{k=1}^{\infty} \mu_k = 1$.

Proof. If $f(z) = \sum_{k=1}^{\infty} \mu_k f_k(z)$ with $\mu_k \geq 0$ and $\sum_{k=1}^{\infty} \mu_k = 1$, then

$$\begin{aligned} & \sum_{k=2}^{\infty} [kC_k\delta(n,k)] \frac{(B-A)(n+1)}{kC_k\delta(n,k)} \mu_k = \sum_{k=2}^{\infty} (B-A)(n+1) \mu_k \\ & = (1 - \mu_1)(B-A)(n+1) \leq (B-A)(n+1). \end{aligned}$$

Hence $f(z) \in VQ(n, \lambda, A, B)$.

Conversely, let the function $f(z)$ defined by (1.1) be in the class $VQ(n, \lambda, A, B)$, define

$$\mu_k = \frac{kC_k\delta(n,k)}{(B-A)(n+1)} |a_k| \quad (k \geq 2)$$

and

$$\mu_1 = 1 - \sum_{k=2}^{\infty} \mu_k.$$

From Theorem 1, $\sum_{k=2}^{\infty} \mu_k \leq 1$ and so $\mu_1 \geq 0$. Since $\mu_k f_k(z) = \mu_k z + a_k z^k$, then

$$\sum_{k=1}^{\infty} \mu_k f_k(z) = z + \sum_{k=2}^{\infty} a_k z^k = f(z).$$

This completes the proof of Theorem 4.

Remarks.

- (i) Putting $\lambda = n = 0$, $A = 2\alpha - 1$ ($0 \leq \alpha < 1$) and $B = 1$ in all the above results, we obtain the corresponding results obtained by Srivastava and Owa [10];
- (ii) Putting $n = 0$ in all the above results, we obtain the corresponding results for the class $VR(\lambda, A, B)$ of which $R(\lambda, A, B)$ is given by (1.9);
- (iii) Putting $\lambda = 0$ in all the above results, we obtain the corresponding results for the class $VQ_n(A, B)$ of which $Q_n(A, B)$ is given by (1.10).

References

- [1] H. S. Al-Amiri, On Ruscheweyh derivatives, *Ann. Polon. Math.*, **38**(1980), 87-94.
- [2] H. S. Al-Amiri, On a subclass of close-to-convex functions with negative coefficients, *Math. (Cluj)*, **31**(1989), 1-7.
- [3] O. Altintas, A subclass of analytic functions with negative coefficients, *Hacettepe Bull. Natur. Sci. Engrg.*, **19**(1990), 15-24.
- [4] A. A. Attiya and M. K. Aouf, A study on certain class of analytic functions defined by Ruscheweyh derivative, *Soochow J. Math.*, **33**(2)(2007), 273-289.
- [5] V. P. Gupta and P. K. Jain, Certain classes of univalent functions with negative coefficients II, *Bull. Austral. Math. Soc.*, **15**(1976), 467-473.
- [6] S. Owa and M. K. Aouf, On subclasses of univalent functions with negative coefficients II, *Pure Appl. Math. Sci.*, **29**(1:2)(1989), 131-139.
- [7] St. Ruscheweyh, New criteria for univalent functions, *Proc. Amer. Math. Soc.*, **49**(1975), 109-115.
- [8] S. M. Sarangi and B. A. Uralegaddi, The radius of convexity and starlikeness for certain classes of analytic functions with negative coefficients I, *Rend. Acad. Naz. Lincei*, **65**(1978), 38-42.
- [9] H. Silverman, Univalent functions with varying arguments, *Houston J. Math.*, **17**(1981), 283-287.
- [10] H. M. Srivastava and S. Owa, Certain classes of analytic functions with varying arguments, *J. Math. Anal. Appl.*, **136**(1)(1988), 217-228.
- [11] B. A. Uralegaddi and S. M. Sarangi, Some classes of univalent functions with negative coefficients, *An. Stiint. Univ. Al. I. Cuza Iasi Sect. I a Mat. (N. S.)*, **34**(1988), 7-11.