

 Open access • Journal Article • DOI:10.1071/WF13058

## Challenges of assessing fire and burn severity using field measures, remote sensing and modelling — [Source link](#)

Penelope Morgan, Robert E. Keane, Gregory K. Dillon, Theresa B. Jain ...+5 more authors

**Institutions:** University of Idaho, United States Forest Service

**Published on:** 24 Dec 2014 - International Journal of Wildland Fire (CSIRO PUBLISHING)

**Topics:** Fire regime and Fire ecology

Related papers:

- [Fire intensity, fire severity and burn severity: a brief review and suggested usage](#)
- [Remote sensing techniques to assess active fire characteristics and post-fire effects](#)
- [Quantifying burn severity in a heterogeneous landscape with a relative version of the delta Normalized Burn Ratio \(dNBR\)](#)
- [A Project for Monitoring Trends in Burn Severity](#)
- [Calibration and validation of the relative differenced Normalized Burn Ratio \(RdNBR\) to three measures of fire severity in the Sierra Nevada and Klamath Mountains, California, USA](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/challenges-of-assessing-fire-and-burn-severity-using-field-46on1c7q5g>

## Challenges of assessing fire and burn severity using field measures, remote sensing and modelling

Penelope Morgan<sup>A,E</sup>, Robert E. Keane<sup>B</sup>, Gregory K. Dillon<sup>B</sup>, Theresa B. Jain<sup>C</sup>, Andrew T. Hudak<sup>C</sup>, Eva C. Karau<sup>B</sup>, Pamela G. Sikkink<sup>C</sup>, Zachary A. Holden<sup>D</sup> and Eva K. Strand<sup>A</sup>

<sup>A</sup>University of Idaho, Department of Forest, Rangeland, and Fire Sciences, 875 Perimeter Drive MS 1133, Moscow, ID 83844, USA.

<sup>B</sup>USDA Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory, Missoula, MT 59807, USA.

<sup>C</sup>USDA Forest Service, Rocky Mountain Research Station, Forestry Sciences Laboratory, Moscow, ID 83843, USA.

<sup>D</sup>USDA Forest Service, Northern Region, Missoula, MT 59807, USA.

<sup>E</sup>Corresponding author. Email: [pmorgan@uidaho.edu](mailto:pmorgan@uidaho.edu)

**Abstract.** Comprehensive assessment of ecological change after fires have burned forests and rangelands is important if we are to understand, predict and measure fire effects. We highlight the challenges in effective assessment of fire and burn severity in the field and using both remote sensing and simulation models. We draw on diverse recent research for guidance on assessing fire effects on vegetation and soil using field methods, remote sensing and models. We suggest that instead of collapsing many diverse, complex and interacting fire effects into a single severity index, the effects of fire should be directly measured and then integrated into severity index keys specifically designed for objective severity assessment. Using soil burn severity measures as examples, we highlight best practices for selecting imagery, designing an index, determining timing and deciding what to measure, emphasising continuous variables measureable in the field and from remote sensing. We also urge the development of a severity field assessment database and research to further our understanding of causal mechanisms linking fire and burn severity to conditions before and during fires to support improved models linking fire behaviour and severity and for forecasting effects of future fires.

**Additional keywords:** fire ecology, fire effects, mapping, remote sensing, retrospective assessment, wildfire environment.

Received 13 April 2013, accepted 14 July 2014, published online 25 November 2014

### Introduction

Wildland fires commonly burn extensive areas in forests and rangelands, often jeopardising homes and municipal watersheds, and other places important to society. Fire as a disturbance drives ecosystem composition and structure from sites to landscapes globally (Bowman *et al.* 2009), thus severity is central to evaluating and predicting ecological conditions before, during and after fire. Fire scientists and managers routinely use field and remotely sensed evaluations of fire and burn severity, defined as the magnitude of change due to fire (Lentile *et al.* 2006; Keeley 2009), to describe fire effects on fuels, vegetation, wildlife habitat and soils (Ryan and Noste 1985; Smith 2000; Carey *et al.* 2003; Key and Benson 2006). This information supports strategic planning before and during fires, prioritising post-fire mitigation to diminish flooding and erosion potential and to foster vegetation recovery post-fire (Robichaud *et al.* 2003, 2007a; Beschta *et al.* 2004; Kuenzi *et al.* 2008),

and making understanding fire and burn severity central to ecologically based fire management.

Despite widespread use, the consistent, objective, repeatable quantification of fire and burn severity remains elusive (Jain *et al.* 2004; Key 2006; Lentile *et al.* 2006; Keeley 2009), and without reliable assessments, the causes and ecological consequences of severity will remain poorly understood. Given the numerous definitions of fire and burn severity, it is important for all users to explain how they define and assess severity (Simard 1991; Lentile *et al.* 2006; Jain *et al.* 2008). As Simard (1991) wrote, ‘no two people interpret fire severity the same’ because observers focus on fire effects selected for a particular set of local objectives or outcomes. These assessments are rarely impartial, standardised, consistent or comprehensive. Moreover, because fire effects can vary in their scale of impact and temporal recovery, one spatial or temporal scale may not fit all objectives, increasing the challenge of objectively and

quantitatively assessing fire or burn severity (Simard 1991; Graham *et al.* 2004). We follow the growing body of literature that differentiates the two severity terms based on temporal and ecological context (Lentile *et al.* 2006; French *et al.* 2008; Veraverbeke *et al.* 2010a, 2010b). We use 'fire severity' to describe the immediate fire effects and 'burn severity' for the longer-term effects of fires on vegetation and soils (Key and Benson 2006; Lentile *et al.* 2006, 2007; French *et al.* 2008; Veraverbeke *et al.* 2010a, 2010b). This is consistent with the US National Wildfire Coordinating Group (2012) definition of fire severity as 'degree to which a site has been altered or disrupted by fire, loosely, a product of fire intensity and residence time', but not with the NWCG definition of burn severity as 'soil heating, large fuel and duff consumption, consumption of the litter and organic layer beneath trees and isolated shrubs, and mortality of buried plant parts'; though these clearly influence vegetation response and soil effects (Chafer *et al.* 2004; Chafer 2008; Neary *et al.* 2008). Similar to French *et al.* (2008), we use 'fire and burn severity' or more simply 'severity' unless we refer specifically to immediate fire effects (fire severity) or fire effects measured in the following year or growing season that include some secondary effects and ecological response (burn severity).

Fire and burn severity are often related to fire behaviour either directly or indirectly, but severity is often confused with fire intensity. One major source of misunderstanding about severity occurs when fire behaviour measures are used to characterise severity (Alexander 1982; Ryan and Noste 1985; Keeley 2009). Including fire behaviour measures in severity descriptions is often problematic given that actual fire behaviour measures are likely lacking. Whereas fire behaviour, especially smouldering combustion, is a critical causal mechanism of fire and burn severity, fire behaviour attributes have often been inappropriately used to describe fire effects (Moreno and Oechel 1991). Oliveras *et al.* (2009) found that fire severity was higher in flanking and head fires than backing fires, and Keeley *et al.* (2008) and Jain *et al.* (2004) emphasised that fire behaviour is a critical element in describing fire severity. Thus, fire and burn severity are intimately linked to fire behaviour, but fire behaviour does not fully describe the effect of fire on the ecosystem, especially if it is focussed only on flaming combustion, because it is missing critical ecological responses from the heat of the fire.

Assessing and predicting severity is challenging because fires have multiple effects that are often assessed in different contexts. Assessments of fire and burn severity are done for post-fire mitigation of erosion potential and invasive species establishment, soil erosion potential (Fox *et al.* 2008; Clark and McKinley 2011), post-fire vegetation recovery including tree seedlings (Turner *et al.* 1999; Díaz-Delgado *et al.* 2003; Miller *et al.* 2003; Pausas *et al.* 2003; Beschta *et al.* 2004; Lentile *et al.* 2006), wildlife habitat (Zarriello *et al.* 1995), species of concern (Kotliar *et al.* 2008) and overall vegetation conditions (Bisson *et al.* 2008; Guay 2011). Soil burn severity can also be used to predict the physical, chemical or biological effects (Jain *et al.* 2012), including water repellency (Lewis *et al.* 2006), erodibility (Pierson *et al.* 2001) and nutrient availability (Belillas and Feller 1998). Severity is also a fire regime attribute (Beukema and Kurz 1998; Morgan *et al.* 2001; Barrett *et al.* 2006; Keane

*et al.* 2006), perhaps the most difficult one to quantify because in this context it is used more conceptually and lacks measurement units; despite this, no other fire regime attribute is as important in fire ecology. Severity is the basis for a national fire atlas for the US (Eidenshink *et al.* 2007; <http://www.mtbs.gov>) and has been used to link landscape patterns and scales of disturbance processes (Turner *et al.* 1994; Chuvieco 1999; Hudak *et al.* 2007a, 2007b). Managers and scientists use fire and burn severity classifications to evaluate prescribed fire success (Ryan and Noste 1985), stratify post-fire vegetation and soil response, and describe burn patterns (Carey *et al.* 2003).

Despite the extensive scientific literature describing fire effects and elements of severity in different ecosystems, there are few widely accepted or standardised measures of severity consistently applicable within such different assessment contexts (Halofsky and Hibbs 2009). Severity is variously used as a concept, a continuous variable, a class and an index. Ideally, metrics of severity should be specific, meaningful and readily interpretable, as well as measureable in the field and remotely, and at multiple spatial and temporal scales (Hudak *et al.* 2007b). Satellite imagery, statistical and simulation modelling, and standardised efficient field sampling have facilitated the generation of quick and inexpensive fire and burn severity maps, minimising the need for extensive resource-intensive, and potentially dangerous, post-fire field sampling. However, all of these advances are limited by ecological, technical and logistical issues.

We address four objectives in this paper. We examine recent advances in predicting and assessing fire and burn severity in the field, remotely and using models. We discuss the numerous factors and associated interactions that influence severity and challenge our ability to consistently assess or predict it. We provide guidance for those who wish to use severity assessments in planning and implementing land management activities, using vegetation and soil burn severity as examples. We propose a strategy for describing fire and burn severity in the future to reduce the confusion and complexity of defining, measuring and assessing severity, while providing the ability to design severity assessments for specific uses. Our focus is on new research building upon previous work conducted by Lentile *et al.* (2006) and Keeley (2009).

## Assessing severity

### *Assessing fire and burn severity in the field*

Many fire effects have been measured to describe fire and burn severity (French *et al.* 2008; Jain *et al.* 2012) (Table 1). Changes in overstorey trees or shrubs are the primary fire effect implicit in most fire and burn severity assessments (Regelbrugge and Smith 1994; Patterson and Yool 1998; Turner *et al.* 1999; Smith *et al.* 2007). However, assessing plant mortality can be somewhat problematic in ecosystems where the dominant plants often re-sprout quickly after fire, such as in many hardwood forests, shrublands and grasslands. Therefore, Keeley (2009) advocated using the amount of biomass consumed instead of plant mortality, and Wang and Glenn (2009) used changes in height to measure severity in shrublands. Surface fuel consumption has also been used as an important indicator of severity (Schimmel and Granstrom 1996; Boby *et al.* 2010;

**Table 1. Quantifying burn severity for forest soils (Jain *et al.* 2012)**

Each subject area includes the literature sources, application, number of categories identified and range of possible post-fire outcomes. Cells with dashes (–) denote an outcome that was not included. Post-fire characteristics most noted had two primary indicators: first, the amount of pre-fire surface organic matter (e.g., litter, humus, rotten wood) present, expressed as abundant, present or absent on the forest floor; and second, whether the litter was scorched (S) and the assessed state of the exposed mineral soil – unburned (U), blackened from combustion (B), gray/white (G) from ash or orange (O) from mineralogical changes and fire residuals. Numbers in the table indicate the level of severity used by that classification, where 0 is unburned and included as a class, and 1–5 represents relative low to high severity.

Source	Application	Number of categories	Post-fire characteristic													
			Pre-fire		Forest Floor Surface Organic Matter Condition (S – scorched, U – unburned, B – black, G – grey/white, O – orange)											
			Unburned	Abundant	Present				Absent							
	S	U	B	G	O	U	B	G	O	U	B	G	O			
<b>Physical effects</b>																
Johansen <i>et al.</i> (2001)	Erosion	2	–	1	1	1	1	1	1	1	1	1	2	2	2	2
Neff <i>et al.</i> (2005)	Erosion	2	0	2	2	2	2	2	2	2	2	2	2	2	2	2
Parsons <i>et al.</i> (2010)	Values at risk	3	–	1	1	1	1	1	3	3	3	3	5	5	5	5
Robichaud and Hungerford (2000)	Water infiltration	2	–	1	1	1	1	1	2	2	2	2	2	2	2	2
Shakesby <i>et al.</i> (2003)	Water repellency	2	–	1	1	1	1	1	1	1	1	1	2	2	2	2
Ulery and Graham (1993)	Physical	4	–	1	–	–	–	–	–	–	–	–	1	2	4	5
<b>Chemical effects</b>																
Arocena and Opio (2003)	Mineralogy	2	0	–	–	–	–	–	–	–	–	–	–	–	2	–
Baird <i>et al.</i> (1999)	Nitrogen	2	0	2	2	2	2	2	2	2	2	2	2	2	2	2
Brais <i>et al.</i> (2000)	Chemistry	2	–	1	1	1	1	1	1	1	1	1	2	2	2	2
Cerri <i>et al.</i> (1991)	Nutrient dynamics	2	0	–	–	–	–	–	–	–	–	–	–	2	2	2
Ellingson <i>et al.</i> (2000)	Nitrogen	2	0	–	–	–	–	–	–	–	–	–	2	2	2	2
Rumpel <i>et al.</i> (2007)	Chemistry	3	0	–	–	–	–	–	3	3	3	3	5	5	5	5
Yeager <i>et al.</i> (2005)	Nitrogen fixation	3	0	–	–	–	–	–	–	–	3	3	–	–	5	–
<b>Biological effects</b>																
Bentley and Fenner (1958)	Seed survival	5	–	1	1	1	1	1	2	2	2	2	3	4	5	–
Bernhardt <i>et al.</i> (2011)	Vegetation	3	–	1	1	1	1	1	3	3	3	3	5	5	5	5
Blank <i>et al.</i> (1994)	Vegetation	2	0	2	2	2	2	2	2	2	2	2	2	2	2	2
Bonnet <i>et al.</i> (2005)	Vegetation	4	0	3	4	5	5	5	4	5	5	5	5	5	5	5
Choromanska and DeLuca (2002)	Microbes	2	0	–	–	–	–	–	–	–	–	–	2	2	2	2
Dyrness and Norum (1983)	Biological review	5	0	2	3	3	3	3	4	4	4	4	–	–	5	–
Jain <i>et al.</i> 2006; Jain and Graham (2007)	Forest structure	4	0	3	3	3	3	3	4	4	4	4	5	5	5	5
Larivière <i>et al.</i> (2005)	Arthropods	2	0	2	2	2	2	2	2	2	2	2	–	–	–	–
Lentile <i>et al.</i> (2005)	Regeneration	3	–	1	3	3	3	3	3	3	3	3	5	5	5	5
Morgan and Neuenschwander (1988)	Shrubs	2	–	1	1	1	1	1	–	–	–	–	2	2	2	2
Schimmel and Granstrom (1996)	Seed survival	3	0	1	1	1	1	1	3	3	3	3	5	5	5	5
Tyler (1995)	Vegetation	2	0	1	1	1	1	1	2	2	2	2	2	2	2	2
Wang and Kembal (2005)	Seed survival	3	–	1	2	2	2	2	–	–	–	–	5	5	5	5
<b>Assessments</b>																
Alexander <i>et al.</i> (2006)	Physical setting	3	–	1	1	1	1	1	3	3	3	3	3	5	5	5
Barkley (2006)	Monitoring	3	–	1	1	1	1	1	3	3	3	3	5	5	5	5
Chafer <i>et al.</i> (2004)	Remote sensing	5	–	1	1	2	2	2	3	3	3	3	4	5	5	5
Key and Benson (2006)	Remote sensing	4	0	2	2	2	2	2	3	3	3	3	5	5	5	5
Lutes <i>et al.</i> (2006)	Monitoring	5	0	1	2	2	2	2	2	3	3	3	3	5	5	5
Miller (2001)	Monitoring	5	0	2	2	2	2	3	4	4	3	4	4	5	5	5
Patterson and Yool (1998)	Remote sensing	4	0	2	2	2	2	2	4	4	4	4	4	5	5	5
Ryan and Noste (1985)	Prescribed fire	4	0	2	2	2	2	3	3	4	4	4	4	5	5	5
US Department of Interior (2003) (forests, shrublands)	Monitoring	5	0	2	3	3	3	3	4	4	4	4	4	5	5	5
US Department of Interior (2003) (grasslands)	Monitoring	5	0	2	3	3	3	3	–	–	–	4	4	5	5	5
White <i>et al.</i> 1996	Remote sensing	3	–	1	1	1	1	1	1	1	1	1	3	5	5	5

**Table 2. Variables to measure in assessing burn severity**

Those variables in bold are most commonly measured and can be more readily inferred from satellite imagery. To measure fire severity for a particular objective, select those fire effects important for assessing the objective. Use continuous variables whenever practical: these can always be collapsed into classes if needed; for example, for interpretation. Name the resulting fire severity index for the objective, such as soil severity index (see examples in Table 1 and Table 3). From Jain *et al.* (2012)

Selected fire effect	Measurement variable	Calculation	Description
<b>Plant mortality</b>	Percentage dead (%)	Trees >5 cm DBH Trees <5 cm DBH	Fire-caused overstorey tree mortality Fire-caused understorey tree mortality
	Reduction in cover (%)	Shrubs Herbs	Shrub cover reduction Herbaceous cover reduction
<b>Fuel consumption</b>	Reduction in loading (%)	Woody Duff	Proportion woody fuel consumed Proportion duff fuel consumed
	Consumption (kg m <sup>-2</sup> )	Woody Duff	Amount of woody fuel consumed Amount of duff fuel consumed
<b>Smoke</b>	Emissions (kg km <sup>-2</sup> )	PM2.5 CO <sub>2</sub>	Amount of particulate matter released Amount of carbon dioxide emitted
<b>Char fraction</b>	Char fraction	Percentage	Amount of char in soil
Soil heating <sup>1</sup>	Depth lethal heating (cm)	Depth >60°C for 1 min	Soil depth of tissue death
	Depth nutrient heating (cm)	Depth >250°C	Soil depth of nutrient changes
	Total heat	Integrated area >60°C under time–temperature curve at 2 cm	Total soil heating
<b>Soil water</b>	Soil infiltration rate (mm hr <sup>-1</sup> )	Rate (mm hr <sup>-1</sup> ) measured with mini disk infiltrometer	Soil infiltration conditions
Repellency <sup>1</sup>			
Nutrients <sup>1</sup>	Reduction in nitrogen (%)	NH <sub>4</sub> concentration before and after fire	Difference between pre- and post-fire NH <sub>4</sub> concentration
Erosion <sup>1</sup>	Exposed mineral soil	Percentage based on visual estimate	Amount of ground area in mineral soil

<sup>1</sup>Not readily inferred from remotely sensed imagery.

Keane *et al.* 2010; Hudak *et al.* 2013), and changes in soil properties, such as water repellency (Lewis *et al.* 2006), erodibility (Pierson *et al.* 2001) and nutrient availability (Belillias and Feller 1998) have been the focus in other studies.

The composite burn index (CBI) is one field measurement that has been widely used in burn severity assessments, especially for vegetation (Key and Benson 2006; Miller and Thode 2007; Holden *et al.* 2009; Miller *et al.* 2009; Soverel *et al.* 2010; Dillon *et al.* 2011; Cansler and McKenzie 2012). CBI was designed to correlate rapid field assessments of fire effects on vegetation and soils with the difference between pre- and post-fire satellite images (Key and Benson 2006). However, CBI is an integrated metric that averages the magnitude of change across five strata from soil to vegetation canopies, with each strata having four or five variables that are visually assessed and assigned a value between zero (unburned) and three (highest severity). As such the specific factors resulting in a given CBI value can become obscured. CBI is heavily weighted to measuring fire effects on vegetation (Miller and Thode 2007) and correlates most closely with changes in the upper canopy structure in forests (Miller and Thode 2007; Miller *et al.* 2009). To improve the correlation between ground measurements and remotely sensed data, De Santis and Chuvieco (2009) developed the GeoCBI, which adjusts the weighting of each stratum according to its fractional cover as viewed from overhead. Although the CBI and GeoCBI have been applied successfully in a wide variety of ecosystems, CBI performs poorly in ecosystems like Alaskan boreal forests (French *et al.* 2008) and California chaparral (Keeley *et al.* 2008), where severity is best represented respectively by depth of burn in soil

organic matter or amount of shrub biomass consumed. CBI and GeoCBI can be correlated with spectral reflectance in a satellite image, but it can be difficult to interpret the specific fire effects that are present using these integrated indices. Further, while visual estimates are the only practical method of estimating most fire effects that comprise the CBI, these estimates have subjective bias across users and site conditions that make consistent and accurate evaluation difficult. Potentially great differences in perceptions and interpretations among observers with varying levels of experience can confound consistent CBI visual evaluations of severity attributes across different ecosystems. Plant species, soil types and fuelbeds are also different across large fires making consistent evaluations problematic.

Burn severity is commonly assessed in the field and then combined with remote sensing (Table 2). Parsons *et al.* (2010) developed a quantitative method for assessing soil burn severity. They focussed on soil effects to assist interpretation of Burned Area Reflectance Classification (BARC) maps (Clark and McKinley 2011; <http://www.fs.fed.us/eng/rsac/baer/barc.html>) in post-fire assessment and rehabilitation. They used five factors to help validate the BARC maps, which include ground cover remaining, ash colour and depth, fine roots remaining, soil structure changes, and water repellency. Like the soil post-fire index developed by Jain *et al.* (2012) (Table 3), the assessments are made in the field with continuous measures of importance to fire effects on soils. A developing database (P.R. Robichaud, pers. comm.) of these continuous measures assessed in the field immediately after fire will be immensely helpful in future evaluations of fire and burn severity from satellite imagery.

**Table 3. Soil post-fire index (PFI) classification key developed by Jain *et al.* (2008) based on (1) the abundance of surface organic matter to create broad categories and (2) mineral soil colour to partition the broad categories**

Surface organic cover can include litter, humus, rotten wood and in some cases a root mat. As with any key, one begins by evaluating the site based on 1a or 1b. If the response to 1b is 'yes' then surface organic cover is evaluated using ocular or grid sampling estimates. For example, if surface organic cover is <40% (3b) than this broad category can be dissected based on mineral soil colour (5a–5d). If the plurality of the soil is charred orange (5d) then the resulting soil PFI is 3.4

Soil characteristics		Soil PFI category
1a	No evidence of a recent fire	0.0
1b	Evidence of recent fire	
2a	Surface organic cover $\geq 85\%$	1.0
2b	Surface organic cover <85%	
3a	Forest floor surface organic cover $\geq 40\%$ and mineral soil appearance has a plurality of:	2.0
4a	Unburned mineral soil	2.1
4b	Black charred mineral soil	2.2
4c	Grey or white charred mineral soil	2.3
4d	Orange charred mineral soil	2.4
3b	Surface organic cover <40% and mineral soil appearance has a plurality of:	3.0
5a	Unburned mineral soil	3.1
5b	Black charred mineral soil	3.2
5c	Grey or white charred mineral soil	3.3
5d	Orange charred mineral soil	3.4
3c	(Forest floor absent) No surface organic matter left and mineral soil appearance has a plurality of:	4.0
6a	Unburned mineral soil	4.1
6b	Black charred mineral soil	4.2
6c	Grey or white charred mineral soil	4.3
6d	Orange charred mineral soil	4.4

Perhaps the greatest challenge to severity evaluations in the field is that without pre-fire measurements, changes due to fire must be inferred retrospectively from post-fire conditions; this is the case in most assessments of severity following wildfires (Hudak *et al.* 2011, 2013). In temperate forests, common fire effects measured to evaluate severity include the amount of surface fuel consumed; percentage mortality in both overstorey trees and understorey plants; percentage tree or shrub volume and cover affected and inferred degree of soil heating. In grasslands and boreal forests, the degree of soil heating and depth of burning, including percentage consumption of soil organic matter, are important fire effects (French *et al.* 2008). However, for most of these effects, both pre- and post-burn measurements are required to accurately and objectively quantify the change caused by fire. Unfortunately, given that wildfire locations are difficult to predict, pre-burn measurements are typically lacking and are at best challenging to acquire or infer (Lentile *et al.* 2007).

Several strategies are used to address the lack of pre-fire measurements. Most often, observers subjectively estimate pre-fire conditions (e.g. canopy cover of understorey and overstorey plants, duff and litter cover, and surface fuel load) based on observations of surrounding areas (Key and Benson 2006). Burned plots can be compared with paired plots in adjacent unburned areas (Diaz-Delgado *et al.* 2003; Karau and Keane 2010; Hudak *et al.* 2011), but these inferences depend on how well the unburned plot represents the pre-fire condition of the burned plot. Sometimes, sites have been sampled pre-burn, such as the network of Forest Inventory and Analysis (FIA) plots across all forested lands in the US (Megown *et al.* 2011), but rarely are the number of burned FIA plots for a given wildfire sufficient for a statistically valid fire effects evaluation.

In some instances, severity can be inferred from post-fire evidence alone. Based on post-fire diameter measurements of all live and dead trees in conifer forests, Miller *et al.* (2009) were able to calculate pre- and post-fire live tree basal area, and subsequently a measure of basal area loss due to fire. In chaparral vegetation, the diameter of the smallest branch remaining on shrub skeletons has been found to be a good indicator of overall biomass loss from fire (Moreno and Oechel 1989; Keeley *et al.* 2008). In other forest studies, the amount of charred surface fuels, soil and trees is sometimes used to evaluate respectively the magnitude of fuel consumption, soil heating and plant mortality. However, this evidence does not provide a complete picture or a true quantitative assessment of the ecological consequences of the fire. Tree bole char, for example, is only partially correlated with tree mortality (Keyser *et al.* 2006; Halofsky and Hibbs 2009; Hudak *et al.* 2011). In rangelands, high fire severity has been associated with reduction of the seedbank, lower species diversity post-fire and increases in exotic species cover (Ghermandi *et al.* 2013).

#### Assessing fire and burn severity with remote sensing

Remotely sensed image data have the great advantage of providing pre-fire information that can be difficult or impossible to retrieve in the field. Surface reflectance changes over the days and weeks following fires (Trigg and Flasse 2000), and fires themselves, change surface reflectance in a wide variety of ecosystems (Landmann 2003, Chafer *et al.* 2004; Cocke *et al.* 2005; Smith *et al.* 2005; Roy *et al.* 2006; Chafer 2008; French *et al.* 2008; Kumar *et al.* 2008; Lee *et al.* 2009; Veraverbeke *et al.* 2011). Many severity assessments use satellite imagery to quantify the magnitude of vegetation change from pre-fire

conditions (e.g. Key and Benson 2006; Miller and Thode 2007; Holden *et al.* 2009; Miller *et al.* 2009; Soverel *et al.* 2010). If the same type of sensor is used to collect imagery both before and after the fire under comparable illumination conditions, then the difference between them can provide an objective means to quantify ecological change induced by the fire, which has a substantial effect on the reflective properties of the scene (Jakubauskas *et al.* 1990; Landmann 2003). This is the basis for using the Normalized Burn Ratio (NBR, Key and Benson 2006), Normalized Difference Vegetation Index (NDVI, Tucker 1979) and similar indices. Many calculate differences between pre- and post-fire indices in absolute (differenced NBR or dNBR) or relative terms as relative differenced Normalized Burn Ratio (RdNBR, Miller and Thode 2007), the Relativized Burn Ratio (RBR), or differenced NDVI (dNDVI, Díaz-Delgado *et al.* 2003; Epting *et al.* 2005), particularly where the pre-burn biomass is low or highly variable. Although the dNBR is used more broadly as a burn severity index, it sometimes performs only marginally better than dNDVI (Hudak *et al.* 2007b; Veraverbeke *et al.* 2010b). Fox *et al.* (2008) found NBR and NDVI are highly correlated. Roy *et al.* (2006) advised caution in relying on NBR and related indices because the resulting burn severity maps may have low accuracy depending on the variable of interest and whether the index was originally developed for detecting burned area, not burn severity. Clearly, each assessment of burn severity must be carefully evaluated to ensure it meets the intended goal with acceptable accuracy.

The Landsat sensors have provided the longest available and most widely interpreted source of image data for assessing severity using the indices detailed above. Fire and burn severity maps derived from Landsat Thematic Mapper (TM) imagery can be used to develop retrospective maps of historical wildland fires back to 1984 (Eidenshink *et al.* 2007). Landsat Multi-Spectral Scanner (MSS) imagery (Russell-Smith *et al.* 1997; Hudak and Brockett 2004) and aerial photographs (Ekstrand 1994; White *et al.* 1996) can extend these records even further back in time. Severity mapping from imagery usually involves relating spectral reflectance characteristics of the post-fire scene to field measures of fire effects or severity indices (e.g. CBI) collected at coincident locations (Cocke *et al.* 2005; Hudak *et al.* 2007b). Strong correlations between the datasets can then be interpreted to indicate the field variable of interest. While 30-m resolution Landsat TM imagery is most commonly used to map fire-induced change, higher spatial and spectral resolution data have obvious potential for quantifying fine-scale post-fire effects.

Other types of imagery are used to quantify severity. Robichaud *et al.* (2007b) used hyperspectral imagery of higher spatial and spectral resolution than Landsat TM imagery to more accurately map post-fire soil and ash cover fractions for assessment of erosion potential. Moreover, they associated the high resolution imagery to directly measured variables instead of indices. However, high resolution multispectral and hyperspectral data are expensive and can be challenging to work with because they contain hundreds of spectral bands, have relatively high densities of pixels and may require extensive image geo-registration. These factors may limit operational use of hyperspectral data for mapping post-fire effects. MODIS imagery, which is freely available like Landsat data but of much

coarser resolution (250–1000 m), can be used to assess severity of large wildfires when other data are unavailable (Kolden and Rogan 2013), but the coarse spatial resolution makes the resulting maps unsatisfying to most managers and scientists.

Active sensor systems, such as Light Detection and Ranging (LiDAR) and radar can also be used to infer fire effects. In contrast to passive sensors measuring the sun's reflected radiation these sensors supply their own power. LiDAR data have demonstrated potential for characterising fire-induced changes in overstorey vegetation characteristics (Wang and Glenn 2009; Wulder *et al.* 2009; Kwak *et al.* 2010; Magnussen and Wulder 2012). For example, LiDAR has been used for post-fire assessments to quantify tree regeneration (Debouk *et al.* 2013), assess how post-fire forest structure varies with burn severity (Kane *et al.* 2013), and estimate post-fire tree height (Magnussen and Wulder 2012). However, operational use of LiDAR remains limited given the sparse coverage of pre-fire data and the expense of acquiring new post-fire data. Radar data may also have potential for severity assessments, but difficulty in data interpretation limits operational utility (Kasischke *et al.* 2007b; Tanase *et al.* 2010a; Tanase *et al.* 2010b; Tanase *et al.* 2010c).

Ultimately, to maximise the utility and efficiency of remote sensing assessments, the spatial, temporal and spectral resolution of the remotely sensed imagery must match the ecological scale of the fire effect of interest. More research into active sensors would likely lead to more physically based severity assessments, which are needed to advance fire science. However, approaches that directly estimate biophysical measures of interest can also be applied to passive optical imagery, which would reduce current overreliance on burn severity indices to infer biophysical attributes (Roy and Landmann 2005; Kasischke *et al.* 2007a). Spectral mixture analysis is an appealing approach to processing post-fire satellite imagery because it estimates the fractional cover of biophysical variables at the sub-pixel level, making them more directly comparable to the same fractional cover variables measured in field plots on the ground. For instance, estimates of char fraction or green vegetation fraction derived from imagery correlate as well as NBR to the same fractional cover variables estimated independently on the ground (Hudak *et al.* 2007b; Smith *et al.* 2007). Others have estimated severity using approaches that incorporate radiative transfer modelling, which takes advantage of specific physical reflectance properties of surfaces to estimate vegetation parameters (Chuvieco *et al.* 2006, 2007). As applied to burn severity mapping, these efforts use a reference spectrum of a range of healthy to damaged vegetation, then invert a radiative transfer model to simulate post-fire spectral response, and finally apply a supervised classification to a post-fire image (De Santis *et al.* 2009; De Santis *et al.* 2010). This method has the capacity to establish scenarios for combinations of vertical strata severities that are less dependent on particular local conditions. It can also simulate the outcome of specific effects of fire, such as changes in leaf colour or leaf area index because they are input variables in the canopy reflectance models (De Santis *et al.* 2009).

#### *Predicting fire and burn severity with modelling*

Statistical and simulation modelling are alternatives to map and assess post-wildfire severity. Statistical relationships to predict severity can provide insight into possible drivers of severity.

Holden *et al.* (2009) created a statistical severity model from field-gathered CBI data, landscape and biophysical spatial data, and NBR from Landsat imagery in New Mexico, USA. Dillon *et al.* (2011) expanded on this statistical approach by adding weather and hydroclimatic indices to predict proportion burned with high severity and to map areas that could burn severely if a wildfire were to occur for the western US. In South Korea, Lee *et al.* (2009) used regression tree analysis to predict severity from landscape characteristics. Simulation models such as FOFEM (Reinhardt *et al.* 1997), and statistical models such as CONSUME (Ottmar *et al.* 1993) are useful for simulating the direct effects of a fire on vegetation, fuels and soils. Keane *et al.* (2010) implemented FOFEM into a spatial computer application called FIREHARM to create severity maps, which have been compared with and integrated with satellite imagery (Karau and Keane 2010; Karau *et al.* 2014). These types of models have several advantages: (1) simulation models can provide biophysically based fire effects estimates, (2) results can be scaled to the resolution most appropriate for describing a specific effect, (3) models allow for rapid assessment because results can be simulated quickly as long as input data are available, and (4) models can be used to predict fire effects, allowing a manager to proactively prioritise resources.


The disadvantage of current empirical and simulation models is that severity predictions are only as good as the input data used to create them (Karau *et al.* 2014), and the widely available spatial data used to develop the simulation and statistical models have high levels of uncertainty (Keane *et al.* 2013). Current

models rarely use data from ongoing severity assessments. Most severity modelling efforts use a completely different set of severity assessments and classifications than remote sensing or field methods. Therefore, a standardised set of ecological metrics to calibrate mapped severity to locally observed fire effects is critical for all three severity methods – empirical, imagery and simulation.

### Challenges common to all assessments of fire and burn severity

#### *Spatial variability*

Fires do not burn homogeneously across landscapes (Fig. 1), nor do they burn similarly in different ecosystems, giving rise to challenges with characterising fire and burn severity. Understanding the causes of this heterogeneity is often a goal of severity assessments. In this context, severity has been linked to many biophysical characteristics and processes (Keeley 2009). Landscape patterns of severity often vary with topography (e.g. Kushla and Ripple 1997; Broncano and Retana 2004; Holden *et al.* 2009) because topography influences the biophysical environment (microclimatic conditions of temperature, precipitation, direct solar radiation, timing of snowmelt, wind exposure) that directly affects both biomass accumulation and the amount of biomass available to burn at the time of fire (Holden and Jolly 2011). Dillon *et al.* (2011) found the distribution of high-severity effects on forests in six regions in western US fires was influenced more strongly by topographic



**Fig. 1.** Post-fire landscape near the lightning ignition start of the 2003 Black Mountain 2 fire near Missoula, Montana, 8 weeks after burning. The heterogeneous pattern of fire severity patches is a result of the interactions of fire with the biotic and physical environment at multiple scales. As time since fire progresses, the relative differences in fire severity, and their spatial pattern, will change.



characteristics than weather or climate, but the specific effects of topography varied geographically. Similarly, Broncano and Retana (2004) found spatial variability in severity to be correlated with elevation and aspect in Spain, and Chafer *et al.* (2004) and Bradstock *et al.* (2010) found that topographic setting, combined with fuel loading and fire weather, influenced severity patterns in Australia.

Unlike topography, which is relatively static and easily quantified pre-fire, quantitative measures of pre-fire fuel and local weather conditions on the ground during a fire are rarely available and difficult to infer from post-fire assessments. Pre-fire character of the vegetation community, such as structure (density, size, height), composition (fire-adapted species *v.* fire-sensitive species), and productivity (biomass, deposition), moisture content and phenology, sets the stage for a range of possible fire behaviours and ecological responses. Severity might be considered low in a grassland but high in a forest; thus the magnitude of severity will vary depending on what is burned. However, there are studies that show a relationship between pre-fire structure and severity. Jain *et al.* (2006) and Jain and Graham (2004) demonstrated close relationships between soil burn severity and both pre- and post-burn forest structure. Hessburg *et al.* (2007) related severity to forest structure across landscapes in the north-western US. Bigler *et al.* (2005) found that severity correlated with forest structure and composition, stand history (previous disturbances) and elevation.

Previous disturbance histories, climates and antecedent weather create vegetation conditions that influence the complex web of fire effects that can be spatially heterogeneous, making it difficult to predict severity (Turner *et al.* 1999; Romme 2005; Thompson and Spies 2010). As a fire burns, the micrometeorology of fire weather interacts with ignition and combustion to leave behind unique patterns of fire effects. Subtle changes between day- and night-time temperatures and humidity can alter fire behaviour and severity. Wind during a fire event can influence severity patterns even in relatively uniform vegetation. The presence or absence of ladder fuels may also determine the severity outcome and forest composition and structures can be influenced by harvest, disease and insect infestations. Tree mortality, for example, is a result of a complex set of interactions between instantaneous fire behaviour, topography, ambient weather conditions and tree characteristics including size, crown position, adjacency of neighbouring trees, bark thickness and other factors influencing the ability of the tree to survive the fire – all of which vary at different scales. These interacting processes and scales make severity estimation difficult because the elements that dictate fire and burn severity may vary at scales that differ by ecosystem, fire and biophysical environment.

Ecological responses occur at multiple time scales and this can confound fire and burn severity assessments (Lentile *et al.* 2006). Pre- and post-fire precipitation patterns, wind events, human interventions, and plant reproductive strategies all influence long-term ecosystem response to fire yet they act at different temporal scales, which adds uncertainty to assessments of fire severity. Changes in post-fire soil nitrogen, for example, last for a shorter time (years) than changes in forest structure (decades). Soil water repellency usually lasts only a few years (Doerr *et al.* 2000; Robichaud and Hungerford 2000), but recovery of vegetation can take decades.

### *Multiple interacting fire effects*

The major factors used to assess severity, such as plant mortality and soil heating, are not independent ecological processes but are interrelated through mutual feedback mechanisms that vary greatly between fuel and vegetation types. Grass fires can have high intensities, rapid rates of spread and nearly complete fuel consumption, yet soil heating and plant mortality are usually low and vegetation conditions 1-year post-fire can be similar to that found before the fire. In contrast, a low-intensity surface fire burning in a stand of fire-intolerant spruce and fir trees can kill many trees through cambial damage on stems and roots through soil heating. Tree mortality, for example, often increases when insects and disease agents attack trees weakened by fire (Hood *et al.* 2007), sometimes enough to alter post-fire assessments of severity. Conversely, soil erosion potential after fire may be mitigated by needles from scorched trees that fall on severely burned, highly erodible soils (Pannkuk *et al.* 2000). High consumption of aboveground biomass in perennial grasslands and shrublands may be short lived with sprouting species recovering relatively quickly post-fire. Fires that increase non-native plant species cover may be more severe than fires where only native plants are present. In some instances, it may be necessary, and even desirable, to account for some secondary fire effects in fire and burn severity assessments (Veraverbeke *et al.* 2010a; 2010b, 2011; Dillon *et al.* 2011).

### *Spatial and temporal scale*

Scale considerations are essential for appropriate assessment of fire effects (Simard 1991). Wildland fire acts across multiple temporal and spatial scales responding to factors that control both fire behaviour (e.g. fuel moisture, wind) and the characteristics of the biological elements that are burned (e.g. species, size, loadings) (King *et al.* 2008). These interactions in turn influence severity. As the factors that control fire and vegetation act across different scales, it follows that fire and burn severity must be described across multiple time and space scales. Further, spatial pattern influences vegetation response – if patches of high severity are very large, recovery of vegetation dependent on dispersal of seeds from surviving plants will be slower than in a fine-scale mosaic (Turner *et al.* 1999; Bonnet *et al.* 2005; Donato *et al.* 2009). Thus, metrics of the spatial distribution of fire effects are needed to fully quantify severity.

The timing of specific fire effects and ecosystem responses can be dramatically different within a single fire. Vegetation recovery, for example, can take only a few years in low-elevation grasslands, but may take decades in upper subalpine forests (Keane and Parsons 2010). Similarly, tree regeneration may occur quickly after fire in productive mesic forests, but may be slower in xeric, cold upper subalpine environments (Agee and Smith 1984). This difference in response timing serves to complicate many burn severity assessments.

### **Recommended best practices**

Instead of collapsing complex fire–biota–environmental interactions and responses down to a generalised classification, we recommend directly measuring the actual fire effect, be it tree mortality, fuel consumption, soil water repellency or any other

important measurable fire effect. As Jain *et al.* (2004) emphasised, researchers should simply quantify severity with what they are actually measuring (e.g. see Table 3). The soil burn severity index is a good example that is widely applied in assessing post-fire effects on soils with a focus on soil erosion potential. Another is the soil PFI based on post-fire characteristics that relate to nutrient availability, seed availability and other soil characteristics (Jain *et al.* 2012) (Table 3). These are good examples that use physically based fire effects variables of interest that can then be input into other fire effects applications (erosion modelling, wildlife habitat evaluations) for a more tailored assessment of severity, and they can be predicted from simulation models to expand the use of the severity index from operational to planning and from only retrospective to predictive. A common database of severity assessments will be immensely helpful for improving inferences beyond local applications. Further, we urge use of continuous variables for measurement whenever practical; these can always be collapsed to classes if needed for interpretation.

Every assessment will require addressing questions of which imagery, which indices, what timing and what to measure, as illustrated in Table 4. We outline best practices as these choices are made for assessing soil burn severity and vegetation effects. Imagery choice is often dependent on what is available. For burn severity assessments, Landsat TM and ETM+ sensors are often used because of the 30-m spatial resolution, and the global availability of the imagery every 16 days and large catalogue of free images dating back to 1984, all of which are important for rapid post-fire assessments needed for mitigating erosion potential. Clearly, soil effects

vary at scales finer than 30 m (Hudak *et al.* 2007b) and degree of soil charcoal and organic content of soils can complicate satellite-inferred burn severity (Epting *et al.* 2005; Smith *et al.* 2010; Picotte and Robertson 2011). For vegetation effects, Landsat imagery is commonly used, but other imagery products with finer spatial and spectral resolution is available. NBR, dNBR and RdNBR are most common, but RBR and char fraction (Lentile *et al.* 2009) may be better suited depending on the specific fire effects of interest. Relativised measures, including both RdNBR (Miller and Thode 2007) and RBR are better for detecting high-severity effects across a wide range of pre-fire conditions, including those with low total biomass.

Timing of imagery depends on the purpose of assessment. When choosing pre- and post-fire images, it is important to consider that vegetation that burns often does so in a drought-stressed state. Therefore, the pre-fire image must be collected as close to the fire date as possible to isolate the effects of fire from the effects of drought. Timing of the post-fire image depends on several considerations, including ecological context and the specific purpose of the assessment. For instance, how quickly the vegetation will respond or recover from the fire must be considered. Is it important to capture immediate, same-season effects before any recovery or is it desirable to allow time for some second-order effects and initial recovery? Has the vegetation senesced or did snow fall immediately post-fire, making change detection impossible? If next growing season imagery is required, when will phenology most closely match the pre-fire image? In rangelands, remotely sensed reflectance is highly variable with phenology throughout the

**Table 4. Guidance for assessing burn severity for vegetation effects and soil burn severity using field and remote sensing methods**

For each application, users need to decide which imagery and index, the timing, and what to assess in the field: see text, Eidenshink *et al.* (2007) for further discussion and references for further information. The resulting severity indices would differ from each other and from those developed for assessing fire effects on habitat for invasive species, wildlife or other purposes

Question	Soil burn severity	Vegetation severity
What imagery?	<ul style="list-style-type: none"> <li>● LANDSAT most commonly used due to availability, spatial resolution and cost.</li> <li>● Quickbird or other high spatial resolution imagery useful when higher resolution is needed, but costs more.</li> <li>● MODIS over larger extents where lower spatial resolution is acceptable.</li> </ul>	
What index?	<ul style="list-style-type: none"> <li>● NBR (one image immediately post-fire), dNBR or RdNBR or RBR</li> <li>● Adjust based on field assessments.</li> </ul>	<ul style="list-style-type: none"> <li>● Relativised measures (e.g. RdNBR or RBR) commonly used, especially for areas with relatively low or heterogeneous vegetation cover; dNDVI useful</li> </ul>
Timing of imagery?	<ul style="list-style-type: none"> <li>● Immediately post-fire to support planning for rehabilitation and recovery</li> </ul>	<ul style="list-style-type: none"> <li>● Usually extended with pre-fire image as close to fire date as possible, post-fire image 1 year post-fire at same phenology, but with rapid vegetation recovery use imagery immediately post-fire</li> <li>● For non-forests, often immediately pre- and immediately post-fire</li> <li>● Depends on purpose of assessment (see Table 2)</li> <li>● If field measures will be used in combination with remote sensing, then only measure variables that can be readily inferred from imagery and match the spatial and temporal scale</li> <li>● CBI or GeoCBI commonly used but we recommend measuring the actual effect(s) of interest directly using quantitative, continuous measures where possible. Examples include tree mortality, fuel consumption, proportion of foliar biomass burned and reduction in canopy cover</li> </ul>
Field measures	<ul style="list-style-type: none"> <li>● Focus on direct measures such as soil colour and exposure, and water repellency. Indirect measures include fuel consumption and amount of ash</li> </ul>	

growing season, which must be considered when selecting images for burn severity assessments.

It is important to think carefully about field measures, especially if these are to be inferred from satellite imagery or linked to predictive models. If field measurements are to be correlated with remote sensing data, the variables measured in the field must have a logical and mechanistic connection to properties the sensor can detect. For instance, soil heating by fire, although ecologically important, cannot be inferred directly from pre- and post-fire satellite imagery comparisons. Correlations between field and remotely sensed variables say nothing about causation; remotely sensed indices of severity are only indices, and therefore should not be interpreted as direct measures of fire or burn severity.


**Future directions**

We suggest that the first step for improving severity assessments is to move towards a unified, physically based, hierarchical terminology (Table 2). Fire and burn severity are general concepts to qualitatively or quantitatively describe the magnitude of the myriad immediate and longer-term fire effects at a point, plot, stand and across a landscape (Fig. 2). Remote sensing indices (Fig. 2) such as the RdNBR, are not direct measures of severity *per se*, but are useful for inferring severity when the fire effect(s) of interest can be meaningfully interpreted from imagery (e.g. Miller et al. 2009). Much of the confusion associated with terminology noted by Keeley (2009)

and Lentile et al. (2006) can be alleviated simply by clearly articulating two factors suggested by Jain et al. (2004): (1) the element or aspect of severity being assessed or inferred (Holden et al. 2007; Miller et al. 2009), and (2) the specific timing of the post-fire assessment.

We suggest that recording actual fire effects measurements, such as percentage tree mortality or pre- and post-fire live tree basal area in forested areas (Miller et al. 2009), or average diameter of the smallest remaining branches in shrublands, is preferable to collapsing these measures into an index like CBI. We recognise that there will always be utility in composite metrics like CBI and GeoCBI, but without specific, ecologically meaningful measurements, it will remain difficult to directly relate ordinal severity class values to specific ecological characteristics or fire effects. These measures can be summarised as CBI or GeoCBI *ex post facto*, as appropriate. Severity classifications based only on relationships to composite measures may have little predictive power to describe potential severity before a site actually burns.

Moving towards more ecologically based severity classifications will require major improvements in the measurement of the direct effects of wildland fires. Developing meaningful relationships between individual fire effects or composite severity metrics and the conditions before, during and after fire will require studies with detailed quantitative descriptions of pre-fire conditions, fire behaviour and the post-fire environment at different time periods. Novel methods for assessing pre-burn



**Fig. 2.** Fire and burn severity, defined as the immediate and longer-term ecological effects of fire, can be assessed in the field using any one or a combination of metrics (top). Severity can also be inferred from individual remote sensing indices (side); this is only effective and interpretable when the index is correlated with fire effects on the ground.

conditions and for describing fire behaviour will be useful. Until we understand the causes and consequences of severity well enough to predict them, proactive, effective pre- and during- fire assessment and management to alter fire effects will continue to be challenging. Intensive spatial field surveys could support improved linkages between remotely sensed map products and field data, especially when fire effects vary greatly at fine spatial scales. One example of this in the US is the Accelerated Remeasurement and Evaluation of Burned Areas (or AREBA) project (Megown *et al.* 2011). The FIA program of the USDA Forest Service now measures both surface and crown fuels, along with many other ecosystem characteristics useful for quantifying fire effects, but the sparse distribution of plots established only in forested ecosystems will limit its operational use in severity mapping efforts, especially in rangelands.

Fire and burn severity mapping projects will continue to depend on remotely sensed imagery and field measurements. Therefore, it is critical that ecological advances in field assessments of severity be matched with the most appropriate imagery (Fig. 2). It is important that key fire effects are related to image signatures at appropriate scales. With the use of more advanced remote sensing technologies, such as hyperspectral imagery, LiDAR and radar, important fire effects may be more accurately and consistently inferred from imagery with higher spectral, spatial and temporal resolutions. It is exciting to see the many different research and management applications of severity, some of which have been prompted by the availability of MTBS data in the US. We look forward to learning as much about the causes and consequences of fire and burn severity as we know about fire behaviour. We also urge the development of a severity field assessment database and research to further our understanding of causal mechanisms linking fire and burn severity to conditions before and during fires to support improved models linking fire behaviour and severity and for forecasting effects of future fires. Understanding where, why and how fires burn severely will be greatly enhanced by efforts to: (1) relate severity to climate, weather, topography, fuels and land use (e.g. Dillon *et al.* 2011; Miller and Safford 2012), (2) explain temporal trends (Dillon *et al.* 2011; Miller *et al.* 2012; Mallek *et al.* 2013), and (3) develop tools that effectively link conditions before fire to flaming and glowing combustion, soil heating, biomass consumption and vegetation mortality. Better understanding will support better and proactive management of fire and fire effects.

One of the grand challenges for fire science remains to link conditions before, during and after fires together based on understanding of how fire behaviour causes fire and burn severity (Kremens *et al.* 2010). Without examining these linkages, it will be difficult to predict the ways in which pre-fire fuels and vegetation influences fire effects and vegetation response, yet that is key to proactive fuels and vegetation management. An important step towards meeting this challenge is a common base of terminology for severity that builds on measurable, physically based metrics linked to conditions before, during and after fires to characterise fire effects across multiple scales and applications. Only by taking this approach will the confusion and ambiguity be reduced and, more importantly, will our understanding of the ecological role of fire be enhanced. Ultimately, we need to more fully understand the

causal mechanisms of severity, such as the multiple ecological interactions, scales of variability and fire behaviour drivers if we are to predict the consequences of alternative pre-, during and post-fire management strategies focused on influencing fire and burn severity outcomes.

### Acknowledgements

We based this paper, in part, on the lessons we learned during several Joint Fire Sciences Program and National Fire Plan-funded projects that dealt with fire and assessing burn severity and creating severity classification systems. This research was supported in part by the USDA Forest Service Rocky Mountain Research Station, the University of Idaho (09-JV-11221637-270), the National Fire Plan and the Joint Fire Science Program (JFSP-09-1-07-4). We appreciate constructive reviews from Pete Robichaud and anonymous reviewers.

### References

- Agee JK, Smith L (1984) Subalpine tree reestablishment after fire in the Olympic Mountains, Washington. *Ecology* **65**, 810–819. doi:10.2307/1938054
- Alexander ME (1982) Calculating and interpreting forest fire intensities. *Canadian Journal of Botany* **60**(4), 349–357. doi:10.1139/B82-048
- Alexander JD, Seavy NE, Ralph JC, Hogoboom B (2006) Vegetation and topographical correlates of fire severity from two fires in the Klamath-Siskiyou region of Oregon and California. *International Journal of Wildland Fire* **15**, 237–245. doi:10.1071/WF05053
- Arocena JM, Opio C (2003) Prescribed fire-induced changes in properties of sub-boreal forest soils. *Geoderma* **113**(1–2), 1–16. doi:10.1016/S0016-7061(02)00312-9
- Baird M, Zabowski D, Everett RL (1999) Wildfire effects on carbon and nitrogen in inland coniferous forests. *Plant and Soil* **209**(2), 233–243. doi:10.1023/A:1004602408717
- Barkley YC (2006) After the burn: assessing and managing your forestland after a wildfire. University of Idaho Extension, Idaho Forest, Wildlife and Range Experiment Station, Bulletin No. 76. (Moscow, ID)
- Barrett SW, DeMeo T, Jones JL, Zeiler JD, Hutter LC (2006) Assessing ecological departure from reference conditions with the Fire Regime Condition Class (FRCC) mapping tool. In 'Fuels Management – How to Measure Success'. (Eds PL Andrews, BW Butler) USDA Forest Service, Rocky Mountain Research Station, Proceedings RMRS-P-41 pp. 575–585. (Fort Collins, CO).
- Belillias CM, Feller MC (1998) Relationships between fire severity and atmospheric and leaching nutrient losses in British Columbia's coastal Western Hemlock zone forests. *International Journal of Wildland Fire* **8**, 87–101. doi:10.1071/WF9980087
- Bentley JR, Fenner RL (1958) Soil temperatures during burning related to postfire seedbeds on woodland range. *Journal of Forestry* **56**, 737–740.
- Bernhardt EL, Hollingsworth TN, Chapin IFS (2011) Fire severity mediates climate-driven shifts in understory community composition of black spruce stands of interior Alaska. *Journal of Vegetation Science* **22**, 32–44. doi:10.1111/J.1654-1103.2010.01231.X
- Beschta RL, Rhodes JJ, Kauffman JB, Gresswell RE, Minshall GW, Karr JR, Perry DA, Hauer FR, Frissell C (2004) Postfire management on forested public lands of the western United States. *Conservation Biology* **18**, 957–967. doi:10.1111/J.1523-1739.2004.00495.X
- Beukema SJ, Kurz WA (1998) 'Vegetation dynamics development tool – Users Guide Version 3.0.' (ESSA Technologies: Vancouver, BC).
- Bigler C, Kulakowski D, Veblen TT (2005) Multiple disturbance interactions and drought influence fire severity in rocky mountain subalpine forests. *Ecology* **86**, 3018–3029. doi:10.1890/05-0011
- Bisson M, Fornaciai A, Coli A, Mazzarini F, Pareschi MT (2008) The vegetation resilience after fire (VRAF) index: development, implementation and an illustration from central Italy. *International Journal of*

- Applied Earth Observation and Geoinformation* **10**, 312–329. doi:10.1016/J.JAG.2007.12.003
- Blank RR, Allen F, Young JA (1994) Extractable anions in soils following wildfire in a sagebrush–grass community. *Soil Science Society of America Journal* **58**, 564–570. doi:10.2136/SSSAJ1994.03615995005800020045X
- Boby LA, Schuur EAG, Mack MC, Verbyla D, Johnstone JF (2010) Quantifying fire severity, carbon, and nitrogen emissions in Alaska's boreal forest. *Ecological Applications* **20**, 1633–1647. doi:10.1890/08-2295.1
- Bonnet VH, Schoettle AW, Shepperd WD (2005) Postfire environmental conditions influence the spatial pattern of regeneration for *Pinus ponderosa*. *Canadian Journal of Forest Research* **35**, 37–47. doi:10.1139/X04-157
- Bowman DM, Balch JK, Artaxo P, Bond WJ, Carlson JM, Cochrane MA, D'Antonio CM, DeFries RS, Doyle JC, Harrison SP, Johnston FH, Keeley JE, Krawchuk MA, Kull CA, Marston JB, Moritz MA, Prentice IC, Roos CI, Scott AC, Swetnam TW, van der Werf GR, Pyne SJ (2009) Fire in the Earth system. *Science* **324**(5926), 481–484. doi:10.1126/SCIENCE.11663886
- Bradstock RA, Hammill KA, Collins L, Price O (2010) Effects of weather, fuel and terrain on fire severity in topographically diverse landscapes of south-eastern Australia. *Landscape Ecology* **25**, 607–619. doi:10.1007/S10980-009-9443-8
- Brais S, Pare D, Ouimet R (2000) Impacts of wild fire severity and salvage harvesting on the nutrient balance of jack pine and black spruce boreal stands. *Forest Ecology and Management* **137**, 231–243. doi:10.1016/S0378-1127(99)00331-X
- Broncano MJ, Retana J (2004) Topography and forest composition affecting the variability in fire severity and post-fire regeneration occurring after a large fire in the Mediterranean basin. *International Journal of Wildland Fire* **13**, 209–216. doi:10.1071/WF03036
- Cansler CA, McKenzie D (2012) How robust are burn severity indices when applied in a new region? Evaluation of alternate field-based and remote sensing methods. *Remote Sensing* **4**, 456–483. doi:10.3390/RS4020456
- Carey A, Evans M, Hann P, Lintermans M, MacDonald T, Ormay P, Sharp S, Shorthouse D, Webb N (2003) Wildfires in the ACT 2003: Report on initial impacts on natural ecosystems. Australian Capital Territory, Technical Report 17, Environment ACT, Canberra, Australia. 123 pp.
- Cerri CC, Volkoff B, Andreaux F (1991) Nature and behaviour of organic matter in soils under natural forest, and after deforestation, burning, and cultivation near Manaus. *Forest Ecology and Management* **38**, 247–257. doi:10.1016/0378-1127(91)90146-M
- Chafer CJ (2008) A comparison of fire severity measures: an Australian example and implications for predicting major areas of soil erosion. *Catena* **74**, 235–245. doi:10.1016/J.CATENA.2007.12.005
- Chafer CJ, Noonan M, Macnaught E (2004) The post-fire measurement of fire severity and intensity in the Christmas 2001 Sydney wildfires. *International Journal of Wildland Fire* **13**, 227–240. doi:10.1071/WF03041
- Choromanska U, DeLuca TH (2002) Microbial activity and nitrogen mineralization in forest mineral soils following heating: evaluation of post-fire effects. *Soil Biology & Biochemistry* **34**, 263–271. doi:10.1016/S0038-0717(01)00180-8
- Chuvieco E (1999) Measuring changes in landscape pattern from satellite images: short-term effects of fire on spatial diversity. *International Journal of Remote Sensing* **20**, 2331–2346. doi:10.1080/014311699212056
- Chuvieco E, Riaño D, Danson F, Martin P (2006) Use of a radiative transfer model to simulate the postfire spectral response to burn severity. *Journal of Geophysical Research* **111**, G04S09. doi:10.1029/2005JG000143
- Chuvieco E, De Santis A, Riano D, Halligan K (2007) Simulation approaches for burn severity estimation using remotely sensed images. *Fire Ecology* **3**, 129–150. doi:10.4996/FIREECOLOGY.0301129
- Clark J, McKinley R (2011) Remote sensing and geospatial support to Burned Area Emergency Response Teams. *Fire Management Today* **71**(2), 15–18.
- Cocke AE, Fule PZ, Crouse JE (2005) Comparison of burn severity assessments using Differenced Normalized Burn Ratio and ground data. *International Journal of Wildland Fire* **14**, 189–198. doi:10.1071/WF04010
- De Santis A, Chuvieco E (2009) GeoCBI: a modified version of the Composite Burn Index for the initial assessment of the short-term burn severity from remotely sensed data. *Remote Sensing of Environment* **113**, 554–562. doi:10.1016/J.RSE.2008.10.011
- De Santis A, Chuvieco E, Vaughn P (2009) Short-term assessment of burn severity using the inversion of PROSPECT and GeoSail models. *Remote Sensing of Environment* **113**, 126–136. doi:10.1016/J.RSE.2008.08.008
- De Santis A, Asner GP, Vaughan PJ, Knapp DE (2010) Mapping burn severity and burning efficiency in California using simulation models and Landsat imagery. *Remote Sensing of Environment* **114**, 1535–1545. doi:10.1016/J.RSE.2010.02.008
- Debouk H, Riera-Tatche R, Vega-Garcia C (2013) Assessing post-fire regeneration in a Mediterranean mixed forest using LiDAR data and artificial neural networks. *Photogrammetric Engineering and Remote Sensing* **79**(12), 1121–1130. doi:10.14358/PERS.79.12.1121
- Díaz-Delgado R, Lloret F, Pons X (2003) Influence of fire severity on plant regeneration through remote sensing imagery. *International Journal of Remote Sensing* **24**, 1751–1763. doi:10.1080/01431160210144732
- Dillon GK, Holden ZA, Morgan P, Crimmins MA, Heyerdahl EK, Luce CH (2011) Both topography and climate affected forest and woodland burn severity in two regions of the western US, 1984 to 2006. *Ecosphere* **2**, art130. doi:10.1890/ES11-00271.1
- Doerr SH, Shakesby RA, Walsh RPD (2000) Soil water repellency: its causes, characteristics and hydro-geomorphological significance. *Earth-Science Reviews* **51**, 33–65. doi:10.1016/S0012-8252(00)00011-8
- Donato DC, Fontaine JB, Campbell JL, Robinson WD, Kauffman JB, Law BE (2009) Conifer regeneration in stand-replacement portions of a large mixed-severity wildfire in the Klamath-Siskiyou Mountains. *Canadian Journal of Forest Research* **39**, 823–838. doi:10.1139/X09-016
- Dyrness CT, Norum RA (1983) The effects of experimental fires on black spruce forest floors in interior Alaska. *Canadian Journal of Forest Research* **13**, 879–893. doi:10.1139/X83-118
- Eidenshink J, Schwind B, Brewer K, Zhu Z, Quayle B, Howard S (2007) A project for monitoring trends in burn severity. *Fire Ecology* **3**, 3–21. doi:10.4996/FIREECOLOGY.0301003
- Ekstrand S (1994) Assessment of forest damage with Landsat TM: correction for varying forest stand characteristics. *Remote Sensing of Environment* **47**, 291–302. doi:10.1016/0034-4257(94)90097-3
- Ellingson LJ, Kauffman JB, Cummings DL, Sanford RL, Jr, Jaramillo VJ (2000) Soil N dynamics associated with deforestation, biomass burning, and pasture conversion in a Mexican tropical dry forest. *Forest Ecology and Management* **137**, 41–51. doi:10.1016/S0378-1127(99)00311-4
- Epting J, Verbyla D, Sorbel B (2005) Evaluation of remotely sensed indices for assessing burn severity in interior Alaska using Landsat TM and ETM+. *Remote Sensing of Environment* **96**, 328–339. doi:10.1016/J.RSE.2005.03.002
- Fox DM, Maselli F, Carrega P (2008) Using SPOT images and field sampling to map burn severity and vegetation factors affecting post forest fire erosion risk. *Catena* **75**, 326–335. doi:10.1016/J.CATENA.2008.08.001
- French NHF, Kasischke ES, Hall RJ, Murphy KA, Verbyla DL, Hoy EE, Allen JL (2008) Using Landsat data to assess fire and burn severity in the North American boreal forest region: an overview and summary of results. *International Journal of Wildland Fire* **17**, 443–462. doi:10.1071/WF08007

- Ghermandi L, Gonzalez S, Lescano MN, Oddi F (2013) Effects of fire severity on early recovery of Patagonian steppes. *International Journal of Wildland Fire* **22**, 1055–1062. doi:10.1071/WF12198
- Graham RT, McCaffrey S, Jain TB (2004) Science basis for changing forest structure to modify wildfire behavior and severity. USDA Forest Service, Rocky Mountain Research Station, General Technical Report RMRS-GTR-120. (Fort Collins, CO).
- Guay T (2011) Rapid assessment of vegetation condition after wildfire. *Fire Management Today* **71**(2), 5–8.
- Halofsky JE, Hibbs DE (2009) Relationships among indices of fire severity in riparian zones. *International Journal of Wildland Fire* **18**, 584–593. doi:10.1071/WF07050
- Hessburg PF, Salter RB, James KM (2007) Re-examining fire severity relations in pre-management era mixed conifer forests: inferences from landscape patterns of forest structure. *Landscape Ecology* **22**, 5–24. doi:10.1007/S10980-007-9098-2
- Holden ZA, Jolly WM (2011) Modeling topographic influences on fuel moisture and fire danger in complex terrain for improved wildland fire management decision support. *Forest Ecology and Management* **262**, 2133–2141. doi:10.1016/J.FORECO.2011.08.002
- Holden ZA, Morgan P, Crimmins M, Steinhorst RK, Smith AMS (2007) Fire season precipitation variability influences fire extent and severity in a large southwestern wilderness area, United States. *Geophysical Research Letters* **34**, L16708. doi:10.1029/2007GL030804
- Holden ZA, Morgan P, Evans JS (2009) A predictive model of burn severity based on 20-year satellite-inferred burn severity data in a large southwestern US wilderness area. *Forest Ecology and Management* **258**, 2399–2406. doi:10.1016/J.FORECO.2009.08.017
- Hood S, Bentz BJ, Gibson K, Ryan KC, DeNitto G (2007) Assessing post-fire Douglas-fir mortality and Douglas-fir beetle attacks in the northern Rocky Mountains. USDA Forest Service, Rocky Mountain Research Station, General Technical Report RMRS-GTR-199. (Fort Collins, CO).
- Hudak AT, Brockett BH (2004) Mapping fire scars in a southern African savanna using Landsat imagery. *International Journal of Remote Sensing* **25**, 3231–3243. doi:10.1080/01431160310001632666
- Hudak AT, Morgan P, Bobbitt M, Lentile L (2007a) Characterizing stand-replacing harvest and fire disturbance patches in a forested landscape: a case study from Cooney Ridge, Montana. In 'Understanding Forest Disturbance and Spatial Patterns: Remote Sensing and GIS Approaches'. (Eds MA Wulder, SE Franklin) pp. 209–231. (Taylor and Francis: London, UK).
- Hudak AT, Morgan P, Bobbitt MJ, Smith AMS, Lewis SA, Lentile LB, Robichaud PR, Clark JT, McKinley RA (2007b) The relationship of multispectral satellite imagery to immediate fire effects. *Fire Ecology* **3**, 64–90. doi:10.4996/FIREECOLOGY.0301064
- Hudak AT, Rickert I, Morgan P, Strand E, Lewis SA, Robichaud PR, Hoffman C, Holden ZA (2011) Review of fuel treatment effectiveness in forests and rangelands and a case study from the 2007 megafires in central, Idaho, USA. USDA Forest Service, Rocky Mountain Research Station RMRS-GTR-252. (Fort Collins, CO).
- Hudak AT, Ottmar RD, Vihnanek BRE, Brewer NW, Smith AMS, Morgan P (2013) The relationship of post-fire white ash cover to surface fuel consumption. *International Journal of Wildland Fire* **22**, 780–785. doi:10.1071/WF12150
- Jain TB, Graham RT (2004) Is forest structure related to fire severity? Yes, no, and maybe: methods and insights in quantifying the answer. In 'Silviculture in Special Places: Proceedings of the 2003 National Silviculture Workshop' 8–11 Sept 2003, Granby, CO. (Eds WD Shepperd, LG Eskew) USDA Forest Service, Rocky Mountain Research Station Proceedings RMRS-P-34, pp. 217–234. (Fort Collins, CO).
- Jain TB, Graham RT (2007) The relation between tree burn severity and forest structure in the Rocky Mountains. In 'Restoring Fire-adapted Ecosystems: Proceedings of the 2005 National Silviculture Workshop', 6–10 June 2005. (Ed. RF Powers) USDA Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-203, pp. 213–250. (Albany, CA).
- Jain TB, Graham RT, Pilliod DS (2004) Tongue-tied. *Wildfire* **4**, 22–26.
- Jain TB, Graham RT, Pilliod DS (2006) The relation between forest structure and soil burn severity. In 'Fuels Management – How to Measure Success', 28–30 March 2006, Portland, OR. (Eds PL Andrews, B Butler) USDA Forest Service, Rocky Mountain Research Station, RMRS-P-41, pp. 615–631. (Fort Collins, CO).
- Jain TB, Gould WA, Graham RT, Pilliod DS, Lentile LB, Gonzalez G (2008) A soil burn severity index for understanding soil–fire relations in tropical forests. *Ambio* **37**, 563–568. doi:10.1579/0044-7447-37.7.563
- Jain TB, Pilliod DS, Graham RT, Lentile LB, Sandquist JE (2012) Index for characterizing post-fire soil environments in temperate coniferous forests. *Forests* **3**, 445–466. doi:10.3390/F3030445
- Jakubauskas ME, Lulla KP, Mausell PW (1990) Assessment of vegetation change in a fire-altered forest landscape. *Photogrammetric Engineering and Remote Sensing* **56**, 371–377.
- Johansen MP, Hakonson TE, Breshears DD (2001) Post-fire runoff and erosion from rainfall simulation: contrasting forests with shrublands and grasslands. *Hydrological Processes* **15**, 2953–2965. doi:10.1002/HYP.384
- Kane VR, Lutz JA, Roberts SL, Smith DF, McGaughey RJ, Povak NA, Brooks ML (2013) Landscape-scale effects of fire severity on mixed-conifer and red fir forest structure in Yosemite National Park. *Forest Ecology and Management* **287**, 17–31. doi:10.1016/J.FORECO.2012.08.044
- Karau EC, Keane RE (2010) Burn severity mapping using simulation modelling and satellite imagery. *International Journal of Wildland Fire* **19**, 710–724. doi:10.1071/WF09018
- Karau EC, Sikkink PG, Keane RE, Dillon G (2014) Integration of satellite imagery with simulation modeling improves fire severity mapping. *Environmental Management* **54**, 98–111. doi:10.1007/S00267-014-0279-X
- Kasischke E, Hoy EE, French NHF, Turetsky MR (2007a) Post-fire evaluation of the effects of fire on the environment using remotely-sensed data. In 'Towards an Operational Use of Remote Sensing in Forest Fire Management'. (Eds I Gitas, C Carmona) pp. 34–52. (Office for Official Publications of the European Communities: Luxembourg).
- Kasischke ES, Bourgeau-Chavez LL, Johnstone JF (2007b) Assessing spatial and temporal variations in surface soil moisture in fire-disturbed black spruce forests in Interior Alaska using spaceborne synthetic aperture radar imagery – implications for post-fire tree recruitment. *Remote Sensing of Environment* **108**, 42–58. doi:10.1016/J.RSE.2006.10.020
- Keane RE, Parsons RA (2010) A management guide to ecosystem restoration treatments: whitebark pine forests of the Northern Rocky Mountains. USDA Forest Service, Rocky Mountain Research Station, RMRS-GTR-232. (Fort Collins, CO).
- Keane RE, Holsinger L, Pratt S (2006) Simulating historical landscape dynamics using the landscape fire succession model LANDSUM version 4.0. USDA Forest Service, Rocky Mountain Research Station, RMRS-GTR-171CD. (Fort Collins, CO).
- Keane RE, Drury SA, Karau EC, Hessburg PF, Reynolds KM (2010) A method for mapping fire hazard and risk across multiple scales and its application in fire management. *Ecological Modelling* **221**, 2–18. doi:10.1016/J.ECOLMODEL.2008.10.022
- Keane RE, Herynk JM, Toney C, Urbanski SP, Lutes DC, Ottmar RD (2013) Evaluating the performance and mapping of three fuel classification systems using Forest Inventory and Analysis surface fuel measurements. *Forest Ecology and Management* **305**, 248–263. doi:10.1016/J.FORECO.2013.06.001
- Keeley JE (2009) Fire intensity, fire severity and burn severity: a brief review and suggested usage. *International Journal of Wildland Fire* **18**, 116–126. doi:10.1071/WF07049

- Keeley JE, Brennan T, Pfaff AH (2008) Fire severity and ecosystem responses following crown fires in California shrublands. *Ecological Applications* **18**, 1530–1546. doi:10.1890/07-0836.1
- Key CH (2006) Ecological and sampling constraints on defining landscape fire severity. *Fire Ecology* **2**, 34–59. doi:10.4996/FIREECOLOGY.0202034
- Key CH, Benson NC (2006) Landscape assessment: ground measure of severity, the Composite Burn Index; and remote sensing of severity, the Normalized Burn Ratio. In 'FIREMON: Fire Effects Monitoring and Inventory System'. (Eds DC Lutes, RE Keane, JF Caratti, CH Key, NC Benson, LJ Gangi) USDA Forest Service, Rocky Mountains Research Station General Technical Report RMRS-GTR-164-CD, pp. 219–279. (Fort Collins, CO).
- Keyser TL, Smith FW, Lentile LB, Shepperd WD (2006) Modeling postfire mortality of ponderosa pine following a mixed-severity wildfire in the Black Hills: the role of tree morphology and direct fire effects. *Forest Science* **52**, 530–539.
- King KJ, Bradstock RA, Cary GJ, Chapman J, Marsden-Smedley JB (2008) The relative importance of fine-scale fuel mosaics on reducing fire risk in south-west Tasmania, Australia. *International Journal of Wildland Fire* **17**, 421–430. doi:10.1071/WF07052
- Kolden CA, Rogan J (2013) Mapping wildfire burn severity in the arctic tundra from downsampled MODIS data. *Arctic, Antarctic, and Alpine Research* **45**(1), 64–76. doi:10.1657/1938-4246-45.1.64
- Kotliar NB, Reynolds EW, Deutschman DH (2008) American Three-toed Woodpecker response to burn severity and prey. *Fire Ecology Special Issue* **4**, 26–45. doi:10.4996/FIREECOLOGY.0402026
- Kremens RL, Smith AMS, Dickinson MB (2010) Fire metrology: current and future directions in physics-based measurements. *Fire Ecology* **6**, 13–25. doi:10.4996/FIREECOLOGY.0602013
- Kuenzi AM, Fulé PZ, Sieg CH (2008) Effects of fire severity and pre-fire stand treatment on plant community recovery after a large wildfire. *Forest Ecology and Management* **255**, 855–865. doi:10.1016/J.FORECO.2007.10.001
- Kumar L, Clarke P, Munoz C, Knox K (2008) Mapping of fire severity and comparison of severity indices across vegetation types in Gibraltar Range National Park, Australia. *International Archives of the Photogrammetry, Remote Sensing, and Spatial Information Sciences* **37**, 1477–1482.
- Kushla JD, Ripple WJ (1997) The role of terrain in a fire mosaic of a temperate coniferous forest. *Forest Ecology and Management* **95**, 97–107. doi:10.1016/S0378-1127(97)82929-5
- Kwak DA, Chung J, Lee WK, Kafatos M, Lee SY, Cho HK, Lee SH (2010) Evaluation for damaged degree of vegetation by forest fire using Lidar and a digital aerial photograph. *Photogrammetric Engineering and Remote Sensing* **76**, 277–287. doi:10.14358/PERS.76.3.277
- Landmann T (2003) Characterizing sub-pixel Landsat ETM+ fire severity on experimental fires in the Kruger National Park, South Africa. *South African Journal of Science* **99**, 357–360.
- Larrivé M, Fahrig L, Drapeau P (2005) Effects of a recent wildfire and clearcuts on ground-dwelling boreal forest spider assemblages. *Canadian Journal of Forest Research* **35**, 2575–2588. doi:10.1139/X05-169
- Lee SW, Lee MB, Lee YG, Won MS, Kim JJ, Hong SK (2009) Relationship between landscape structure and burn severity at the landscape and class levels in Samchuck, South Korea. *Forest Ecology and Management* **258**, 1594–1604. doi:10.1016/J.FORECO.2009.07.017
- Lentile LB, Smith FW, Shepperd WD (2005) Patch structure, fire-scar formation, and tree regeneration in a large mixed-severity fire in the South Dakota Black Hills, USA. *Canadian Journal of Forest Research* **35**, 2875–2885. doi:10.1139/X05-205
- Lentile LB, Holden ZA, Smith AMS, Falkowski MJ, Hudak AT, Morgan P, Lewis SA, Gessler PE, Benson NC (2006) Remote sensing techniques to assess active fire characteristics and post-fire effects. *International Journal of Wildland Fire* **15**, 319–345. doi:10.1071/WF05097
- Lentile LB, Morgan P, Hudak AT, Bobbitt MJ, Lewis SA, Smith AM, Robichaud PR (2007) Post-fire burn severity and vegetation response following eight large wildfires across the western United States. *Fire Ecology* **3**, 91–108. doi:10.4996/FIREECOLOGY.0301091
- Lentile LB, Smith AMS, Hudak AT, Morgan P, Bobbitt MJ, Lewis SA, Robichaud PR (2009) Remote sensing for prediction of 1-year post-fire ecosystem condition. *International Journal of Wildland Fire* **18**, 594–608. doi:10.1071/WF07091
- Lewis SA, Wu JQ, Robichaud PR (2006) Assessing burn severity and comparing soil water repellency, Hayman Fire, Colorado. *Hydrological Processes* **20**, 1–16. doi:10.1002/HYP.5880
- Lutes DC, Keane RE, Caratti JF, Key CH, Benson NC, Sutherland S, Gangi LJ (2006) FIREMON: fire effects monitoring and inventory system. USDA Forest Service, Rocky Mountain Research Station, RMRS-GTR-164-CD. (Fort Collins, CO).
- Magnussen S, Wulder MA (2012) Post-fire canopy height recovery in Canada's boreal forests using Airborne Laser Scanner (ALS). *Remote Sensing* **4**, 1600–1616. doi:10.3390/RS4061600
- Mallek C, Safford H, Viers J, Miller J (2013) Modern departures in fire severity and area vary by forest type, Sierra Nevada and southern Cascades, California, USA. *Ecosphere* **4**(12), art153. doi:10.1890/ES13-00217.1
- Megown K, Finco M, Brewer K, Schwind B (2011) Accelerated re-measurement and evaluation of burned areas. *Fire Management Today* **71**(2), 9–11.
- Miller M (2001) Fire behavior and characteristics. In 'Fire Effects Guide'. (Ed. M Miller), pp. 12–38. (National Interagency Fire Center: Boise, ID).
- Miller JD, Safford HD (2012) Trends in wildfire severity 1984–2010 in the Sierra Nevada, Modoc Plateau, and southern Cascades, California, USA. *Fire Ecology* **8**, 41–57. doi:10.4996/FIREECOLOGY.0803041
- Miller JD, Thode AE (2007) Quantifying burn severity in a heterogeneous landscape with a relative version of the delta Normalized Burn Ratio (dNBR). *Remote Sensing of Environment* **109**(1), 66–80. doi:10.1016/J.RSE.2006.12.006
- Miller JD, Nyhan JW, Yool SR (2003) Modeling potential erosion due to the Cerro Grande Fire with a GIS-biased implementation of the Revised Universal Soil Loss Equation. *International Journal of Wildland Fire* **12**, 85–100. doi:10.1071/WF02017
- Miller JD, Knapp EE, Key CH, Skinner CN, Isbell CJ, Creasy RM, Sherlock JW (2009) Calibration and validation of the relative differenced Normalized Burn Ratio (RdNBR) to three measures of fire severity in the Sierra Nevada and Klamath Mountains, California, USA. *Remote Sensing of Environment* **113**, 645–656. doi:10.1016/J.RSE.2008.11.009
- Miller JD, Collins BM, Lutz JA, Stephens SL, van Wagendonk JW, Yasuda DA (2012) Differences in wildfires among ecoregions and land management agencies in the Sierra Nevada region, California, USA. *Ecosphere* **3**, art80. doi:10.1890/ES12-00158.1
- Moreno JM, Oechel WC (1989) A simple method for estimating fire intensity after a burn in California chaparral. *Oecologia Plantarum* **10**, 57–68.
- Moreno JM, Oechel WC (1991) Fire intensity effects on germination of shrubs and herbs in southern California chaparral. *Ecology* **72**, 1993–2004. doi:10.2307/1941554
- Morgan P, Neuenschwander LF (1988) Seed-bank contributions to regeneration of shrub species after clear-cutting and burning. *Canadian Journal of Botany* **66**, 169–172. doi:10.1139/B88-026
- Morgan P, Hardy CC, Swetnam TW, Rollins MG, Long DG (2001) Mapping fire regimes across time and space: understanding coarse and fine-scale fire patterns. *International Journal of Wildland Fire* **10**, 329–342. doi:10.1071/WF01032
- National Wildfire Coordinating Group (2012) Glossary of Wildland Fire Terminology. Available at <http://www.nwcg.gov/pms/pubs/glossary/> [Verified 22 February 2014]

- Neary DG, Ryan KC, DeBano LF (Eds) (2008) Wildland fire in ecosystems: effects of fire on soils and water. USDA Forest Service, Pacific Northwest Research Station, RMRS-GTR-42-vol.4. (Ogden, UT)
- Neff JC, Harden JW, Gleixner G (2005) Fire effects on soil organic matter content, composition, and nutrients in boreal interior Alaska. *Canadian Journal of Forest Research* **35**, 2178–2187. doi:10.1139/X05-154
- Oliveras I, Gracia M, Mora G, Retana J (2009) Factors influencing the pattern of fire severities in a large wildfire under extreme meteorological conditions in the Mediterranean basin. *International Journal of Wildland Fire* **18**, 755–764. doi:10.1071/WF08070
- Ottmar RD, Burns MF, Hall JN, Hanson AD (1993) CONSUME user's guide. USDA Forest Service, Pacific Northwest Research Station, PNW-GTR-304. (Portland, OR)
- Parsons A, Robichaud PR, Lewis SA, Napper C, Clark JT (2010) Field guide for mapping post-fire soil burn severity. USDA Forest Service, Rocky Mountain Research Station, Report RMRS-GTR-243. (Fort Collins, CO)
- Patterson MW, Yool SR (1998) Mapping fire-induced vegetation mortality using Landsat thematic mapper data: a comparison of linear transformation techniques. *Remote Sensing of Environment* **65**, 132–142. doi:10.1016/S0034-4257(98)00018-2
- Pausas JG, Ouadah N, Ferran A, Gimeno T, Vallejo R (2003) Fire severity and seedling establishment in *Pinus halepensis* woodlands, eastern Iberian Peninsula. *Plant Ecology* **169**, 205–213. doi:10.1023/A:1026019528443
- Picotte JJ, Robertson KM (2011) Validation of remote sensing of burn severity in south-eastern US ecosystems. *International Journal of Wildland Fire* **20**, 453–464. doi:10.1071/WF10013
- Pierson FB, Robichaud PR, Spaeth KE (2001) Spatial and temporal effects of wildfire on the hydrology of a steep rangeland watershed. *Hydrological Processes* **15**, 2905–2916. doi:10.1002/HYP.381
- Regelbrugge JC, Smith DW (1994) Postfire tree mortality in relation to wildfire severity in mixed oak forests in the Blue Ridge of Virginia. *Northern Journal of Applied Forestry* **11**, 90–97.
- Reinhardt ED, Keane RE, Brown JK (1997) First Order Fire Effects Model: FOFEM 4.0, user's guide. USDA Forest Service, Intermountain Research Station, INT-GTR-344. (Ogden, UT)
- Robichaud PR, Hungerford RD (2000) Water repellency by laboratory burning of four northern Rocky Mountain forest soils. *Journal of Hydrology* **231–232**, 207–219. doi:10.1016/S0022-1694(00)00195-5
- Robichaud PR, MacDonald L, Freeouf J, Neary D, Martin D, Ashmun L (2003) Postfire rehabilitation of the Hayman Fire. In 'Hayman Fire Case Study'. (Ed. RT Graham) pp. 293–313. USDA Forest Service, Rocky Mountain Research Station RMRS-GTR-114. (Ogden, UT)
- Robichaud PR, Elliot WJ, Pierson FB, Hall DE, Moffet CA (2007a) Predicting postfire erosion and mitigation effectiveness with a web-based probabilistic erosion model. *Catena* **71**, 229–241. doi:10.1016/J.CATENA.2007.03.003
- Robichaud PR, Lewis SA, Laes DYM, Hudak AT, Kokaly RF, Zamudio JA (2007b) Postfire soil burn severity mapping with hyperspectral image unmixing. *Remote Sensing of Environment* **108**, 467–480. doi:10.1016/J.RSE.2006.11.027
- Romme WH (2005) The importance of multiscale spatial heterogeneity in wildland fire management and research. In 'Ecosystem Function in Heterogeneous Landscapes'. (Eds GM Lovett, CG Jones, MG Turner, KC Weathers) pp. 253–266. (Springer: New York)
- Roy D, Landmann T (2005) Characterizing the surface heterogeneity of fire effects using multi-temporal reflective wavelength data. *International Journal of Remote Sensing* **26**, 4197–4218. doi:10.1080/01431160500112783
- Roy DP, Boschetti L, Trigg SN (2006) Remote sensing of fire severity: assessing the performance of the normalized burn ratio. *Geoscience and Remote Sensing Letters, IEEE* **3**(1), 112–116. doi:10.1109/LGRS.2005.858485
- Rumpel C, Gonzalez-Perez JA, Bardoux G, Largeau C, Gonzalez-Vila FJ, Valentin C (2007) Composition and reactivity of morphologically distinct charred materials left after slash-and-burn practices in agricultural tropical soils. *Organic Geochemistry* **38**, 911–920. doi:10.1016/J.ORGGEOCHEM.2006.12.014
- Russell-Smith J, Ryan PG, Durieu R (1997) A Landsat MSS-derived fire history of Kakadu National Park, monsoonal northern Australia, 1980–94: seasonal extent, frequency and patchiness. *Journal of Applied Ecology* **34**, 748–766. doi:10.2307/2404920
- Ryan KC, Noste NV (1985) Evaluating prescribed fires. In 'Wilderness Fire Symposium Missoula, MT', 22–26 September 1983, Missoula, MT. (Eds JE Lotan, BM Kilgore, WC Fischer, RW Mutch) pp. 230–237. USDA Forest Service, Intermountain Research Station, INT-182. (Ogden, UT)
- Schimmel J, Granstrom A (1996) Fire severity and vegetation response in the boreal Swedish forest. *Ecology* **77**, 1436–1450. doi:10.2307/2265541
- Shakesby RA, Chafer CJ, Doerr SH, Blake WH, Wallbrink P, Humphreys GS, Harrington BA (2003) Fire severity, water repellency characteristics and hydrogeomorphological changes following the Christmas 2001 Sydney forest fires. *The Australian Geographer* **34**, 147–175. doi:10.1080/00049180301736
- Simard AJ (1991) Fire severity, changing scales, and how things hang together. *International Journal of Wildland Fire* **1**, 23–34. doi:10.1071/WF9910023
- Smith JK, ed. (2000) Wildland fire in ecosystems: effects of fire on fauna. USDA Forest Service, Rocky Mountain Research Station, RMRS-GTR-42-vol 1. (Ogden, UT)
- Smith AMS, Wooster MJ, Drake NA, Dipotso FM, Falkowski MJ, Hudak AT (2005) Testing the potential of multi-spectral remote sensing for retrospectively estimating fire severity in African savannahs. *Remote Sensing of Environment* **97**(1), 92–115. doi:10.1016/J.RSE.2005.04.014
- Smith AMS, Lentile LB, Hudak AT, Morgan P (2007) Evaluation of linear spectral unmixing and dNBR for predicting post-fire recovery in a North American ponderosa pine forest. *International Journal of Remote Sensing* **28**(22), 5159–5166. doi:10.1080/01431160701395161
- Smith AMS, Eitel JUH, Hudak AT (2010) Spectral analysis of charcoal on soils: implications for wildland fire severity mapping methods. *International Journal of Wildland Fire* **19**, 976–983. doi:10.1071/WF09057
- Soverel NO, Perrakis DDB, Coops NC (2010) Estimating burn severity from Landsat dNBR and RdNBR indices across western Canada. *Remote Sensing of Environment* **114**, 1896–1909. doi:10.1016/J.RSE.2010.03.013
- Tanase MA, Perez-Cabello F, de la Riva J, Santoro M (2010a) TerraSAR-X data for burn severity evaluation in Mediterranean forests on sloped terrain. *IEEE Transactions on Geoscience and Remote Sensing* **48**, 917–929. doi:10.1109/TGRS.2009.2025943
- Tanase MA, Santoro M, de la Riva J, Perez-Cabello F, Le Toan T (2010b) Sensitivity of X-, C-, and L-band SAR backscatter to burn severity in Mediterranean pine forests. *IEEE Transactions on Geoscience and Remote Sensing* **48**, 3663–3675. doi:10.1109/TGRS.2010.2049653
- Tanase MA, Santoro M, Wegmuller U, de la Riva J, Perez-Cabello F (2010c) Properties of X-, C- and L-band repeat-pass interferometric SAR coherence in Mediterranean pine forests affected by fires. *Remote Sensing of Environment* **114**, 2182–2194. doi:10.1016/J.RSE.2010.04.021
- Thompson JR, Spies TA (2010) Factors associated with crown damage following recurring mixed-severity wildfires and post-fire management in southwestern Oregon. *Landscape Ecology* **25**, 775–789. doi:10.1007/S10980-010-9456-3
- Trigg S, Flasse S (2000) Characterising the spectral-temporal response of burned savanna using *in situ* spectroradiometry and infrared thermometry. *International Journal of Remote Sensing* **21**, 3161–3168. doi:10.1080/01431160050145045



- Tucker CJ (1979) Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment* **8**(2), 127–150. doi:10.1016/0034-4257(79)90013-0
- Turner MG, Hargrove WW, Gardner RH, Romme WH (1994) Effects of fire on landscape heterogeneity in Yellowstone National Park, Wyoming. *Journal of Vegetation Science* **5**, 731–742. doi:10.2307/3235886
- Turner MG, Romme WH, Gardner RH (1999) Prefire heterogeneity, fire severity, and early postfire plant reestablishment in subalpine forests of Yellowstone National Park, Wyoming. *International Journal of Wildland Fire* **9**, 21–36. doi:10.1071/WF99003
- Tyler CM (1995) Factors contributing to postfire seedling establishment in chaparral: direct and indirect effects of fire. *Journal of Ecology* **83**, 1009–1020. doi:10.2307/2261182
- Ulery AL, Graham RC (1993) Forest fire effects on soil color and texture. *Soil Science Society of America Journal* **57**, 135–140. doi:10.2136/SSSAJ1993.03615995005700010026X
- US Department of Interior (2003) Fire monitoring handbook. (Fire Management Program Center, National Interagency Fire Center: Boise, ID)
- Veraverbeke S, Lhermitte S, Verstraeten WW, Goossens R (2010a) The temporal dimension of differenced Normalized Burn Ratio (dNBR) fire/burn severity studies: the case of the large 2007 Peloponnese wildfires in Greece. *Remote Sensing of Environment* **114**, 2548–2563. doi:10.1016/J.RSE.2010.05.029
- Veraverbeke S, Verstraeten W, Lhermitte S, Goossens R (2010b) Evaluating Landsat Thematic Mapper spectral indices for estimating burn severity of the 2007 Peloponnese wildfires in Greece. *International Journal of Wildland Fire* **19**, 558–569. doi:10.1071/WF09069
- Veraverbeke S, Lhermitte S, Verstraeten WW, Goossens R (2011) Evaluation of pre/post-fire differenced spectral indices for assessing burn severity in a Mediterranean environment with Landsat Thematic Mapper. *International Journal of Remote Sensing* **32**, 3521–3537. doi:10.1080/01431161003752430
- Wang C, Glenn NF (2009) Estimation of fire severity using pre- and post-fire LiDAR data in sagebrush steppe rangelands. *International Journal of Wildland Fire* **18**, 848–856. doi:10.1071/WF08173
- Wang GG, Kembell KJ (2005) Effects of fire severity on early development of understory vegetation. *Canadian Journal of Forest Research* **35**, 254–262. doi:10.1139/X04-177
- White JD, Ryan KC, Key CC, Running SW (1996) Remote sensing of forest fire severity and vegetation recovery. *International Journal of Wildland Fire* **6**, 125–136. doi:10.1071/WF9960125
- Wulder MA, White JC, Alvarez F, Han T, Rogan J, Hawkes B (2009) Characterizing boreal forest wildfire with multi-temporal Landsat and LIDAR data. *Remote Sensing of Environment* **113**, 1540–1555. doi:10.1016/J.RSE.2009.03.004
- Yeager CM, Northup DE, Grow CC, Barns SM, Kuske CR (2005) Changes in nitrogen-fixing and ammonia-oxidizing bacterial communities in soil of a mixed conifer forest after wildfire. *Applied and Environmental Microbiology* **71**, 2713–2722. doi:10.1128/AEM.71.5.2713-2722.2005
- Zarriello TJ, Knick ST, Rotenberry JT (1995) Producing a burn/disturbance map for the Snake River Birds of Prey National Conservation Area. In ‘Snake River Birds of Prey National Conservation Area Research and Monitoring Annual Report’. (Boise, ID)