

Changes in Alternative Pathway and Mitochondrial Respiration in Avocado in Response to Elevated Carbon Dioxide Levels

Diana L. Lange¹ and Adel A. Kader

Department of Pomology, University of California, Davis, CA 95616

ADDITIONAL INDEX WORDS. *Persea americana*, alternative oxidase, cyanide-resistant respiration, respiratory control, residual respiration

ABSTRACT. Partially ripened avocado [*Persea americana* (Mill.) cv. Hass] fruit harvested in either June or Aug. 1994 were kept at 10 °C in air (21% O₂), 20% CO₂ (17% O₂, balance N₂), or 40% CO₂ (13% O₂, balance N₂) for 7 to 12 days and then were transferred to air at 10 °C for 2 to 3 days. Mitochondrial respiration was stimulated in response to elevated CO₂ treatments at 10 °C. A shift to alternative pathway (Alt) respiration occurred on day 4 in experiments using avocados from both harvest dates, with a return to initial levels in only the 20% CO₂-treated fruit (June-harvested fruit after return to air). Elevated CO₂ at 20 °C decreased the in vitro O₂ consumption of isolated mitochondria compared to mitochondria kept in air. The Alt pathway contributed less to the total O₂ uptake of CO₂-treated mitochondria compared to mitochondria kept in air. The respiratory control ratios of the CO₂-treated fruit and mitochondria were higher and lower, respectively, than the air controls. Induction of 33 to 37 kD proteins (corresponding to the size of the alternative oxidase proteins) occurred in avocados after 4 days in 40% CO₂. These results indicate that elevated CO₂ has various effects depending on concentration, duration and temperature of exposure, and mitochondrial function of avocado fruit, such as increased and altered respiratory oxidation and up-regulation of alternative oxidase proteins.

The alternative respiratory (Alt) pathway, which is cyanide resistant, is present in a wide range of plant tissues (Latices, 1982; Solomos, 1977; Solomos and Latices, 1976), including avocados (Latices, 1982), cucumbers (*Cucumis sativus* L.) (Morohashi et al., 1991), oranges [*Citrus sinensis* (L.) Osbeck] (Bruemmer, 1989), apples (*Malus domestica* Borkh.), and peppers (*Capsicum annuum* L.) (Lurie and Klein, 1989). While the function of this nonphosphorylating pathway is apparent in thermogenic species, its role in nonthermogenic tissue is unclear (McIntosh, 1994). The Alt pathway may provide a way to disperse excess reducing power when cytochrome oxidase is inhibited (Lambers, 1982). Collier and Cummins (1991) reported that the Alt pathway may have been used during *Saxifraga cernua* L. petal unfolding as an inefficient energy source. During fruit ripening, the energy charge of the tissue is high and the rate of cytochrome (Cyt) pathway oxidation is coupled tightly to oxidative phosphorylation. The Alt pathway may provide a means of oxidizing the large electron flow that occurs during fruit ripening without producing as large amounts of ATP as occurs when the Cyt pathway is operating alone.

The ability of CO₂ to stimulate the Alt pathway has been reported previously, but most of these studies focused on the effects of CO₂ in combination with ethylene (Day et al., 1978; Latices, 1987). Latices (1982) concluded that CO₂ appeared to enhance ethylene-induced Alt respiration. Carbon dioxide alone has sustained the cyanide resistant pathway in wheat (*Triticum aestivum* L.) seedlings (McCaig and Hill, 1977), carnation (*Dian-*

thus caryophyllus L.) callus (Palet et al., 1991), and Jerusalem artichoke (*Helianthus tuberosus* L.) tubers (Stegink and Siedow, 1986). To our knowledge, the ability of CO₂ to stimulate the Alt pathway in fruit tissue has not been demonstrated. One of our objectives was to determine if, and to what degree, CO₂ stimulates the Alt pathway.

Elthon and McIntosh (1987) identified alternative oxidase (AltOx) proteins (molecular weights = 35 to 37 kD) in the thermogenic spadix of *Sauromatum guttatum* Schott and later developed monoclonal antibodies to these proteins (Elthon et al., 1989). AltOx proteins subsequently have been identified in a number of nonthermogenic plant species, including avocado fruit, etiolated mung bean (*Vigna radiata* L.) hypocotyls, fresh potato (*Solanum tuberosum* L.) tubers, and tobacco (*Nicotiana tabacum* L.) callus (Elthon et al., 1989). Rhoads and McIntosh (1993) reported that increased Alt pathway activity correlated with an accumulation of a 35-kD protein in tobacco suspension cultures and de novo transcription and translation were necessary to cause the maximum accumulation of the 35-kD protein. Yoshimoto et al. (1989) reported that the Alt pathway was induced simultaneously with the appearance of a 36-kD protein in yeast. Although the AltOx proteins are present in fruit tissue, there is still some question of whether or not they are constitutive and always present or synthesized de novo in coordination with developmental changes. AltOx in pear (*Pyrus communis* L.) was constitutive, and changes in its activity did not involve transcriptional or translational events (R.J. Romani, unpublished data).

Huang and Romani (1991) reported that avocado mitochondria have an intrinsic homeostatic capacity in which disrupted energy-linked functions are self-restored. In addition, the storage life of avocados can be extended by exposure to elevated CO₂ atmospheres (Spalding and Reeder, 1974; Truter and Eksteen, 1987). Therefore, we used avocado fruit and their mitochondria to study the direct and indirect effects of CO₂ stress on enhancement and partitioning of respiration in fruit tissue.

Received for publication 5 Mar. 1996. Accepted for publication 19 Sept. 1996. Research supported in part by U.S. Dept. of Agriculture research agreement no. 58-319R-3-004. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Extension specialist; current address, Dept. of Horticulture, Southwest Purdue Agricultural Research and Extension Center, R.R. 6, Box 139A, Vincennes, IN 47591.

Materials and Methods

PLANT MATERIAL AND TREATMENTS. 'Hass' avocado fruit were harvested at commercial maturity ($\geq 24\%$ dry mass) from an orchard in Santa Barbara County, Calif., and were transported to our laboratory in Davis, Calif., where they were stored at 10°C for up to 1 week until initiation of the experiments. Fruit uniform in size and free from defects were selected. Before initiating experiments, fruit were partially ripened at 20°C for 4 d with a continuous flow ethylene at $10\ \mu\text{L}\cdot\text{L}^{-1}$ in humidified air at a flow rate of $400\ \text{mL}\cdot\text{min}^{-1}$.

Experiments were conducted at 10°C for up to 15 d. Nine fruit were placed in a 4-L glass jar and ventilated with humidified air ($21\% \text{O}_2 + 0.3\% \text{CO}_2 + \text{balance N}_2$) or a specified gas mixture at a continuous flow rate of $100\ \text{mL}\cdot\text{min}^{-1}$. The gas mixtures included $20\% \text{CO}_2$ mixed with air ($20\% \text{CO}_2 + 17\% \text{O}_2 + \text{balance N}_2$) or $40\% \text{CO}_2$ mixed with air ($40\% \text{CO}_2 + 13\% \text{O}_2 + \text{balance N}_2$). Avocado fruit were kept in air or CO_2 -enriched atmospheres at 10°C for 12 (June 1994) or 7 (Aug. 1994) d and then transferred to air for 3 (June) or 2 (August) d.

Individual fruit were selected for each experiment based on screening for uniform C_2H_4 production rates. Partially ripened avocados were screened after 4 d of ripening. For every sampling date, treatments were replicated three times, each replicate having three fruit. All of the fruit for one sampling date for each treatment were contained in the same jar.

Intact avocados were exposed to air or CO_2 -enriched atmospheres at 10°C and then were used in mitochondrial assays unless otherwise stated.

MITOCHONDRIAL ISOLATION AND PARTIAL PURIFICATION. Avocado fruit mitochondria were isolated using the method of Moreau and Romani (1982b). All extraction procedures were performed over ice. Homogenization of the fruit tissue was conducted in a 10°C storage room. For the mitochondrial O_2 uptake assays, $100\ \text{g}$ of tissue were homogenized in $300\ \text{mL}$ isolation medium using a fine wire-mesh screen submerged in medium. The isolation medium consisted of $0.25\ \text{M}$ sucrose, $50\ \text{mM}$ potassium phosphate ($\text{pH} = 7.2$), $5\ \text{mM}$ EDTA, $5\ \text{mM}$ β -mercaptoethanol, 0.2% (w/v) soluble ($40,000\ \text{MW}$) PVP, and 0.1% BSA. In all experiments, the homogenate was filtered through four layers of cheesecloth and centrifuged at $2000\times g$ for 10 min. The supernatant was filtered through four layers of cheesecloth and centrifuged at $10,000\times g$ for 15 min. This supernatant was discarded, and the pellet was resuspended in $3\ \text{mL}$ wash medium and homogenized in the presence of $11\ \text{mL}$ additional wash medium. The wash medium consisted of $0.25\ \text{M}$ sucrose, $50\ \text{mM}$ potassium phosphate ($\text{pH} = 7.2$), $5\ \text{mM}$ β -mercaptoethanol, and 0.1% BSA. This homogenate was recentrifuged at $2000\times g$ for 5 min to pelletize remaining chloroplasts out of the mitochondrial supernatant.

Partial purification of the mitochondria was conducted in the following manner. An $8,000\times g$ centrifugation step was substituted with a $20,000\times g$ centrifugation step through $20\ \text{mL}$ of 25% Percoll (Sigma Chemical Co., St. Louis) "pad" for 10 min [as described in Day and Hanson (1977) and modified by Romani (1994) personal communication]. This partial purification was necessary to eliminate most of the nonmitochondrial organelles that would interfere with the oxidase assays. A full Percoll purification system was not used (Moreau and Romani, 1982a) because a more rapid cleanup was necessary. The pad medium consisted of 25% Percoll, $0.25\ \text{M}$ sucrose, and $50\ \text{mM}$ potassium phosphate ($\text{pH} = 7.2$). The resulting mitochondrial pellet was washed with wash medium and centrifuged at $2000\times g$ for 5 min several times to remove the Percoll with the supernatant, and then the mitochondrial pellet was resuspended

in 0.3 to $0.5\ \text{mL}$ wash medium. Between the addition of wash medium and centrifugation steps, the supernatant was removed by vacuum aspiration and discarded. Protein content of the purified mitochondrial preparation was determined by the Bradford (1976) method using BSA as the standard.

POLAROGRAPHIC MEASUREMENT OF O_2 UPTAKE AND RESPIRATORY CONTROL. Mitochondrial functions were monitored, including the measurement of O_2 uptake, the contributions of different oxidase pathways toward total O_2 consumption, and respiratory control ratios that indicate the degree of oxidative phosphorylation. One milliliter of reaction media [$0.25\ \text{M}$ sucrose, 0.1% BSA, $50\ \text{mM}$ phosphate ($\text{pH} 7.2$), $1\ \text{mM}$ MgCl_2 , $10\ \mu\text{M}$ CoA, $100\ \mu\text{M}$ TPP (thiamin pyrophosphate), and $100\ \mu\text{M}$ NAD^+] and an appropriate aliquot of mitochondria (0.2 to $0.4\ \text{mg}$ protein) were placed in a 1-mL chamber fashioned from plexiglass and equipped with an O_2 electrode (Yellow Springs Instruments, Yellow Springs, Ohio). The electrode was maintained at a constant temperature of 25°C . The reaction mixture was constantly stirred with a micro stir bar. The O_2 content of air-saturated water was estimated according to Estabrook (1987). Oxygen consumption was measured following the addition of $10\ \text{mM}$ malate as substrate. The contribution of the Cyt and Alt pathways to total respiration was determined using $1\ \text{mM}$ KCN and $3\ \text{mM}$ SHAM (salicylhydroxamic acid), respectively. KCN was always added first to avoid stimulation of O_2 uptake by SHAM through CN-sensitive peroxidases (Møller et al., 1988). Respiratory control ratios (RCR) were determined by adding stoichiometric amounts of ADP ($\approx 0.15\ \text{mM}$) and measuring the state 3 : state 4 O_2 consumption rates ratio.

IN VITRO EXPOSURE OF ISOLATED MITOCHONDRIA TO ELEVATED CO_2 ATMOSPHERES. To measure the effects of in vitro exposure to stress levels of CO_2 , mitochondria were extracted as previously described from avocado fruit stored in air for <2 weeks at 10°C and were ripened partially with a continuous flow of C_2H_4 at $10\ \mu\text{L}\cdot\text{L}^{-1}$ in humidified air at a flow rate of $400\ \text{mL}\cdot\text{min}^{-1}$. The fruit were ripened partially to facilitate tissue homogenization. The mitochondria were suspended in reaction mixture and separated into three equal aliquots (one per treatment) of $10\ \text{mL}$ each in a 50-mL Erlenmeyer flask. The mitochondrial suspensions were held at 20°C on a rotary shaker on a low setting. Each flask was sealed with a serum cap supplied with inlet and outlet flow lines to the headspace. Air or the CO_2 -enriched atmosphere ($20\% \text{CO}_2 + 17\% \text{O}_2$ or $40\% \text{CO}_2 + 13\% \text{O}_2$) was flushed through the headspace at $20\ \text{mL}\cdot\text{min}^{-1}$. Before each assay, the headspace was aerated momentarily to provide necessary O_2 as a substrate for oxidase reactions. At the designated times, a 1-mL aliquot of the mitochondrial mixture was removed and tested for respiratory control and different oxidase pathway contributions.

DETERMINING ALTERNATIVE OXIDASE (ALT Ox) PROTEIN ABUNDANCE. Mitochondria were isolated and partially purified as previously described with minor modifications. No BSA was used in the isolation, wash, or purification medium to avoid overestimation of protein in the samples. Partially purified samples were dialyzed overnight at 4°C in $5\ \text{mM}$ phosphate buffer ($\text{pH} 7.2$) and 10% glycerol to remove sucrose and salts, frozen in liquid N_2 , and stored at -80°C for ≤ 5 months. Samples were prepared as previously described by Elthon and McIntosh (1987) with the following modifications: β -mercaptoethanol concentration was increased to 10% (v/v), and the samples were boiled for 5 min before adding the tracking dye 0.04% bromophenol blue. Electrophoresis was conducted with the buffer system of Laemmli (1970) using a 5% stacking and a 15% polyacrylamide separating gel. The gels were stained with Coomassie Blue to verify equal loading of protein. Bio-Rad low molecular-mass protein standards, either unstained

or biotinylated for gels or immunoblots, respectively, were used to estimate molecular mass. Protein blotting followed the protocol of Blake et al. (1984) except that antibody incubations were for 2 h at room temperature. In accordance with Elthon and McIntosh (1987), the AltOx proteins were detected with a 1 monoclonal antibody : 10 AltOx dilution from *Sauromatum guttatum* (courtesy of T. Elthon).

Results and Discussion

The O₂ uptake of mitochondria extracted from partially ripe 'Hass' avocados that were kept at 10 °C in air (Fig. 1) followed a similar pattern to that of the intact fruit (Lange and Kader, 1996). Uptake of O₂ by mitochondria from June-harvested fruit kept in air gradually declined over the 15-d storage period at 10 °C, whereas O₂ uptake by mitochondria from August-harvested fruit peaked on day 4. Avocado fruit exposed to 20% CO₂ had an increase in the O₂ uptake of extracted mitochondria on day 4 (Fig. 1 A and B). The O₂ uptake of 40% CO₂-stored mitochondria from June-harvested avocado fruit was 3-fold greater than the air control on treatment days 4 and 9 (Fig. 1A), whereas with August-harvested fruit, mitochondrial O₂ uptake was 2-fold greater than the air control on day 4 (Fig. 1B). After transfer to air, the mitochondrial O₂ uptake of 20% CO₂- (Fig. 1A) and 40% CO₂-treated (Fig. 1 A and B) avocados increased again. Mathooko et al. (1995) also found that elevated CO₂ concentrations increased the mitochondrial activity




Fig. 1. Changes in the O₂ uptake (protein mass basis) of mitochondria extracted from (A) June- and (B) August-harvested 'Hass' avocados that were kept at 10 °C in air (21% O₂), 20% CO₂ + 17% O₂, or 40% CO₂ + 13% O₂ before transfer to air.

of cucumber fruit exposed to 30% or 60% CO₂. Perez-Trejo et al. (1981) found that only 2 to 3 h of exposure to 10% to 30% CO₂ were required to cause a 6-fold rise in respiration of potato tubers. This increased whole potato tuber respiration may be a result of increased mitochondrial respiration. Lange and Kader (1997) found that 40% CO₂-stored avocado fruit had increased total fruit respiration, just as we observed in this fruit study with mitochondrial respiration.

Mitochondrial respiration of June-harvested, air-stored avocado fruit was relatively low and gradually declined, regardless of the pathway contributing to O₂ uptake (Fig. 2A). In 20% CO₂-stored avocado fruit, the primary terminal oxidase pathway was the cytochrome oxidase (Cyt) pathway on day 4. However, on day 9, the predominant terminal oxidase pathway was the alternative (Alt) pathway (Fig. 2B). Treatment of avocado fruit with 40% CO₂




Fig. 2. Oxygen uptake (protein mass basis) due to the activities of the cytochrome alternative and residual oxidase pathways of mitochondria extracted from June-harvested 'Hass' avocados that were stored at 10 °C in (A) air, (B) 20% CO₂, or (C) 40% CO₂ before transfer to air on day 12.

elicited a similar effect as that of 20% CO₂, but after transfer to air for 2 d, the Alt pathway became predominant again (Fig. 2C). The August-harvested avocados had a similar pattern of response due to terminal oxidase pathways with most of the differences occurring on day 4 (Fig. 3). The Cyt pathway was favored in air-stored fruit (Fig. 3A), whereas the Alt pathway was favored in avocados exposed to either CO₂ concentration (Fig. 3 B and C). Residual respiration (i.e., respiration in the presence of KCN and SHAM) usually accounts for at most 10% to 20% of the total O₂ uptake in fruit (Romani, personal communication). In our study, residual respiration was substantial (often >30%) and appeared to be stimulated by elevated CO₂ exposure (Figs. 2 and 3).

Laties (1982) reported that exposure of potato tubers to 10% CO₂ for 72 h yielded CN-resistant tissue as well as a significant rise

in residual respiration. Residual respiration, at least in part, may be due to monooxygenases, which probably contribute to the total O₂ consumed by plant tissues (Day et al., 1980). There also may be a contribution of α -oxidation to residual respiration.

The direct in vitro exposure to CO₂ caused partial inhibition of O₂ uptake of mitochondria extracted from 'Hass' avocados (Fig. 4). Within 2 h, the rates of O₂ uptake of mitochondria kept in air, 20% CO₂, or 40% CO₂ were all reduced by \approx 50%. Rates remained at a constant level in mitochondria kept in air, whereas the respiration rates of CO₂-treated mitochondria decreased to 20% of the initial rate after 20 h at 20 °C. Mathooko et al. (1995) also found that elevated CO₂ concentrations (up to 60%) inhibited the in vitro O₂ uptake rates of mitochondria from cucumber fruit, broccoli (*Brassica oleracea* var. *italica* Plenck) buds, and carrots (*Daucus carota* L.).

Most of the initial O₂ uptake of extracted avocado mitochondria was contributed by the Alt pathway (Fig. 5), but this decreased to 50%, 40%, and 20% of the initial activity after only 2 h in air, 20% CO₂, and 40% CO₂, respectively. For the rest of the experiment, the mitochondria kept in air maintained an Alt O₂ uptake that was 50% of the initial rate, whereas the CO₂-treated mitochondria had a continued decrease in O₂ uptake resulting from the Alt pathway. The relative contribution of the Cyt pathway was higher than the other pathways in the CO₂-treated mitochondria after 2 and 4 h at 20 °C (Figs. 5 B and C).

Since the CO₂ treatments included slightly lower levels of O₂ (17% for the 20% CO₂ treatment and 13% for the 40% CO₂ treatment) than the air treatment (21% O₂), partial inhibition of the Alt pathway by CO₂ may be due to the higher K_m of alternative oxidase for O₂, which is estimated to be 10-times higher than the K_m for cytochrome oxidase (Solomos, 1977). For example, Serald and Sisler (1972) observed a K_m of 11 to 14 μ M for AltOx compared with 1.2 to 1.4 μ M for CytOx. However, the levels of O₂ in the CO₂ treatments are too high to have a major effect on the affinities of these terminal oxidase pathways for O₂. Other direct effects of CO₂ on the mitochondria, such as the acidification of the intracellular spaces or structural changes in proteins or lipid membranes, may have played a role as well (Romo-Parada et al., 1991; Shipway and Bramlage, 1973). Moriguchi and Romani (1995) concluded that




Fig. 3. Oxygen uptake (protein mass basis) due to the activities of the cytochrome, alternative and residual oxidase pathways of mitochondria extracted from August-harvested 'Hass' avocados which were stored at 10 °C in (A) air, (B) 20% CO₂, or (C) 40% CO₂ before transfer to air on day 7.




Fig. 4. Changes in the O₂ uptake (protein mass basis) of isolated 'Hass' avocado fruit mitochondria treated at 20 °C with air (21% O₂), 20% CO₂ + 17% O₂, or 40% CO₂ + 13% O₂ for up to 20 h.

exposure of avocado fruit to CO₂-rich atmospheres enhanced the capacity of their mitochondria to restore energy-linked functions.

The respiratory control ratio (RCR) is defined as the state 3 O₂ uptake rate (in the presence of added ADP) divided by the state 4 O₂ uptake rate (where ADP supply has been depleted). RCR values for freshly isolated avocado mitochondria are in the range of 2 to 5 (Moreau and Romani, 1982a). The indirect effect of CO₂ on the RCR of treated avocado fruit was to maintain or to protect the tissue from a loss of respiratory control (Fig. 6). After transfer to air for 2 to 3 d, the RCR levels dropped to near 1, regardless of the previous treatment. The effect of CO₂ on the RCR levels of either June- or August-harvested 'Hass' avocado fruit was similar. Elevated CO₂ atmospheres have slowed senescence of a wide range of horticultural commodities (Kader, 1986; Wang, 1990). This slowing may be due partially to the protective role that CO₂ may play in maintaining membrane integrity (Day et al., 1978).

The RCR levels of *in vitro* CO₂-treated avocado mitochondria were slightly lower than mitochondria kept in air after 2 and 4 h of CO₂ exposure (Fig. 7). RCR dropped to 1 in all treatments, indicating the total loss of respiratory control, after 20 h. (RCR = 1). Although it was not a strong uncoupler, the direct effect of CO₂ appears to be through an uncoupling of oxidative phosphorylation, as was described previously by Shipway and Bramlage (1973).

Western blot (immunoblot) analysis, using a monoclonal antibody to the AltOx proteins, revealed the presence of three or more proteins in the 33- to 41-kDa range (Figs. 8–10). Elthon et al. (1989) reported the presence of alternative oxidase (AltOx) proteins in avocado fruit (range = 35 to 37 kDa). The Alt pathway is stimulated by ethylene produced during climacteric fruit ripening (Laties, 1982). Since partially ripe avocado fruit tissue that was producing large amounts of ethylene was used in these experiments, there was already an abundance of AltOx proteins on day 0 (Figs. 8 and 9) (Lange and Kader, 1996). On days 9 and 15 (labeled AT-Air) of air storage of June-harvested avocado fruit at 10 °C, the AltOx proteins accumulated to higher than initial levels (Fig. 8). Exposure to CO₂-enriched atmospheres decreased the amount of AltOx proteins until after transfer to air for 3 d at 10 °C, after which there was no effect of previous treatment with CO₂ on the abundance of AltOx proteins. In August-harvested avocado fruit stored at 10 °C, an accumulation of AltOx proteins only occurred in fruit stored in 40% CO₂ for 4 d (Fig. 9). Again, after transfer to air for 2 d, there were no differences in the levels of AltOx proteins. In a second experiment to determine the effects of all treatments on August-harvested avocados on day 4 (Fig. 10), the air-stored fruit had small amounts of AltOx proteins, the 20% CO₂-stored fruit had intermediate levels, and the 40% CO₂-stored fruit had a high abundance of AltOx proteins.

Expression of the Alt pathway in nonthermogenic tissues is often correlated with increased metabolic activity, such as that observed in fruit ripening or wound stress responses (Day et al., 1980; Laties, 1982; Palmer, 1976; Solomos, 1977). One reason for Alt pathway expression during fruit ripening or stress responses may be to allow for the continued generation of synthetic intermediates by the mitochondria when the energy charge is high (Lambers, 1982). In addition, expression of the Alt pathway allows for the recycling of mitochondrial matrix and cytoplasmic NADH by bypassing normal respiratory control. Possibly, the increased expression of the Alt pathway during elevated CO₂ exposure was due to a physiological response that triggered activation of the Alt pathway. During conditions of Cyt pathway inhibition (such as the use of stress levels of CO₂), the fermentative and Alt pathways are stimulated. Induction of the Alt pathway, compared to induction of the fermentative pathway, results in an 8-fold increase in ATP




Fig. 5. Changes in the O₂ uptake (protein mass basis) of the cytochrome, alternative, and residual oxidase pathways of 'Hass' avocado fruit mitochondria treated at 20 °C with air (21% O₂), 20% CO₂ + 17% O₂, or 40% CO₂ + 13% O₂ for up to 20 h.

production, less substrate consumption, less disturbance to normal metabolism by maintaining the Krebs cycle, and, therefore, greater tolerance of the commodity to high CO₂ stress. Recently, the importance of the Alt pathway *in vivo* has become more clear with the advent of transgenic tobacco plants that have reduced or overexpressed AltOx (Vanlerberghe et al., 1994). The antisense-AltOx tobacco suspension cells did not survive when they were grown under conditions that inhibited the Cyt pathway, while the wild type cells were able to grow due to the functioning of AltOx. Cells with overexpressed AltOx did not have increased partitioning to the Alt pathway, suggesting that this partitioning may be subject to additional regulatory factors *in vivo*, such as posttranslational modifications to AltOx protein that limits AltOx pathway activity.

Mitz (1979) suggested that the mechanism of CO₂ action in disrupting typical cellular function may be due to changes in protein configuration and membrane permeability. Taking into




Fig. 6. Changes in respiratory control ratio (RCR) of mitochondria extracted from (A) June-harvested and (B) August-harvested, partially ripe 'Hass' avocados that were kept at 10 °C in air (21%), 20% CO₂ + 17% O₂, or 40% CO₂ + O₂, and transferred to air.




Fig. 7. Changes in the respiratory control ratio (RCR) of isolated 'Hass' avocado fruit mitochondria treated at 20 °C with air (21% O₂), 20% CO₂ + 17% O₂, or 40% CO₂ + 13% O₂ for up to 20 h.

account these broad changes in plant cells, the use of stress levels of CO₂ to control disease, insects, or physiological disorders in plant tissues may be a feasible alternative to using chemicals, provided that specific plant tissues and treatment conditions (time, temperature, relative humidity, and O₂ and CO₂ concentrations) are thoroughly tested before applying these treatments.

Literature Cited

- Blake, M.S., K.H. Johnston, G.J. Russell-Jones, and E.C. Gotschlich. 1984. A rapid, sensitive method for detection of alkaline phosphatase-conjugated antibody on western blots. *Anal. Biochem.* 136:175-179.
- Bradford, M. 1976. A rapid and sensitive method for the quantitation of protein utilizing the principle of protein dye binding. *Anal. Biochem.* 72:248-254.
- Bruemmer, J.H. 1989. Terminal oxidase activity during ripening of Hamlin orange. *Phytochemistry* 8:2901-2902.
- Collier, D.E. and W.R. Cummins. 1991. Respiratory shifts in developing petals of *Saxifraga cernua*. *Plant Physiol.* 95:324-328.
- Day, D.A., G.P. Arron, R.E. Christoffersen, and G.G. Laties. 1978. Effect of ethylene and carbon dioxide on potato metabolism: Stimulation of tuber and mitochondrial respiration, and inducement of the alternative path. *Plant Physiol.* 62:820-825.
- Day, D.A., G.P. Arron, and G.G. Laties. 1980. Nature and control of respiratory pathways in plants: The interaction of cyanide-resistant respiration with the cyanide-sensitive pathway, p. 197-241. In: D.D. Davies (ed.). *Biochemistry of plants*, vol. 2. Academic, New York.
- Day, D.A. and J.B. Hanson. 1977. On methods for the isolation of mitochondria from etiolated corn shoots. *Plant Sci. Lett.* 11:99-104.
- Elthon, T.E. and L. McIntosh. 1987. Identification of the alternative terminal oxidase of higher plant mitochondria. *Proc. Natl. Acad. Sci. U.S.A.* 84:8399-8403.
- Elthon, T.E., R.L. Nickels, and L. McIntosh. 1989. Monoclonal antibodies to the alternative oxidase of higher plant mitochondria. *Plant Physiol.* 89:1311-1317.
- Estabrook, R.W. 1987. Mitochondrial respiratory control and the polarographic measurement of ADP:O ratios. *Methods Enzymol.* 10:41-47.
- Huang, L.S. and R.J. Romani. 1991. Metabolically driven self-restoration of energy-linked functions by avocado mitochondria. *Plant Physiol.* 95:1096-1105.
- Kader, A.A. 1986. Biochemical and physiological basis for effects of controlled and modified atmospheres on fruits and vegetables. *Food Technol.* 40:99-100, 102-104.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 227:680-685.
- Lambers, H. 1982. Cyanide-resistant respiration: a non-phosphorylating electron transport pathway acting as an energy overflow. *Physiol. Plant.* 55: 478-485.
- Lange, D.L. and A.A. Kader. 1997. Effects of elevated carbon dioxide on key mitochondrial respiratory enzymes in 'Hass' avocado fruit and fruit disks. *J. Amer. Soc. Hort. Sci.* 122:238-244.
- Laties, G.G. 1982. The cyanide-resistant, alternative path in higher plant respiration. *Annu. Rev. Plant Physiol.* 33:519-555.
- Laties, G.G. 1987. Ethylene and wound-induced cyanide-resistant respiration climaxes in potato: The synergistic role of CO₂ and the selective role of lycorine. *Physiol. Plant.* 69:305-312.
- Lurie, S. and J.D. Klein. 1989. Cyanide metabolism in relation to ethylene production and cyanide insensitive respiration in climacteric and non-climacteric fruits. *J. Plant Physiol.* 135:518-521.
- Mathooko, F.M., T. Fukuda, Y. Kubo, A. Inaba, and R. Nakamura. 1995. Regulation of mitochondrial activity in cucumber fruit, broccoli buds and carrot by carbon dioxide. *Acta Hort.* 398:71-79.
- McCaig, T.N. and R.D. Hill. 1977. Cyanide-insensitive respiration in wheat: cultivar differences and effects of temperature, carbon dioxide and oxygen. *Can. J. Bot.* 55:549-555.
- McIntosh, L. 1994. Molecular biology of the alternative oxidase. *Plant Physiol.* 105:781-786.
- Mitz, M.A. 1979. CO₂ biodynamics: A new concept of cellular control. *J. Theor. Biol.* 80:537-551.

Møller, I.M., A. Bérczi, L.W.H. van der Plas, and H. Lambers. 1988. Measurement of the activity of and capacity of the alternative pathway in intact plant tissue: Identification of problems and possible solutions. *Physiol. Plant.* 72:642-649.

Moreau, F. and R. Romani. 1982a. Preparation of avocado mitochondria using self-generated percoll density gradients and changes in buoyant density during ripening. *Plant Physiol.* 70:1380-1384.

Moreau, F. and R. Romani. 1982b. Malate oxidation and cyanide-insensitive respiration in avocado mitochondria during the climacteric cycle. *Plant Physiol.* 70:1385-1390.

Moriguchi, T. and R.J. Romani. 1995. Mitochondrial self-restoration as an index to the capacity of avocado fruit to sustain atmospheric stress at two climacteric states. *J. Amer. Soc. Hort. Sci.* 120:643-649.

Morohashi, Y., T. Seto, and H. Matsushima. 1991. Appearance of alternative respiration in cucumber cotyledon mitochondria after treatment with cycloheximide. *Physiol. Plant.* 83:640-646.

Palet, A., M. Ribas-Carbó, J.M. Argilés, and J. Azcón-Bieto. 1991. Short-term effects of carbon dioxide on carnation callus cell respiration. *Plant Physiol.* 96:467-472.

Palmer, J.M. 1976. The organization and regulation of electron transport in plant mitochondria. *Annu. Rev. Plant Physiol.* 27:133-157.

Perez-Trejo, M.S., H.W. Janes, and C. Frenkel. 1981. Mobilization of respiratory metabolism in potato tubers by carbon dioxide. *Plant Physiol.* 67:514-517.

Rhoads, D.M. and L. McIntosh. 1993. Cytochrome and alternative pathway respiration in tobacco: Effects of salicylic acid. *Plant Physiol.* 103:877-883.

Romo-Parada, L., L.-P. Vézina, P.M. Charest, F. Castaigne, and C. Willemot. 1991. Effect of modification of storage atmosphere on phospholipids and ultrastructure of cauliflower mitochondria. *Physiol. Plant.* 83:664-674.

Sherald, J.L. and H.D. Sisler. 1972. Selective inhibition of antimycin A-insensitive respiration in *Ustilago maydis* and *Ceratocystis ulmi*. *Plant Cell Physiol.* 13:1039-1052.

Shipway, M.R. and W.J. Bramlage. 1973. Effects of carbon dioxide on activity of apple mitochondria. *Plant Physiol.* 51:1095-1098.

Solomos, T. 1977. Cyanide-resistant respiration in higher plants. *Annu. Rev. Plant Physiol.* 28:279-297.

Solomos, T. and G.G. Laties. 1976. Effects of cyanide and ethylene on the respiration of cyanide-sensitive and cyanide-resistant plant tissues. *Plant Physiol.* 58:47-50.

Spalding, D.H. and W.F. Reeder. 1974. Low-oxygen, high-carbon diox-




Fig. 8. Immunoblot analysis of the alternative oxidase proteins (see arrows) from mitochondria of June-harvested 'Hass' avocado fruit which were kept at 10 °C in air, 20% CO₂, or 40% CO₂ for up to 12 d (day 0 and 9 shown only) and after transfer (AT) to air for 3 d. Purified mitochondrial protein (100 µg) were loaded into each sample lane.




Fig. 9. Immunoblot analysis of the alternative oxidase proteins (see arrows) from mitochondria of August-harvested 'Hass' avocado fruit that were kept at 10 °C in air, 20% CO₂, or 40% CO₂ for 7 d (day 0 and 4 shown only) and after transfer (AT) to air for 2 d. Purified mitochondrial protein (60 µg) were loaded into each sample lane.

ide controlled atmosphere storage for control of anthracnose and chilling injury of avocados. *Phytopathology* 65:458-460.

Stegink, S.J. and J.N. Siedow. 1986. Binding of butyl gallate to plant mitochondria. II. Relationship to the presence or absence of the alternative pathway. *Plant Physiol.* 62:232-237.

Truter, A.B. and G.J. Eksteen. 1987. Controlled and modified atmospheres to extend storage life of avocados. *South African Avocado Growers' Assn. Yrbk.* 10:151-153.

Vanlerberghe, G.C., A.E. Vanlerberghe, and L. McIntosh. 1994. Molecular genetic alteration of plant respiration: Silencing and overexpression of alternative oxidase in transgenic tobacco. *Plant Physiol.* 106:1503-1510.

Wang, C.Y. 1990. Physiological and biochemical effects of controlled atmosphere on fruits and vegetables, p. 197-223. In: M. Calderon and R. Barkai-Golan (eds.). *Food preservation by modified atmospheres*. CRC Press, Boca Raton, Fla.

Yoshimoto, A., S. Shigeru, N. Minagawa, and T. Komiyama. 1989. Possible role of a 36 kDa protein induced by respiratory inhibitors in cyanide-resistant respiration in *Hansenula anomala*. *J. Biochem.* 105:864-866.




Fig. 10. Immunoblot analysis of the alternative oxidase proteins (see arrows) from mitochondria of August-harvested 'Hass' avocado fruit that were kept at 10 °C in air, 20% CO₂, or 40% CO₂ for 4 d. One-hundred micrograms of purified mitochondrial protein was loaded into each lane.