

Changes in yield and seed quality traits in rapeseed genotypes by sulphur fertilization

Cem Ömer Egesel · M. Kemal Gül · Fatih Kahrıman

Published online: 5 July 2009
© Springer-Verlag 2009

Erratum to: Eur Food Res Technol (2009) 229:505–513
DOI 10.1007/s00217-009-1067-3

Unfortunately, the printed version of article contains Turkish words in the column 1 of Table 1. Corrected version of Table 1 is given below with English words.

Table 1 Soil characteristics for the experimental area

Parameter	2006	2007
pH	6.4	7.8
Lime (%)	2.6	18.6
Salt (%)	0.025	0.020
Sand (%)	44	54.1
Org matter (%)	2.1	1.4
Total N (%)	0.106	0.090
S (ppm)	55.6	12.74

The online version of the original article can be found under
doi:[10.1007/s00217-009-1067-3](https://doi.org/10.1007/s00217-009-1067-3).

C. Ö. Egesel (✉) · F. Kahrıman
Department of Field Crops,
Çanakkale Onsekiz Mart University,
17020 Çanakkale, Turkey
e-mail: cegesel@comu.edu.tr

M. K. Gül
COMPO GmbH Co. KG Turkey Liaison Office,
Büyükdere Street, No: 7 GIZ 2000 Plaza,
7th Floor, 34398 Maslak, Istanbul, Turkey