

G OPEN ACCESS

Citation: Mullan PC, Vazifedan T (2022) Changing temporal trends in patient volumes in a pediatric emergency department during a COVID-19 pandemic lockdown: A retrospective cohort study. PLoS ONE 17(9): e0271708. https://doi.org/ 10.1371/journal.pone.0271708

Editor: Yong-Hong Kuo, University of Hong Kong, HONG KONG

Received: July 5, 2022

Accepted: August 29, 2022

Published: September 12, 2022

Copyright: © 2022 Mullan, Vazifedan. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its <u>Supporting Information</u> files.

Funding: The authors received no specific funding for this work. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: No. The authors have declared that no competing interests exist.

RESEARCH ARTICLE

Changing temporal trends in patient volumes in a pediatric emergency department during a COVID-19 pandemic lockdown: A retrospective cohort study

Paul C. Mullan^{1,2}*, Turaj Vazifedan³

1 Division of Pediatric Emergency Medicine, Department of Pediatrics, Children's Hospital of the King's Daughters, Norfolk, Virginia, United States of America, 2 Eastern Virginia Medical School, Norfolk, Virginia, United States of America, 3 Department of Pediatrics, Children's Hospital of the King's Daughters, Norfolk, Virginia, United States of America

* paul.mullan@chkd.org

Abstract

Objective

Emergency department (ED) teams have had to adjust limited staffing resources to meet the fluctuating levels of patient volume and acuity during the COVID-19 pandemic. Historically, Mondays have had the highest reported ED volumes. We are unaware of any studies reporting on the change of this Monday effect during the COVID-19 pandemic.

Methods

This retrospective, observational study of a single pediatric ED compared a pandemic lockdown period (3/23/2020-11/1/2020) with a seasonally comparative period (3/25/2019-11/3/ 2019). We compared the mean number of patients who arrived on Monday versus any other specific weekday (Tuesday, Wednesday, Thursday, or Friday) and the aggregate of other weekdays (Tuesday to Friday) for both study periods. Secondary analyses investigated overall mean volumes, admission rates, and differences in triage acuity levels.

Results

There were 31,377 and 18,098 patients in the comparative and pandemic periods. The mean number of ED visits on Mondays in the comparative period was significantly more than any other weekday and the aggregate of weekdays (latter p<0.001). In contrast, there were no significant differences in the mean number of ED visits on Mondays in the pandemic period relative to any other weekday and the aggregate of weekdays (all p>0.05). The pandemic period had significantly lower mean volumes, higher admission rates, and more patients with higher acuity levels.

Conclusion

The previously experienced Monday effect of increased relative ED patient volumes was not seen during the pandemic period. This change has operational implications for scheduling

Abbreviations: ED, Emergency Department; ESI, Emergency Severity Index; GLM, Generalize Linear Model; SARS-CoV-2, Severe Acute Respiratory Syndrome Coronavirus. ED staffing resources. Larger database studies are needed to determine the generalizability of these findings.

Introduction

Understanding temporal trends in the volume of patients arriving at emergency departments (EDs) is critical in the operational planning of resources during this period of the coronavirus disease (COVID-19) pandemic as well as during non-pandemic periods. By predicting the anticipated ED staffing needs according to the hour of the day, day of the week, or month of the year, leadership can appropriately determine staff rosters, offer surge moonlighting opportunities, flex rooming areas, and plan financial budgets. Optimal planning of these resources can support ED care that is timely, efficient, safe, and effective. Prior studies have identified other temporal factors that were associated with changes in ED patient volumes including seasonality, day of the week, time of day, and holidays [1, 2].

The initial spread of the novel severe acute respiratory syndrome coronavirus (SARS-CoV-2) to the United States (U.S.) in early 2020, and the resulting pandemic that followed, led to lockdowns with the closure of schools and daycare facilities. These lockdowns were associated with significant decreases in patient volumes in U.S. adult and pediatric emergency departments [3–5]. In one U.S. based, multi-center ED study, this lockdown period was associated with significant ED revenue decreases that led to a 15% reduction in ED physician hours when compared to a pre-pandemic period [6, 7] Given the economic vulnerability of U.S. EDs, which are primarily reimbursed on a fee-for-service model, it is critical that ED staff scheduling can anticipate temporal changes in patient volumes.

Higher volumes of ED patients have been historically been reported to occur on Mondays [8–10]. Given this trend, we have historically scheduled additional physician and nurse staffing coverage on Mondays to manage the workload. This study aims to characterize the weekday variability in patient volume in a U.S. pediatric emergency department during a pandemic period and a comparative pre-pandemic period. We hypothesize that there was a significantly higher proportion of ED weekday patient volumes on Mondays in the pre-pandemic period and that this difference was no longer present during the pandemic period.

Materials and methods

Study design and setting

This study was a retrospective, observational cohort study of ED visits at a 31-bed freestanding, academic children's hospital. This hospital is the state's only dedicated Level-I pediatric trauma center and the only freestanding children's hospital in the state, with a catchment area population of approximately two million people. This report was written according to the Strengthening The Reporting of Observational Studies in Epidemiology guideline statement [11]. The research project was exempted from our institutional review board given that it was deemed not to be human subjects research (IRB #20-12-NH-0281).

Methods and measurements

The patients were identified with a patient report generated from our electronic health record. Each patient was designated as arriving on a particular day of the week based on the patient's ED arrival time.

The study period was evaluated based on historical events. The first reported SARS-CoV-2 infection in Virginia was on March 7, 2020 and the governor declared a state of emergency on March 12, 2020 to start public health interventions aimed to change behaviors and limit the infectious spread of SARS-CoV-2 [12]. The governor closed schools statewide on Monday, March 23, 2020 for the academic school year through to June 2020; the majority of schools in the region and state remained fully remote in the Fall of 2020 [13]. The first surge in COVID-19 cases in the region occurred in July and August 2020; the second surge started in November 2020. This study's pandemic period of analysis was a period of 32 weeks (3/23/2020-11/1/2020) from the initial school closures until the second surge when more in-person schooling started to become available. This pandemic period was compared with a seasonally comparative period in 2019 (3/25/2019-11/3/2019). All ED patients from both periods were eligible for inclusion.

Outcomes and analyses

The primary outcome was to determine if there was a difference in the proportion of weekday patients who arrived on a Monday versus another weekday in each study period. For the primary outcome, Generalized Linear Model (GLM) was conducted for comparing the mean number of all ED visits on Mondays versus the mean number of all ED visits on each of other weekdays (that is, Monday versus Tuesday; Monday versus Wednesday; Monday versus Thursday; Monday versus Friday) and in aggregate (that is, Monday versus Tuesday-through-Friday) during the two study periods in 2019 and 2020. A box plot was created to graphically display all ED visit numbers in each period.

Secondary analyses used GLM for comparing the Emergency Severity Index (ESI) triage level proportions between the 2019 and 2020 periods; any patients without an assigned ESI level were excluded from this analysis. Continuous variables were presented as mean and Standard Deviation (SD). Categorical variables were presented as frequency and percentage. A Chi-square test was used for comparing the percentage of ED visits that arrived on a weekday versus a weekend day of the week between 2019 and 2020. All statistical tests were performed using SPSS 26 (Chicago, IL). All statistical tests were two-sided, and p < 0.05 was considered statistically significant.

Results

There were 31,377 and 18,098 patients included in the 2019 comparative period and 2020 pandemic period, respectively (Table 1 and S1 File). In the 2020 pandemic period, the mean number of patients per week significantly declined versus the 2019 comparative period. The 2020 pandemic period also had a significantly higher admission rate. There were no differences in the percentage of patients who arrived on weekdays (Monday to Friday) in the 2019 comparative period (73.4%) and 2020 (73.5%) pandemic period. There were also no differences in the percentage of patients who arrived on either of the weekend days in the 2019 comparative period and 2020 pandemic period.

For the primary outcome, the mean number of ED visits on Monday was significantly more than any other weekday in the 2019 comparative period or in aggregate (Tuesday to Friday; p<0.001; Table 2). In contrast, there was not any significant difference in the mean number of ED visits on Monday versus any other weekday during the 2020 pandemic period or in aggregate (p = 0.22). The box plot of the number of visits on each weekday is displayed graphically (Fig 1).

For the secondary analyses, 80 patients from the study periods did not have a triage acuity assigned and the proportion of these unassigned patients did not vary between periods. There

	2019 (3/25/2019-11/3/2019) n = 31,377	2020 P (3/23/2020-11/1/2020) n = 18,098	Difference (95% Confidence Interval)	P value
Mean Number of patients per week (standard deviation)	980.5	565.6	414.9	< 0.001
Number of admissions (%)	3,805 (12.13%)	2,867 (15.84%)	3.71% (3.07, 4.36)	< 0.001
Weekday arrivals (%)	23,042 (73.4%)	13,297 (73.5%)	0.1% (-0.01, 0.01)	0.93
Weekend arrivals (%)	8,335 (26.6%)	4,801 (26.5)	0.1% (-0.01, 0.01)	0.93
Number (%) of Monday arrivals	5,126 (16.34%)	2,791 (15.42%)	0.92% (0.25, 1.58)	0.007
Number (%) of Tuesday arrivals	4,699 (14.98%)	2,665 (14.73%)	0.25% (-0.40, 0.9)	0.45
Number (%) of Wednesday arrivals	4,577 (14.59%)	2,637 (14.57%)	0.02% (-0.63, 0.66)	0.96
Number (%) of Thursday arrivals	4,407 (14.05%)	2,614 (14.44%)	0.4% (-1.04, 0.24)	0.22
Number (%) of Friday arrivals	4,233 (13.49%)	2,590 (14.31%)	0.82% (-1.46, -0.19)	0.01
Number (%) of Saturday arrivals	4,055 (12.92%)	2,362 (13.05%)	0.13% (-0.74, 0.49)	0.68
Number (%) of Sunday arrivals	4,280 (13.64%)	2,439 (13.48%)	0.16% (-0.46, 0.79)	0.61
Number (%) without a triage acuity level assigned	55 (0.16%)	25 (0.14%)	0.03% (-0.04, 0.10)	0.45
Triage Emergency Severity Index Level 1	260 (0.83%)	217 (1.20%)	0.37% (0.18, 0.56)	< 0.001
Triage Emergency Severity Index Level 2	2981 (9.50%)	2370 (13.10%)	3.59% (3.00, 4.19)	< 0.001
Triage Emergency Severity Index Level 3	5277 (16.82%)	5255 (29.04%)	12.22% (11.43, 13.0)	< 0.001
Triage Emergency Severity Index Level 4	18844 (60.06%)	9299 (51.38%)	8.68% (-9.59, -7.76)	< 0.001
Triage Emergency Severity Index Level 5	3960 (12.62%)	932 (5.15%)	7.47% (-7.96, -6.98)	< 0.001
Triage Emergency Severity Index Level Not documented	55 (0.18%)	25 (0.14%)	0.037% (-0.11, 0.04)	0.38

Table 1. Description of Emergency Department (ED) patients in the 2019 comparative and 2020 pandemic period.

https://doi.org/10.1371/journal.pone.0271708.t001

was a significant difference in the distribution of ESI per day between the ED visits in each study period, such that the likelihood of a greater proportion of high (that is, sicker) acuity level patients was higher in 2020 than in 2019; this significant difference (p<0.001) was present on the weekly levels (that is, all 7 days combined; p<0.001) and at the level of each individual day of the week (p<0.001) (Fig 2).

Year	Weekday Period	Mean (SD) number of all ED visits	Estimate of Difference	95% CI	р
2019	Monday ^a	160.2 (25.5)	-	-	-
	Tuesday	146.8 (26.5)	-13.3	(-23.7, -3.0)	0.011
	Wednesday	143.0 (25.7)	-17.2	(-27.5, -6.8)	0.001
	Thursday	137.7 (24.7)	-22.5	(-32.8, -12.2)	< 0.001
	Friday	132.3 (21.5)	-27.9	(-38.2, -17.6)	< 0.001
	Tuesday to Friday	140.0 (25.0)	-20.2	(-28.5, -11.9)	< 0.001
2020	Monday ^a	87.2 (16.8)	-	-	-
	Tuesday	83.3 (17.6)	-3.9	(-14.2, 6.4)	0.45
	Wednesday	82.4 (19.2)	-4.8	(-15.1, 5.5)	0.36
	Thursday	81.7 (16.9)	-5.5	(-15.8, 4.8)	0.29
	Friday	80.9 (15.6)	-6.3	(-16.6, 4.0)	0.23
	Tuesday to Friday	82.1 (17.2)	-5.1	(-13.4, 3.1)	0.22

Table 2. Mean number of all Emergency Department (ED) visits by weekday in both study periods.

^aMonday as reference Level

Abbreviations: CI (Confidence Interval), SD (Standard Deviation)

https://doi.org/10.1371/journal.pone.0271708.t002





Fig 1. Box plot of the number of visits by day of the week during the pandemic period (3/23/2020-11/1/2020) and comparative period (3/25/2019-11/3/2019).


https://doi.org/10.1371/journal.pone.0271708.g001

Discussion

In this single-center study, the historical increase in the relative number of ED patient visits on Mondays was not seen during our pandemic lockdown period. The overall number of ED visits was dramatically decreased in this pandemic period compared to a seasonally comparative period in the prior year. The 14% relative increase in the proportion of weekday visits on Mondays that we reported during the 2019 comparative period, relative to other weekdays (i.e., Tuesday through Friday), was comparable to the 12% to 17% relative increases reported in other ED studies [8–10]. We are unaware of any other ED studies that have reported this disappearance of the historical Monday effect associated with a pandemic lockdown period. One United Kingdom database study has reported that there is no longer an increase in the number of ED fracture visits presenting on weekends during the pandemic period as compared to a pre-pandemic period [14]. We described no change in the relative percentage of weekend ED arrivals between the two periods, but future studies could investigate for differences in the frequency of specific patient diagnoses that present on weekends during the pandemic and prepandemic periods.

During the pandemic, the greatest decrease in pediatric ED visits has been reported in communicable conditions (76% reduction) while non-infectious diagnoses have seen a smaller reduction of only 36% [15]. Given that 28% of U.S. pediatric emergency department diagnoses are related to infectious conditions, this dramatic decrease in overall ED volume is expected with children having fewer opportunities to spread contagious infections to one another while out of school and daycare during the periods of lockdown [16]. The specific decrease in the proportion of Monday ED visits, versus other weekdays, might be due to various factors and is potentially linked to the start of the workweek, as an Israeli ED study has reported Sunday (i.e.,





https://doi.org/10.1371/journal.pone.0271708.g002

the first day of their traditional workweek) as the busiest day of ED visits [2]. One possibility is that part of the Monday increase has historically been attributed to the return of families from a weekend trip away who then seek care the following day [17]. With the lockdown period and its associated reductions in traveling, this effect might have been dampened [18]. Another contributor to the Monday effect has been attributed to the increased number of Monday referrals from primary care providers who return to the office after the weekend off. The significant decrease in primary care visits during the pandemic might have diminished this phenomenon as well [19]. Additionally, the pandemic's shift to the virtual workplace for many workers and the muddled schedules of each workday in the week might have also contributed to the decrease in this Monday effect. More workers are working from home or after-hours to afford flexibility in balancing their workload demands at-home virtual schooling, job site closures, and other pandemic-related challenges [20]. The increase in the availability and access to virtual healthcare visits during the pandemic, as a more convenient source of care for caregivers, might have also contributed to the decrease in the Monday effect for ED visits.

Whether the explanation for this vanishing Monday effect is due to one or multiple factors, the trend has implications for appropriately scheduling ED staffing resources optimally, a goal with heightened importance in the face of pandemic-related staffing shortages [21]. The

appropriate balancing of workloads across shifts is one strategy that might be effective in addressing the emotional burnout of healthcare providers in the pandemic [6]. With optimal staffing, both the mental health of workers as well as the financial health of an ED can be supported [7]. In our own ED, we typically scheduled an additional ED physician shift on Monday (as well as Tuesday) evenings, in addition to the other 35 weekly shifts. These two shifts, early on in the week, represent an additional 5.7% of scheduled weekly physicians hours, which might be necessary during traditional periods but might be avoidable in the future if a similar, extended lockdown period occurs.

The absolute number of ED patients is only part of the equation when determining the staffing needs for an ED. In this study, we also saw a significant increase in the triage acuity of patients and an increase in our admission rates. Caring for higher acuity patients is associated with longer ED length of stay for each patient and can require more staffing per patient to match the increased resource utilization for this sicker population. Similar increases in acuity and admission rates have been described in other ED settings across the globe [15, 22]. This higher proportion of high acuity patients has contributed to a decline in the performance of many traditional ED throughput measures of quality including longer length of stay, longer time to perform lab and radiological studies, and delayed time to antibiotic administration [23].

This study had several limitations. First, as a single-center, pediatric ED, our findings might not be generalizable to other settings. Second, these findings might not be applicable to future pandemics or future surges within the current pandemic, as the various factors (e.g. the degree of school, daycare, and work closures) that impact this decrease might be present to differing degrees. Lastly, there were some missing data on the triage acuity level assignments; however this was a relatively small (<1%) proportion, and the proportion did not change between the two periods.

Conclusions

In conclusion, our ED visit volume did not show a statistically significant increase in the proportion of Monday visits, compared to the other weekdays, during the lockdown period of the COVID-19 pandemic. The absence of a significant increase in the proportion of weekday ED visits on Monday was a change compared to the pre-pandemic period. In this single center study, this change only represented a 0.92% decrease in the proportion of Monday visits in the pandemic period. Future studies are needed to determine if this observed decrease in the Monday effect is present more widely using national database studies. If we can increase our understanding of the impact of societal factors, such as pandemic lockdown periods, on ED volumes, then we can continue to plan the judicious use of our limited ED resources [7].

Supporting information

S1 Checklist. (DOCX)

S1 File. Dataset. (XLSX)

Acknowledgments

The authors thank Dr. Faiqa Qureshi, MD for her review and feedback on the manuscript.

Author Contributions

Conceptualization: Paul C. Mullan.

Data curation: Paul C. Mullan.

Formal analysis: Paul C. Mullan, Turaj Vazifedan.

Investigation: Paul C. Mullan, Turaj Vazifedan.

Methodology: Paul C. Mullan, Turaj Vazifedan.

Project administration: Paul C. Mullan.

Resources: Paul C. Mullan.

Supervision: Paul C. Mullan.

Validation: Paul C. Mullan.

Visualization: Paul C. Mullan, Turaj Vazifedan.

Writing – original draft: Paul C. Mullan, Turaj Vazifedan.

Writing – review & editing: Paul C. Mullan, Turaj Vazifedan.

References

- Sun Y, Heng BH, Seow YT, Seow E. Forecasting daily attendances at an emergency department to aid resource planning. *BMC Emerg Med.* 2009; 9:1. <u>https://doi.org/10.1186/1471-227X-9-1</u> PMID: 19178716
- Rotstein Z, Wilf-Miron R, Lavi B, Shahar A, Gabbay U, Noy S. The dynamics of patient visits to a public hospital ED: a statistical model. *Am J Emerg Med.* 1997; 15(6):596–599. https://doi.org/10.1016/s0735-6757(97)90166-2 PMID: 9337370
- Haddadin Z, Blozinski A, Fernandez K, et al. Changes in Pediatric Emergency Department Visits During the COVID-19 Pandemic. *Hosp Pediatr.* 2021; 11(4):e57–e60. <u>https://doi.org/10.1542/hpeds.2020-005074 PMID</u>: 33436415
- Pines JM, Zocchi MS, Black BS, et al. Characterizing pediatric emergency department visits during the COVID-19 pandemic. *Am J Emerg Med.* 2021; 41:201–204. https://doi.org/10.1016/j.ajem.2020.11.037 PMID: 33257144
- DeLaroche AM, Rodean J, Aronson PL, et al. Pediatric Emergency Department Visits at US Children's Hospitals During the COVID-19 Pandemic. *Pediatrics*. 2021; 147(4):e2020039628.
- Sharifi M, Asadi-Pooya AA, Mousavi-Roknabadi RS. Burnout among Healthcare Providers of COVID-19; a Systematic Review of Epidemiology and Recommendations. *Arch Acad Emerg Med.* 2021; 9(1): e7. https://doi.org/10.22037/aaem.v9i1.1004 PMID: 33490964
- Pines JM, Zocchi MS, Black BS, et al. The Effect of the COVID-19 Pandemic on the Economics of United States Emergency Care. Ann Emerg Med. 2021; 78(4):487–499. https://doi.org/10.1016/j. annemergmed.2021.04.026 PMID: 34120751
- Erkamp NS, van Dalen DH, de Vries E. Predicting emergency department visits in a large teaching hospital. *Int J Emerg Med.* 2021; 14(1):34. https://doi.org/10.1186/s12245-021-00357-6 PMID: 34118866
- Castner J, Yin Y, Loomis D, Hewner S. Medical Mondays: ED Utilization for Medicaid Recipients Depends on the Day of the Week, Season, and Holidays. *J Emerg Nurs.* 2016; 42(4):317–324. https:// doi.org/10.1016/j.jen.2015.12.010 PMID: 26972368
- McHale P, Wood S, Hughes K, Bellis MA, Demnitz U, Wyke S. Who uses emergency departments inappropriately and when—a national cross-sectional study using a monitoring data system. *BMC Med.* 2013; 11(1):258.
- Vandenbroucke JP, von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Int J Surg.* 2014; 12(12):1500–1524. https:// doi.org/10.1016/j.ijsu.2014.07.014 PMID: 25046751
- 12. Documenting Virginia's path to recovery from the coronavirus (COVID-19) pandemic, 2020–2021—Ballotpedia. Accessed January 20, 2022. https://ballotpedia.org/Documenting_Virginia%27s_path_to_recovery_from_the_coronavirus_(COVID-19)_pandemic, 2020-2021

- 13. COVID-19 pandemic in Virginia. In: *Wikipedia*.; 2022. Accessed January 20, 2022. <u>https://en.wikipedia.</u> org/w/index.php?title=COVID-19_pandemic_in_Virginia&oldid=1064752543
- Ferraro CF, Findlater L, Morbey R, et al. Describing the indirect impact of COVID-19 on healthcare utilisation using syndromic surveillance systems. *BMC Public Health*. 2021; 21:2019. https://doi.org/10. 1186/s12889-021-12117-5 PMID: 34740346
- Kruizinga MD, Peeters D, van Veen M, et al. The impact of lockdown on pediatric ED visits and hospital admissions during the COVID19 pandemic: a multicenter analysis and review of the literature. *Eur J Pediatr.* 2021; 180(7):2271–2279. https://doi.org/10.1007/s00431-021-04015-0 PMID: 33723971
- Hasegawa K, Tsugawa Y, Cohen A, Camargo CA. Infectious Disease-related Emergency Department Visits Among Children in the US. *Pediatr Infect Dis J.* 2015; 34(7):681–685. https://doi.org/10.1097/INF. 0000000000000704 PMID: 25860534
- Wargon M, Guidet B, Hoang TD, Hejblum G. A systematic review of models for forecasting the number of emergency department visits. *Emerg Med J.* 2009; 26(6):395–399. <u>https://doi.org/10.1136/emj.2008.</u> 062380 PMID: 19465606
- Abdullah M, Dias C, Muley D, Shahin Md. Exploring the impacts of COVID-19 on travel behavior and mode preferences. *Transportation Research Interdisciplinary Perspectives*. 2020; 8:100255. https:// doi.org/10.1016/j.trip.2020.100255 PMID: 34173481
- Chatterji P, Li Y. Effects of the COVID-19 Pandemic on Outpatient Providers in the United States. Med Care. 2021; 59(1):58–61. https://doi.org/10.1097/MLR.00000000001448 PMID: 33136711
- Dunatchik A, Gerson K, Glass J, Jacobs JA, Stritzel H. Gender, Parenting, and The Rise of Remote Work During the Pandemic: Implications for Domestic Inequality in the United States. *Gender & Society*. 2021; 35(2):194–205.
- CDC. Healthcare Workers. Centers for Disease Control and Prevention. Published February 11, 2020. Accessed January 20, 2022. <u>https://www.cdc.gov/coronavirus/2019-ncov/hcp/mitigating-staff-shortages.html</u>
- Finkelstein Y, Maguire B, Zemek R, et al. Effect of the COVID-19 Pandemic on Patient Volumes, Acuity, and Outcomes in Pediatric Emergency Departments: A Nationwide Study. *Pediatr Emerg Care*. 2021; 37(8):427–434. https://doi.org/10.1097/PEC.00000000002484 PMID: 34074990
- Bae SJ, Chung HS, Namgung M, Choi YH, Min JH, Lee DH. Comparison of the Clinical Process and Outcomes in Patients after Coronavirus Infection 2019 Outbreak. *Medicina (Kaunas)*. 2021; 57 (10):1086. https://doi.org/10.3390/medicina57101086 PMID: 34684122