



Characteristics and sources of submicron aerosols above the urban canopy (260 m) in Beijing, China, during the 2014 APEC summit

C. Chen^{1,2,3}, Y. L. Sun^{1,2}, W. Q. Xu¹, W. Du^{1,4}, L. B. Zhou¹, T. T. Han¹, Q. Q. Wang¹, P. Q. Fu¹, Z. F. Wang¹, Z. Q. Gao^{1,2}, Q. Zhang⁵, and D. R. Worsnop⁶

¹State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

²Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science & Technology, Nanjing, China

³College of Applied Meteorology, Nanjing University of Information Science and Technology, Nanjing, China

⁴Department of Resources and Environment, Air Environmental Modeling and Pollution Controlling Key Laboratory of Sichuan Higher Education Institutes, Chengdu University of Information Technology, Chengdu, China

⁵Department of Environmental Toxicology, University of California, 1 Shields Ave., Davis, CA, USA

⁶Aerodyne Research, Inc., Billerica, MA, USA

Correspondence to: Y. L. Sun (sunyele@mail.iap.ac.cn)

Received: 3 July 2015 – Published in Atmos. Chem. Phys. Discuss.: 26 August 2015

Revised: 10 November 2015 – Accepted: 11 November 2015 – Published: 19 November 2015

Abstract. The megacity of Beijing has experienced frequent severe fine particle pollution during the last decade. Although the sources and formation mechanisms of aerosol particles have been extensively investigated on the basis of ground measurements, real-time characterization of aerosol particle composition and sources above the urban canopy in Beijing is rare. In this study, we conducted real-time measurements of non-refractory submicron aerosol (NR-PM₁) composition at 260 m at the Beijing 325 m meteorological tower (BMT) from 10 October to 12 November 2014, by using an aerosol chemical speciation monitor (ACSM) along with synchronous measurements of size-resolved NR-PM₁ composition near ground level using a high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS). The NR-PM₁ composition above the urban canopy was dominated by organics (46 %), followed by nitrate (27 %) and sulfate (13 %). The high contribution of nitrate and high NO₃⁻ / SO₄²⁻ mass ratios illustrates an important role of nitrate in particulate matter (PM) pollution during the study period. The organic aerosol (OA) was mainly composed of secondary OA (SOA), accounting for 61 % on an average. Different from that measured at the ground site, primary OA (POA) correlated moderately with SOA, likely suggesting a high contribution from regional transport above the urban

canopy. The Asia–Pacific Economic Cooperation (APEC) summit with strict emission controls provides a unique opportunity to study the impacts of emission controls on aerosol chemistry. All aerosol species were shown to have significant decreases of 40–80 % during APEC from those measured before APEC, suggesting that emission controls over regional scales substantially reduced PM levels. However, the bulk aerosol composition was relatively similar before and during APEC as a result of synergetic controls of aerosol precursors. In addition to emission controls, the routine circulations of mountain–valley breezes were also found to play an important role in alleviating PM levels and achieving the “APEC blue” effect. The evolution of vertical differences between 260 m and the ground level was also investigated. Our results show complex vertical differences during the formation and evolution of severe haze episodes that are closely related to aerosol sources and boundary-layer dynamics.

1 Introduction

Beijing (39°56' N, 116°20' E), the capital of China, is one of the largest megacities in the world with more than 21 million residents and 5.4 million vehicles in operation by the

end of 2013 (Beijing Municipal Bureau of Statistics, 2014). In the west, north, and northeast, the city is surrounded by the Taihang and Yanshan mountains at approximately 1000–1500 m above sea level. The fan-shaped topography in addition to the rapid urbanization has caused frequent severe haze pollution episodes in Beijing. These conditions have received a significant amount of attention from atmospheric scientists, the government, and the general public (Sun et al., 2006, 2012a, 2013b, 2014; Guo et al., 2014). For in-depth elucidation of severe urban haze formation and particulate matter (PM) characteristics, extensive studies have been conducted in Beijing including real-time online measurements and filter sampling with subsequent offline analyses (Sun et al., 2006; Pope III. et al., 2009; Zhao et al., 2013). Aerosol mass spectrometers (AMS), which are capable of determining size-resolved aerosol compositions with high sensitivity, have been widely deployed in Beijing and other cities in China since 2006 (Huang, X.-F. et al., 2012; Zhang et al., 2014; Li et al., 2015). Numerous conclusions and findings have been obtained since then, which have greatly improved our understanding of aerosol composition, formation mechanisms, and evolution processes (Sun et al., 2010; Xiao et al., 2011; Zhang et al., 2012, 2014; Hu et al., 2013; Huang et al., 2013; Guo et al., 2014; Li et al., 2015). However, most previous AMS studies include short-term measurements of generally less than 2 months, because of the high cost and maintenance of the instrument. The recently developed Aerodyne Aerosol Chemical Speciation Monitor (ACSM) (Ng et al., 2011) has been used in some studies for examining the chemical composition, sources, and processes of atmospheric aerosols in China. The advantage of the ACSM is its robustness for real-time long-term measurements of aerosol particle composition with little attendance (Ng et al., 2011; Sun et al., 2012a, 2013b, 2014; Budisulistiorini et al., 2014; Jiang et al., 2015; Parworth et al., 2015; Petit et al., 2015). The first ACSM measurements in Beijing highlighted the important role of nitrate in PM pollution in summer, which was mainly attributed to the partitioning of nitric acid into liquid ammonium nitrate particles (Sun et al., 2012a). The PM pollution characteristics also dramatically differed between summer and winter. Agricultural burning and photochemical production play major roles in PM pollution in summer (Li et al., 2010; Huang, K. et al., 2012; Sun et al., 2012a; Zhang et al., 2015), whereas coal combustion is the dominant source of PM in winter (Sun et al., 2013b). A more detailed analysis of a severe haze pollution episode that occurred in January 2013 suggested that stagnant meteorological conditions, source emissions, secondary production, and regional transport are four major factors driving the formation and evolution of haze pollution in Beijing during winter (Sun et al., 2013b, 2014; Guo et al., 2014; Zhang et al., 2014).

Despite extensive efforts for the characterization of fine particle pollution in Beijing, most studies are conducted at ground sites, which are subject to significant influences of local emission sources such as traffic, cooking, and biomass

burning. In comparison, measurements obtained above the urban canopy with much less influence of local source are more representative for a large scale, which is of great importance for characterizing regional transport. However, such studies in Beijing are rare due to the absence of high platforms. The Beijing 325 m meteorological tower (BMT) is a unique platform for measuring aerosol and gaseous species at various heights in Beijing megacity. Moreover, this platform is beneficial for studying the interactions of the lower boundary layer (< 300 m) and air pollution, particularly during autumn and winter when the nocturnal planetary boundary height is often below 300 m (Ting et al., 2008; Zhang et al., 2013). Based on the BMT measurements, Sun et al. (2009, 2013) reported that the SO₂ concentration reached its maximal value at 50 m during heating periods, whereas PM_{2.5} showed a “higher top and lower bottom” vertical pattern due to the inversions of temperature (*T*) and relative humidity (RH) during summer hazy days. Guinot et al. (2006) and Meng et al. (2008) also determined that local concentration peaks at 50 to 100 m were likely related to the urban canopy. However, real-time characterization of aerosol particle composition above the urban canopy has been performed only once (Sun et al., 2015). The 2-week study found substantially different aerosol compositions between ground level and 260 m. In addition, the compositional differences at the two heights were found to be strongly associated with source emissions, the vertical mixing mechanism, and RH/*T*-dependent secondary production. Because these measurements only lasted 2 weeks, the aerosol characteristics and sources above the urban canopy remain poorly understood.

The 2014 Asia–Pacific Economic Cooperation (APEC) summit was hosted in Beijing during 5–11 November 2014, when strict emission control measures were implemented in Beijing and surrounding regions to ensure good air quality. During 3–12 November emission controls such as reducing the number of vehicles in operation by approximately 50 %, shutting down factories, stopping construction activities, and enhancing the cleanliness of urban roads were gradually implemented (http://www.gov.cn/xinwen/2014-11/14/content_2778635.htm, in Chinese). The neighboring provinces such as Hebei, Tianjin, and Shandong implemented the same emission controls during APEC (<http://www.bjepb.gov.cn/bjepb/324122/412670/index.html>, in Chinese). As a result, the PM levels in Beijing during the summit were significantly reduced, leading to “APEC blue”, a phrase commonly used to refer to the good air quality. However, the response of aerosol chemistry to emission controls over a regional scale has not been investigated. Measurements above the urban canopy are ideal for evaluating the roles of emission controls in reducing PM levels under the condition of minimizing the influences of local point sources.

In this study, we conduct real-time measurements of non-refractory submicron aerosol (NR-PM₁) composition including organics (Org), sulfate (SO₄²⁻), nitrate (NO₃⁻), ammo-

nium (NH_4^+), and chloride (Cl^-) at 260 m at the BMT before and during APEC, 10 October–2 November and 3–12 November 2014, respectively, by using an ACSM. The aerosol composition, diurnal variation, and sources above the urban canopy are investigated in detail. The responses of aerosol composition, particle acidity, and sources of organic aerosol (OA) to emission controls are elucidated by comparing the changes before and during APEC, and the roles of meteorological conditions in PM reduction during APEC are discussed. In addition, the vertical differences of aerosol composition and its interactions with boundary-layer dynamics are also examined.

2 Experimental methods

2.1 Sampling site and measurements

All of the measurements in this study were conducted at the same site as that reported by Sun et al. (2013b), which is an urban site at the Institute of Atmospheric Physics, Chinese Academy of Sciences, between North 3rd and 4th Ring Road from 10 October to 12 November 2014. The ACSM and gas measurement instruments were mounted inside a container at 260 m on the BMT. The ACSM sampling setup used in this study is similar to that described by Sun et al. (2012a). Briefly, aerosol particles were first sampled into the container with a $\text{PM}_{2.5}$ cyclone to remove coarse particles larger than $2.5\ \mu\text{m}$. After passing through a diffusion silica-gel dryer, aerosol particles were sampled into the ACSM at a flow rate of $\sim 0.1\ \text{L}\ \text{min}^{-1}$. The ACSM was operated by alternating ambient air and filtered air with a mass spectrometer at a scanning rate of $500\ \text{ms}\ \text{amu}^{-1}$ from m/z 10 to 150. The data were saved every two cycles, leading to a time resolution of approximately 5 min. The detailed principles of the ACSM can be found elsewhere (Ng et al., 2011; Sun et al., 2012a). An Aerodyne high-resolution time-of-flight AMS (HR-ToF-AMS) was simultaneously deployed near ground level at the same location to measure the size-resolved NR- PM_{10} aerosol composition. Details of the sampling and operation procedures of the HR-ToF-AMS are given in Xu et al. (2015).

Meteorological variables including wind speed (WS), wind direction (WD), RH, and T at 15 heights of 8, 15, 32, 47, 65, 100, 120, 140, 160, 180, 200, 280, and 320 m were obtained from the BMT. In addition, a Doppler wind lidar (Windcube 200, Leosphere, Orsay, France) was deployed at the same location to obtain the wind profiles from 100 to 5000 m with a spatial resolution of 50 m and a time resolution of 10 min. All of the data in this study are reported in Beijing Standard Time (BST), which is equal to Coordinated Universal Time (UTC) plus 8 h.

2.2 Data analysis

The ACSM data were analyzed for the mass concentration and chemical composition of NR- PM_{10} species including or-

ganics, sulfate, nitrate, ammonium, and chloride by using ACSM standard data analysis software (v. 1.5.3.0). Detailed analytical procedures have been reported by Ng et al. (2011) and Sun et al. (2012a). Similar to that of previous studies in Beijing (Sun et al., 2011, 2012a, 2013b, 2014), an empirical and constant collection efficiency (CE) of 0.5 was applied during the entire campaign to compensate for the particle loss due mainly to particle bounce at the vaporizer (Matthew et al., 2008). The CE of 0.5 is rational for this study because aerosol particles were dried, and the mass fraction of ammonium nitrate was overall below the threshold value (40 %) that affects CE (Middlebrook et al., 2012). The average ratio of measured NH_4^+ ($\text{NH}_{4\ \text{meas}}^+$) versus predicted NH_4^+ ($\text{NH}_{4\ \text{pred}}^+$) was 0.56, suggesting that the aerosol particles were acidic. Although the particle acidity would have a slightly higher CE than 0.5 (~ 0.59) if the equation $\text{CE}_{\text{dry}} = \max(0.45, 1.0 - 0.73 \times (\text{NH}_{4\ \text{meas}}^+ / \text{NH}_{4\ \text{pred}}^+))$ recommended by Middlebrook et al. (2012) were used, no effect on CE is present if using the parameterization reported by Quinn et al. (2006). For consistency with our previous studies and with the HR-ToF-AMS measurements at the ground site, we maintained $\text{CE} = 0.5$ in this study. The default relative ionization efficiency (RIE) values were 1.4 for organics, 1.1 for nitrate, 1.2 for sulfate, and 1.3 for chloride, except ammonium (6.5) which was determined from pure ammonium nitrate particles. Note that the ACSM measurements were compared with those of HR-AMS at the same location before the campaign. All submicron aerosol species measured by the ACSM were highly correlated with those by the HR-AMS ($r^2 > 0.97$). Although the total NR- PM_{10} mass measured by the ACSM agreed well with that by HR-AMS ($r^2 = 0.99$, slope = 0.99), the regression slopes of ACSM against HR-AMS varied from 0.61 to 1.24 for different aerosol species. Because ACSM was found to have a larger uncertainty in the quantification of submicron aerosol species, particularly in the determination of relative ionization efficiency, the mass concentrations of aerosol species measured by the ACSM at 260 m were further corrected using the regression slopes of ACSM/HR-AMS obtained from the intercomparison study.

Positive matrix factorization (PMF) with the PMF2.exe algorithm (Paatero and Tapper, 1994) was performed on the ACSM OA mass spectra to resolve potential OA components with different sources and processes. Only m/z s < 125 were included in the PMF analysis due to the large interferences of naphthalene signals on several larger m/z s (e.g., m/z 127–129) (Sun et al., 2012a, 2013b, 2014). The PMF results were then evaluated using an Igor Pro-based PMF Evaluation Tool (PET, v. 2.06) (Ulbrich et al., 2009) following procedures detailed by Zhang et al. (2011). After careful evaluation of the mass spectra and time series of OA factors, a 2-factor solution, i.e., an oxygenated OA (OOA) and a hydrocarbon-like OA (HOA) with $f_{\text{peak}} = 0.4$, was chosen. More detailed PMF diagnostics are presented in Figs. S1, S2, and Table S1 in the Supplement. While the 3-factor solution resolved an




Figure 1. Evolution of vertical profiles of (a) wind speed (WS) and (b) wind direction (WD) from the measurements of the Doppler wind lidar. The time series of NR-PM₁ (i.e., Org + SO₄²⁻ + NO₃⁻ + NH₄⁺ + Cl⁻) is shown as the black line in (a). The shaded area refers to the APEC period (same for following figures).

unrealistic factor with unexpectedly high m/z 12 and m/z 15, the 2-factor solution at $f_{\text{peak}} = 0$ showed much higher m/z 44 in the HOA spectrum, which is generally a characteristic of OOA (Fig. S3).

2.3 Air mass trajectory analyses

The 3-day (72 h) back trajectories were calculated every hour at 500 m height using the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT, NOAA) 4.9 model (Draxler and Hess, 1997; Li et al., 2015). The trajectories were then grouped into four clusters before and during APEC using the algorithm of cluster analysis. The clustering of trajectories is based on the total spatial variance (TSV) method (Draxler et al., 2012). This method minimizes the inter-cluster differences among trajectories while maximizing the inter-cluster differences, which has been widely used in previous studies (Sun et al., 2014; Zhang et al., 2014; Li et al., 2015).

3 Results and discussion

3.1 General description

3.1.1 Submicron aerosol and meteorology

The NR-PM₁ mass concentration varied significantly from 0.7 to 254 $\mu\text{g m}^{-3}$, with an average of 53.5 $\mu\text{g m}^{-3}$. As indicated in Fig. 1, the variations of NR-PM₁ were strongly associated with WD and WS. The formation of severe haze episodes was generally initiated by a WD change from northerly to southerly and a decrease of WS to less than 5 m s^{-1} below 1 km. The southern air flow and low WS were then dominant most of the time during the evolution of the haze episode; subsequently, the air masses changed from the south to the north/northwest, leading to a rapid decrease of

PM level in a few hours. Haze episodes with such life cycle driven by meteorological conditions have also been observed many times in Beijing (Jia et al., 2008; Sun et al., 2013b, 2014; Guo et al., 2014). Note that a mountain–valley breeze lasting approximately half a day was frequently observed throughout the study, which reduced the daytime PM levels to a certain degree. As shown in Fig. 1, most of the cleaning processes were similar, all driven by the switch of air masses from south/southwest to north/northwest associated with high WS across the entire vertical layer ($> 5 \text{ m s}^{-1}$). However, the cleaning process that occurred on 20–21 October was different. As the WD changed from the south to the northwest/northeast, the NR-PM₁ concentration remained high. This phenomenon can be explained by the low WS ($< 4 \text{ m s}^{-1}$) below 500 m and the high RH (Figs. 2, S4). The NR-PM₁ began to decrease at $\sim 20:00$ as WD shifted to the south associated with a decrease in RH. This result indicates that a cleaner and drier air mass was located to the south of Beijing during this stage. Such a cleaning process by southern air flow is not common and is generally weaker than that by northern/northwestern flow. This observation is supported by the higher NR-PM₁ concentration of $\sim 20 \mu\text{g m}^{-3}$ on 21 October compared to during other cleaning periods at $\sim < 5 \mu\text{g m}^{-3}$. The average mass concentration of NR-PM₁ during APEC was 24.1 $\mu\text{g m}^{-3}$, which is significantly lower than the 65.1 $\mu\text{g m}^{-3}$ recorded before APEC, indicating a large reduction of PM during APEC. In addition, the southern air mass occurred less frequently and had a shorter duration during APEC. These results manifest that meteorology, in addition to emission controls, might have played an important role in reducing PM levels during APEC.

The NR-PM₁ species showed similar and dramatic variations to the total NR-PM₁ mass (Fig. 2). In particular, three haze episodes before APEC (Ep1, Ep2, and Ep3 in Fig. 2d) and two episodes during the summit (APEC1 and APEC2 in Fig. 2d) were observed in this study. The three episodes before APEC were all characterized by high RH at 48–70 % and low WS at 2.3–3.4 m s^{-1} , elucidating the important roles of stagnant meteorological conditions in severe haze formation. In comparison, the RH in the two episodes during APEC was lower at 34–38 %, and the WS was comparably higher at 3.1–3.8 m s^{-1} (Table 1). These results suggest that the meteorological conditions during APEC appeared to be more favorable for the dispersion of pollutants. Indeed, clear accumulation processes of aerosol species were observed for three episodes before APEC, yet they were much weaker during the summit. However, the two episodes during APEC showed obvious temperature inversions, which inhibited the vertical convection of pollutants. The meteorological conditions during haze episodes differed substantially from those during clean periods, which were characterized by high WS at $> 5 \text{ m s}^{-1}$ and low RH at $< 20 \%$.

The NR-PM₁ was dominated by organics, accounting for on average 46 % of the total mass, followed by nitrate at 27 %, sulfate at 13 %, ammonium at 9 %, and chloride at




Figure 2. Time series of (a) T , (b) RH, (c) WS and WD, and (d) concentrations of NR-PM₁ species (Org, SO₄²⁻, NO₃⁻, NH₄⁺, and Cl⁻), and (e) mass fraction of each species in NR-PM₁. Two clean periods and five haze episodes are marked in (d) for further discussions. The meteorological parameters in this figure were all from the tower measurements.

Table 1. Summary of average meteorological variables for different periods and the mass differences of aerosol species between the ground site and 260 m (i.e., ground – 260 m).

	Before APEC				During APEC		
	Entire	Ep1	Ep2	Ep3	Entire	APEC1	APEC2
Meteorological variables							
RH (%)	47.1	48.4	69.7	56.7	29.8	34.2	38.5
T (°C)	13.3	16.7	12.5	10.9	9.0	11.5	8.1
WS (m s ⁻¹)	4.0	3.4	2.3	2.3	4.9	3.8	3.1
Mass differences (µg m ⁻³)							
Org	0.7	0.3	4.5	-5.2	9.6	14.6	13.6
SO ₄ ²⁻	3.4	3.0	8.8	1.3	1.3	1.6	1.9
NO ₃ ⁻	4.3	4.5	10.9	0.8	0.7	1.0	1.0
NH ₄ ⁺	3.9	4.2	9.0	2.3	1.6	2.9	2.3
Cl ⁻	-0.1	0.0	-0.4	-0.2	1.0	1.7	1.5
NR-PM ₁	12.1	12.0	32.8	-1.1	14.1	21.8	20.2

5%. The nitrate contribution ranged from 27 to 28% during the three episodes before APEC and from 29 to 31% in the two episodes during APEC, which is significantly higher than the sulfate contribution of 10–15 and 8–11%, respectively (Fig. 6). Although the dominance of organics in PM₁ was consistent with that in previous studies in Beijing (Sun et al., 2012a, 2013b, 2014; Guo et al., 2014; Zhang et al., 2014), the nitrate contribution in this study was approximately twice that of sulfate and significantly higher than previously reported values of 16% in 2011 (Sun et al., 2013b) and 13–14% in 2013 (Sun et al., 2014; Zhang et al., 2014). The mass ratio of NO₃⁻ / SO₄²⁻ can be used to indicate the relative im-

portance of mobile and stationary sources (Arimoto et al., 1996). Therefore, higher NO₃⁻ / SO₄²⁻ in this study likely indicates the predominance of mobile sources rather than stationary sources. Because of the continuous increase of NO_x emissions associated with a decrease in SO₂ (Wang et al., 2013), nitrate is expected to play a more important role in PM pollution in the future. Our results highlight that NO_x emission control should be a priority in mitigating air pollution, particularly in non-heating seasons with low SO₂ precursors.

Figure 3 further shows the time series of NO₃⁻ / SO₄²⁻ mass ratio and sulfur oxidation ratio (SOR) calculated as the molar fraction of sulfate in total sulfur (i.e., sulfate and




Figure 3. Time series of (a) sulfur oxidation ratio (SOR), (b) ratio of $\text{NO}_3^-/\text{SO}_4^{2-}$, and (c) NR-PM₁. The SOR and $\text{NO}_3^-/\text{SO}_4^{2-}$ were color-coded by RH.

SO_2) (Sun et al., 2014). The $\text{NO}_3^-/\text{SO}_4^{2-}$ was ubiquitously greater than 1 during five haze episodes, indicating the importance of nitrate in the formation of severe haze pollution. Interestingly, we observed a rapid increase in $\text{NO}_3^-/\text{SO}_4^{2-}$ during the formation stage of a pollution episode, followed by a decrease in $\text{NO}_3^-/\text{SO}_4^{2-}$ during the subsequent evolution stage. The variations of $\text{NO}_3^-/\text{SO}_4^{2-}$ illustrates that two different formation mechanisms might drive the formation and evolution of haze episodes. During the early stage of haze formation, the RH was relatively low and the formation rate of sulfate was correspondingly low, which is supported by the low SOR values. Consequently, the nitrate formation played a dominant role during this stage. The SO_4^{2-} concentration remained consistently low when the nitrate began to increase (Fig. 2d). As the RH continued to increase, the SOR showed a corresponding increase, indicating that more SO_2 was oxidized to form sulfate, most likely via aqueous-phase processing (Zhang and Tie, 2011; Sun et al., 2013a). The SO_4^{2-} concentration then showed a substantial increase, and the $\text{NO}_3^-/\text{SO}_4^{2-}$ ratio decreased as a result. For example, during Ep2, the hourly $\text{NO}_3^-/\text{SO}_4^{2-}$ increased from ~ 1.1 to 4.0 during the formation stage and then decreased to ~ 1.8 during the evolution stage. These results indicate that SO_4^{2-} played an enhanced role in PM pollution during the evolution stage of haze episodes with high RH. Moreover, the $\text{NO}_3^-/\text{SO}_4^{2-}$ ratios during clean periods (~ 0.3) were much lower than those during haze episodes. One explanation is that the nitrate in clean air masses from the north/northwest is significantly lower than that of sulfate.

3.1.2 Sources and composition of OA

Two OA factors, HOA and OOA, were identified in this study. The HOA spectrum was similar to those determined at other urban sites (Huang, K. et al., 2012; Sun et al., 2012a, b), which is characterized by prominent hydrocarbon ion peaks of m/z 27, 29, 41, 43, 55, and 57 (Fig. 4a). The HOA spectrum showed a higher m/z 55/57 ratio compared with that of exhaust aerosols from diesel trucks and gasoline vehicles (Mohr et al., 2009), yet it had characteristics similar to those resolved in urban Beijing (Sun et al., 2010, 2012a). The high m/z 55/57 ratio and the two visible peaks at meal times in diurnal variations (Fig. 4b) indicate the impact of local cooking activities (Sun et al., 2011, 2012a, 2013b). However, the two HOA peaks were much smaller than those observed at the ground site (Xu et al., 2015), indicating a significantly smaller impact of local cooking emissions on OA at 260 m. Moreover, the HOA spectrum showed a considerable m/z 60 peak, a marker m/z for biomass burning (Aiken et al., 2009; Huang et al., 2011; Zhang et al., 2015). The fraction of m/z 60 was 0.9%, which is much higher than $\sim 0.3\%$ in the absence of biomass burning. All these results suggest that HOA was a primary OA factor combined with traffic, cooking, and biomass burning emissions. Limited by the ACSM spectra and PMF analysis, we were not able to separate the different primary OA factors in this study. HOA correlated well with chloride ($r^2 = 0.61$) and moderately well with secondary inorganic species ($r^2 = 0.42\text{--}0.65$), indicating that a major fraction of HOA shared similar sources to secondary species at 260 m, which likely came from regional transport. HOA on average contributed 39% of total organics, which is less than the 57% observed at the ground site during the same study period (Xu et al., 2015). This result indicates a




Figure 4. (a) Mass spectra of HOA and OOA, (b) diurnal variations of the mass concentration and mass fraction of HOA and OOA, (c) time series of HOA, OOA, and inorganic species (SO_4^{2-} , NO_3^- , Cl^-). The correlations of HOA and OOA with inorganic species are also shown in the figure.

smaller impact of primary sources above the urban canopy. The diurnal cycle of HOA was relatively flat, with two visible peaks occurring at noon and night. The HOA contribution to OA was relatively constant throughout the day, ranging from 36 to 43 %. This result further supports the theory that HOA above the urban canopy came dominantly from regional transport and was well mixed with regional secondary OA (SOA). Indeed, the correlation of HOA with OOA in this study was quite high ($r^2 = 0.76$), supporting that HOA and OOA might have some common sources (e.g., regional transport) at 260 m.

The mass spectrum of OOA resembles that identified in 2012 in summer in Beijing (Sun et al., 2012a) in addition to the spectra resolved at other urban sites (Ulbrich et al., 2009), in that it is characterized by a prominent m/z 44 peak (mainly CO_2^+). OOA dominated the OA composition throughout the day, ranging from 57 to 64 %. The average OOA contribution to OA was 61 %, which is close to that previously reported in Beijing (Huang et al., 2010; Sun et al., 2012a, 2013b). The diurnal cycle of OOA was relatively flat, yet a gradual increase during the day was also observed despite the rising planetary boundary layer, suggesting daytime photochemical processing. OOA is often considered as a good surrogate of SOA (Zhang et al., 2005; Jimenez et al., 2009; Ng et al.,

2011). In this study, OOA correlated well with secondary inorganic species such as NO_3^- and SO_4^{2-} ($r^2 = 0.72$ – 0.90), which is consistent with previous conclusions that OOA is a secondary species in nature (Zhang et al., 2005; Sun et al., 2012a).

3.2 Response of aerosol chemistry to emission controls

3.2.1 Aerosol composition

Figure 5 shows the variations of aerosol composition as a function of NR-PM₁ mass loading before and during APEC. The organics contribution showed a notable decrease from 62 to 32 % as the NR-PM₁ mass concentration increased from < 10 to $> 200 \mu\text{g m}^{-3}$ before APEC. In contrast, the sulfate contribution showed a corresponding increase from 8 to 22 %. Except for low values at NR-PM₁ $< 10 \mu\text{g m}^{-3}$, nitrate and ammonium constituted relatively constant fractions of NR-PM₁ across different NR-PM₁ loadings and varied at 21–31 and 8–12 %, respectively. These results highlighted the enhanced roles of secondary inorganic species in severe PM pollution before APEC. This observation is further supported by a comparison of average chemical composition between three pollution episodes and a clean event (Fig. 6). The secondary inorganic aerosol (SIA; $\text{SO}_4^{2-} + \text{NO}_3^- + \text{NH}_4^+$)




Figure 5. Submicron aerosol composition as a function of NR-PM₁ mass loadings (a) before APEC and (b) during APEC. The solid line shows the probability of NR-PM₁ mass.

on average contributed 46–51 % of the total NR-PM₁ mass during the three episodes before APEC, which is significantly higher than the 40 % reported during the clean event (Fig. 6). The NR-PM₁ mass-loading-dependent aerosol composition showed different behavior during APEC. As shown in Fig. 5b, all aerosol species had relatively constant contributions to NR-PM₁ at 10–100 µg m⁻³. The contribution of organics ranged from 43 to 58 %, which is higher overall than before APEC. This result indicates an enhanced role of organics during APEC, particularly during severe PM pollution periods. Similarly, nitrate contributed the largest fraction of NR-PM₁, varying from 23 to 32 %. Figure 5 also shows a very broad range of NR-PM₁ mass concentration with the maximum concentration over 200 µg m⁻³ before APEC. In contrast, the range of NR-PM₁ was much narrower during APEC, suggesting a significantly lower amount of severe haze pollution during APEC. Indeed, 93 % of the time during APEC, the NR-PM₁ level was lower than 60 µg m⁻³, whereas 49 % of the time before APEC, it exceeded such a concentration level. These results indicate that the air pollution was substantially more severe before APEC. The average mass concentration of NR-PM₁ was 24.1 µg m⁻³ during

APEC, which is 63 % lower than the 65.1 µg m⁻³ recorded before APEC (Fig. 6). This result demonstrates a significant reduction of PM during APEC due to emission controls and better weather conditions, including higher WS and lower RH. However, the bulk NR-PM₁ composition was rather similar before and during APEC, both of which were dominated by organics, 46 % versus 47 %, followed by nitrate at 27 % versus 29 %, and sulfate at 14 % versus 10 % (Fig. 6). The lower sulfate contribution during APEC might be due to the lower RH associated with lower liquid water content, leading to less production of sulfate. These results highlight that the emission controls during APEC did not significantly affect the regional aerosol bulk composition, although the mass concentrations of precursors and aerosol species were reduced substantially. One possible explanation is the synergetic control of various precursors such as SO₂, NO_x, and volatile organic compounds (VOCs) over a regional scale during APEC. Our results clearly imply that synergetic controls of the emissions of precursors over a regional scale are efficient for mitigating air pollution in north China.

3.2.2 Diurnal variations

The diurnal variations of meteorological variables, NR-PM₁ species, and OA components before and during APEC are presented in Fig. 7. The diurnal cycles of meteorological conditions were overall similar before and during APEC except for lower temperatures and RH during APEC. The WS during APEC was consistently higher than that before APEC, particularly in the morning (04:00–12:00) and evening (18:00–22:00). Although the WD during APEC came dominantly from the northwest at night and shifted to the south during the day, it came mainly from the south before APEC (Fig. 2c).

The total NR-PM₁ showed pronounced diurnal variation with two peaks in the early afternoon (12:00–14:00) and late evening (20:00–22:00) that were dominantly influenced by organics. By checking the diurnal cycles of the OA factors, we concluded that the two peaks occurring at meal times are mainly attributed to primary emissions such as cooking-related activities and traffic emissions (Allan et al., 2010; Sun et al., 2011, 2012a). Compared with the diurnal cycles of OA previously observed at the ground site in Beijing (Sun et al., 2012a), the two peaks of organics were considerably smaller. This result indicates that local source emissions can be vertically mixed above the urban canopy but at substantially reduced concentrations. Our results also demonstrate that sampling above the urban canopy is less influenced by local source emissions and can be more representative over a regional scale.

SIA and OOA showed similar diurnal patterns before and during APEC, all of which were characterized by gradual increases during the day. These results indicate that their diurnal cycles were driven by similar formation mechanisms before and during APEC, such as photochemical process-




Figure 6. Average chemical composition of NR-PM₁ before and during APEC, and also that of five haze episodes and two clean events marked in Fig. 2.




Figure 7. Diurnal variations of meteorological variables (T , RH , WS , and WD), NR-PM₁ species, and OA factors before and during APEC. The change rates during APEC (i.e., $(\text{before APEC} - \text{APEC}) / \text{before APEC} \times 100$) are also marked as light gray in the figure.

ing and daytime vertical mixing. Higher concentrations of secondary species were also observed at night, which might have been associated with a more shallow boundary-layer height (Sun et al., 2012a). It should be noted that all secondary species showed relatively constant background concentrations, indicating that a major fraction was likely from regional transport. SIA and OOA during APEC showed substantial reductions (45–74 %) throughout the day compared with those before APEC, indicating that regional emission controls played a significant role in reducing secondary species during APEC, although the lower RH and higher WS were also important. Moreover, a higher reduction percent-

age was observed between 04:00 and 12:00, when higher mountain–valley breezes occurring routinely during APEC cleaned the air pollutants more efficiently.

The diurnal cycles of chloride showed some differences before and during APEC. Although it was relatively flat during APEC, chloride showed a clear decrease in the afternoon before APEC, likely due to the evaporative loss and dilution effects associated with higher T and the elevated boundary layer (Sun et al., 2012a). The diurnal cycle of HOA showed a lower overall concentration during the day except for a pronounced noon peak before and during APEC. Considering that the peak time corresponds to lunchtime, we concluded




Figure 8. Variations of NR-PM₁ species and OA factors as a function of (a) RH and (b) WS before and during APEC. The RH and WS were from the tower measurements at 280 m.

that it was attributed mainly to local cooking sources. In addition, a more significant reduction in the evening peak of HOA was observed during APEC. One explanation is that controls of heavy-duty vehicles (HDV) and heavy-duty diesel trucks (HDDT) decreased the HOA emissions at night during APEC.

3.2.3 Meteorological effects

Meteorological parameters contribute the largest uncertainties to the evaluation of the effects of emission controls on PM reduction. Here we compared the variations of aerosol species as a function of RH and WS before and during APEC (Fig. 8). At low RH levels (< 40%), all aerosol species appeared to increase linearly as a function of RH in both periods at similar rates of increase. Moreover, the mass concentrations of aerosol species were slightly lower during APEC than before the summit, indicating small reductions in aerosol species during APEC. By checking the air mass trajectories (Fig. S5), we determined that the low RH periods were mainly associated with the air masses from the north/northwest where fewer emission controls were implemented during APEC. This finding explains the small reductions in aerosol species ($\sim 22\%$) during APEC under the same RH conditions. However, the variations in aerosol

species showed substantially different behaviors as a function of RH at high RH levels (> 40%) before and during APEC. Whereas most aerosol species continued to linearly increase as a function of RH before APEC, they remained relatively constant and even showed decreases during APEC. As a result, significant reductions in aerosol species at high RH levels were observed during APEC. The air masses during high RH periods were found to be dominantly from the south/southeast where strict emission controls were implemented such as in Hebei, Tianjin, and Shandong provinces. These results clearly indicate that emission controls played a major role in PM reduction during APEC and that the control effects tended to be more efficient under higher RH periods. The primary HOA and chloride showed decreases when the RH was > 60%, indicating that humidity has a significantly lower impact on primary aerosols than secondary components at high RH levels.

The mass concentrations of aerosol species showed a strong dependence on WS before and during APEC. For example, the total NR-PM₁ mass was decreased by $\sim 80\%$ from ~ 100 to $< 20 \mu\text{g m}^{-3}$ as WS increased to 7 m s^{-1} before APEC. These results indicate that wind is efficient in cleaning air pollutants in Beijing, which is consistent with previous conclusions (Han et al., 2009; Sun et al., 2013b). In comparison, the decreasing rates of aerosol species as a func-




Figure 9. Wind rose plots (a) before APEC and (b) during APEC.

tion of WS were lower during APEC. As a result, aerosol species showed the largest concentration differences before and during APEC in periods with low WS. As indicated by the wind increase plots in Fig. 9, low and high WS levels were mainly associated with southern/southeastern and northern/northwestern winds, respectively. These results further indicate that larger reductions of aerosol species occurred in Beijing when air masses were from the south.

3.2.4 Back trajectory analysis

Figure 10 presents the average chemical composition of NR-PM₁, corresponding to four clusters before and during APEC, determined from the cluster analysis of back trajectories (Draxler and Hess, 1997). The air masses before APEC were predominantly from the south/southeast at 54 % of the time (C1 in Fig. 10a), and the aerosol loading was the highest (96.7 $\mu\text{g m}^{-3}$) among the clusters. Comparatively, the northwesterly clusters (C3 and C4 in Fig. 10a) presented significantly lower aerosol loadings at 8.3 and 3.5 $\mu\text{g m}^{-3}$, respectively, with fewer frequencies of 14 and 11 %, respectively. Such large differences in aerosol loadings between the northerly and southerly air masses are consistent with the spatial distributions of anthropogenic emissions such as SO₂, NO_x, and BC (Zhang et al., 2007; Lu et al., 2011). Although the areas to the north/northwest of Beijing are relatively clean with low emissions of anthropogenic primary pollutants, the south/southeast regions are characterized by substantially higher emissions. In addition, 21 % of the air masses originated from the west and showed moderately high NR-PM₁ mass at 55.4 $\mu\text{g m}^{-3}$. It should be noted that the air masses from the south were often stagnant, as indicated by their shorter trajectories that played an important role in facilitating the accumulation of pollutants. The aerosol composition varied significantly among four clusters, reflecting the variety in chemical characteristics of aerosol particles from different source regions. The aerosol particle composition from the southeastern and western clusters (C1 and C2) were dominated by nitrate at 27 and 30 % and by OOA at 26 and 32 %, respectively, with considerable contribution from sulfate at 14 and 10 %, respectively. These results elucidate the dominant roles of nitrate and OOA in severe PM pollu-




Figure 10. The average NR-PM₁ composition for each cluster (a) before and (b) during APEC. The numbers on the pie charts refer to the average total NR-PM₁ mass for each cluster. In addition, the number of trajectories and corresponding percentages of the total trajectories are also shown in the legends.

tion before APEC, which differs significantly from previous studies that reported that sulfate was generally more prevalent than nitrate (Huang et al., 2014; Sun et al., 2014). These results also highlight very different pollution characteristics during the late fall season compared to winter. In comparison, the nitrate contributions were significantly lower, at 17 and 8 %, in the two northwestern clusters (C3 and C4), associated with an enhanced contribution of sulfate at 19 and 21 %, respectively. Moreover, the cleanest cluster (C4) showed a dominant contribution of organics at 64 %, indicating the important role of organics during clean periods (Sun et al., 2010, 2013b).

The air masses during APEC showed changes, particularly the increases in frequency of two northwestern clusters




Figure 11. Comparisons of time series of total NR-PM₁ mass and NR-PM₁ species between 260 m and ground level.

(C1 and C4), which was 40 % of the time compared with 25 % before APEC (Fig. 10b). These two clusters showed similar bulk aerosol compositions to those before APEC yet with reductions of the total NR-PM₁ mass loading at nearly 40–50 %. The air masses during APEC were dominated by cluster 3 (C3 in Fig. 10b). Although C3 originated from the north of Beijing, it circulated around the south of Beijing including Baoding, a polluted city in the Hebei province, before arriving at the sampling site. As a result, C3 presented the highest aerosol mass loading, at $44.0 \mu\text{g m}^{-3}$, composed primarily of nitrate and OOA at 30 and 29 %, respectively. Moreover, cluster 2 (C2 in Fig. 10b), originating from the northwest, showed a similar aerosol composition, yet had a ~ 50 % decrease in total mass compared to C3. One explanation is that air masses in C2 passed through western Beijing, which is relatively clean compared to the southeastern regions. As shown in Fig. 10, similar clusters before and during APEC showed ubiquitous reductions in NR-PM₁ mass during APEC, indicating that emission controls played an important role in PM reduction. Moreover, the decreases in frequency of southern/southeastern air masses during APEC also helped to alleviate the PM level for the entire period, thus achieving the “APEC blue” effect. Emission controls in surrounding regions south of Beijing should be taken as a priority for the mitigation of air pollution in Beijing.

3.3 Vertical differences: insights into emission controls and boundary-layer dynamics

Figure 11 shows a comparison of the time series of NR-PM₁ species between 260 m and the ground level for the entire study. All submicron species showed overall similar variations at the two different heights, indicating their relatively similar sources and evolution processes. However, large vertical differences in aerosol composition were also frequently observed, illustrating complex vertical gradients of aerosol species caused by multiple factors such as local emissions, regional transport, and boundary-layer dynamics. The average compositional differences before and during APEC are shown in Fig. 12. Although the concentration difference in NR-PM₁ was similar before and during APEC, at 12.1 and $14.1 \mu\text{g m}^{-3}$, respectively, the composition differed significantly. SIA dominated the compositional difference before APEC, together accounting for 95 % of the total NR-PM₁ mass. In comparison, organics and chloride showed minor vertical differences (< 5 %). These results indicate different sources and formation mechanisms between SIA and organic aerosol. During APEC, the compositional difference was dominated by organics, accounting for 68 % on average, and the contribution of SIA was largely reduced to 25 %. These results suggest that emission controls over regional scales affect the composition differences between ground level and the urban canopy. As discussed in Sect. 3.2 and by Xu et al. (2015), secondary species including SIA and SOA showed significant reductions at both ground level and 260 m dur-




Figure 12. Average chemical composition of the difference between ground level and 260 m (a) before APEC and (b) during APEC. The “1 %” in the box indicates a lower concentration of chloride at the ground site compared to at 260 m.

ing APEC as a result of emission controls. Although primary OA showed similar reductions as those of SOA above the urban canopy, the changes remained small near ground level. Thus, the largest organic difference during APEC was mainly caused by local primary source emissions.

The vertical differences in aerosol composition also varied largely among different haze episodes. As indicated in Fig. 11 and Table 1, Ep3 presented the smallest vertical differences for all aerosol species, indicating a well-mixed layer below 260 m. The WS was consistently low at $< 2.5 \text{ m s}^{-1}$ across the different heights, and the WD was predominantly from the south during Ep3. Moreover, the vertical profiles of extinction showed an evident reduction in pollution from $\sim 2 \text{ km}$ to the ground on 28 October, leading to the formation of Ep3 (Fig. S6). Such boundary-layer dynamics would produce a well-mixed layer in the lower atmosphere, leading to minor chemical differences between the ground level and 260 m.

Comparatively, the vertical evolution of Ep2 differed significantly (Fig. 13a). The mass concentrations of all aerosol species between the ground level and 260 m were similar during the formation stage of Ep2, from 23 October to 09:00 24 October. However, although aerosol species near ground level showed large increases after 09:00 on 24 October, they remained relatively constant at 260 m, leading to the largest vertical concentration gradients among five episodes. The average NR-PM₁ at 260 m was $143.4 \mu\text{g m}^{-3}$, which is 38 % lower than that at the ground site. By checking the vertical profiles of meteorological variables, we observed a clear temperature inversion between 120 and 160 m that formed during 00:00–09:00 on 24 October. Such a temperature inversion formed a stable layer below $\sim 200 \text{ m}$ and inhibited the vertical mixing of air pollutants between the ground and 260 m. In addition, the stagnant meteorological conditions as indicated by low WS and high RH further facilitated the accumulation of ground pollution. It should be noted that the aqueous-phase processing, most likely fog processing under the high RH conditions (often $> 90 \%$) during this stage, also played an important role in the increase of SIA, particularly




Figure 13. Evolution of vertical profiles of meteorological variables (WD, WS, RH, and T) and NR-PM₁ concentration at 260 m and at the ground site during two pollution episodes (a) Ep2 and (b) APEC2. The vertical profiles of wind speed and wind direction were from the measurements of the Doppler wind lidar, and those of RH and T were from the tower measurements. The white areas in the figure indicate that the data were not available.

sulfate. This finding is also supported by the significant increase of SOR during this stage (Fig. 3).

The evolution of the severe Ep2 was terminated at approximately 00:00 on 26 October when the WD changed from south to northwest. Although the mass concentrations of aerosol species at 260 m began to show rapid decreases at that time, the concentration at the ground site decreased significantly after 4 h. The different cleaning processes between 260 m and the ground level are closely linked to the vertical profiles of meteorological variables. As indicated in Fig. 13a, a strong temperature inversion below 320 m was observed during the cleaning period, which resulted in a significantly higher WS and lower RH at 260 m than those at

ground level. Indeed, both WS and RH showed clear shears during the cleaning period, suggesting a gradual interaction between the northern air mass and boundary pollution from top to bottom. Such an interacting mechanism resulted in a time lag of approximately 4 h in cleaning the pollutants at ground level compared to that at 260 m. Similar interactions between boundary-layer dynamics and aerosol pollution were also observed on 1, 5, and 11 November.

The evolution of vertical differences during APEC differed from those in three episodes before APEC. As shown in Fig. 13b, frequent mountain–valley breezes were observed during 8–11 November (APEC2). The northwest mountain–valley breeze began routinely at approximately midnight and dissipated at approximately noon. The NR-PM₁ aerosol species showed direct responses to the mountain–valley breeze, which was characterized by similar routine diurnal cycles. All aerosol species began to decrease at midnight because the cleaning effects of mountain–valley breeze reached minimum concentrations at noon, then increased continuously when the WD changed to south. The mountain–valley breeze also caused a unique diurnal cycle of vertical differences. As shown in Fig. 13b, aerosol species were well mixed within the lower boundary layer between 12:00 and 16:00, and the concentrations between 260 m and the ground level were similar. However, the differences in concentration began to increase when the boundary-layer height decreased after sunset at ~18:00, and the differences were maximum at midnight when the NR-PM₁ mass approached 100 μg m⁻³. A detailed investigation of the evolution of aerosol species showed that such vertical differences in NR-PM₁ were caused mainly by organics from local primary sources (Xu et al., 2015; Fig. 11). These results indicate that local source emissions played a more important role in PM pollution near ground level during APEC. The concentration differences in NR-PM₁ began to decrease with the occurrence of the mountain–valley breeze and reached a minimum at noon. Our results revealed the important role of mountain–valley breeze in affecting the boundary-layer structure and reducing the daytime PM levels during APEC. It was estimated that the mountain–valley breeze caused a reduction in NR-PM₁ concentration of approximately 50 μg m⁻³ at the ground site during the day on 10–11 November (Fig. 13b). Therefore, our results illustrated that the achievement of “APEC blue” was also due partly to meteorological effects, particularly the mountain–valley breeze, in addition to emission controls.

4 Conclusions

We have presented a detailed characterization of aerosol particle composition and sources above the urban canopy in Beijing from 10 October to 12 November 2014. This study is unique because it examines strict emission controls implemented during the 2014 APEC summit and synchronous

real-time measurements of aerosol particle composition at 260 m and near ground level obtained by two aerosol mass spectrometers. The NR-PM₁ composition above the urban canopy was dominated by organics at 46 %, followed by nitrate at 27 %, and sulfate at 13 %. The high contribution of nitrate and high NO₃⁻ / SO₄²⁻ mass ratios illustrates the important role of nitrate in PM pollution during the study period. This result has significant implications, namely, that NO_x emission controls should be prioritized for the mitigation of air pollution in Beijing, particularly in non-heating seasons with low SO₂ precursors. The OA above the urban canopy was dominated by OOA at 61 % and included HOA at 39 %. Different from that at the ground site, HOA correlated moderately with OOA above the urban canopy, indicating similar sources, likely through regional transport.

With the implementation of emission controls, the mass concentrations of aerosol species were shown to have decreased significantly by 40–80 % during APEC, whereas the bulk aerosol composition was relatively similar before and during APEC. Organics were dominant before and during the summit, at 46 % versus 47 %, respectively, followed by nitrate at 27 % versus 29 %, and sulfate at 14 % versus 10 %, respectively. Our results suggest that synergistic controls of various precursors such as SO₂, NO_x, and VOCs over a regional scale would not significantly affect regional aerosol bulk composition, although the mass concentrations would be reduced substantially. By linking aerosol compositions and sources to meteorological conditions, we determined that meteorological parameters, particularly mountain–valley breezes, played an important role in suppressing PM growth and hence reducing PM levels during APEC. Our results elucidated that the good air quality in Beijing during APEC was the combined result of emission controls and meteorological effects, with the former playing the dominant role. We further investigated the vertical evolution of aerosol particle composition by comparing the aerosol chemistry between the ground level and 260 m. We observed very complex vertical differences during the formation and evolution of severe haze episodes which were closely related to aerosol sources (local versus regional) and boundary-layer dynamics. Although a stable *T* inversion layer between 120 and 160 m associated with stagnant meteorology caused higher concentrations of aerosol species at the ground site, the interaction of boundary-layer dynamics and aerosol chemistry during the cleaning processes resulted in a lag time of approximately 4 h in cleaning pollutants near ground level compared to processes that occur above the urban canopy.

Acknowledgements. This work was supported by the National Key Project of Basic Research (2014CB447900), the Strategic Priority Research Program (B) of the Chinese Academy of Sciences (XDB05020501), the Key Research Program of the Chinese Academy of Sciences (KJZD-EW-TZ-G06-01-0), and the Special Fund for Environmental Protection Research in the Public Interest (201409001).

Edited by: J. Ma

References

- Aiken, A. C., Salcedo, D., Cubison, M. J., Huffman, J. A., DeCarlo, P. F., Ulbrich, I. M., Docherty, K. S., Sueper, D., Kimmel, J. R., Worsnop, D. R., Trimborn, A., Northway, M., Stone, E. A., Schauer, J. J., Volkamer, R. M., Fortner, E., de Foy, B., Wang, J., Laskin, A., Shutthanandan, V., Zheng, J., Zhang, R., Gaffney, J., Marley, N. A., Paredes-Miranda, G., Arnott, W. P., Molina, L. T., Sosa, G., and Jimenez, J. L.: Mexico City aerosol analysis during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite (T0) – Part 1: Fine particle composition and organic source apportionment, *Atmos. Chem. Phys.*, 9, 6633–6653, doi:10.5194/acp-9-6633-2009, 2009.
- Allan, J. D., Williams, P. I., Morgan, W. T., Martin, C. L., Flynn, M. J., Lee, J., Nemitz, E., Phillips, G. J., Gallagher, M. W., and Coe, H.: Contributions from transport, solid fuel burning and cooking to primary organic aerosols in two UK cities, *Atmos. Chem. Phys.*, 10, 647–668, doi:10.5194/acp-10-647-2010, 2010.
- Arimoto, R., Duce, R., Savoie, D., Prospero, J., Talbot, R., Cullen, J., Tomza, U., Lewis, N., and Ray, B.: Relationships among aerosol constituents from Asia and the North Pacific during PEM – West A, *J. Geophys. Res.-Atmos.*, 101, 2011–2023, 1996.
- Budisulistiorini, S. H., Canagaratna, M. R., Croteau, P. L., Baumann, K., Edgerton, E. S., Kollman, M. S., Ng, N. L., Verma, V., Shaw, S. L., Knipping, E. M., Worsnop, D. R., Jayne, J. T., Weber, R. J., and Surratt, J. D.: Intercomparison of an Aerosol Chemical Speciation Monitor (ACSM) with ambient fine aerosol measurements in downtown Atlanta, Georgia, *Atmos. Meas. Tech.*, 7, 1929–1941, doi:10.5194/amt-7-1929-2014, 2014.
- Draxler, R. R. and Hess, G.: Description of the HYSPLIT4 modeling system, Air Resources Laboratory, Silver Spring, Maryland, 1997.
- Draxler, R., Stunder, B., Rolph, G., Stein, A., and Taylor, A.: HYSPLIT4 user's guide, version 4, report, NOAA, Silver Spring, Md, 2012.
- Guinot, B., Roger, J.-C., Cachier, H., Pucal, W., Jianhui, B., and Tong, Y.: Impact of vertical atmospheric structure on Beijing aerosol distribution, *Atmos. Environ.*, 40, 5167–5180, 2006.
- Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., and Zeng, L.: Elucidating severe urban haze formation in China, *P. Natl. Acad. Sci. USA*, 111, 17373–17378, 2014.
- Han, S., Kondo, Y., Oshima, N., Takegawa, N., Miyazaki, Y., Hu, M., Lin, P., Deng, Z., Zhao, Y., and Sugimoto, N.: Temporal variations of elemental carbon in Beijing, *J. Geophys. Res.-Atmos.*, 114, D23202, doi:23210.21029/22009JD012027, 2009.
- Hu, W. W., Hu, M., Yuan, B., Jimenez, J. L., Tang, Q., Peng, J. F., Hu, W., Shao, M., Wang, M., Zeng, L. M., Wu, Y. S., Gong, Z. H., Huang, X. F., and He, L. Y.: Insights on organic aerosol aging and the influence of coal combustion at a regional receptor site of central eastern China, *Atmos. Chem. Phys.*, 13, 10095–10112, doi:10.5194/acp-13-10095-2013, 2013.
- Huang, K., Zhuang, G., Lin, Y., Fu, J. S., Wang, Q., Liu, T., Zhang, R., Jiang, Y., Deng, C., Fu, Q., Hsu, N. C., and Cao, B.: Typical types and formation mechanisms of haze in an Eastern Asia megacity, Shanghai, *Atmos. Chem. Phys.*, 12, 105–124, doi:10.5194/acp-12-105-2012, 2012a.
- Huang, R. J., Zhang, Y., Bozzetti, C., Ho, K. F., Cao, J. J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Brun, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., El Haddad, I., and Prevot, A. S.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, doi:10.1038/nature13774, 2014.
- Huang, X.-F., He, L.-Y., Hu, M., Canagaratna, M. R., Sun, Y., Zhang, Q., Zhu, T., Xue, L., Zeng, L.-W., Liu, X.-G., Zhang, Y.-H., Jayne, J. T., Ng, N. L., and Worsnop, D. R.: Highly time-resolved chemical characterization of atmospheric submicron particles during 2008 Beijing Olympic Games using an Aerodyne High-Resolution Aerosol Mass Spectrometer, *Atmos. Chem. Phys.*, 10, 8933–8945, doi:10.5194/acp-10-8933-2010, 2010.
- Huang, X.-F., He, L.-Y., Hu, M., Canagaratna, M. R., Kroll, J. H., Ng, N. L., Zhang, Y.-H., Lin, Y., Xue, L., Sun, T.-L., Liu, X.-G., Shao, M., Jayne, J. T., and Worsnop, D. R.: Characterization of submicron aerosols at a rural site in Pearl River Delta of China using an Aerodyne High-Resolution Aerosol Mass Spectrometer, *Atmos. Chem. Phys.*, 11, 1865–1877, doi:10.5194/acp-11-1865-2011, 2011.
- Huang, X.-F., He, L.-Y., Xue, L., Sun, T.-L., Zeng, L.-W., Gong, Z.-H., Hu, M., and Zhu, T.: Highly time-resolved chemical characterization of atmospheric fine particles during 2010 Shanghai World Expo, *Atmos. Chem. Phys.*, 12, 4897–4907, doi:10.5194/acp-12-4897-2012, 2012b.
- Huang, X.-F., Xue, L., Tian, X.-D., Shao, W.-W., Sun, T.-L., Gong, Z.-H., Ju, W.-W., Jiang, B., Hu, M., and He, L.-Y.: Highly time-resolved carbonaceous aerosol characterization in Yangtze River Delta of China: Composition, mixing state and secondary formation, *Atmos. Environ.*, 64, 200–207, 2013.
- Jia, Y., Rahn, K. A., He, K., Wen, T., and Wang, Y.: A novel technique for quantifying the regional component of urban aerosol solely from its sawtooth cycles, *J. Geophys. Res.*, 113, D21309, doi:10.1029/2008jd010389, 2008.
- Jiang, Q., Sun, Y. L., Wang, Z., and Yin, Y.: Aerosol composition and sources during the Chinese Spring Festival: fireworks, secondary aerosol, and holiday effects, *Atmos. Chem. Phys.*, 15, 6023–6034, doi:10.5194/acp-15-6023-2015, 2015.
- Jimenez, J., Canagaratna, M., Donahue, N., Prevot, A., Zhang, Q., Kroll, J., DeCarlo, P., Allan, J., Coe, H., and Ng, N.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.
- Li, W. J., Shao, L. Y., and Buseck, P. R.: Haze types in Beijing and the influence of agricultural biomass burning, *Atmos. Chem. Phys.*, 10, 8119–8130, doi:10.5194/acp-10-8119-2010, 2010.
- Li, Y. J., Lee, B. P., Su, L., Fung, J. C. H., and Chan, C. K.: Seasonal characteristics of fine particulate matter (PM) based on high-resolution time-of-flight aerosol mass spectrometric (HR-ToF-AMS) measurements at the HKUST Supersite in Hong Kong, *Atmos. Chem. Phys.*, 15, 37–53, doi:10.5194/acp-15-37-2015, 2015.
- Lu, Z., Zhang, Q., and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010,

- Atmos. Chem. Phys., 11, 9839–9864, doi:10.5194/acp-11-9839-2011, 2011.
- Matthew, B. M., Middlebrook, A. M., and Onasch, T. B.: Collection efficiencies in an Aerodyne Aerosol Mass Spectrometer as a function of particle phase for laboratory generated aerosols, *Aerosol Sci. Tech.*, 42, 884–898, 2008.
- Meng, Z., Ding, G., Xu, X., Xu, X., Yu, H., and Wang, S.: Vertical distributions of SO₂ and NO₂ in the lower atmosphere in Beijing urban areas, China, *Sci. Total Environ.*, 390, 456–465, 2008.
- Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of composition-dependent collection efficiencies for the aerodyne aerosol mass spectrometer using field data, *Aerosol Sci. Tech.*, 46, 258–271, 2012.
- Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of primary organic aerosol emissions from meat cooking, trash burning, and motor vehicles with high-resolution aerosol mass spectrometry and comparison with ambient and chamber observations, *Environ. Sci. Technol.*, 43, 2443–2449, 2009.
- Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., and Sun, Y.: An Aerosol Chemical Speciation Monitor (ACSM) for routine monitoring of the composition and mass concentrations of ambient aerosol, *Aerosol Sci. Tech.*, 45, 780–794, 2011.
- Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, 111–126, 1994.
- Parworth, C., Fast, J., Mei, F., Shippert, T., Sivaraman, C., Tilp, A., Watson, T., and Zhang, Q.: Long-term measurements of submicrometer aerosol chemistry at the Southern Great Plains (SGP) using an Aerosol Chemical Speciation Monitor (ACSM), *Atmos. Environ.*, 106, 43–55, 2015.
- Petit, J.-E., Favez, O., Sciare, J., Crenn, V., Sarda-Estève, R., Bonnaire, N., Mocnik, G., Dupont, J.-C., Haeffelin, M., and Leoz-Garziandia, E.: Two years of near real-time chemical composition of submicron aerosols in the region of Paris using an Aerosol Chemical Speciation Monitor (ACSM) and a multi-wavelength Aethalometer, *Atmos. Chem. Phys.*, 15, 2985–3005, doi:10.5194/acp-15-2985-2015, 2015.
- Pope III, C. A., Ezzati, M., and Dockery, D. W.: Fine-particulate air pollution and life expectancy in the United States, *New Engl. J. Med.*, 360, 376–386, 2009.
- Quinn, P., Bates, T., Coffman, D., Onasch, T., Worsnop, D., Baynard, T., De Gouw, J., Goldan, P., Kuster, W., and Williams, E.: Impacts of sources and aging on submicrometer aerosol properties in the marine boundary layer across the Gulf of Maine, *J. Geophys. Res.-Atmos.*, 111, D23S36, doi:10.1029/2006jd007582, 2006.
- Sun, J., Zhang, Q., Canagaratna, M. R., Zhang, Y., Ng, N. L., Sun, Y., Jayne, J. T., Zhang, X., Zhang, X., and Worsnop, D. R.: Highly time- and size-resolved characterization of submicron aerosol particles in Beijing using an Aerodyne Aerosol Mass Spectrometer, *Atmos. Environ.*, 44, 131–140, 2010.
- Sun, Y., Wang, Y., and Zhang, C.: Measurement of the vertical profile of atmospheric SO₂ during the heating period in Beijing on days of high air pollution, *Atmos. Environ.*, 43, 468–472, 2009.
- Sun, Y., Zhang, Q., Schwab, J. J., Demerjian, K. L., Chen, W.-N., Bae, M.-S., Hung, H.-M., Hogrefe, O., Frank, B., Rattigan, O. V., and Lin, Y.-C.: Characterization of the sources and processes of organic and inorganic aerosols in New York city with a high-resolution time-of-flight aerosol mass spectrometer, *Atmos. Chem. Phys.*, 11, 1581–1602, doi:10.5194/acp-11-1581-2011, 2011.
- Sun, Y., Song, T., Tang, G., and Wang, Y.: The vertical distribution of PM_{2.5} and boundary-layer structure during summer haze in Beijing, *Atmos. Environ.*, 74, 413–421, 2013.
- Sun, Y., Du, W., Wang, Q., Zhang, Q., Chen, C., Chen, Y., Chen, Z., Fu, P., Wang, Z., and Gao, Z.: Real-Time Characterization of Aerosol Particle Composition above the Urban Canopy in Beijing: Insights into the Interactions between the Atmospheric Boundary Layer and Aerosol Chemistry, *Environ. Sci. Technol.*, 49, 11340–11347, 2015.
- Sun, Y. L., Zhuang, G., Tang, A., Wang, Y., and An, Z.: Chemical characteristics of PM_{2.5} and PM₁₀ in haze-fog episodes in Beijing, *Environ. Sci. Technol.*, 40, 3148–3155, 2006.
- Sun, Y. L., Wang, Z., Dong, H., Yang, T., Li, J., Pan, X., Chen, P., and Jayne, J. T.: Characterization of summer organic and inorganic aerosols in Beijing, China with an Aerosol Chemical Speciation Monitor, *Atmos. Environ.*, 51, 250–259, 2012a.
- Sun, Y. L., Zhang, Q., Schwab, J. J., Yang, T., Ng, N. L., and Demerjian, K. L.: Factor analysis of combined organic and inorganic aerosol mass spectra from high resolution aerosol mass spectrometer measurements, *Atmos. Chem. Phys.*, 12, 8537–8551, doi:10.5194/acp-12-8537-2012, 2012b.
- Sun, Y. L., Wang, Z., Fu, P., Jiang, Q., Yang, T., Li, J., and Ge, X.: The impact of relative humidity on aerosol composition and evolution processes during wintertime in Beijing, China, *Atmos. Environ.*, 77, 927–934, 2013a.
- Sun, Y. L., Wang, Z. F., Fu, P. Q., Yang, T., Jiang, Q., Dong, H. B., Li, J., and Jia, J. J.: Aerosol composition, sources and processes during wintertime in Beijing, China, *Atmos. Chem. Phys.*, 13, 4577–4592, doi:10.5194/acp-13-4577-2013, 2013b.
- Sun, Y. L., Jiang, Q., Wang, Z., Fu, P., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013, *J. Geophys. Res.-Atmos.*, 119, 4380–4398, 2014.
- Ting, M., Yue-Si, W., Jie, J., Fang-Kun, W., and Mingxing, W.: The vertical distributions of VOCs in the atmosphere of Beijing in autumn, *Sci. Total Environ.*, 390, 97–108, 2008.
- Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009.
- Wang, Y., Zhang, Q. Q., He, K., Zhang, Q., and Chai, L.: Sulfate-nitrate-ammonium aerosols over China: response to 2000–2015 emission changes of sulfur dioxide, nitrogen oxides, and ammonia, *Atmos. Chem. Phys.*, 13, 2635–2652, doi:10.5194/acp-13-2635-2013, 2013.
- Xiao, R., Takegawa, N., Zheng, M., Kondo, Y., Miyazaki, Y., Miyakawa, T., Hu, M., Shao, M., Zeng, L., Gong, Y., Lu, K., Deng, Z., Zhao, Y., and Zhang, Y. H.: Characterization and source apportionment of submicron aerosol with aerosol mass spectrometer during the PRIDE-PRD 2006 campaign, *Atmos.*

- Chem. Phys., 11, 6911–6929, doi:10.5194/acp-11-6911-2011, 2011.
- Xu, W. Q., Sun, Y. L., Chen, C., Du, W., Han, T. T., Wang, Q. Q., Fu, P. Q., Wang, Z. F., Zhao, X. J., Zhou, L. B., Ji, D. S., Wang, P. C., and Worsnop, D. R.: Aerosol composition, oxidative properties, and sources in Beijing: results from the 2014 Asia-Pacific Economic Cooperation Summit study, *Atmos. Chem. Phys. Discuss.*, 15, 23407–23455, doi:10.5194/acpd-15-23407-2015, 2015.
- Zhang, J. K., Sun, Y., Liu, Z. R., Ji, D. S., Hu, B., Liu, Q., and Wang, Y. S.: Characterization of submicron aerosols during a month of serious pollution in Beijing, 2013, *Atmos. Chem. Phys.*, 14, 2887–2903, doi:10.5194/acp-14-2887-2014, 2014.
- Zhang, Q., Worsnop, D. R., Canagaratna, M. R., and Jimenez, J. L.: Hydrocarbon-like and oxygenated organic aerosols in Pittsburgh: insights into sources and processes of organic aerosols, *Atmos. Chem. Phys.*, 5, 3289–3311, doi:10.5194/acp-5-3289-2005, 2005.
- Zhang, Q., Streets, D. G., He, K., Wang, Y., Richter, A., Burrows, J. P., Uno, I., Jang, C. J., Chen, D., Yao, Z., and Lei, Y.: NO_x emission trends for China, 1995–2004: The view from the ground and the view from space, *J. Geophys. Res.*, 112, D22306, doi:10.1029/2007jd008684, 2007.
- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Anal. Bioanal. Chem.*, 401, 3045–3067, 2011.
- Zhang, Q. and Tie, X.: High solubility of SO_2 : evidence in an intensive fog event measured in the NCP region, China, *Atmos. Chem. Phys. Discuss.*, 11, 2931–2947, doi:10.5194/acpd-11-2931-2011, 2011.
- Zhang, Q., Meng, J., Quan, J., Gao, Y., Zhao, D., Chen, P., and He, H.: Impact of aerosol composition on cloud condensation nuclei activity, *Atmos. Chem. Phys.*, 12, 3783–3790, doi:10.5194/acp-12-3783-2012, 2012.
- Zhang, W., Zhang, Y., Lv, Y., Li, K., and Li, Z.: Observation of atmospheric boundary layer height by ground-based LiDAR during haze days, *J. Remote Sens.*, 17, 981–992, 2013.
- Zhang, Y. J., Tang, L. L., Wang, Z., Yu, H. X., Sun, Y. L., Liu, D., Qin, W., Canonaco, F., Prévôt, A. S. H., Zhang, H. L., and Zhou, H. C.: Insights into characteristics, sources, and evolution of submicron aerosols during harvest seasons in the Yangtze River delta region, China, *Atmos. Chem. Phys.*, 15, 1331–1349, doi:10.5194/acp-15-1331-2015, 2015.
- Zhao, X. J., Zhao, P. S., Xu, J., Meng, W., Pu, W. W., Dong, F., He, D., and Shi, Q. F.: Analysis of a winter regional haze event and its formation mechanism in the North China Plain, *Atmos. Chem. Phys.*, 13, 5685–5696, doi:10.5194/acp-13-5685-2013, 2013.