

Characterization and Classification of Soils along the Toposequence of Kindo Koye Watershed in Southern Ethiopia

Mulugeta Demiss^{1*} and Sheleme Beyene²

¹ MoA, P O Box 62347, Addis Ababa, Ethiopia

² Hawassa University, P O Box 05 Hawassa, Ethiopia

Abstract: In developing countries, where research funds are limited, the availability of pedogenic information and proper classification of soils will be of great importance. The soils of Kindo Koye watershed were fully characterized along east and west facing toposequences that formed a catena and classified according to the Soil Taxonomy and the WRB Legend to assess the impact of topography on soil development and characteristics. The morphological and physiochemical properties of seven pedons located at the upper, middle and lower slopes of the two toposequences and at the depression were studied. The study revealed the existence of three different soil orders along the toposequences in an area that was previously mapped as Eutric Nitisols. The pedons on the upper and middle slopes of both east and west-facing toposequences and the pedon on the east-facing lower slope were categorized under Ultisols, whereas the pedons on the foot slope west-facing and the depression were categorized under Inceptisols and Entisols, respectively. The Ultisols, Inceptisols and Entisols were further categorized as Acrisols, Cambisols and Fluvisols major groups according to the WRB Legend, respectively. This detail survey and classification of soils shows that topography has a great influence on soil development and characteristics.

Keywords: Catena; Toposequence; Pedon; Soil Taxonomy; WRB Legend

1. Introduction

In developing countries, where research funds are limited, the availability of pedogenic information and proper classification of soils will be of great importance in adopting well tested management technologies and landscape positions without going through the whole process of time consuming and expensive technology selection trials as this will provide the basic information for sustainable agricultural planning (Fikre, 2003). There has not been a comprehensive compilation on soils of Ethiopia, though the felt need for a process-oriented instruction text and as a suitable reference has long been recognized. Even the limited findings are not easily accessible to those who might wish to utilize them (Mesfin, 1998). Consequently, sustainable soil management practices that are based on the understanding of soil systems are not available for most parts of the country (Fikre, 2003).

Landscape position influences runoff, drainage, soil temperature, soil erosion, soil depth and hence soil formation. Different soil properties encountered along landscapes will affect the patterns of plant production, litter production and decomposition, which will definitely have effects on carbon (C) and nitrogen (N) contents of the soil. Soil properties such as clay content and its distribution with depth, sand content and pH have been shown to be highly correlated with landscape position (Wang *et al.*, 2000) while organic matter has been shown to vary with slope position (Miller *et al.*, 1998).

Soils on steep upper slopes range from moderately deep to shallow. They are well drained with the gravelly and channery silt loam and sandy loam textures commonly associated with rock outcrops. These soils generally have severe erosion potential from exposed or bare soil areas and a greater risk of slope failure. Soils in mid-slope and toe or lower slope positions are usually deep, well-drained, gravelly silt loams, whereas those below prominent sandstone cliffs

are usually sandy loams (www.fs.fed.us/r8/boone/resources/soil/index.shtml 03/08/2009). In the lowest landscape positions, water may saturate the regolith to such a degree that drainage and aeration are restricted. Here, the weathering of some minerals and the decomposition of organic matter are retarded, while the loss of iron and manganese is accelerated. In such low-lying topography, special profile features characteristic of wetland soils may develop (Brady and Weil, 2002).

The Ethiopian Mapping Authority (EMA, 1988) characterized the soils of Wolayita areas as Eutric Nitisols. But the Authority used a very small-scale survey that does not specifically tell about the areas considered in this study. Thus, the present study was initiated to fully characterize and classify the soils of the catena following the Soil Taxonomy (Soil Survey Staff, 1999) and the WRB Legend (FAO/WRB, 2006) systems to assess the impact of topography on soil development and characteristics.

2. Materials and Methods

2.1. Description of the Site

The study was conducted at the Kindo Koye watershed, Damot Woyde Woreda, Wolayita Zone, Southern Nations, Nationalities and Peoples' Regional State (SNNPRS). The watershed is located at the coordinates between 6° 52.82' and 6° 53.41' N and 37° 52.42' and 37° 52.63' E with altitude ranging from 1970 to 2061 meters above sea level (masl). The region has a humid climate with an average annual temperature of 20 °C. The monthly mean temperatures range from 17.2 °C in July to 21.9 °C in February. The average annual precipitation is about 1333 mm with monthly minimum and maximum recorded values of 29 mm and 218 mm in the months of January and July, respectively (FAO, 1984). Eighty three percent of the rainfall falls between April and October every year. The major crops and grasses along the selected toposequence

*Corresponding author. E-mail: Mulugetadem@gmail.com

include maize (*Zea mays*), barley (*Hordem vulgare*), sweet potato (*Ipomoea batatas*), sugarcane (*Saccharum officinarum*), and teff (*Eragrotis teff*) and grasses such as *Digitaria diagonalis*. Besides these, plantations dominated by eucalyptus trees (*Eucalyptus camaldulensis*) were present. The soils of the study area are developed on basaltic parent material. According to the WRB Soil Classification System, Eutric Nitosols are the dominant soil units (EMA, 1988).

2.2. Soil Profile Site Selection, Description and Sampling

A catena was selected along east-west facing slopes (toposequences) encompassing landform components spanning from ridge top to valley bottom. The slopes along the toposequences ranged from 0.5 to 25%. The toposequences were divided into three slope categories namely upper slope, middle slope and foot slope, and

depression at the center. A total of six pedons with 2 m length, 2 m width and 2 m depth were excavated one each on upper slope west-facing (UWF), middle slope west-facing (MWF), foot slope west-facing (FWF), foot slope east-facing (FEF), middle slope east-facing (MEF) and upper slope east-facing (UEF) and the seventh with 3 m x 3 m and 3 m dimension at the depression (DEP) (Table 1). The soil profiles of all the sampling slopes and aspects were described *in situ* following the guidelines for soil profile description (FAO, 1990) and samples were collected from all identified horizons. Freshly excavated sites were used for sampling and profile description. Core sampler was used to collect undisturbed soil samples from each horizon to determine bulk density. The total area of the study site was 65 ha. The east-facing toposequence is 31 ha and the west-facing one is 34 ha.

Table 1. Description of soil profile site characteristics.

Pedon	Slope (%)	Altitude (masl)	Surrounding landform	Physiographic position	Parent material
UEF	13	2010	Gentle slope	Upper slope	Basaltic
MEF	20	1997	Steep slope	Middle slope	Basaltic
FEF	25	1984	Steep slope	Foot slope	Colluvium/Basaltic
FWF	12	1976	Steep slope	Foot slope	Colluvium/Basaltic
MWF	18	2002	Steep slope	Middle slope	Basaltic
UWF	10	2029	Gentle slope	Upper slope	Basaltic
DEP	0.5	1970	Gentle/flat	Depression	Alluvium/colluvium

UWF = Upper slope west-facing; MWF = Middle slope west-facing; FWF = Foot slope west-facing; DEP = Depression; UEF = Upper slope east-facing; MEF = Middle slope east-facing; FEF = Foot slope east-facing

2.3. Soil Analysis

The soil samples collected from every identified horizon were air-dried and ground to pass through 2 mm sieve. For the determinations of total N and organic carbon (OC), a 0.5 mm sieve was used. Analysis of the physico-chemical properties of the soil samples were carried out following standard laboratory procedures.

Bulk density was determined using the core-sampling method (BSI, 1975). Particle size distribution was analyzed by the hydrometer method (Sahlemedhin and Taye, 2000) using hydrogen peroxide (H₂O₂) to oxidize organic matter and sodium hexameta phosphate (Na₆PO₃) as a dispersing agent. Soil pH was determined in H₂O and 0.1M KCl using 1:2.5 soils to solution ratio using a combined glass electrode pH meter (Chopra and Kanwar, 1976) and change in pH was determined by subtracting soil pH (KCl) from soil pH (H₂O). Cation exchange capacity and exchangeable bases were determined using the 1M-ammonium acetate (pH 7) method followed by repeated washing with ethanol (96%) to remove the excess ammonium ions in the soil solution according to the percolation tube procedure (Van Reeuwijk, 1993) and the base saturation (BS) and exchangeable sodium percentages (ESP) were computed.

Total N was analyzed by the Kjeldahl digestion and distillation procedure (Bremner and Mulvaney, 1982), whereas OC was determined following the wet combustion method of Walkley and Black as outlined by Van Ranst *et al.* (1999). Available phosphorus (P) content of the soils was analyzed using the Olsen method as outlined by Van

Reeuwijk (1993). Available micronutrients (Fe, Mn, Zn, and Cu) contents of the soils were extracted by the diethylene triamine pentaacetic acid (DTPA) extraction method (Tan, 1996) all were quantified using atomic absorption spectrophotometer.

Finally, analysis of simple correlation coefficient among the different soil physical and chemical properties was carried out using SAS (1997) software to reveal the magnitude and direction of relationships between each other.

3. Results and Discussion

3.1. Soil Morphological Features

Distinct horizons/layers and argillic B-horizons were observed in the pedons, except for the FWF and depression area. Three of the pedons, MEF, FEF and UWF, had Ap and Bt; the UEF had Ap, BA and Bt; the MWF had Ap, AB and Bt; and the FWF had Ap, AB and B, whereas the pedon at the depression had A, AC and C horizon sequences (Table 2).

The depths of the A-horizons decreased with increasing slope (Table 2). Accordingly, the DEP (0.5% slope) and MEF (20% slope) had the deepest and shallowest A-horizons, respectively. The soils at the shoulders tend to be shallower due to erosion, whereas the soils on the foot slope and toe-slope areas tend to be thicker as a result of deposition. Erosion causes stripping of the soil thus preventing the material to stay in place to develop into a soil. The greatest erodibility was associated with the upper

slope positions where soils tended to be shallow, coarse in texture and low in organic matter (OM), while lower erodibility was observed at the lower slope positions with deep, organic-rich and leached soils (Lawrence, 1992). Irvin (1996) who related landform elements to soil properties stated that generally, an increase in slope is associated with a reduction in: leaching, OM content, clay translocation, mineral weathering, horizon differentiation, and solum thickness.

Surface soil color (moist) ranged from very dusky red (2.5YR 2/2) to very dark brown (7.5YR 2.5/2) except in the pedon of the depression, whereas the color (moist) of the subsurface horizons varied from dark reddish brown (2.5YR 2.5/3) to very dark brown (7.5YR 2.5/3) (Table 2). The moist soil colors of the horizons in the pedon at the depression, however, varied from reddish brown (5YR 4/4) to very dark brown (7.5YR 2.5/3).

The results showed that soil color is highly influenced by soil OM, where the darkness in the A-horizon decreased with depth. Dark colored surface horizons (values ≤ 3) are often enriched with OM, offering many benefits to the soil (soils.missouri.edu/tutorial/page7.asp 03/08/2009). Soils on slopes that were never saturated with water had reddish and brownish subsoil colors, which are indicative of well-drained and aerated conditions. Reddish color is due to the presence of iron compounds in various states of oxidation and hydration (Foth, 1990). The horizons in the pedon at the depression varied in color from the others due to reduction reactions caused by water saturation. Pedons that collect water, and are on poorly drained locations where soils are water saturated much of the time, will tend to have grey-colored B-horizons (Foth, 1990). Topography affects the amount of surface runoff, erosion and deposition. If erosion removes soil from the shoulder or back-slope areas of a hill-slope, thinner and light-colored soils remain where the OM content is low. Soils found on foot-slope or toe-slope areas generally show a higher OM content and thicker A- horizon (grunwald.ifas.ufl.edu/Nat_resources/organic_matter/organic.htm 02/08/2009).

The moist consistence of the soils ranged from friable to extremely firm, whereas the wet consistence ranged from slightly sticky/slightly plastic to very sticky/very plastic (Table 2). Despite high clay contents of up to 81% (Table 3), the soil materials were not extremely sticky (Table 2) probably because of the type of clay mineral present. Many red colored tropical soils have clay particles composed mainly of kaolinite and oxides of iron and aluminum, which have little capacity to develop stickiness and to expand and contract on wetting and drying (Foth, 1990). The very friable and friable consistence observed in the surface soils of the pedons (Table 2) could be attributed to the higher OM contents of the layers (Table 5). Although consistence is an inherent soil characteristic, the presence of high OM in

the surface horizon changes its consistence (Wakene and Heluf, 2004).

3.2. Soil Physical Properties

3.2.1. Particle Size Distribution

The soil texture varied from clay loam to clay in the surface horizons of all pedons (Table 3) and became finer from the upper to the middle of the toposequences which may be due to removal of fine soil particles from steeper slope positions by erosion. According to Lawrence (1992), of the individual soil properties considered, silt and sand contents were the most highly correlated with erodibility. Moore *et al.* (1993) found that slope was one of the topographic factors which was most highly correlated with soil properties. The investigators have reported that slope was positively correlated with sand content and negatively correlated with silt content, and high OM mainly occurred when slopes were less than 2%. The sediment transport was different for each soil particle size. The transport of coarse-sized particles (sand) was lowest, whereas the transport of fine soil particles (clay) and medium-sized particles (silt) was high. If erosion occurs on a hill-slope, the silt content often is higher in the bottom soils compared to the soils on the hill-slope shoulder (grunwald.ifas.ufl.edu/Nat_resources/soil_forming_factors/formation.htm 02/08/2009).

The subsurface horizons of most pedons were finer in texture than their respective surface horizons, indicating that there was a prominent translocation of clay down the profiles forming argillic horizons. However, the texture of the subsoil horizons of the pedon at the west-facing foot slope was more or less similar to that of its surface horizon, whereas the pedon at the depression possessed coarser (sandy clay loam) texture as compared to the clay texture of its surface horizons (Table 3), which could be attributed to successive deposition of contrasting materials.

Change in clay percentages down the soil profile suggests pedogenic eluviation–illuviation processes, particularly in the upper as well as middle slope profiles. The presence of faint to prominent clay coatings in the subsurface horizons of the five pedons also indicates that clay illuviation/translocation was the main factor for the formation of argillic horizon in the pedons. The *in situ* synthesis of secondary clays, the weathering of primary minerals in the B-horizon, or the residual concentration of clays from the selective dissolution of more soluble minerals in the B-horizon could have also contributed to the accumulation of clays in the subsoil horizons (Rust, 1983; Chadwick and Graham, 2000; Buol *et al.*, 2003).

Table 2. Morphological features and physical properties of the soils along the toposequences at Kindo Koye watershed.

Horizon	Depth (cm)	Color (moist)	Field texture	Structure*	Consistence		Horizon boundary
					Moist	Wet	
Upper slope east-facing (UEF) Pedon							
Ap1	0-40	7.5YR 2.5/2	Loam	VW, FI, GR	VFI	SST-SPL	G-S
Ap2	40-64	7.5YR 2.5/2	Clay loam	MO, ME, SB	VFI	ST-PL	C-S
BA	64-90	5YR 3/3	Clay	ST, ME, AB	FI	VST-VPL	C-S
Bt1	90-136	7.5YR 2.5/2	Clay	ST, ME, AB	FI	VS-VP	G-S
Bt2	136+	2.5YR 2.5/4	Clay	ST, ME, AB	VFI	VS-VP	-
Middle slope east-facing (MEF) Pedon							
Ap1	0-14	2.5YR 2/2	Clay	VW, ME, GR	FR	ST-PL	C-S
Ap2	14-40	2.5YR 3/3	Clay	VW, FM, AB,	FR	ST-PL	G-S
Bt1	40-53	2.5YR 2.5/4	Clay	MO, FM, AB	EFI	VST-VPL	G-S
Bt2	53-117	2.5YR 3/3	Clay	VS, FM, SB	EFI	VST-VPL	G-S
Bt3	117-165	2.5YR 3/6	Clay	ST, ME, AB	EFI	VST-VPL	G-S
Bt4	165+	2.5YR 3/4	Clay	ST, ME, AB	EFI	VST-VPL	-
Foot slope east-facing (FEF) Pedon							
Ap1	0-18	7.5YR 2/3	Clay loam	VW, ME, GR	FR	SST-SPL	G-S
Ap2	18-53	5YR 3/2	Clay loam	WE, ME, GR	FR	SST-SPL	G-S
Bt1	53-100	2.5YR 2.5/3	Silty clay	WE, ME, AB	FR	SST-SPL	G-S
Bt2	100-160	2.5YR 2.5/3	Silty clay	MO, ME, AB	FI	SST-SPL	D-S
Bt3	160+	5YR 3/4	Clay	ST, ME, AB	VFI	ST-PL	-
Foot slope west-facing (FWF) Pedon							
Ap1	0-17	7.5YR 2.5/3	Clay	WE, FI, GR	FR	SST-SPL	G-S
Ap2	17-37	7.5YR 3/3	Clay	WE, ME, SB	FR	SST-SPL	G-S
AB	37-54	7.5YR 2.5/3	Clay	MO, FI-ME, AB	FR	SST-SPL	G-S
B1	54-71	7.5YR 2.5/3	Clay	ST, FI-ME, AB	FR	SST-SPL	G-S
B2	71-112	7.5YR 3/3	Clay	ST, FI-ME, AB	FR	SST-SPL	G-S
B3	112-133	7.5YR 3/3	Clay	ST, FI-ME, AB	FR	SST-SPL	G-S
B4	133+	7.5YR 3/3	Clay	ST, FI-ME, AB	FR	SST-SPL	-
Middle slope west-facing (MWF) Pedon							
Ap1	0-22	2.5YR 3/4	Loam	WE, VF, GR	FR	SST-SPL	C-S
Ap2	22-43	2.5YR 3/3	Clay	WE, FI, AB	FR	SST-SPL	A-S
AB	43-75	5YR 3/3	Clay	MO, FM, AB	FR	ST-PL	D-S
Bt1	75-95	5YR 3/3	Clay	ST, ME, AB	FR	ST-PL	D-S
Bt2	95-142	2.5YR 3/6	Clay	ST, ME, SB	FR	ST-PL	D-S
Bt3	142+	2.5YR 3/4	Clay	ST, ME, AB	FR	ST-PL	-
Upper slope west-facing (UWF) Pedon							
A1	0-38	7.5YR 2.5/2	Clay loam	WE, FM,GR	VFR	ST-SS	G-D
A2	38-78	7.5YR 2.5/3	Clay loam	VW, FM, GR	VFR	ST-PL	G-S
Bt1	78-102	5YR 3/2	Clay	WE, FM, SB	VFI	VST-VPL	A-S
Bt2	102-171	5YR 3/3	Clay	FI, FM, AB	VFI	VST-VPL	G-S
Bt3	171+	2.5YR 2.5/4	Clay	FI, ME, AB	VFI	VST-VPL	-
Depression (DEP) Pedon							
A1	0-20	5YR 3/3	Clay loam	WE, FM, G	FR	ST-SPL	G-S
A2	20-50	7.5YR 3/3	Clay loam	WE, FM, SB	FR	ST-SPL	G-S
A3	50-70	7.5YR 2.5/3	Sandy loam	WE, M, SB	FR	ST-SPL	G-S
AC1	70-95	7.5YR 3/4	Sandy loam	VW, C, SB	LO	ST-SPL	G-S
AC2	95-120	5YR 3/2	Silty clay	MS	FR	ST-SPL	G-S
C1	120-150	5YR 3/3	SCL	SG	LO	ST-SPL	G-S
C2	150-170	5YR 3/2	SCL	SG	LO	ST-SPL	A-S
C3	170-200	5YR 3/3	Clay loam	MS	FI	ST-PL	A-S
C4	200-209	5YR 3/3	SCL	MS	LO	ST-PL	A-S
C5	209-230	5YR 3/3	Clay loam	MS	FI	ST-SPL	G-S
C6	230-255	5YR 3/3	Sandy loam	MS	LO	ST-PL	A-S
C7	255+	5YR 3/3	Clay	MS	FI	VST-VPL	-

*WE = Weak; VW = Very weak; FM = Fine and medium; GR = Granular; AB = Angular blocky; SB = Sub angular blocky; FR = Friable; VFR = Very friable; FI = Firm; VFI = Very firm; EFI = Extremely firm; SST-SPL = Slightly sticky and slightly plastic; ST-PL = Sticky and plastic; VST-VPL = Very sticky and very plastic; G-S = Gradual and smooth; C-S = Clear and smooth; D-S = Diffuse and smooth; A-S = Abrupt and smooth; MO = Moderate; SCL = Sandy clay loam; ME = Medium; LO = Loose

Table 3. Particle size distribution and bulk density of the soils in Kindo Koye watershed catena.

Horizon	Depth (cm)	Sand (%)	Silt (%)	Clay (%)	Textural class	Bulk density (g cm ³)
Upper slope East-facing (UEF) Pedon						
Ap1	0-40	31	38	31	Clay loam	1.33
Ap2	40-64	23	30	47	Clay	1.26
BA	64-90	21	28	51	Clay	1.46
Bt1	90-136	11	12	77	Clay	1.42
Bt2	136+	9	10	81	Clay	1.33
Middle slope East-facing (MEF) Pedon						
Ap1	0-14	30	32	38	Clay loam	1.36
Ap2	14-40	24	28	48	Clay	1.31
Bt1	40-53	20	20	60	Clay	1.33
Bt2	53-117	18	16	66	Clay	1.34
Bt3	117-165	16	12	72	Clay	1.66
Bt4	165+	20	12	68	Clay	1.35
Foot slope East-facing (FEF) Pedon						
Ap1	0-18	32	30	38	Clay loam	1.34
Ap2	18-53	26	34	40	Clay	1.21
Bt1	53-100	22	20	58	Clay	1.49
Bt2	100-160	18	20	62	Clay	1.42
Bt3	160+	24	16	60	Clay	1.42
Foot slope West-facing (FWF) Pedon						
Ap1	0-17	30	28	42	Clay	1.35
Ap2	17-37	34	26	40	Clay	1.49
AB	37-54	34	24	42	Clay	1.35
B1	54-71	32	26	42	Clay	1.43
B2	71-112	36	30	34	Clay loam	1.27
B3	112-133	30	34	36	Clay loam	1.38
B4	133+	26	34	40	Clay	1.45
Middle slope West-facing (MWF) Pedon						
Ap1	0-22	20	28	52	Clay	1.37
Ap2	22-43	20	24	56	Clay	1.44
A	43-63	20	24	56	Clay	1.41
AB	63-75	22	22	56	Clay	1.41
Bt1	75-95	20	14	66	Clay	1.44
Bt2	95-142	14	10	76	Clay	1.41
Bt3	142+	14	8	78	Clay	1.38
Upper slope West-facing (UWF) Pedon						
A1	0-38	31	34	35	Clay loam	1.25
A2	38-78	31	36	33	Clay loam	1.36
Bt1	78-102	19	22	59	Clay	1.37
Bt2	102-171	19	22	59	Clay	1.26
Bt3	171+	15	20	65	Clay	1.43
Depression (DEP) Pedon						
A1	0-20	23	32	45	Clay	1.28
A2	20-50	17	40	43	Clay	1.28
A3	50-70	31	28	41	Clay	1.37
AC1	70-95	27	30	43	Clay	1.33
AC2	95-120	31	34	35	Clay loam	1.32
C1	120-150	51	20	29	Sandy clay loam	1.45
C2	150-170	47	22	31	Sandy clay loam	1.59
C3	170-200	45	26	29	Sandy clay loam	1.20
C4	200-209	59	16	25	Sandy clay loam	1.34
C5	209-230	27	36	37	Clay loam	1.20
C6	230-255	36	26	38	Clay loam	1.30
C7	255+	20	36	44	Clay	1.43

3.2.2. Bulk Density

Bulk density was highest in the surface horizon of the Pedon in the MEF followed by that in the MEF while the lowest in the UWF followed by the Depression area. The subsoil bulk density was highest (1.66 g cm^{-3}) in the MEF and second (1.59 g cm^{-3}) as well as lowest (1.20 g cm^{-3}) were observed in the Depression area (Table 3). Generally, bulk density increased with depth primarily because of decrease in soil OM content and soil aggregation, as was indicated by the significant negative correlation between the two properties (Table 6). Soils that are loose, porous, or well-aggregated will have lower bulk densities than soils that are compacted or non-aggregated, as pore space (or air) weighs less than the solid space (soil particles) ([weather.nmsu.edu/teaching material/soil252/Chapt5.htm](http://weather.nmsu.edu/teaching_material/soil252/Chapt5.htm) 05/12/2007).

The values of bulk density in the middle slope positions were relatively high, which might be attributed to cultivation. Secondary tillage (cultivation) generally decrease pore space and thus increases bulk density which stands as a reason for the higher bulk densities of the cropped soils than the uncropped soils. The movement of machinery over the field forces solid particles into spaces once occupied by water or air, resulting in less pore space and increased bulk density ([weather.nmsu.edu/teaching material/soil252/Chapt5.htm](http://weather.nmsu.edu/teaching_material/soil252/Chapt5.htm) 05/12/2007). Bulk density is an indirect measure of pore space and is affected primarily by texture and structure showing that as solid space and clay content increase, bulk density decreases.

3.3. Chemical Properties

3.3.1. Soil pH

The pH (H_2O) of the soil in the surface layers of the pedons was found to be slightly to moderately acidic, as per the rating of Jones (2003), with values ranging from 5.6 to 6.2 (Table 4). In all the soil profiles of the different landscape positions, soil pH measured in water was higher by about 1-2 units than the respective pH values measured in KCl solution. The low soil pH values with KCl determination indicate the presence of substantial quantity of exchangeable hydrogen ion. According to Mekar and Uehara (1972) and Anon (1993), high soil acidity with KCl solution

determination showed the presence of high potential acidity and relatively readily weatherable minerals.

3.3.2. Cation Exchange Capacity and Exchangeable Bases

The cation exchange capacity (CEC) of the soils across the surface and subsurface horizons ranged from 15.4 to 28.8 $\text{cmol}(+) \text{ kg}^{-1}$. The surface soil CEC was the highest in the UEF pedon followed by the Depression while the lowest was observed in the MEF pedon (Table 4). The CEC values of the five pedons (UWF, MWF, FWF, MEF and FEF) could be considered as medium and that of the remaining two pedons (UEF and DEP) as high in accordance with the rating of Landon (1991). The CEC values of the pedons showed inconsistent relationship with depth (Table 4).

The concentrations of the basic exchangeable cations in the upper slopes were in the order of $\text{Ca} > \text{Mg} > \text{K} > \text{Na}$ (Table 4). The exchangeable Ca contents of the upper slope positions were more than quadruple as compared to that of the foot slope positions of the toposequences. The higher Ca content of the soils at the upper slope positions might be due to its strong adsorption to the soil colloids as compared to other cations, particularly Na, because of its higher charge and small hydrated radius (Foth, 1990). However, the order of abundance the basic cations varied markedly along the toposequences which may be due to differences in land use system on the landscape positions. Similar to Ca, the highest Mg contents were obtained in the upper slope positions, although the sub-soil layers of the MWF were rich in Mg. Highest exchangeable K was recorded in the surface horizon of the MWF, and generally the middle and foot slope positions had more K content in surface horizons as compared to the upper slope positions in both east and west-facing sides (Table 4).

Exchangeable Ca and Mg were positively correlated with CEC, whereas the correlations of Na and K with CEC were negative (Table 6). The percent base saturation (52) was found to be highest in the surface horizon of the UEF pedon, whilst the lowest (26) was recorded in the surface horizon of FEF pedon.

Table 4. Cation exchange capacity (CEC), exchangeable bases, ESP, and pH of the soils at the Kindo Koye Watershed.

Depth (cm)	pH (H ₂ O)	pH (KCl)	Na	K	Ca	Mg	*TEB	CEC (cmol(+) kg ⁻¹)	PBS	ESP
			(cmol(+) kg ⁻¹)				(cmol(+) kg ⁻¹)			
Upper slope East-facing (UEF) Pedon										
0-40	6.1	5.2	0.17	0.50	10.98	2.55	14.20	27.40	52	0.62
40-64	5.8	5.3	0.21	0.50	9.18	1.91	12.40	27.60	45	0.76
64-90	6.3	5.2	0.29	0.38	6.99	2.55	10.21	26.60	38	1.09
90-136	6.0	4.8	0.33	0.41	5.49	3.54	9.77	27.60	35	1.19
136+	6.1	4.8	0.37	0.41	5.84	4.03	10.65	28.80	37	1.28
Middle slope East-facing (MEF) Pedon										
0-14	6.2	4.5	0.62	2.03	1.77	0.28	4.70	17.20	27	3.60
14-40	6.1	4.7	0.87	1.55	1.98	0.25	4.65	16.60	28	5.24
40-53	6.1	4.8	0.96	1.97	1.61	0.64	5.18	16.90	31	5.68
53-117	6.6	4.8	0.84	3.69	1.15	0.45	6.13	16.70	37	5.03
117-165	6.7	5.1	0.84	5.11	1.12	0.43	7.50	18.00	42	4.67
165+	6.7	5.1	0.90	5.43	1.18	0.51	8.02	17.20	47	5.23
Foot slope East-facing (FEF) Pedon										
0-18	5.8	4.2	0.73	2.33	1.36	0.69	5.11	19.50	26	3.74
18-53	5.8	4.2	0.96	1.57	1.52	0.38	4.43	16.10	28	5.96
53-100	5.8	4.3	0.97	2.75	1.01	0.41	5.14	18.00	29	5.39
100-160	5.1	4.0	1.75	2.19	0.96	0.43	5.33	17.80	30	9.83
160+	5.5	4.0	1.08	2.68	0.96	0.42	5.14	18.20	28	5.93
Foot slope West-facing (FWF) Pedon										
0-17	5.8	4.4	0.77	3.50	1.04	0.25	5.56	20.60	27	3.74
17-37	5.7	4.3	0.69	2.95	0.81	0.32	4.77	20.20	24	3.41
37-54	5.8	4.3	0.85	2.16	1.01	0.21	4.23	17.60	24	4.83
54-71	5.6	4.2	0.99	1.65	1.19	0.34	4.17	19.50	21	5.08
71-112	5.8	4.3	0.76	1.46	1.17	0.22	3.61	21.00	17	3.62
112-133	5.7	4.3	0.79	1.16	1.28	0.21	3.44	19.50	18	4.05
133+	5.2	4.0	0.87	1.04	1.03	0.30	3.24	19.10	17	4.55
Middle slope West-facing (MWF) Pedon										
0-22	6.1	4.6	0.92	5.93	1.40	0.75	9.00	23.79	38	3.87
22-43	5.8	4.5	0.29	0.26	6.04	2.30	8.89	23.40	38	4.40
43-63	5.7	4.1	0.27	0.21	5.34	3.13	8.94	25.20	35	4.54
63-75	5.7	4.0	0.23	0.16	4.84	2.63	7.87	23.40	34	4.71
75-95	5.8	4.0	0.27	0.17	3.79	3.21	7.45	23.60	32	4.45
95-142	5.9	4.0	0.21	0.16	5.84	3.79	9.99	25.60	39	4.26
142+	6.5	4.1	0.89	3.59	1.16	0.68	6.32	19.50	32	4.56
Upper slope West-facing (UWF) Pedon										
0-38	5.6	4.4	0.15	0.14	6.94	2.47	9.70	23.20	42	0.65
38-78	5.9	4.4	0.20	0.21	7.24	1.88	9.13	24.40	37	0.82
78-102	5.3	4.4	0.44	0.18	6.34	2.88	9.85	26.20	38	1.68
102-171	5.2	4.1	0.52	0.17	3.64	1.23	5.56	25.00	22	2.08
171+	5.6	3.8	0.40	0.17	6.29	1.56	8.42	24.40	35	1.64
Depression (DEP) Pedon										
0-20	6.2	4.4	0.24	0.09	6.64	2.39	9.35	26.60	35	0.90
20-50	5.6	4.3	0.20	0.05	4.14	1.07	5.46	26.00	21	0.77
50-70	6.2	4.2	0.27	0.05	6.89	1.73	8.93	22.00	41	1.23
70-95	6.0	4.3	0.98	1.02	1.51	0.32	3.83	16.40	23	5.98
95-120	5.9	4.4	0.98	1.37	1.60	0.34	4.29	15.40	28	6.36
120-150	6.3	4.4	0.96	1.86	1.22	0.23	4.27	19.50	22	4.92
150-170	6.1	4.4	0.38	0.11	7.49	3.70	11.68	24.60	47	1.54
170-200	6.5	4.4	0.29	0.07	6.09	2.55	9.00	21.20	42	1.37
200-209	6.6	4.5	0.27	0.07	4.69	1.32	6.35	15.60	41	1.73
209-230	6.1	4.4	0.40	0.11	7.98	3.37	11.86	25.40	47	1.57
230-255	6.5	4.6	0.42	0.11	5.49	2.47	8.48	24.40	35	1.72
255+	5.8	4.4	0.38	0.48	8.23	3.70	12.80	28.20	45	1.34

*TEB = Total exchangeable bases; CEC = Cation exchange capacity; PBS = Percent base saturation; ESP = Exchangeable sodium percentage

3.3.3. Organic Carbon and Total Nitrogen

The organic carbon (OC) and total nitrogen (N) contents of the soils decreased with depth in most pedons, and the surface horizons in the pits on the east-facing slope positions (UEF, MEF and FEF) contained relatively higher OC and total N than their respective west-facing pedons (Table 5). Generally, the pedon at the Depression area has relatively higher OC and total N than the other pedons except for OC in the UEF and MEF and for total N in the FEF. The range of soil OC contents in the surface horizons are considered as very low according to the ratings of Metson (1961), whereas the total N content of the surface horizons ranged from 0.115 (MEF) to 0.217% (FEF) (Table 5) and are considered as low to medium according to Havlin *et al.* (1999).

Organic carbon in the surface layer decreased down slope on east-facing toposequence. The highest organic carbon value of 2.16% was recorded in UEF, the front yard area used for cattle tethering, indicating higher content of organic carbon in uncultivated land as compared to its cultivated counterparts. Wakene and Heluf (2004) have also indicated that intensive cultivation aggravates OM oxidation and hence reduces OC content. Similarly, the organic carbon in surface layer decreased from UWF to MWF. However the FWF had higher organic carbon than MWF, which might be attributed to partial accumulation of the material from the upper and middle slopes. The total nitrogen content of the surface layers along toposequences followed similar trend with that of organic matter, except for the FEF pedon. The difference in OC and total N content among the pedons could be attributed to the effect of variation in the land use systems along the toposequences. In addition, higher OC and total N contents were recorded in the surface as compared to subsurface layers indicating strong correlation between them. However, their contents in the Depression pedon did not follow similar trend due to accumulation of contrasting material that add different materials from top parts through erosion in different years and water-logging, which might have affected decomposition and mineralization (Wang *et al.*, 2000). Organic carbon and total nitrogen contents were positively correlated with available P and micronutrients, but negatively with clay (Table 6).

3.3.4. Available Phosphorus

Available phosphorus (P) contents of the soils in the surface horizons was highest in the UEF (7.3 mg kg⁻¹) followed by FWF (4.54 mg kg⁻¹) and MEF (4.2 mg kg⁻¹) while the lowest was recorded in the MWF (0.64 mg kg⁻¹) pedon. Available P generally has inconstant relationship with depth in all pedons and irregularities were observed due to contrasting materials (Table 5). There is a general increase in the distribution of available P from top to bottom along the slope both on the east and west-facing toposequences. This is due to the fact that the relationship of slope position to soil properties is, to a great degree, controlled by erosion processes in that it alters the distribution of soil particles

and water redistribution over the field. Kravchenko and Bullock (2000) found that in more than half of their study sites, slope was negatively correlated to CEC, organic matter, available P, and exchangeable K.

According to Havlin *et al.* (1999), the available P contents of the soils in the surface horizons are considered to be very low to low. The relatively higher available P in the surface horizons of most soil profiles as compared to that of subsurface layers could be attributed to the difference in organic matter contents of the layers. Available P content was positively correlated ($r = 0.41$) with organic carbon (Table 6). High OM content and a good rate of its mineralization could ensure release of phosphate ions adequate for crop production, though most of the phosphate released in this way will be in the topsoil. If not immediately taken up by the plant or by soil organisms, however, it will be converted to non-labile form. Besides, the phosphate ions are most likely to combine with free iron or aluminum ions in acid soils to form iron III phosphates and aluminum phosphates, which are relatively insoluble (Ahn, 1993).

The decrease in available P content with depth in the pedons, except for the Depression pedon, is attributed to the increment of clay, as available P correlates negatively ($r = -0.56$) with clay content (Table 6). Iron and aluminum oxides are intimately associated with the kaolinitic clay fraction of the soil or as coatings on the clay, and thereby increase P fixation. Fixation is thus related to soil texture and would be expected to be greater in clayey soils than in light textured ones. This applies particularly when the clay is kaolinitic (Ahn, 1993). Available P had also a very highly significant correlation with Fe, Zn and Cu, and high correlation with Mn (Table 6).

3.2.5. Available Micronutrients (Fe, Mn, Zn and Cu)

The micronutrients contents in all pedons decreased with increasing soil depth (Table 5). The order of micronutrients concentration in the pedons was Mn > Zn > Fe > Cu, except for east-facing foot slope where Zn > Mn > Fe > Cu was recorded.

The concentrations of Cu and Fe are very low and that of Mn and Zn fall in low to high ranges as compared with the normal ranges of these nutrients in soil (Havlin *et al.*, 1999). The trend of Mn concentration under different slope positions was similar to that of Fe distribution (Table 5) indicating that these two elements have similar chemical behavior in tropical soils (Kravskoof, 1972).

The distribution of CU was consistently decreased from the surface to the subsurface horizons, which might be attributed to the strong association of Cu with soil organic matter. There was also a positive correlation between copper and organic matter showing that copper is strongly complexed with organic matter as was also described earlier by Wakene and Heluf (2004). Moreover, the Fe and Mn contents were also decreased consistently with depth in the profiles of every landscape positions.

Table 5. Total nitrogen, organic carbon (OC) and available phosphorus and micronutrients contents of the soils at the Kindo Koye Watershed.

Depth (cm)	Total N (%)	OC (%)	C/N ratio	Available P (mg kg ⁻¹)	Fe (mg kg ⁻¹)	Mn (mg kg ⁻¹)	Zn (mg kg ⁻¹)	Cu (mg kg ⁻¹)
Upper slope East-facing (UEF) Pedon								
0-40	0.178	2.163	12	7.30	1.39	4.51	10.23	0.18
40-64	0.140	1.344	10	1.40	0.59	0.53	3.94	0.11
64-90	0.113	0.789	7	0.92	1.06	0.40	3.41	0.31
90-136	0.070	0.481	7	0.92	0.59	0.95	0.79	0.18
136+	0.090	0.585	7	1.66	0.37	0.44	0.48	0.15
Middle slope East-facing (MEF) Pedon								
0-14	0.115	1.948	17	4.20	1.19	3.83	6.09	0.18
14-40	0.185	1.318	7	1.34	0.79	4.80	2.49	0.24
40-53	0.132	1.127	9	1.34	0.37	0.88	1.91	0.18
53-117	0.110	0.879	8	1.34	0.66	0.84	1.17	0.07
117-165	0.123	0.813	7	2.34	0.37	0.29	0.40	0.13
165+	0.070	0.530	8	2.20	0.55	0.20	0.66	0.13
Foot slope East-facing (FEF) Pedon								
0-18	0.217	1.743	8	2.30	1.03	8.78	7.96	0.20
18-53	0.153	1.743	11	1.30	2.57	7.55	7.52	0.24
53-100	0.137	0.905	7	0.34	0.66	0.81	1.76	0.15
100-160	0.112	0.813	7	0.30	0.51	0.33	0.62	0.18
160+	0.098	0.720	7	0.30	0.18	0.15	0.70	0.07
Foot slope West-facing (FWF) Pedon								
0-17	0.127	1.555	12	4.54	2.20	7.22	6.40	0.20
17-37	0.108	1.113	10	2.48	1.36	7.11	5.30	0.15
37-54	0.148	0.966	7	1.98	1.30	5.30	4.36	0.13
54-71	0.141	1.325	9	1.98	1.72	4.60	4.29	0.15
71-112	0.160	1.452	9	1.90	1.85	11.97	7.15	0.29
112-133	0.140	1.662	12	2.56	ND	6.27	8.07	0.24
133+	0.143	1.489	10	2.56	3.48	9.77	8.36	0.22
Middle slope West-facing (MWF) Pedon								
0-22	0.116	1.069	9	0.64	0.70	0.59	3.85	0.07
22-43	0.109	1.051	10	0.46	0.73	0.20	3.15	0.13
43-63	0.092	0.879	10	1.18	0.66	0.33	2.64	0.13
63-75	0.095	0.668	7	1.12	0.66	0.81	2.53	0.11
75-95	0.070	0.554	8	1.00	0.37	0.62	2.02	0.07
95-142	0.049	0.513	10	1.00	0.29	0.24	1.14	0.13
142+	0.028	0.358	13	1.00	0.31	0.24	0.51	0.04
Upper slope West-facing (UWF) Pedon								
0-38	0.146	1.572	11	1.38	2.22	8.16	8.58	0.22
38-78	0.115	1.407	12	1.20	2.09	3.21	8.07	0.24
78-102	0.085	0.880	10	0.56	0.57	0.92	2.73	0.13
102-171	0.092	0.657	7	0.56	0.44	0.37	2.24	0.13
171+	0.070	0.491	7	0.28	0.33	0.68	1.23	0.18
Depression (DEP) Pedon								
0-20	0.189	1.890	10	1.50	1.43	9.79	6.62	0.31
20-50	0.185	1.638	9	1.50	1.45	6.89	6.25	0.22
50-70	0.168	1.134	7	1.24	1.63	6.93	7.04	0.11
70-95	0.143	1.239	9	1.80	1.19	5.43	5.48	0.31
95-120	0.169	1.155	7	2.42	1.52	4.93	4.84	0.31
120-150	0.112	0.819	7	3.60	1.87	7.85	5.90	0.29
150-170	0.126	0.945	8	4.30	1.32	6.23	5.52	0.29
170-200	0.115	1.176	10	4.18	1.83	6.18	6.09	0.24
200-209	0.147	0.990	7	3.70	2.42	12.56	5.52	0.33
209-230	0.112	1.197	11	5.20	1.01	2.79	4.51	0.24
230-255	0.113	0.882	8	4.60	1.08	1.36	3.98	0.22
255+	0.147	1.113	8	6.36	4.93	2.95	5.30	0.40

3.4. Classification of the Soils

The dominant soils of the area were previously mapped as Eutric Nitisols (EMA, 1988). The soil profiles, except that of the depression, had thick (37 to 78) surface horizons, having moist color of 7.5YR 5/2 and dark and very weak to weak structure. The organic carbon content of the surface horizons of the pedons ranged from 1.05 to 1.95% with percent base saturation values of less than 50 using 1M NH₄OAc at pH 7. Thus, the six pedons possessed umbric epipedons, whereas the pedon on the depression failed to meet criteria for any other epipedons except Ochric.

In the subsurface horizons of all, but FWF pedon, thick horizons (133 to 171 cm) with clay contents ranging from 51 to 81% were observed. The clay contents of the subsurface horizons were found to be more than 1.2 times greater than their respective surface layers, the subsurface horizons contained 8% more clay than the horizon above where the horizon above had more than 40% clay and these clay increments were found within distances less than 15 cm (Table 4). The apparent cation exchange capacities of the horizons ranged between 25 and 88 cmol(+) kg⁻¹. Besides, faint and prominent pedfaces, argillians, were observed in the horizons. These properties would therefore qualify the horizons of the five pedons as argillic subsurface diagnostic horizons as described by Buol *et al.* (2003). Considering these features, the five pedons were grouped under Ultisols. The subsurface layers of the FWF pedon did not show clay increment, although there was evidence of color alteration indicating that it possesses cambic horizons. In addition, it has base saturation less than 50% using 1M NH₄OAc at pH 7 between the umbric epipedon and a depth of 180 cm. Thus, it was classified as Inceptisols, whereas the pedon at the Depression without any diagnostic horizon was classified as Entisols.

The region is characterized by isothermic temperature and ustic moisture regimes based on the estimates made using the mean annual and monthly temperature and moisture distributions of the region, respectively (Van Wambeke, 2003). Thus, the five pedons were classified as Ustults on the bases of soil moisture regime at the suborder level. Further, the pedons did not have a densic, lithic, paralithic or petroferic contact within 150 cm; and did not have a clay decrease of 20% or more from the maximum clay content. Hence, they were classified as Paleustults and Typic Paleustults at great groups and subgroups, respectively (Table 7).

Considering the ustic soil moisture regime, absence of free carbonates within 200 cm of the mineral soil surface and base saturation (1M NH₄OAc) of less than 60% in all horizons at a depth between 25 and 75 cm from the mineral soil surface and the presence of umbric epipedon, the FWF pedon was further classified as Ustepts, Dystrustepts and Humic Dystrustepts at suborder, great groups and subgroup levels, respectively (Table 7).

The pedon at the Depression was classified as Fluvents based on the features such as < 25% slope, 0.2% or more

OC and irregular decrease in OC content from a depth of 25 cm to 125 cm. This Depression area pedon was further classified as Aquic Usticfluvents due to ustic soil moisture regime with seasonal aquatic conditions.

The classification of the soils of the five pedons as Ultisols, and that of FWF and DEP as Inceptisols and Entisols, respectively, was a major deviation from the previous soil map (EMA, 1988). The position of the FWF and DEP pedons on the catena makes these orders different from the rest of the pedons in the area and showing the influence of topography on soil development.

The five pedons, which were grouped under Ultisols following Soil Taxonomy, were classified as Acrisols due to the presence of an argillic B horizon; having a base saturation which is less than 50% in, at least, some part of the B horizon within 125 cm of the surface. They were further grouped under Ferric Acrisols at the unit level because of their ferric properties (Table 7)

The FWF pedon was categorized as Dystric Cambisols due to the presence of an umbric A horizon, which is more than 25 cm thick a base saturation of less than 50% in at least some part of the B horizon. The pedon at the depression was categorized under Dystric Fluvisols as it developed from recent alluvial deposits, and had no diagnostic horizons other than an ochric A and base saturation of less than 50%, at least in some part of the soil between 20 and 50 cm from the surface (Table 7).

Contrary to the report of the EMA (1988), the present study revealed the existence of three soil orders or mapping units within the catena. This is a result of detail survey and classification of soils and shows that topography has a great influence on soil development.

4. Conclusions

The soils of a catena could differ as a result of erosion, transport and deposition of surface materials as well as leaching, translocation and deposition of chemicals and particulate constituents in the soil. Topography plays a major role in these processes and thereby influences the development and characteristics of the soils along the toposequences. Most of the important soil quality indicators such as bulk density, structure, OC, soil pH, CEC, total N, available P, exchangeable bases, and available micronutrients were influenced by the different landscape positions, particularly at the surface horizon. Continuous intensive cultivation without appropriate soil management practices has contributed to the degradation of the important soil quality indicators. Therefore, reducing intensive cultivation, and integrated use of inorganic and organic fertilizers could replenish the degraded soil quality parameters for sustainable productivity. However, further study of the areas is recommended especially with respect to soil landscape - land management relationships so as to give sound conclusion for the sustainable use of the land.

Table 6. Correlation between properties of the soils in Kindo Koye watershed catena.

	BD	Na	K	Ca	Mg	CEC	Total N	OC	Av. P	Fe	Mn	Zn	Cu
Clay	0.296	0.107	0.263	-0.123	0.184	0.151	-0.660**	-0.693**	-0.564**	-0.586**	-0.753**	-0.862**	-0.618**
BD		0.157	0.263	-0.165	0.000	-0.044	-0.269	-0.387	-0.095	-0.121	-0.259	-0.323	-0.152
Na			0.664**	-0.826**	-0.756**	-0.731**	0.049	-0.062	-0.182	-0.093	-0.028	-0.228	-0.092
K				-0.696**	-0.633**	-0.529*	-0.097	-0.102	-0.097	-0.176	-0.165	-0.277	-0.377
Ca					0.828**	0.797**	-0.009	0.069	0.284	0.122	-0.115	0.191	0.193
Mg						0.815**	-0.300	-0.252	0.158	0.005	-0.282	-0.108	0.070
CEC							-0.188	-0.083	0.088	-0.004	-0.260	0.015	0.004
TN								0.782**	0.283	0.416*	0.662**	0.673**	0.497*
OC									0.414*	0.428*	0.627**	0.840**	0.390
Av. P										0.526*	0.352	0.501**	0.478*
Fe											0.588**	0.606**	0.646*
Mn												0.782**	0.585**
Zn													0.517*

** = Correlation is significant at $P \leq 0.0$; * = Correlation is significant at $P \leq 0.05$; BD = Bulk density; CEC = Cation exchange capacity; OC = Organic carbon; Av. P = Available phosphorus

Table 7. Classification of the soils of Kindo Koye watershed catena following Soil Taxonomy and the WRB Systems.

Pedon*	Soil Taxonomy				The WRB legend	
	Order	Suborder	Great group	Subgroup	Major Groups	Units
UEF	Ultisols	Ustults	Paleustults	Typic Paleustults	Acrisols	Ferric Acrisols
MEF	Ultisols	Ustults	Paleustults	Typic Paleustults	Acrisols	Ferric Acrisols
FEF	Ultisols	Ustults	Paleustults	Typic Paleustults	Acrisols	Ferric Acrisols
DEP	Entisols	Fluvents	Usticfluvents	Aquic Ustifluvents	Fluvisols	Dystric Fluvisols
FWF	Inceptisols	Ustepts	Dystrustepts	Humic Dystrustepts	Cambisols	Dystric Cambisols
MWF	Ultisols	Ustults	Paleustults	Typic Paleustults	Acrisols	Ferric Acrisols
UWF	Ultisols	Ustults	Paleustults	Typic Paleustults	Acrisols	Ferric Acrisols

*UWF = Upper slope west-facing; MWF = Middle slope west-facing; FEF = Foot slope east-facing; DEP = Depression; FWF = Foot slope west-facing; MEF = Middle slope east-facing; UEF = Upper slope east-facing

5. Acknowledgments

We would like to thank B.B. Mishra, Alemayehu Kiflu, Dereje Tsegaye, Fitsum Tekly, Gifole Gidago and Tigist Yimer for their kind assistance in fieldwork. We also extend our gratitude to the CIDA/UPCD Project for providing the financial support for the fieldwork and analytical service costs of the study.

6. References

- Ahn, P.M. 1993. *Tropical Soils and Fertilizer Use*. Longman Group UK Limited, UK.
- Anon. 1993. *Soil Chemical Analysis*. Part I, Vol. 1. Improvement of soil services for Agricultural development.
- Brady, N.C. and Weil, R.R. 2002. *The Nature and Properties of Soils*. 13th edition. Saurabh Printers Pvt. Ltd., India.
- Bremner, J.M. and Mulvaney, C.S. 1982. Nitrogen-total. In: Page, A.L. (ed.). *Methods of Soil Analysis*. Part 2. Chemical and microbiological properties. SSSA, Madison, Wisconsin. pp. 595-641.
- BSI (British Standards Institution). 1975. *Methods of testing soils for civil engineering purpose*. British standards 1377. British standards institution, London.
- Buol, S.W., Southard, R.J. Graham, R.C. and McDaniel, P.A. 2003. *Soil Genesis and Classification*, Fifth edition. Iowa state University Press. Ames, USA.
- Chadwick, O.A. and Graham, R.C. 2000. Pedogenic processes. In: Sumner, M.E. (ed.). *Handbook of soil science*. CRC Press, Boca Raton. pp. 41-75.
- Chopra, S.H. and Kanwar, J.S. 1976. Analytical agricultural chemistry. Kalyani Publisher Ludhiana, New Delhi, Indian. p. 518. En. wikipedia.org/wiki/Soil_pH. (Accessed on 25/12/2007)
- EMA (Ethiopian Mapping Authority). 1988. National Atlas of Ethiopia, Addis Ababa, Ethiopia.
- FAO (Food and Agricultural Organization). 1984. Agro climatic data for Africa.V.1. Countries North of the equator. FAO Plant Production and Protection series No.22. Rome.
- FAO (Food and Agricultural Organization). 1990. *Guidelines for Soil Profile Description*. 3rd edition. Soil Resources, Management and Conservation Service, Land and water Development Division, FAO, Rome.
- FAO/WRB (Food and Agricultural Organization/World Resource Base for Soil Resources). 2006. World Resources Report No. 103. 2nd Edition. FAO, Rome, Italy.
- Fikre, M. 2003. Pedogenesis of major volcanic soils of the south central rift valley Region, Ethiopia. MSc Thesis. University of Saskatchewan. Saskatoon, Canada. pp. 128.
- Foth, H.D. 1990. *Fundamentals of Soil Science*. 12th edition. John Willey and Sons .New York. Grunwald.ifas.ufl.edu/Nat_resources/organic_matter/organic.htm (Accessed on 02/08/2009).
- Havlin, J.L., Beaton, J.D. Tisdale, S.L. and Nilson, W.L. 1999. *Soil Fertility and Fertilizers: An Introduction to Nutrient Management*. 6th edition. Prentice hall. Upper Saddle River, New Jersey.
- Irvin, B.J. 1996. Spatial Information Tools for Delineating Landform Elements to Support Soil/Landscape Analysis. PhD Thesis, University of Wisconsin-Madison. grunwald.ifas.ufl.edu/Nat_resources/soil_formation_factors/formation.htm (Accessed on 05/12/2007).
- Jones, J.B. 2003. *Agronomic Handbook: Management of Crops, Soils and Their Fertility*. CRC Press LLC, Bxa Ratan, FL, USA.
- Kravchenko, A.N. and Bullock, D.G. 2000. Correlation of corn and soybean grain yield with topography and soil properties. *Argon. J.* 92: 75-83.
- Kravskoof, K.B. 1972. Geochemistry of micronutrients. In: Mortued, J.J. (ed.). *Soil Sci. Soc. Am. Proc*, Madison, Wisconsin, USA. pp. 7-40.
- Landon, J.R. 1991. Booker tropical soil manual: a handbook for soil survey and agricultural land evaluation in the tropics and subtropics. Long man scientific and technical. Booker Tate Ltd. John Wiley and Sons, Inc., New York.
- Lawrence, W. M.1992. The variation of soil erodibility with slope position in a cultivated Canadian prairie landscape. *Earth Surface Processes and Landforms*. Vol. 17, Issue 6, Wiley and Sons, Ltd. pp. 543-556 .
- Mekaru, T. and Uehara, G. 1972. Anion adsorption in ferruginous tropical soils. *Soil Sci. Soc. Am. Proc.* 36: 296-300.
- Mesfin, A. 1998. Nature and management of Ethiopian soils. Alemaya University of Agriculture. Alemaya, Ethiopia.
- Metson, A.J. 1961. *Methods of Chemical Analysis for Soil Survey Samples*. New Zealand DSIR Soil Bur Bull 12. Govt Printer, Wellington, New Zealand.
- Miller, P.M., Singer, M.J. and Nielsen, D.R. 1998. Spatial variability of wheat yields and soil properties on complex hills. *Soil Sci. Soc. Am. J.* 52: 1133-1141.
- Moore, I.D., Gessler, P.E., Nielsen, G.A. and Peterson, G.A. 1993. Soil attributes prediction using terrain analysis. *Soil Sci. Soc. Am. J.* 57: 443-452.
- Rust, R.H. 1983. Alfisols. In: Wilding, L.P., Smeck, N.E. and Hall, G.F. (eds.). *Pedogenesis and Soil Taxonomy II*. The soil orders. Elsevier Science Publishers, Amsterdam. pp. 253-281.
- Sahlemedhin, S. and Taye, B. 2000. Procedure for soil and plant analysis. National Soil Research Centre, Ethiopian Agricultural Research Organization, Addis Ababa, Ethiopia. p. 110.
- SAS .1997. SAS Institute Inc., Cary. NC. USA.
- Soil Survey Staff. 1999. *Soil Taxonomy: a basic system of classification for making and Interpreting soil surveys*, USDA-NRCS. US Government printing office, Washington, DC.
- Tan, K.N. 1996. *Soil Sampling, Preparation and Analysis*. Marcel Dekker, Inc, New York.
- Van Ranst, E., Verloo, M., Demeyer, A. and Pauwels, J.M. 1999. *Manual for the soil chemistry and fertility laboratory: analytical methods for soils and plants, equipment and management of consumables*. University of Gent, Belgium.

- Van Reeuwijk. 1993. *Procedures for Soil Analysis*. 4th Edition. International soil reference and Information center, Netherlands. [Weather.nmsu.edu/teaching_Material/soil252/Chapt5.htm](http://www.weather.nmsu.edu/teaching_Material/soil252/Chapt5.htm) (Accessed on 05/12/2007).
- Van Wambeke, A. 2003. Properties and Management of Soils of the Tropics. Food and Agriculture Organization (FAO) of the United Nations.
- Wakene, N. and Heluf, G. 2004. The impact of different land use systems on soil quality of western Ethiopia Alfisols. International Research on Food Security: Natural Resource Management and Rural Poverty Reduction through Research for Development and Transformation. Deutcher Tropentage-Berlin 5-7 October 2004. pp. 1-7. <http://www.Tropentage.de/2004/abstracts/full/265.pdf>
- Wang, J., Bojie, F., Yang, Q. and Liding, C. 2000. Soil nutrients in relation to land use and landscape position in the semi-arid small catchments on the loess plateau in China. Department of Systems Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, China. *J. Arid. Environ.* Online at <http://www.ideallibrary.com>,48:537-550.