

Characterization of a novel wood mouse virus related to murid herpesvirus 4

David J. Hughes¹, Anja Kipar², Steven G. Milligan³, Charles Cunningham³, Mandy Sanders⁴, Michael A. Quail⁴, Marie-Adele Rajandream⁴, Stacey Efstathiou⁵, Rory J. Bowden³, Claude Chastel⁶, Malcolm Bennett², Jeffery T. Sample⁷, Bart Barrell⁴, Andrew J. Davison³ and James P. Stewart^{1*}

¹ School of Infection and Host Defence, University of Liverpool, Liverpool L69 3GA, UK

²Department of Veterinary Pathology, University of Liverpool, Liverpool, L69 7ZJ, UK

³MRC Virology Unit, Institute of Virology, University of Glasgow, Church Street, Glasgow G11 5JR, UK

⁴The Wellcome Trust Sanger Institute, The Wellcome Trust Genome Campus, Cambridge, CB10 1SA, UK

⁵Division of Virology, Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge CB2 1QP, UK

⁶Laboratoire de Virologie, Faculté de Médecine, 29285 Brest, France

⁷Department of Microbiology and Immunology, The Pennsylvania State University College of Medicine, Hershey, PA 17033, USA

* **Corresponding author:** James P. Stewart, School of Infection and Host Defence, University of Liverpool, Liverpool L69 3GA, UK. Tel: +44 151 706 4381. E-mail: j.p.stewart@liv.ac.uk.

Present address: David J. Hughes, Department of Microbiology and Immunology, The Pennsylvania State University College of Medicine, Hershey, PA 17033, USA

Present address: Steven G. Milligan, Infection and Immunity Section, Level 9, University of Glasgow Dental School, 378 Sauchiehall Street, Glasgow G2 3JZ, UK

Present address: Rory J. Bowden, Oxford Centre for Gene Function, Department of Statistics, 1 South Parks Road, Oxford, OX1 3TG, UK

Words in summary: 206

Words in main text: 5425

Figures and Tables: 8 + 1 Supplementary Table

Contents category: Animal Viruses – Large DNA

Running title: Wood mouse herpesvirus

The DNA sequences of WMHV and BRHV were deposited in GenBank under accession numbers GQ169129 and EF495130, respectively.

1 **SUMMARY**

2 We have isolated two novel gammaherpesviruses, one from a field vole (*Microtus agrestis*)
3 and the other from wood mice (*Apodemus sylvaticus*). The genome of the latter, wood
4 mouse herpesvirus (WMHV), was completely sequenced. WMHV had the same genome
5 structure and predicted gene content as murid herpesvirus 4 (MuHV4; murine
6 gammaherpesvirus 68). Overall nucleotide sequence identity between WMHV and MuHV4
7 was 85 % and most of the 10 kbp region at the left end of the unique region is particularly
8 highly conserved, especially the viral *tRNA*-like sequences and the coding regions of genes
9 *M1* and *M4*. The partial sequence (71913 bp) of another gammaherpesvirus, Brest
10 herpesvirus (BRHV), which was isolated ostensibly isolated from a white-toothed shrew
11 (*Crocidura russula*), was also determined. The BRHV sequence was 99.2 % identical to the
12 corresponding portion of the WMHV genome. Thus, WMHV and BRHV appear to be strains
13 of a new virus species. Biological characterization of WMHV indicated that it grows with
14 similar kinetics to MuHV4 in cell culture. The pathogenesis of WMHV in wood mice was also
15 extremely similar to that of MuHV4, except for the absence of inducible bronchus-associated
16 lymphoid tissue at day 14 post infection and a higher load of latently infected cells at 21 days
17 post infection.

18

19 INTRODUCTION

20 The most extensively characterized members of the family *Herpesviridae* that have hosts in
21 the family Muridae are mouse cytomegalovirus (MCMV) and rat cytomegalovirus (RCMV),
22 which are classified in the genus *Muromegalovirus* of the subfamily *Betaherpesvirinae*, and
23 murid herpesvirus 4 (MuHV4; also known as murine gammaherpesvirus 68, often
24 abbreviated to MHV-68 or γ HV-68; species *Murid herpesvirus 4*), classified in the genus
25 *Rhadinovirus* of the subfamily *Gammaherpesvirinae* (Davison *et al.*, 2009; Efstathiou *et al.*,
26 1990). Species in the genus *Rhadinovirus* also represent five herpesviruses of primates and
27 one of ungulates. However, these viruses are not closely related to MuHV4, and the best
28 estimate is that the lineages within the genus diverged approximately 60 million years ago
29 (McGeoch *et al.*, 2005). At least three other murid herpesviruses have been reported,
30 though these are unclassified at present (Davison *et al.*, 2009).

31
32 MuHV4 was originally isolated from bank voles (*Myodes glareolus*) and yellow-necked field
33 mice (*Apodemus flavicollis*) in Slovakia (Blaskovic *et al.*, 1980); reviewed in (Nash *et al.*,
34 2001). An epidemiological survey of MuHV4 infection in free-living rodents in the UK
35 (Blasdell *et al.*, 2003) showed that MuHV4 is endemic in wood mice (but not bank voles),
36 indicating that the wood mouse is a major natural host for this virus. Recent definitive
37 molecular data has also shown that MuHV4 is present in free-living yellow-necked field and
38 wood mice (Ehlers *et al.*, 2007).

39
40 In consideration of these observations, a wood mouse infection model was developed as an
41 alternative to a model utilizing the laboratory (house) mouse (*Mus musculus*), which has
42 been used to date for MuHV4 studies (Hughes *et al.*, submitted). In comparison with the
43 BALB/c laboratory mouse, the features of MuHV4 infection in the wood mouse are: (i) after
44 intranasal inoculation, viral titres achieved in the lung are approximately 1000-fold lower; (ii)
45 replication is restricted to scattered alveolar epithelial cells and macrophages within focal
46 granulomatous infiltrations, rather than being evident as a diffuse, T-cell dominated

47 interstitial pneumonitis; (iii) latently infected lymphocytes are abundant in inducible
48 bronchus-associated lymphoid tissue (iBALT); (iv) the spleens of wood mice show reduced
49 splenomegaly and leukocytosis; (v) well-delineated secondary follicles with classical
50 germinal centres are formed; and (vi) titres of neutralizing antibody to MuHV4 are
51 significantly higher.

52

53 The present study focuses on the isolation and genetic and biological characterization of a
54 novel, MuHV4-like virus. Two independent strains were examined, one isolated in the
55 present study from wood mice in Cheshire, UK and the other from a white-toothed shrew
56 (*Crocidura russula*) in Brest, France (Chastel *et al.*, 1994).

57

58 RESULTS

59 Genomic characterization of WMHV and BRHV

60 Three distinct viruses were obtained from free-living murids captured in Cheshire, UK. The
61 isolation and TEM results are summarized in Table 1.

62 The TEM-positive samples gave rise to PCR products from the DPOL gene, and the TEM-
63 negative samples did not (data not shown). The sequences of the 213-bp amplicons
64 originating from WM1, WM2, WM7 and WM8 DPOL were identical to each other, regardless
65 of the tissue from which the viruses were isolated. These sequences (minus the primers;
66 160 bp) exhibited 89 % nucleic acid identity and 94 % predicted amino acid sequence
67 identity to the corresponding region of MuHV4 DPOL. The FV1 DPOL amplicon was also
68 213 bp in size, and the 160-bp sequence (minus primers) was more closely related to
69 MuHV4 than any other herpesvirus, at 61 % nucleic acid and 54 % amino acid sequence
70 identity. The HM4 virus DNA yielded a DPOL PCR product of 231 bp (178 bp minus primers)
71 that was closely related to MCMV (strain Smith) DPOL, at 99 % nucleic acid and 100 %
72 amino acid sequence identity. These results confirm the identification of two novel
73 gammaherpesviruses. The WM isolates were designated WMHV, and the FV1 isolate was
74 designated field vole herpesvirus (FVHV).

75

76 An initial analysis of the coding regions of genes *M1*, *M2* and *M3*, which had been identified
77 hitherto only in MuHV4, showed that cognate PCR products were detected in all WMHV
78 isolates using MuHV4-specific primers, thus confirming the close relatedness of these
79 isolates to MuHV4 (Fig. 1). In contrast, none of these genes were amplified from FVHV,
80 consistent with its more distant relationship to MuHV4. The DNA sequences obtained from
81 these PCR products for WMHV *M1* and *M3* were 97 % and 94 % identical to MuHV4 *M1* and
82 *M3*, respectively, while WMHV and MuHV4 *M2* sequences were more divergent, sharing
83 only 83 % identity.

84

85 The complete nucleotide sequence of the WMHV genome was then determined. The

86 genome structure deduced from the WMHV genome sequence is the same as that of
87 MuHV4, consisting of a unique region (U) flanked at both ends by multiple direct repeats of a
88 terminal repeat (TR). In WMHV and MuHV4, the size of U is 118864 and 118211 bp,
89 respectively, and that of TR is 1244 and 1240 bp. Overall nucleotide sequence identity is 85
90 %. The predicted gene content of WMHV is the same as that for MuHV4, as represented by
91 the most up-to-date annotation (NC_001826). A 71913-bp segment of the BRHV genome
92 was sequenced. This represented TR (1265 bp; plus a partial copy) linked to the left portion
93 of U (70439 bp) terminating within *ORF53*. The BRHV sequence is 86 % and 99.2 %
94 identical to the corresponding portion of the MuHV4 and WMHV genomes, respectively. The
95 information used to annotate the genome sequences is shown in Supplementary Table S1.

96

97 Fig. 2 shows a representation of DNA sequence identity along the entire WMHV and MuHV4
98 genomes, and Fig. 3 provides detail on amino acid sequence identity between WMHV,
99 BRHV and MuHV4 protein-coding regions. The most highly conserved regions between
100 WMHV and MuHV4 include two sets of internal tandem repeats and the region from the left
101 end of U to the end of the *M4* coding region, which includes the eight viral tRNA-like genes
102 (*vtRNAs*) (Bowden *et al.*, 1997). Although the *vtRNAs* are well conserved, there are
103 functionally relevant differences in the sequences of micro-RNAs (*miRNAs*) 1, 2, 5, 6 and 9
104 that are derived from *vtRNA* primary transcripts (Pfeffer *et al.*, 2004)(Fig. 4). The most highly
105 conserved coding regions at the amino acid sequence level are *M4* (98.3 %), *ORF43* (97.2
106 %), *M1* (96.9 %) and *ORF60* (96.7 %), and the least conserved are *ORF73* (67.0 %), *ORF51*
107 (68.0%) and *M2* (72.8 %). In the comparable sequences, the most highly conserved coding
108 regions at the amino acid sequence level between WMHV and BRHV are *ORF8*, *ORF28*,
109 *ORF29*, *ORF34*, *ORF43*, *ORF44* and *ORF46* (each 100 %), and the least conserved are
110 *ORF45* (95.2 %), *ORF51* (95.8 %) and *M2* (97.4 %). Fig. 5 shows the amino acid sequence
111 alignments for *M1*, *M3* and *M4* (which are related), and also *M2*, *ORF51* and *ORF73*.

112

113 **Biological characterization of WMHV**

114 The relative rates of growth of MuHV4 and WMHV in NIH3T3 cells were compared by
115 determining a one-step growth curve as described previously (Macrae *et al.*, 2001), and
116 were not significantly different (data not shown).

117

118 Fig. 6 shows the features of infection of wood mice by WMHV in comparison with MuHV4.
119 Infectious virus was detected in the lungs of wood mice infected with WMHV or MuHV4 at 7
120 d PI, but not at 14 d PI, and a significantly greater amount of infectious virus was recovered
121 from MuHV4-infected wood mice at 7 d PI (Fig. 6a). The numbers of leukocytes per spleen
122 isolated from infected wood mice were similar for WMHV and MuHV4 at all time points PI
123 (Fig. 6b). There was an increase in the number of leukocytes at 14 d PI with both viruses,
124 but this was marginal and transient, and infection with neither virus induced significant
125 splenomegaly. In WMHV- and MuHV4-infected wood mice, the number of latently infected
126 cells per spleen increased dramatically from 7 d PI, peaking at 14 d PI (Fig. 6c); the mean
127 number of latently infected cells then declined approximately six-fold by 21 d PI in WMHV-
128 infected animals and twenty-five-fold in the MuHV4-infected mice, and was largely
129 unchanged at 28 d PI in both infections. The difference observed at 21 d PI was statistically
130 significant ($P < 0.05$).

131

132 Histological examination identified broadly similar changes in both WMHV- and MuHV4-
133 infected wood mice and similar to those reported in detail previously (D.J. Hughes *et al.*
134 Submitted). On day 7 PI, mild to moderate perivascular or peribronchial, B cell dominated
135 lymphocyte infiltration with evidence of B cell emigration from blood vessels was seen
136 together with multifocal, predominantly perivascular macrophage and lymphocyte (i.e.,
137 granulomatous) infiltrates. There was a mild to moderate increase in the number of
138 disseminated T and B cells in the interstitium. Viral antigen was scarce and seen in
139 occasional alveolar epithelial cells (type I and II pneumocytes) and in macrophages within
140 the granulomatous infiltrates. The mediastinal lymph nodes and spleens of these animals

141 contained primary and secondary follicles and unaltered T cell zones. Rare virus antigen-
142 positive macrophages were seen in the lymph nodes. At 14 d PI, wood mice infected with
143 MuHV4 displayed intense perivascular and peribronchial, B cell rich lymphocyte infiltration
144 with evidence of lymphatic follicle formation (Fig. 7a, b). This has been described previously
145 as being iBALT (D.J. Hughes, submitted). In contrast, in WMHV-infected animals moderate
146 multifocal perivascular and peribronchial B cell infiltration and emigration was seen, but
147 without distinct evidence of follicle formation (Fig. 7c, d). Granulomatous infiltrates were still
148 observed in both groups; these contained macrophages exhibiting viral antigen. Large, well
149 delineated secondary follicles were observed in the spleens, and virus antigen-positive
150 macrophages were detected in the red pulp. At 20 d PI, both granulomatous infiltrates and
151 perivascular and peribronchial lymphocyte infiltrations were still observed in the lungs.
152 However, the follicle formation that was previously seen in MuHV4-infected wood mice had
153 subsided. Spleens exhibited smaller secondary follicles than at day 14 PI. Virus antigen-
154 laden macrophages were seen in the spleen. In the lung, however, viral antigen expression
155 was restricted to one individual macrophage in a granulomatous infiltrate in a MuHV4-
156 infected animal. By day 28 PI, the granulomatous infiltrates were few in number and small,
157 but a mild to moderate perivascular and peribronchial lymphocyte infiltration remained. This
158 persisted until day 46 d PI, to a mild degree. Thus, the changes observed were extremely
159 similar except for the less intense B cell infiltration and absence of iBALT in the lungs of
160 WMHV-infected mice at day 14 PI.

161

162

163

164 **DISCUSSION**

165 This study demonstrated the abundance of herpesviruses in natural populations of wood
166 mice in Cheshire. Two novel gammaherpesviruses (FVHV and WMHV) were isolated.
167 Previous analyses have concluded that herpesvirus genomes of less than 95 % nucleotide
168 sequence identity may represent different species (Ehlers *et al.*, 2007). Thus, the degree of
169 divergence between WMHV and MuHV4, both overall (85 %) and within specific loci (e.g.,
170 *M2* and *ORF73*) (Figs. 2, 3), is probably sufficient to warrant classification of WMHV as a
171 new species. Under the current taxonomic scheme, in which murid herpesvirus species are
172 named after the host family, this would be *Murid herpesvirus 7*. Although an epidemiological
173 study of free-living rodents in the UK was unable to distinguish between the two viruses
174 (Blasdell *et al.*, 2003), a PCR-based study of mice trapped in Germany (Ehlers *et al.*, 2007)
175 indicated that MuHV4 is present predominantly in yellow-necked field mice (*Apodemus*
176 *flavicollis*), whereas WMHV is present in wood mice (*A. sylvaticus*). However, MuHV4 was
177 detected in some wood mice. Thus, it is possible that the two viruses normally infect different
178 *Apodemus* species, but that there is some crossover. The biological characteristics of the
179 two viruses in the wood mouse model exhibit significant similarities. However, the viruses do
180 differ in their ability to grow in the lungs, in the development of iBALT, and also perhaps in
181 the efficiency of reactivation from splenic leukocytes. Interestingly, WMHV was isolated from
182 trigeminal ganglia as well as spleens, suggesting that this virus may be neurotropic during a
183 natural infection. This hypothesis warrants further investigation.

184

185 The analysis of the sequence of a large portion of the genome of the BRHV genome showed
186 that the relationship of this virus to WMHV is sufficiently close (99 % identity) to warrant the
187 consideration of WMHV and BRHV as strains of the same virus. Given that herpesviruses
188 are thought generally to have evolved with their hosts (McGeoch *et al.*, 2006), this
189 relationship was unanticipated, since the wood mouse and white-toothed shrew are
190 classified in different mammalian orders, *Rodentia* (family *Muridae*) and *Insectivora* (family
191 *Soricidae*), respectively. Thus, the claimed insectivore source of BRHV must be viewed as

192 questionable. It is possible that the virus actually originated from a rodent, either by cross-
193 infection in the wild or by laboratory contamination, since BRHV was isolated by passage in
194 suckling mouse brains (Chastel *et al.*, 1994).

195

196 Other viruses related to MuHV4 have been characterized, but none thus far has been shown
197 to be sufficiently divergent from MuHV4 to form a new species. Viruses isolated from bank
198 voles or yellow-necked field mice at the same time as MuHV4 (MHV-76, MHV-72, MHV-60
199 and MHV-78) are considered to be strains of MuHV4. MHV-76, although originally
200 characterized as a novel alphaherpesvirus due to its cytopathic effect *in vitro* (Ciampor *et al.*,
201 1981; Svobodova *et al.*, 1982) and then as a betaherpesvirus (Hamelin & Lussier, 1992),
202 was conclusively demonstrated to be a gammaherpesvirus (Macrae *et al.*, 2001). MHV-76
203 proved to be equivalent to MuHV4 with a 9538-bp deletion at the left end of U, which
204 probably arose during passage of the virus *in vivo* or *in vitro*. MHV-72 *ORF21* (encoding
205 thymidine kinase) is identical in sequence to the corresponding MuHV4 gene (Raslova *et al.*,
206 2000), and *ORF51* (encoding gp150) differs by five nucleotide substitutions (Macakova *et al.*
207 *et al.*, 2003). Analysis of 12 other loci has shown that MHV-72 is more divergent from MuHV4
208 than MHV-76, and that *M1*, *M2* and *M3* are absent; nonetheless, MHV-72 and MuHV4 are
209 highly related (Oda *et al.*, 2005). It seems likely that uncharacterized herpesviruses (MHV-60
210 and MHV-78) isolated at the same time as MuHV4 may also be strains of MuHV4
211 (Mistriková *et al.*, 2000; Nash *et al.*, 2001).

212

213 The WMHV genome is co-linear with that of MuHV4, and the two viruses have the same
214 predicted gene content (Figs. 2, 3). The reason for the generally higher degree of
215 conservation of sequences near the left end of U is not known. Speculative explanations
216 could centre on selective sweeps in this region of the genome or recombination between a
217 WMHV-like virus and a virus more closely related to MuHV4. The noncoding sequences in
218 this region, including the *vtRNA*-like transcripts, are generally highly conserved. However,
219 there are differences in *miRNAs* 1, 2, 5, 6 and 9 that are derived from the primary transcripts

220 of these *vtRNAs* (Pfeffer *et al.*, 2004) (Fig. 4) . The targets and exact functions of these
221 *miRNAs* are not currently known (Pfeffer *et al.*, 2004), but these differences could have
222 functional consequences and the comparative data could be informative. In addition to
223 noncoding regions in this locus, the M1 and M4 proteins are highly conserved (Fig. 5). It has
224 been proposed that the most likely function for the M4 protein is as a modulator of the innate
225 immune system. *M4* is expressed *in vitro* with kinetics similar to immediate-early genes
226 (Ebrahimi *et al.*, 2003), and *in vivo* it is expressed during productive infection but not during
227 latency (Virgin *et al.*, 1999). M4 does not appear to have a role during the initial stages of
228 infection *in vivo*, but is important during establishment of latency in the spleen (Evans *et al.*,
229 2006; Geere *et al.*, 2006). M1 has been shown to stimulate a V β 4⁺ CD8⁺ T cell in a way
230 reminiscent of a superantigen and by doing this facilitate latent infection (Evans *et al.*, 2008).

231

232 The M3 protein, which is related to M1 and M4, is also well conserved, but somewhat less
233 so than the M1 and M4 proteins, particularly towards the N terminus (Fig. 2). The secreted
234 M3 protein is expressed strongly during lytic infection and probably to a lesser extent during
235 latency (Simas *et al.*, 1999; Usherwood *et al.*, 2000; van Berkel *et al.*, 1999; Virgin *et al.*,
236 1999). *In vitro*, the M3 protein selectively binds chemokines associated with the antiviral
237 inflammatory response (Parry *et al.*, 2000; van Berkel *et al.*, 2000). In the laboratory mouse,
238 M3 was found to have a role in enhancing the amplification of latently-infected B cells by
239 affecting the CD8⁺ T cell response (Bridgeman *et al.*, 2001), although this function was not
240 seen in an independent study (van Berkel *et al.*, 2002). In the wood mouse model, M3 has a
241 critical role in the amplification of latently-infected B cells in the lung and the formation of
242 iBALT containing these cells (Hughes *et al.*, submitted for publication). Differences in M3
243 may therefore account for the lack of iBALT in WMHV-infected wood mice.

244

245 The M2 protein is the most divergent of the four proteins encoded by the left end of the
246 genome, and is associated with latency (Husain *et al.*, 1999). Numerous reports largely
247 agree that *M2* is dispensable for long-term persistence, although MuHV4 recombinants

248 lacking a functional *M2* gene are less efficient in the establishment of latency following
249 intranasal infection of mice (Clambey *et al.*, 2002; Jacoby *et al.*, 2002; Macrae *et al.*, 2003;
250 Simas *et al.*, 2004). It has also been postulated that *M2* is required for efficient colonization
251 of follicle B cells and the development of these cells into memory B cells, a cell type
252 exploited by MuHV4 for long-term latency (Simas *et al.*, 2004). Given the relationship, and
253 possible overlap, between the hosts of WMHV and MuHV4, the divergence of the *M2* gene
254 in a region of low overall variation might reflect strong immune selection. Indeed, it has been
255 shown that an H2-Kd–restricted CD8+ T cell epitope present in *M2* (Husain *et al.*, 1999) sets
256 the latent load during persistent infection of laboratory mice (*M. musculus*) (Marques *et al.*,
257 2008). However, this epitope is not conserved between MuHV4 and WMHV (Fig. 5b),
258 suggesting that it may not be functional in the *Apodemus* hosts. The generation of greater
259 numbers of infective centres (a measure of latency) in the spleens of the WMHV-infected
260 wood mice at 21 d PI (Fig. 6c) raises the possibility that *M2* may have evolved in this virus to
261 augment the expansion of latently infected cells during the acute phase of latency.
262 Experiments to address this hypothesis could involve replacing MuHV4 *M2* with WMHV *M2*
263 and testing the phenotype in wood mice. Furthermore, numerous PXXP motifs are found
264 throughout MuHV4 *M2* (labelled P1-9, Fig.6c), some of which have been shown to
265 functionally bind SH3-domain containing proteins, such as Vav1 (Madureira *et al.*, 2005;
266 Rodrigues *et al.*, 2006). Of these, P3, P4 and P5 have not been conserved in WMHV or
267 BRHV. Recent *in vivo* analysis showed that mutations of P3, P4 or P5 had no affect on the
268 establishment of, or reactivation from, splenic latency (Herskowitz *et al.*, 2008). Taken
269 together, these motifs are unlikely to be important for *M2*'s signalling function. In a similar
270 vein, the tyrosine residues at positions 120 and 129 of *M2*, which have been proven to be
271 functional (Herskowitz *et al.*, 2008; Pires de Miranda *et al.*, 2008), are conserved in both
272 WMHV and BRHV, highlighting their importance for *M2*'s signalling function.

273

274 The second most divergent protein in WMHV and MuHV4 is the virion glycoprotein gp150,
275 which is encoded by *ORF51*. It seems likely that, in addition to exhibiting extensive

276 differences in amino acid sequence, these proteins are predicted to be N-glycosylated
277 differently in the two viruses (Fig. 5c). Gp150 is a major target for the host antibody
278 response, so it is likely to be under strong selective pressure (Gillet *et al.*, 2007). However, it
279 is not clear why this membrane glycoprotein is more variable than others encoded by the
280 two viruses.

281

282 The most divergent protein in WMHV and MuHV4 is encoded by *ORF73* (Fig. 5d). *In vivo*
283 analyses of an MuHV4 mutant have shown that *ORF73* is essential for the establishment
284 and maintenance of latency (Fowler *et al.*, 2003), and preliminary characterization of *ORF73*
285 mRNAs suggests that their transcription is similar to that of KSHV *ORF73* encoding the
286 protein LANA (Coleman *et al.*, 2005). In a similar way to KSHV-LANA, the MuHV4 *ORF73*
287 protein interacts with cellular bromodomain-containing BET proteins leading to activation of
288 the promoters of G₁/S cyclins (Ottinger *et al.*, 2009). The reason for the sequence variability
289 in *ORF73* is not clear. However, EBV EBNA1 (the functional analogue of rhadinovirus
290 *ORF73* proteins) shows considerable variability between strains (Wright *et al.*, 1995),
291 and this has consequences for EBV-associated disease (Mai *et al.*, 2007; Wang *et al.*,
292 2003), the function of EBNA1 as a transcriptional transactivator (Do *et al.*, 2008) and the
293 CD8⁺ T cell response (Bell *et al.*, 2008).

294

295 In summary, WMHV is a novel MuHV4-like virus whose study will give further insight into
296 gammaherpesvirus biology, especially in comparative terms alongside MuHV4.

297

298

299 **METHODS**

300 **Cheshire herpesviruses**

301 **Isolation and growth.** Eight wood mice (WM1 to WM8), a bank vole (BV1), a field vole
302 (*Microtus agrestis*; FV1) and six house mice (HM1 to HM6) were captured in Cheshire
303 during 2002. The animals were killed by cervical dislocation and trigeminal ganglia, lungs
304 and spleens were removed for virus reactivation. Virus was reactivated from trigeminal
305 ganglia by explant culture as described previously (Efstathiou *et al.*, 1986). Virus was
306 reactivated from the spleen by using an infectious centre assay (Sunil-Chandra *et al.*, 1992).
307 Lung tissue was homogenized and virus recovered as described previously (Stewart *et al.*,
308 1998). Mouse NIH3T3 cells (Todaro & Green, 1963) were used for all virus isolation
309 experiments. Supernatants were examined as negatively stained preparations by
310 transmission electron microscopy (TEM).

311

312 **Preliminary sequence analysis.** Samples were tested for the presence of herpesvirus DNA
313 polymerase gene (DPOL) sequences by PCR. Whole-cell DNA from NIH3T3 cells was
314 purified at 18-24 h post infection (PI) using a QIAamp DNA Mini Kit (Qiagen). PCR was
315 carried out using the degenerate, deoxyinosine-substituted primers 5'-
316 TGTAACCTCGGTGTAYGGITTYACIGGIGT-3' and 5'-CACAGAGTCCGTRTCICCRTAIAT-3'
317 (Ehlers *et al.*, 1999). PCR products were inserted into pCR2.1topo (Invitrogen Life
318 Technologies) and inserts were sequenced from three individual clones per product by Lark
319 Technologies Inc., UK. Amino acid sequences deduced from the sequences of the PCR
320 products were compared to known herpesvirus DPOL sequences using BLAST (Altschul *et*
321 *al.*, 1997).

322 To amplify protein-coding DNA from genes *M1*, *M2* and *M3*, the samples were subjected to
323 PCR using primers M1-f/M1-r (5'-TCATTGAGCAGCGGCGAC-3' and 5'-
324 GTATTCAGGCTTAGGACTG-3'; 1292 bp), M2-f/M2-r (5'-ATGGCCCCAACACCCCCAC-3'
325 and 5'-ACTCCTCGCCCCACTCCAC-3'; 577 bp) and M3-f/M3-r (5'-
326 CTCTGGGAGAGCGTCAG-3' and 5'-GTTACTGAGTATCAATGATCC-3'; 1251 bp),

327 respectively. PCR products were sequenced as described above. The sequences obtained,
328 minus those of the primers, accounted for the entire protein-coding region of each gene but
329 for a few codons at one or both ends.

330

331 **Genome sequence analysis.** Virus isolated from the WM8 spleen was plaque-purified three
332 times from infected NIH3T3 cells overlaid with agarose, and a master stock of cell-
333 associated virus was prepared and titrated. For the purposes of the present study, this virus
334 was designated wood mouse herpesvirus (WMHV). WMHV was found to be primarily cell-
335 associated in culture. To prepare virions for DNA extraction, WMHV was grown on a mouse
336 cell line ($\alpha\beta$ SV1) deficient in the response to α/β interferon. This line was derived by first
337 generating mouse embryonic fibroblasts (Todaro & Green, 1963) from interferon α/β
338 receptor knockout mice (Muller *et al.*, 1994). These cells were then transformed by
339 transfection with a plasmid expressing SV40 T antigen (pVU0) (Kalderon *et al.*, 1982) to
340 generate an immortal cell line. The resulting cell line was found in preliminary experiments to
341 release a much higher level of cell-free virus.

342

343 Twenty 150-cm² tissue culture flasks of sub-confluent $\alpha\beta$ SV1 cells were infected with WMHV
344 at an MOI of 0.01 for 7 d. Virus DNA was then purified as described (Baldick *et al.*, 1997)
345 and its integrity confirmed by agarose gel electrophoresis.

346

347 The DNA was sequenced at the Wellcome Trust Sanger Institute by a standard random
348 shotgun approach to an average coverage of 12 reads per nucleotide. Tandem repeat
349 regions in the genome were determined using the program MREPS (Kolpakov *et al.*, 2003),
350 and the genome ends were inferred by comparison with the MuHV4 sequence (U97553;
351 (Virgin *et al.*, 1997). The main computer programs used to analyse the sequence were: for
352 sequence annotation, Artemis (Rutherford *et al.*, 2000), ACT (Carver *et al.*, 2005), and
353 Sequin (NCBI); for sequence alignment, ClustalW (Thompson *et al.*, 1994) and Mafft (Katoh

354 & Toh, 2008); for DNA sequence analyses, GCG (Accelrys) and EMBOSS (Rice *et al.*,
355 2000); for amino acid sequence analysis, GCG, ExPASy (Gasteiger *et al.*, 2003), PTrans
356 (Taylor, 1986) and Philius (Reynolds *et al.*, 2008); and for similarity searches, BLAST and
357 FASTA and its relatives (Pearson & Lipman, 1988).

358

359 **Biological characterization.** The growth properties of WMHV were compared with those of
360 MuHV4 in laboratory-bred wood mice using the procedures described (Sunil-Chandra *et al.*,
361 1992). All animal work was performed under UK Home Office Project Licence number
362 40/2483 and Personal Licence number 60/6501.

363

364 Wood mice (*Apodemus sylvaticus*) were obtained from an out-bred colony established at the
365 Faculty of Veterinary Science, University of Liverpool (Bennett *et al.*, 1997; Feore *et al.*,
366 1997). This colony was obtained from Dr. J. Clarke in 1995, and was derived from captive-
367 bred colonies that had been maintained for several decades in the Department of Zoology,
368 University of Oxford, with only occasional introductions of new stock from the wild. Their
369 general housing and maintenance has been described elsewhere (Clarke, 1998), and at
370 Liverpool they are maintained under semi-barrier conditions. The Liverpool colony has
371 suffered no clinical disease, and, although not specified pathogen free (SPF) in the sense
372 used for most laboratory rodents, all samples tested for the major infections of laboratory
373 rodents have so far been negative. Of particular relevance to this study, no evidence of
374 MuHV4 infection has been detected by serology and PCR analysis (Blasdell *et al.*, 2003).
375 Both male and female wood mice of 5 – 8 weeks of age were used. They were infected
376 intranasally with 4×10^5 PFU of virus, and the lungs, spleens and bronchial lymph nodes
377 were harvested at various times PI. Lung tissue was homogenized, and the lysate was
378 freeze-thawed three times and used in plaque assays. Leukocytes were purified from the
379 spleens and counted, and virus reactivation was monitored using an infective centre assay.
380 Tissues from infected wood mice were routinely processed for histopathological
381 examination, including immunohistology.

382

383 **Brest herpesvirus**

384 ***Isolation and growth.*** The herpesvirus (Brest/AN711) isolated from a white-toothed shrew
385 (Chastel *et al.*, 1994) was grown and titrated on baby hamster kidney (BHK) cells as
386 described (Bridgeman *et al.*, 2001). For the purposes of the present study, this virus was
387 named Brest herpesvirus (BRHV). To prepare viral DNA, confluent monolayers of cells in
388 175-cm² flasks were infected at an MOI of 0.01. When CPE was complete at approximately
389 4 d PI, virions were purified from the medium by Ficoll gradient ultracentrifugation as
390 described (Lopes *et al.*, 2004). Banded virus was diluted with PBS to a total volume of 30 ml,
391 and pelleted at 30000 x *g* for 90 min. The pelleted virus was resuspended in TE buffer
392 containing 0.5 % (w/v) SDS and 50 µg ml⁻¹ proteinase K. The mixture was incubated
393 overnight at 37 °C and extracted with phenol, and the DNA precipitated in ethanol and
394 dissolved in a small volume of TE, as described above.

395

396 ***Preliminary sequence analysis.*** Initial cloning involved the generation of a small library of
397 bacteriophage M13 recombinants containing BRHV AluI fragments, using standard methods.
398 The inserts in three recombinants were sequenced, and found by BLAST similarity search to
399 be most closely related to the MuHV4 genome. Respectively, the insert sizes were 148, 145
400 and 156 bp and exhibited 89.2, 96.6 and 90.4 % nucleotide sequence identity to ORF18,
401 ORF31 and ORF60.

402

403 ***Partial genome sequence analysis.*** A cosmid library was generated from BRHV DNA as
404 described (Cunningham & Davison, 1993). Three overlapping cosmid clones constituting
405 approximately 70 kbp of the genome were sequenced, the first by a standard random
406 shotgun approach, and the other two by iterative primer-walking on both strands, based
407 initially on data generated from the first cosmid or arising from the preliminary sequence
408 analysis described above. The computer programs used for analysis are listed above.

ACKNOWLEDGEMENTS

This work was funded by the Medical Research Council (United Kingdom), the Wellcome Trust, U.S. Public Health Service grant CA090208, and the Penn State Hershey Cancer Institute. JPS was funded by a Royal Society (London) University Research Fellowship. Initial isolation experiments were funded by a Royal Society (London) Summer Studentship to DJH. TEM was performed by Brian Getty (School of Infection and Host Defence, Liverpool), and histology and immunohistology by the Histology Laboratory, Veterinary Laboratory Services, Faculty of Veterinary Science, University of Liverpool.

REFERENCES

- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997).** Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**, 3389-3402.
- Baldick, C. J., Jr., Marchini, A., Patterson, C. E. & Shenk, T. (1997).** Human cytomegalovirus tegument protein pp71 (ppUL82) enhances the infectivity of viral DNA and accelerates the infectious cycle. *J Virol* **71**, 4400-4408.
- Bell, M. J., Brennan, R., Miles, J. J., Moss, D. J., Burrows, J. M. & Burrows, S. R. (2008).** Widespread sequence variation in Epstein-Barr virus nuclear antigen 1 influences the antiviral T cell response. *J Infect Dis* **197**, 1594-1597.
- Bennett, M., Crouch, A. J., Begon, M., Duffy, B., Feore, S., Gaskell, R. M., Kelly, D. F., McCracken, C. M., Vicary, L. & Baxby, D. (1997).** Cowpox in British voles and mice. *J Comp Pathol* **116**, 35-44.
- Blasdell, K., McCracken, C., Morris, A., Nash, A. A., Begon, M., Bennett, M. & Stewart, J. P. (2003).** The wood mouse is a natural host for Murid herpesvirus 4. *J Gen Virol* **84**, 111-113.
- Blaskovic, D., Stancekova, M., Svobodova, J. & Mistrikova, J. (1980).** Isolation of five strains of herpesviruses from two species of free living small rodents. *Acta Virol* **24**, 468.
- Bowden, R. J., Simas, J. P., Davis, A. J. & Efstathiou, S. (1997).** Murine gammaherpesvirus 68 encodes tRNA-like sequences which are expressed during latency. *J Gen Virol* **78**, 1675-1687.
- Bridgeman, A., Stevenson, P. G., Simas, J. P. & Efstathiou, S. (2001).** A secreted chemokine binding protein encoded by murine gammaherpesvirus- 68 is necessary for the establishment of a normal latent load. *J Exp Med* **194**, 301-312.

- Carver, T. J., Rutherford, K. M., Berriman, M., Rajandream, M. A., Barrell, B. G. & Parkhill, J. (2005).** ACT: the Artemis Comparison Tool. *Bioinformatics* **21**, 3422-3423.
- Chastel, C., Beaucournu, J. P., Chastel, O., Legrand, M. C. & Le Goff, F. (1994).** A herpesvirus from an European shrew (*Crocidura russula*). *Acta Virol* **38**, 309.
- Ciampor, F., Stancekova, M. & Blaskovic, D. (1981).** Electron microscopy of rabbit embryo fibroblasts infected with herpesvirus isolates from *Clethrionomys glareolus* and *Apodemus flavicollis*. *Acta Virol* **25**, 101-107.
- Clambey, E. T., Virgin, H. W. t. & Speck, S. H. (2002).** Characterization of a spontaneous 9.5-kilobase-deletion mutant of murine gammaherpesvirus 68 reveals tissue-specific genetic requirements for latency. *J Virol* **76**, 6532-6544.
- Clarke, J. R. (1998).** Voles. In *UFAW Handbook on the Care and Management of Laboratory Animals*. Edited by T. B. Poole. Oxford: Blackwell Scientific.
- Coleman, H. M., Efstathiou, S. & Stevenson, P. G. (2005).** Transcription of the murine gammaherpesvirus 68 ORF73 from promoters in the viral terminal repeats. *J Gen Virol* **86**, 561-574.
- Cunningham, C. & Davison, A. J. (1993).** A cosmid-based system for constructing mutants of herpes simplex virus type 1. *Virology* **197**, 116-124.
- Davison, A. J., Eberle, R., Ehlers, B., Hayward, G. S., McGeoch, D. J., Minson, A. C., Pellett, P. E., Roizman, B., Studdert, M. J. & Thiry, E. (2009).** The order Herpesvirales. *Arch Virol* **154**, 171-177.
- Do, N. V., Ingemar, E., Phi, P. T., Jenny, A., Chinh, T. T., Zeng, Y. & Hu, L. (2008).** A major EBNA1 variant from Asian EBV isolates shows enhanced transcriptional activity compared to prototype B95.8. *Virus Res* **132**, 15-24.
- Ebrahimi, B., Dutia, B. M., Roberts, K. L., Garcia-Ramirez, J. J., Dickinson, P., Stewart, J. P., Ghazal, P., Roy, D. J. & Nash, A. A. (2003).** Transcriptome profile of murine gammaherpesvirus-68 lytic infection. *J Gen Virol* **84**, 99-109.

- Efstathiou, S., Ho, Y. M., Hall, S., Styles, C. J., Scott, S. D. & Gompels, U. A. (1990).** Murine herpesvirus 68 is genetically related to the gammaherpesviruses Epstein-Barr virus and herpesvirus saimiri. *J Gen Virol* **71**, 1365-1372.
- Efstathiou, S., Minson, A. C., Field, H. J., Anderson, J. R. & Wildy, P. (1986).** Detection of herpes simplex virus-specific DNA sequences in latently infected mice and in humans. *J Virol* **57**, 446-455.
- Ehlers, B., Borchers, K., Grund, C., Frolich, K., Ludwig, H. & Buhk, H. J. (1999).** Detection of new DNA polymerase genes of known and potentially novel herpesviruses by PCR with degenerate and deoxyinosine-substituted primers. *Virus Genes* **18**, 211-220.
- Ehlers, B., Kuchler, J., Yasmum, N., Dural, G., Voigt, S., Schmidt-Chanasit, J., Jakel, T., Matuschka, F. R., Richter, D., Essbauer, S., Hughes, D. J., Summers, C., Bennett, M., Stewart, J. P. & Ulrich, R. G. (2007).** Identification of novel rodent herpesviruses, including the first gammaherpesvirus of *Mus musculus*. *J Virol* **81**, 8091-8100.
- Evans, A. G., Moorman, N. J., Willer, D. O. & Speck, S. H. (2006).** The M4 gene of gammaHV68 encodes a secreted glycoprotein and is required for the efficient establishment of splenic latency. *Virology* **344**, 520-531.
- Evans, A. G., Moser, J. M., Krug, L. T., Pozharskaya, V., Mora, A. L. & Speck, S. H. (2008).** A gammaherpesvirus-secreted activator of Vbeta4+ CD8+ T cells regulates chronic infection and immunopathology. *J Exp Med* **205**, 669-684.
- Feore, S. M., Bennett, M., Chantrey, J., Jones, T., Baxby, D. & Begon, M. (1997).** The effect of cowpox virus infection on fecundity in bank voles and wood mice. *Proceedings of the Royal Society of London, Series B - Biological Sciences* **264**, 1457-1461.
- Fowler, P., Marques, S., Simas, J. P. & Efstathiou, S. (2003).** ORF73 of murine herpesvirus-68 is critical for the establishment and maintenance of latency. *J Virol* **84**, 3405-3416.

- Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D. & Bairoch, A. (2003).** ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res* **31**, 3784-3788.
- Geere, H. M., Ligertwood, Y., Templeton, K. M., Bennet, I., Gangadharan, B., Rhind, S. M., Nash, A. A. & Dutia, B. M. (2006).** The M4 gene of murine gammaherpesvirus 68 modulates latent infection. *J Gen Virol* **87**, 803-807.
- Gillet, L., May, J. S., Colaco, S. & Stevenson, P. G. (2007).** The murine gammaherpesvirus-68 gp150 acts as an immunogenic decoy to limit virion neutralization. *PLoS One* **2**, e705.
- Hamelin, C. & Lussier, G. (1992).** Characterization of the DNA of rodent herpesviruses by restriction endonuclease analysis and hybridization. *Lab Anim Sci* **42**, 142-145.
- Herskowitz, J. H., Siegel, A. M., Jacoby, M. A. & Speck, S. H. (2008).** Systematic mutagenesis of the murine gammaherpesvirus 68 M2 protein identifies domains important for chronic infection. *J Virol* **82**, 3295-3310.
- Husain, S. M., Usherwood, E. J., Dyson, H., Coleclough, C., Coppola, M. A., Woodland, D. L., Blackman, M. A., Stewart, J. P. & Sample, J. T. (1999).** Murine gammaherpesvirus M2 gene is latency-associated and its protein a target for CD8(+) T lymphocytes. *Proc Natl Acad Sci USA* **96**, 7508-7513.
- Jacoby, M. A., Virgin, H. W. t. & Speck, S. H. (2002).** Disruption of the M2 gene of murine gammaherpesvirus 68 alters splenic latency following intranasal, but not intraperitoneal, inoculation. *J Virol* **76**, 1790-1801.
- Kalderon, D., Oostra, B. A., Ely, B. K. & Smith, A. E. (1982).** Deletion loop mutagenesis: a novel method for the construction of point mutations using deletion mutants. *Nucleic Acids Res* **10**, 5161-5171.
- Katoh, K. & Toh, H. (2008).** Recent developments in the MAFFT multiple sequence alignment program. *Brief Bioinform* **9**, 286-298.
- Kolpakov, R., Bana, G. & Kucherov, G. (2003).** mreps: Efficient and flexible detection of tandem repeats in DNA. *Nucleic Acids Res* **31**, 3672-3678.

- Lopes, F. B., Colaco, S., May, J. S. & Stevenson, P. G. (2004).** Characterization of murine gammaherpesvirus 68 glycoprotein B. *J Virol* **78**, 13370-13375.
- Macakova, K., Matis, J., Rezuchova, I., Kudela, O., Raslova, H. & Kudelova, M. (2003).** Murine gammaherpesvirus (MHV) M7 gene encoding glycoprotein 150 (gp150): difference in the sequence between 72 and 68 strains. *Virus Genes* **26**, 89-95.
- Macrae, A. I., Dutia, B. M., Milligan, S., Brownstein, D. G., Allen, D. J., Mistrikova, J., Davison, A. J., Nash, A. A. & Stewart, J. P. (2001).** Analysis of a Novel Strain of Murine Gammaherpesvirus Reveals a Genomic Locus Important for Acute Pathogenesis. *J Virol* **75**, 5315-5327.
- Macrae, A. I., Usherwood, E. J., Husain, S. M., Flano, E., Kim, I. J., Woodland, D. L., Nash, A. A., Blackman, M. A., Sample, J. T. & Stewart, J. P. (2003).** Murid herpesvirus 4 strain 68 M2 protein is a B-cell-associated antigen important for latency but not lymphocytosis. *J Virol* **77**, 9700-9709.
- Madureira, P. A., Matos, P., Soeiro, I., Dixon, L. K., Simas, J. P. & Lam, E. W. (2005).** Murine gamma-herpesvirus 68 latency protein M2 binds to Vav signaling proteins and inhibits B-cell receptor-induced cell cycle arrest and apoptosis in WEHI-231 B cells. *J Biol Chem* **280**, 37310-37318.
- Mai, S. J., Ooka, T., Li, D. J., Zeng, M. S., Jiang, R. C., Yu, X. J., Zhang, R. H., Chen, S. P. & Zeng, Y. X. (2007).** Functional advantage of NPC-related V-val subtype of Epstein-Barr virus nuclear antigen 1 compared with prototype in epithelial cell line. *Oncol Rep* **17**, 141-146.
- Marques, S., Alenquer, M., Stevenson, P. G. & Simas, J. P. (2008).** A single CD8+ T cell epitope sets the long-term latent load of a murid herpesvirus. *PLoS Pathog* **4**, e1000177.
- McGeoch, D. J., Gatherer, D. & Dolan, A. (2005).** On phylogenetic relationships among major lineages of the Gammaherpesvirinae. *J Gen Virol* **86**, 307-316.
- McGeoch, D. J., Rixon, F. J. & Davison, A. J. (2006).** Topics in herpesvirus genomics and evolution. *Virus Res* **117**, 90-104.

- Mistríková, J., Raslova, H., Mrmusova, M. & Kudelova, M. (2000).** A murine gammaherpesvirus. *Acta Virol* **44**, 211-226.
- Muller, U., Steinhoff, U., Reis, L. F., Hemmi, S., Pavlovic, J., Zinkernagel, R. M. & Aguet, M. (1994).** Functional role of type I and type II interferons in antiviral defense. *Science* **264**, 1918-1921.
- Nash, A. A., Dutia, B. M., Stewart, J. P. & Davison, A. J. (2001).** Natural history of murine gamma-herpesvirus infection. *Philos Trans R Soc Lond B Biol Sci* **356**, 569-579.
- Oda, W., Mistríkova, J., Stancekova, M., Dutia, B. M., Nash, A. A., Takahata, H., Jin, Z., Oka, T. & Hayashi, K. (2005).** Analysis of genomic homology of murine gammaherpesvirus (MHV)-72 to MHV-68 and impact of MHV-72 on the survival and tumorigenesis in the MHV-72-infected CB17 scid/scid and CB17+/+ mice. *Pathol Int* **55**, 558-568.
- Ottinger, M., Pliquet, D., Christalla, T., Frank, R., Stewart, J. P. & Schulz, T. F. (2009).** The interaction of the gammaherpesvirus 68 orf73 protein with cellular BET proteins affects the activation of cell cycle promoters. *J Virol* **83**, 4423-4434.
- Parry, C. M., Simas, J. P., Smith, V. P., Stewart, C. A., Minson, A. C., Efstathiou, S. & Alcami, A. (2000).** A broad spectrum secreted chemokine binding protein encoded by a herpesvirus. *J Exp Med* **191**, 573-578.
- Pearson, W. R. & Lipman, D. J. (1988).** Improved tools for biological sequence comparison. *Proc Natl Acad Sci USA* **85**, 2444-2448.
- Pfeffer, S., Zavolan, M., Grasser, F. A., Chien, M., Russo, J. J., Ju, J., John, B., Enright, A. J., Marks, D., Sander, C. & Tuschl, T. (2004).** Identification of virus-encoded microRNAs. *Science* **304**, 734-736.
- Pires de Miranda, M., Alenquer, M., Marques, S., Rodrigues, L., Lopes, F., Bustelo, X. R. & Simas, J. P. (2008).** The Gammaherpesvirus m2 protein manipulates the Fyn/Vav pathway through a multidocking mechanism of assembly. *PLoS One* **3**, e1654.

Raslova, H., Matis, J., Rezuchova, I., Macakova, K., Berebbi, M. & Kudelova, M. (2000).

The bystander effect mediated by the new murine gammaherpesvirus 72-- thymidine kinase/5'-fluoro-2'-deoxyuridine (MHV72-TK/5-FUdR) system in vitro [In Process Citation]. *Antivir Chem Chemother* **11**, 273-282.

Reynolds, S. M., Kall, L., Riffle, M. E., Bilmes, J. A. & Noble, W. S. (2008).

Transmembrane topology and signal peptide prediction using dynamic bayesian networks. *PLoS Comput Biol* **4**, e1000213.

Rice, P., Longden, I. & Bleasby, A. (2000). EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet* **16**, 276-277.

Rodrigues, L., Pires de Miranda, M., Caloca, M. J., Bustelo, X. R. & Simas, J. P. (2006).

Activation of Vav by the gammaherpesvirus M2 protein contributes to the establishment of viral latency in B lymphocytes. *J Virol* **80**, 6123-6135.

Rutherford, K., Parkhill, J., Crook, J., Horsnell, T., Rice, P., Rajandream, M. A. &

Barrell, B. (2000). Artemis: sequence visualization and annotation. *Bioinformatics* **16**, 944-945.

Simas, J. P., Marques, S., Bridgeman, A., Efstathiou, S. & Adler, H. (2004). The M2

gene product of murine gammaherpesvirus 68 is required for efficient colonization of splenic follicles but is not necessary for expansion of latently infected germinal centre B cells. *J Gen Virol* **85**, 2789-2797.

Simas, J. P., Swann, D., Bowden, R. & Efstathiou, S. (1999). Analysis of murine

gammaherpesvirus-68 transcription during lytic and latent infection. *J Gen Virol* **80**, 75-82.

Stewart, J. P., Usherwood, E. J., Ross, A., Dyson, H. & Nash, T. (1998). Lung epithelial

cells are a major site of murine gammaherpesvirus persistence. *J Exp Med* **187**, 1941-1951.

Sunil-Chandra, N. P., Efstathiou, S., Arno, J. & Nash, A. A. (1992). Virological and

pathological features of mice infected with murine gamma-herpesvirus 68. *J Gen Virol* **73**, 2347-2356.

- Svobodova, J., Blaskovic, D. & Mistrikova, J. (1982).** Growth characteristics of herpesviruses isolated from free living small rodents. *Acta Virol* **26**, 256-263.
- Taylor, P. (1986).** A computer program for translating DNA sequences into protein. *Nucleic Acids Res* **14**, 437-441.
- Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994).** CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**, 4673-4680.
- Todaro, G. J. & Green, H. (1963).** Quantitative studies of the growth of mouse embryo cells in culture and their development into established lines. *J Cell Biol* **17**, 299-313.
- Usherwood, E. J., Roy, D. J., Ward, K., Surman, S. L., Dutia, B. M., Blackman, M. A., Stewart, J. P. & Woodland, D. L. (2000).** Control of gammaherpesvirus latency by latent antigen-specific CD8(+) T cells. *J Exp Med* **192**, 943-952.
- van Berkel, V., Barrett, J., Tiffany, H. L., Fremont, D. H., Murphy, P. M., McFadden, G., Speck, S. H. & Virgin, H. W. (2000).** Identification of a gammaherpesvirus selective chemokine binding protein that inhibits chemokine action. *J Virol* **74**, 6741-6747.
- van Berkel, V., Levine, B., Kapadia, S. B., Goldman, J. E., Speck, S. H. & Virgin, H. W. t. (2002).** Critical role for a high-affinity chemokine-binding protein in gamma-herpesvirus-induced lethal meningitis. *J Clin Invest* **109**, 905-914.
- van Berkel, V., Preiter, K., Virgin, H. W. & Speck, S. H. (1999).** Identification and initial characterization of the murine gammaherpesvirus 68 gene M3, encoding an abundantly secreted protein. *J Virol* **73**, 4524-4529.
- Virgin, H. W., Latreille, P., Wamsley, P., Hallsworth, K., Weck, K. E., Dal Canto, A. J. & Speck, S. H. (1997).** Complete sequence and genomic analysis of murine gammaherpesvirus 68. *J Virol* **71**, 5894-5904.
- Virgin, H. W., Presti, R. M., Li, X. Y., Liu, C. & Speck, S. H. (1999).** Three distinct regions of the murine gammaherpesvirus 68 genome are transcriptionally active in latently infected mice. *J Virol* **73**, 2321-2332.

Wang, J. T., Sheeng, T. S., Su, I. J., Chen, J. Y. & Chen, M. R. (2003). EBNA-1 sequence variations reflect active EBV replication and disease status or quiescent latency in lymphocytes. *J Med Virol* **69**, 417-425.

Wright, M. N., Stewart, J. P., Janjua, N. J., Pepper, S. D., Sample, C., Rooney, C. M. & Arrand, J. R. (1995). Antigenic and sequence variation in the C-terminal unique domain of the Epstein-Barr virus nuclear antigen EBNA-1. *Virology* **208**, 521-530.

Table 1. Isolation of herpesviruses from free-living rodents in Cheshire.

| Animal* | Tissues harvested† | Tissues yielding CPE† | Herpesvirus particles‡ |
|---------|--------------------|-----------------------|------------------------|
| WM1 | TG, S, L | TG, S | + |
| WM2 | TG, S, L | S | + |
| WM3 | TG, S, L | – | – |
| WM4 | TG, S, L | – | – |
| WM5 | TG, S, L | – | – |
| WM6 | TG, S, L | – | – |
| WM7 | TG, S, L | TG | + |
| WM8 | TG, S, L | TG, S | + |
| BV1 | TG, S, L | – | – |
| FV1 | S, L | S | + |
| HM1 | TG, S, L | – | – |
| HM2 | TG, S, L | – | – |
| HM3 | TG, S, L | – | – |
| HM4 | TG, S, L | S | + |
| HM5 | TG, S, L | – | – |
| HM6 | TG, S, L | – | – |

* WM, wood mouse; BV, bank vole; FV, field vole; HM, house mouse; followed by a number for each animal.

† TG, trigeminal ganglia; S, spleen; L, lungs; –, no CPE.

‡ +, particles observed by EM; –, particles not observed by Transmission Electron Microscopy on negatively stained preparations.

FIGURE LEGENDS

Fig. 1. PCR amplification of the coding regions of genes (a) *M1*, (b) *M2* and (c) *M3* from viruses isolated from FV1, WM1, WM2, WM7 and WM8, in comparison with MuHV4. TG, trigeminal ganglia; S, spleen.

Fig. 2. Variation between the genome sequences of WMHV and MuHV4. The lower part of the panels represent the genome, commencing in the first panel at the start of U and ending in the last panel with one copy of TR, which is shown in a thicker format. Protein-coding regions are depicted by shaded arrows, with connecting introns indicated by white horizontal bars, and genes encoding the tRNA-like genes (1-8) are shown as arrowheads. Internal tandem repeats are represented by black horizontal bars. The upper part of each panel shows the nucleotide divergence (nd) calculated for a 100-nucleotide window, shifted by increments of 3 nucleotides. A nucleotide position was counted as divergent if it differed between the two sequences; insertions or deletions were not scored.




Fig. 3. Divergence between the amino acid sequences of predicted protein-coding regions in WMHV, BRHV and MuHV4. The histogram illustrates sequence divergence (% non-identity) between the amino acid sequence of predicted protein-coding regions in WMHV and MuHV4 (grey bars, all coding regions) and BRHV and WMHV (black bars, coding regions to *ORF52*)





Fig. 4. Alignments of the predicted nucleotide sequences of the *tRNA*-like genes and *miRNAs* from MuHV4 and WMHV. A diagrammatic representation of the genomic region showing the relative positions of these non-coding RNAs is shown at the top. The positions of the *M1-M3* ORFs and viral *tRNA*-like transcripts (t1–t8) are shown by arrows. The positions of the *miRNAs* (miR-M1-1 through M1-9) derived from primary transcripts of the *tRNA*-like RNAs are shown by vertical lines. Sequence alignments of the *tRNA/miRNA* transcripts are shown below. The sections processed to generate the *tRNA*-like molecules are shaded grey and pre-*miRNAs* are shaded blue. The positions of the A and B box of the RNA Polymerase III promoters are shown by open boxes, as are the positions of the processed *miRNAs*. The positions of the anti-codons in the *tRNAs* are shown in blue type. Differences between MuHV4 and WMHV are shown in red type. Data for MuHV4 are from (Bowden *et al.*, 1997) and (Pfeffer *et al.*, 2004).


Fig. 5. Alignments of the predicted amino acid sequences of (a) M1, M3 and M4, (b) M2, (c) ORF51, and (d) ORF73. Each individual alignment consists of the sequences from MuHV4, WMHV and BRHV, with residues that differ from the consensus (or from each other in the case of ORF73) shaded grey. In (a), the alignments for M1, M3 and M4 are aligned with each other because these three proteins are related via the residues in bold type; each sequence contains a predicted signal peptide (lower case). In (b), the positions of PXXP motifs (P1–P9), tyrosine residues 120 and 129 and the CD8 CTL epitope (CTL) are indicated above the M2 sequence. In (c), the bold residues indicate potential N-linked glycosylation sites in gp150 encoded by *ORF51*. In (d), the positions of the Brd4- and Brd2-interacting domains of the ORF73 protein are shown above the sequences.

Fig. 6. Virological analyses of WMHV infection of wood mice. Wood mice (three per time-point) were infected intranasally with 4×10^5 PFU of MuHV4 or WMHV. Bars represent standard deviation from the mean; the asterisk represents statistically significant differences between species ($p < 0.05$). (a) Infectious virus recovered from the lung at 7 and 14 d PI. Titres were measured by plaque assay on NIH3T3 cells. (b) Mean leukocyte numbers per spleen. (c) Infective centre assay of the level of latency in splenocytes. Infectious virus titres in the samples were analysed in parallel and were subtracted from the total infectious centres.

Fig. 7. Cellular response to WMHV and MuHV4 infection in the lungs of wood mice at 14 d PI. (a & b), Infection with MuHV4; (a) Intense peribronchial focal lymphocyte infiltration with evidence of lymphatic follicle formation (F). B, bronchiole. HE stain. Bar = 50 μ m. (b) Focal perivascular B cell infiltration with lymphatic follicle formation (F). A, artery. Staining for the B cell marker CD45R, avidin biotin peroxidase complex method, Papanicolaou's haematoxylin counterstain. Bar = 20 μ m. (c & d) Infection with WMHV; (c) Moderate peribronchiolar focal lymphocyte infiltration (arrows). B, bronchiole. HE stain. Bar = 20 μ m. (d) Artery with focal B cell-dominated (CD45R-positive) perivascular lymphocyte infiltration (arrows). There is evidence of B cell rolling and emigration (arrowheads). Bar = 20 μ m.







(a)

| | |
|----------|--|
| MuHV4 M1 | mglatlclllscilggg.....iaWPSVVNIETYPPEDDDTKEDMRDYLFVQNCLLQDNFNATYCSDSFEKLDKRSHTLPTDTCNVKTTFLVNVY |
| WMHV M1 | mwlatlclllscilggg.....iaWPSVVNIETYPPEDDDTKEDMRDYLFVQNCLLQDNFNATYCSDSFEKLDKRSHTLPTDTCNVKTTFLVNVY |
| BRHV M1 | mwlatlclllscilggg.....iaWPSVVNIETYPPEDDDTKEDMRDYLFVQNCLLQDNFNATYCSDSFEKLDKRSHTLPTDTCNVKTTFLVNVY |
| MuHV4 M3 | maf1stsvylikocicillagglaes1t1g1LAPALSTHSSGVSTQSVDSLQIKRGAETQAHCLTTPAETEVTECASILKDVLAONLHLELQGLCNVKNKMGAPW |
| WMHV M3 | maf1stpsmlikocicillagglvvaesTSIGLRPALTYTSSGVTTQSVDSLQIKRGAETQAHCLTTPAETEVTECASILKDVLAONLHLELQGLCNVKNKMGAPW |
| BRHV M3 | maf1stpsmlikocicillagglvvaesTSIGLRPALTYTSSGVTTQSVDSLQIKRGAETQAHCLTTPAETEVTECASILKDVLAONLHLELQGLCNVKNKMGAPW |
| MuHV4 M4 | mgplgrpwaasfgffflavislattptgEDDDIPVKIHRLTFVKGVLVDVTGGS.SYVVCVYPSRKLFCPTTRWKDLSRFLNSETLQVCSARTIYSVVP |
| WMHV M4 | mgplgrpwaasfgffflavislattptgEDDDIPVKIHRLTFVKGVLVDVTGGS.SYVVCVYPSRKLFCPTTRWKDLSRFLNSETLQVCSARTIYSVVP |
| BRHV M4 | mgplgrpwaasfgffflavislattptgEDDDIPVKIHRLTFVKGVLVDVTGGS.SYVVCVYPSRKLFCPTTRWKDLSRFLNSETLQVCSARTIYSVVP |

| | |
|----------|---|
| MuHV4 M1 | N.RHKYGIKFKFESRLPLPTMASATSGRVIKVLVAEAGR....PWKRHWANLAMVYTSNVVRLTDLNNAKFRTRFSRIWVSVTLDRHEVDLDTF...AGFL |
| WMHV M1 | N.RHKYGMFKFESRLPLPTMASATSGRVIKVLVAEAGR....PWKRHWANLAMVYTSNVVRLTDLNNAKFRTRFSRIWVSVTLDRHEVDLDTF...AGFL |
| BRHV M1 | N.RHKYGMFKFESRLPLPTMASATSGRVIKVLVAEAGR....PWKRHWANLAMVYTSNVVRLTDLNNAKFRTRFSRIWVSVTLDRHEVDLDTF...AGFL |
| MuHV4 M3 | VSVEELGQEIITGRLPFPVGGTTPVNDLVRVLVAESNTPPEETPEEFYAYVELOQTELYTFLGLDNDVVTSDYMTIWMIDIPKSYVDVGMLT...RATF |
| WMHV M3 | VSVEELGQEIITGRLPFPVGGTTPVNDLVRVLVAESNTPPEETPEEFYAYVELOQTELYTFLGLDNDVVTSDYMTIWMIDIPKSYVDVGMLT...RATF |
| BRHV M3 | VSVEELGQEIITGRLPFPVGGTTPVNDLVRVLVAESNTPPEETPEEFYAYVELOQTELYTFLGLDNDVVTSDYMTIWMIDIPKSYVDVGMLT...RATF |
| MuHV4 M4 | VEMLRIISLPLPEIKS.IVGGIRSQYVSFPTTISLHPAKLP1WKSFANLKVSMKFRSSQWEVAFSVVSKTDYAITIYWAEIPGFLIHESATINLINQPL |
| WMHV M4 | VEMLRIISLPLPEIKS.IVGGIRSQYVSFPTTISLHPAKLP1WKSFANLKVSMKFRSSQWEVAFSVVSKTDYAITIYWAEIPGFLIHESATINLINQPL |
| BRHV M4 | VEMLRIISLPLPEIKS.IVGGIRSQYVSFPTTISLHPAKLP1WKSFANLKVSMKFRSSQWEVAFSVVSKTDYAITIYWAEIPGFLIHESATINLINQPL |

| | |
|----------|---|
| MuHV4 M1 | FAAPESVQLTLLMDVYPTFTWCGQI.SLNDPDLVPSFOAIRTLVPMCFP...MWRYLNGQDFHHDGCHQESK.WWNPTHIIFRLNPGTES...HNIT |
| WMHV M1 | FAAPESVQLTLLMDVYPTFTWCGQI.SLNDPDLVPSFOAIRTLVPMCFP...MWRYLNGQDFHHDGCHQESN.WWNPTHIIFRLNPGTES...HNIT |
| BRHV M1 | FAAPESVQLTLLMDVYPTFTWCGQI.SLNDPDLVPSFOAIRTLVPMCFP...MWRYLNGQDFHHDGCHQESN.WWNPTHIIFRLNPGTES...HNIT |
| MuHV4 M3 | LEQWPGAKVTVMIPYSSFTTWCGEIGAISEESAPQPSLSAR...SPVCKN...SARYSTS.KFCEVDGCTAETG.MEKMSLLTP...FGGPP...QQAQ |
| WMHV M3 | LEQWPGAKVTVMIPYSSFTTWCGEIGAISEESAPQPSLSAR...SPVCKN...SARYSTS.KFCEVDGCTAETG.MEKMSLLTP...FGGPP...QQAQ |
| BRHV M3 | LEQWPGAKVTVMIPYSSFTTWCGEIGAISEESAPQPSLSAR...SPVCKN...SARYSTS.KFCEVDGCTAETG.MEKMSLLTP...FGGPP...QQAQ |
| MuHV4 M4 | LALYADLHVDMVRLTDFKFIYCQTY.TLQQKNLTDPRGTGRKPTSSVILPSPHVKNQIIRRNETHFVETCSSAWDNYTSEAHNISRNSSSRGNSVQLVN |
| WMHV M4 | LALYADLHVDMVRLTDFKFIYCQTY.TLQQKNLTDPRGTGRKPTSSVILPSPHVKNQIIRRNETHFVETCSSAWDNYTSEAHNISRNSSSRGNSVQLVN |
| BRHV M4 | LALYADLHVDMVRLTDFKFIYCQTY.TLQQKNLTDPRGTGRKPTSSVILPSPHVKNQIIRRNETHFVETCSSAWDNYTSEAHNISRNSSSRGNSVQLVN |

| | |
|----------|--|
| MuHV4 M1 | LNTCVCHVKYNDLQELDAAHRIKILTIISNFFGFYKPLVVLVYFSGSDVNEGQAPPLQYCVVFIHRGNYGFFRTRQRGDPCPCFSLG..RDEIVLVG |
| WMHV M1 | LNTCVCHVKYNDLQELDAAHRIKILTIISNFFGFYKPLVVLVYFSGSDVNEGQAPPLQYCVVFIHRGNYGFFRTRQRGDPCPCFSLG..RDEIVLVG |
| BRHV M1 | LNTCVCHVKYNDLQELDAAHRIKILTIISNFFGFYKPLVVLVYFSGSDVNEGQAPPLQYCVVFIHRGNYGFFRTRQRGDPCPCFSLG..RDEIVLVG |
| MuHV4 M3 | MNTCPCYKYK.SVSLPAMPDHLILADLAGLDSLTSFVYVMAAYFDSTHENPVRPSSKLYHICALQMTS.HDGVWTSTS...SECCPIRLVEGQSRNVLQVLV |
| WMHV M3 | MNTCPCYKYK.SVSLPAMPDHLILADLAGLDSLTSFVYVMAAYFDSTHENPVRPSSKLYHICALQMTS.HDGVWTSTS...SECCPIRLVEGQSRNVLQVLV |
| BRHV M3 | MNTCPCYKYK.SVSLPAMPDHLILADLAGLDSLTSFVYVMAAYFDSTHENPVRPSSKLYHICALQMTS.HDGVWTSTS...SECCPIRLVEGQSRNVLQVLV |
| MuHV4 M4 | ITANPCTLPT.LWDNWPCTNYRSSVPVEIVIHENILLEGRAIYIYHQIGLFDQPRLCVATFWMSK.....EETLLMQLDYPCEVSVKKGKFLIKS |
| WMHV M4 | ITANPCTLPT.LWDNWPCTNYRSSVPVEIVIHENILLEGRAIYIYHQIGLFDQPRLCVATFWMSK.....EETLLMQLDYPCEVSVKKGKFLIKS |
| BRHV M4 | ITANPCTLPT.LWDNWPCTNYRSSVPVEIVIHENILLEGRAIYIYHQIGLFDQPRLCVATFWMSK.....EETLLMQLDYPCEVSVKKGKFLIKS |

| | |
|----------|---|
| MuHV4 M1 | HYVDVKRIVGITIFFDQGEHRI SYLGLKLSRAAVVGDdTNNKIIFPGQOS |
| WMHV M1 | HYVDVKRIVGITIFFDQGEHRI SYLGLKLSRAAVVGDdTNNKIIFPGQOS |
| BRHV M1 | HYVDVKRIVGITIFFDQGEHRI SYLGLKLSRAAVVGDdTNNKIIFPGQOS |
| MuHV4 M3 | APTSMPKLVGVSMLLEGQYRLEYFGDH |
| WMHV M3 | APTSMPKLVGVSMLLEGQYRLEYFGDH |
| BRHV M3 | APTSMPKLVGVSMLLEGQYRLEYFGDH |
| MuHV4 M4 | IVSTYHAISMVTFIWEYGIETDFLE |
| WMHV M4 | IVSTYHAISMVTFIWEYGIETDFLE |
| BRHV M4 | IVSTYHAISMVTFIWEYGIETDFLE |

(b)

| | | | | | |
|----------|---|-----------|-------------|------------|-----------|
| MuHV4 M2 | <u>P1</u> MAPTPPQGIKIPNPPGGSSQNPVLWGDGTGNYRPSPEWILGOVPCDQRFHPHSGNKNSSSTSGGRPORPPLPRTREPKTIRRGFNKLRSTLKSPPKPRP | <u>P2</u> | <u>P3/4</u> | <u>CTL</u> | <u>P5</u> |
| WMHV M2 | MAPTPPQGIKIPNPPGGSSQNPVLWGDGTGHNPKPEWILGOVPCDHRGGHPAGNKNSSSTSGGKPHRRSWPRWRVPSVKARFKFRSTIRSPRNIET. | | | | |
| BRHV M2 | MAPTPPQGIKIPNPPGGSSQNPVLWGDGTGHNPKPEWILGOVPCDHRGGHPAGNKNSSSTSGGKPHRRSWPRWRVPSVKARFKFRSTIRSPRNIET. | | | | |

| | |
|----------|---|
| MuHV4 M2 | SPVSPPEEVPNAPGSPPEENIYETANSEPVYIQPISTRSLMMLDSGSTSEENLGPPTRPPLKLPNQHPMNPDIPLIPIPPSKCHKGFVEWGEE |
| WMHV M2 | NPGRPEENGNAPGSPPEENIYDALNNEPLYIQPICNSMMLDSGSTGSTEISLAPTRPLKLPNQHPMNPDIPLIPIPPSKCHKGFVEWGEE |
| BRHV M2 | NPGRPEENGNAPGSPPEENIYDALNNEPLYIQPICNSMMLDSGSTGSTEISLAPTRPLKLPNQHPMNPDIPLIPIPPSKCHKGFVEWGEE |

(c)

| | |
|-------------|---|
| MuHV4 ORF51 | MCGVKSIAKCFLLFQIISFLGNHNLVWVPGAALGAETVEGITSREMEINATKAPSSGATSELLVTLNPNPTIIMRPPVAQNGESVHKDARASASDPT |
| WMHV ORF51 | MCGVKPIVCKFLLFHIINFLGTYNVGVWPGTPLCAAQAVDGTISREMEINATLAPSSGATSLLVTLNPNPTIIMRPPVAQNGESVIDNSAPASDPT |
| BRHV ORF51 | MCGVKPIVCKFLLFHIINFLGNVNVGVWPGTLLCAAQAVDGTISREMEINATLAPSSGATSLLVTLNPNPTIIMRPPVAQNGESVIDNSAPASDPT |

| | |
|-------------|--|
| MuHV4 ORF51 | TSEPTSPGEEPTFA.....DPKAAEASAGHVGETESESPTPLPAITPKPSSQEDNPTMTPTTAEPTSNADVSTEHVDETEPESPTFLPTPEPDTPTTE |
| WMHV ORF51 | TSNPSSPEEAPTAAPITPITPTAAQSAEHVGETDEAPTLPITPKPSSQEDPTMTSPETPTTAAISTEQDDETEPESAPPPTAPE.....E |
| BRHV ORF51 | TSNPSSPEEAPTAAPITPITPTAAQSAEHVGETDEAPTLPITPKPSSQEDPTMTSPETPTTAAISTEQDDETEPESAPPPTAPE.....E |

| | |
|-------------|--|
| MuHV4 ORF51 | TTTPTKNQDEPTLITS.....SSDAPADTS.DTSPKQEDDPVKPTESKQOAEFKDNSPSDVPETADSDTDPASPTVELTPEPTTETVTPADSPV |
| WMHV ORF51 | TTTPTKNQDEPTLINTSDQDDSSSDIPAGTPGPTTPPKQETETTKFVDSKQVEFNDSAPSDIPETSDSTPTPV.....TDPTSPSVEETSAPAEPT |
| BRHV ORF51 | TTTPTKNQDEPTLINTSDQDDSSSDIPAGTPGPTTPPKQETETTKFVDSKQVEFNDSAPSDIPETSDSTPTPA.....TDPTSPSVEETSAPAEPT |

| | |
|-------------|---|
| MuHV4 ORF51 | FOPTAPAEPSKPEPTTPVDPATEPNTFADPSTPESTPTDPPAPQPTPAEAPSNEPEPTPPVDPATPPNIPADPSTPESTPADPAEQPTPAEPSTP |
| WMHV ORF51 | PDSTPADPPAPQPTPAEPSTPDSTPQDEPSTPDSTPADPPAPQPTPAEAPSTPESTPADPPAAQPTPAEPSTPDSTPADPAEQPTPAEPSTP |
| BRHV ORF51 | PDSTPADPPAPQPTPAEPSTPDSTPQDEPSTPDSTPADPPAPQPTPAEAPSTPESTPADPPAAQPTPAEPSTPDSTPADPAEQPTPAEPSTP |

| | |
|-------------|---|
| MuHV4 ORF51 | EPSTPAKAPAEPTTPEPSGPMTEATPSTACGAEETETPDGDTTTPASPOTTAPMHEVVDISTLLWIRPTVAIILIFLLMIFHIMYCVCLHE |
| WMHV ORF51 |PSPEPMTQSGPAITPEIATPSTTEPGAGDASMGATTTQAASALTTKPMRVVVDVSTMLWIRPTVAIIVLIFLLMIFHIMYCVCLHE |
| BRHV ORF51 |PSPEPMTQSGPAITPEIATPSTTESGAGKEASMGDTTQAASALTTKPMRVVVDVSTMLWIRPTVAIIVLIFLLMIFHIMYCVCLHE |


(d)

| | |
|-------------|---|
| MuHV4 ORF73 | MPTSPPTTRNTTSGKTRSGCKRRCFNKPAAAMPKRRAAPKRPAPPPPPGCCDEESS-----OGTQTPNPPSPVPPS-----S |
| WMHV ORF73 | MPTSPPTTRNTTSGKTRSGCKRRCYKFKQASMPKRRRVPKRPAPPPPPPPGQDEESSLLGESSHTEERPEQSEERQSPKQPIPPSPPTPLKKTPT |

| | |
|-------------|---|
| MuHV4 ORF73 | PTLPSPPVPPSPVHEPSPSPAPPSPD---VDVEGLDVGETDDPGPPPKR--YSRYQKPHNPSDELPKKYQGMRRHLQVTAAPRLFDPEGHPETHF |
| WMHV ORF73 | SDPPSPAPPSTVPVPPQSPSPAPPSPDSSDVEGLDQGGDDGPPAKKAPSPRYQKPYNEKSTLPAKYRGMRRNHMVIAPRLFDPERPAPTHF |

| | |
|-------------|--|
| MuHV4 ORF73 | KSAMFSSHTPYTLNKLKHKIQSKHVLSTPVSLCLPVPVGTTOCCVYIYLLSFVEDKKQAKLKRVLVAYCEKYHSSVEGTIVKAKYFPLPEPTEPTD |
| WMHV ORF73 | KSAMFSTKPYSLTLKLVKQSKQVVTSTVSLCLPVPVGGQPTLVKYIILSFVENKKLAKMLRKKVVLVYCAKFKHSNIDGTIVKAKYFPLPG---TPTD |

| | |
|-------------|--|
| MuHV4 ORF73 | PEQPSTSTQASGTOHGPTASLDAGAEQATGSPGSSPGQQGQSQT |
| WMHV ORF73 | PEQPSTSAQVSGTQSPASPEASTEQEATV----- |







Table S1. Information used to annotate the WMHV and MuHV4 genome sequences

| | Protein name* | Family | Description | EC no. | Activity |
|---------|--|--------|---|-----------|---|
| M1 | protein M1 | M1 | contains a signal peptide; secreted protein | | unknown |
| M2 | protein M2 | | interacts with Vav oncoprotein; involved in B cell proliferation and differentiation | | immune regulation; latency |
| M3 | chemokine-binding protein M3 | M1 | contains a signal peptide; secreted protein | | immune regulation |
| M4 | chemokine-binding protein M4 | M1 | contains a signal peptide; secreted protein | | immune regulation |
| ORF4 | complement control protein | CCP | type 1 envelope glycoprotein; contains four SCR domains | | immune regulation |
| ORF6 | single-stranded DNA-binding protein | | contains a zinc-finger | | DNA replication; possibly gene regulation |
| ORF7 | DNA packaging terminase subunit 2 | | | | DNA encapsidation |
| ORF8 | envelope glycoprotein B | | type 1 membrane protein; possible membrane fusogen; binds cell surface heparan sulphate | | cell entry; cell-to-cell spread |
| ORF9 | DNA polymerase catalytic subunit | | | 2.7.7.7 | DNA replication |
| ORF10 | protein G10 | DURP | | | unknown |
| ORF11 | virion protein G11 | DURP | | | unknown |
| ORF12 | E3 ubiquitin ligase MIR1 | MIR | type 3 membrane protein; 2 transmembrane domains; contains a PHD finger; downregulates MHC-I | | immune regulation |
| 13M | protein 13M | | hypothetical protein encoded by 5'-region of ORF12 transcript | | unknown |
| ORF17.5 | capsid scaffold protein | | clipped near C terminus | | capsid morphogenesis |
| ORF17 | capsid maturation protease | | serine protease (N-terminal region); minor scaffold protein (remainder of protein, clipped near C terminus) | 3.4.21.97 | capsid morphogenesis |
| ORF18 | protein UL79 | | required for expression of late genes | | gene regulation |
| ORF19 | DNA packaging tegument protein UL25 | | located on capsid near vertices; possibly stabilizes the capsid and retains the genome | | DNA encapsidation |
| ORF20 | nuclear protein UL24 | | | | unknown |
| ORF21 | thymidine kinase | | | 2.7.1.21 | nucleotide metabolism |
| ORF22 | envelope glycoprotein H | | type 1 membrane protein; possible membrane fusogen; complexed with envelope glycoprotein L | | cell entry; cell-to-cell spread |
| ORF23 | protein UL88 | | | | unknown |
| ORF24 | virion protein UL87 | | | | unknown |
| ORF25 | major capsid protein | | 6 copies form hexons, 5 copies form pentons | | capsid morphogenesis |
| ORF26 | capsid triplex subunit 2 | | complexed 2:1 with capsid triplex subunit 1 to connect capsid hexons and pentons | | capsid morphogenesis |
| ORF27 | envelope glycoprotein 48 | | type 2 membrane protein | | cell-to-cell spread |
| ORF28 | envelope glycoprotein 150 | | type 1 membrane protein | | immune regulation |
| ORF29 | DNA packaging terminase subunit 1 | | contains an ATPase domain | | DNA encapsidation |
| ORF30 | protein UL91 | | | | unknown |
| ORF31 | protein UL92 | | | | unknown |
| ORF32 | DNA packaging tegument protein UL17 | | capsid-associated | | DNA encapsidation; capsid transport |
| ORF33 | tegument protein UL16 | | | | possibly virion morphogenesis |
| ORF34 | protein UL95 | | | | unknown |
| ORF35 | tegument protein UL14 | | | | virion morphogenesis |
| ORF36 | tegument serine/threonine protein kinase | PK | | 2.7.11.1 | protein phosphorylation |
| ORF37 | deoxyribonuclease | | | | DNA processing |
| ORF38 | myristylated tegument protein | | envelope-associated | | virion morphogenesis |
| ORF39 | envelope glycoprotein M | | type 3 membrane protein; 8 transmembrane domains; complexed with envelope glycoprotein N | | virion morphogenesis; membrane fusion |
| ORF40 | helicase-primase subunit | | | | DNA replication |
| ORF42 | tegument protein UL7 | | | | virion morphogenesis |
| ORF43 | capsid portal protein | | dodecamer located at one capsid vertex in place of a penton | | DNA encapsidation |

| | | | | | | |
|--------|--|--|--|--------|----------|---|
| ORF44 | helicase-primase helicase subunit | | | | | DNA replication |
| ORF45 | tegument protein G45 | | | | | unknown |
| ORF46 | uracil-DNA glycosylase | | | | 3.2.2.3 | DNA repair |
| ORF47 | envelope glycoprotein L | | | | | cell entry; cell-to-cell spread |
| ORF48 | tegument protein G48 | | | | | unknown |
| ORF49 | protein G49 | | | | | gene regulation |
| ORF50 | protein Rta | | | | | gene regulation; latency |
| ORF51 | envelope glycoprotein 350 | | | | | cell attachment |
| ORF52 | virion protein G52 | | | | | unknown |
| ORF53 | envelope glycoprotein N | | | | | virion morphogenesis; membrane fusion |
| ORF54 | deoxyuridine triphosphatase | | | DURP | 3.6.1.23 | nucleotide metabolism |
| ORF55 | tegument protein UL51 | | | | | virion morphogenesis |
| ORF56 | helicase-primase subunit | | | | | DNA replication |
| ORF57 | multifunctional expression regulator | | | MER | | gene regulation; RNA metabolism and transport |
| ORF58 | envelope protein UL43 | | | | | possibly membrane fusion |
| ORF59 | DNA polymerase processivity subunit | | | | | DNA replication |
| ORF60 | ribonucleotide reductase subunit 2 | | | | 1.17.4.1 | nucleotide metabolism |
| ORF61 | ribonucleotide reductase subunit 1 | | | | 1.17.4.1 | nucleotide metabolism |
| ORF62 | capsid triplex subunit 1 | | | | | capsid morphogenesis |
| ORF63 | tegument protein UL37 | | | | | virion morphogenesis |
| ORF64 | large tegument protein | | | | | capsid transport |
| ORF65 | small capsid protein | | | | | capsid morphogenesis; possibly capsid transport |
| ORF66 | protein UL49 | | | | | unknown |
| ORF67 | nuclear egress type 2 membrane protein | | | | | nuclear egress |
| ORF67A | DNA packaging protein UL33 | | | | | DNA encapsidation |
| ORF68 | DNA packaging protein UL32 | | | | | DNA encapsidation; possibly capsid transport |
| ORF69 | nuclear egress lamina protein | | | | | nuclear egress |
| ORF72 | cyclin | | | | | cell cycle regulation |
| M11 | apoptosis regulator M11 | | | Bcl-2 | | apoptosis |
| ORF73 | nuclear antigen LANA | | | | | latency |
| ORF74 | membrane protein G74 | | | GPCR | | intracellular signalling |
| ORF75C | tegument protein G75C | | | FGARAT | | unknown |
| ORF75B | protein G75B | | | FGARAT | | unknown |
| ORF75A | protein G75A | | | FGARAT | | unknown |

* Protein names are a provisional standard for all herpesviruses



- Encoded by a core gene (i.e. inherited from an ancestor of alpha-, beta- and gammaherpesviruses)
- Encoded by a betagamma gene (i.e. inherited from an ancestor of beta- and gammaherpesviruses)
- Encoded by a gamma gene (i.e. inherited from an ancestor of gammaherpesviruses)
- Encoded by a gene specific to a subset of gammaherpesviruses

This is an author manuscript that has been accepted for publication in *Journal of General Virology*, copyright Society for General Microbiology, but has not been copy-edited, formatted or proofed. Cite this article as appearing in *Journal of General Virology*. This version of the manuscript may not be duplicated or reproduced, other than for personal use or within the rule of 'Fair Use of Copyrighted Materials' (section 17, Title 17, US Code), without permission from the copyright owner, Society for General Microbiology. The Society for General Microbiology disclaims any responsibility or liability for errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final copy-edited, published article, which is the version of record, can be found at <http://vir.sgmjournals.org>, and is freely available without a subscription.