

SCIENTIFIC REPORTS



OPEN

Characterization of AiiK, an AHL lactonase, from *Kurthia huakui* LAM0618^T and its application in quorum quenching on *Pseudomonas aeruginosa* PAO1

Weiwei Dong^{1,2}, Jie Zhu³, Xiang Guo², Delong Kong², Qi Zhang³, Yiqing Zhou², Xiaoyang Liu², Shumiao Zhao¹ & Zhiyong Ruan²

N-Acyl homoserine lactones (AHLs) act as the key quorum sensing (QS) signal molecules in gram-negative bacteria, which coordinates gene expression and then activates various processes, including biofilm formation and production of virulence factors in some pathogens. Quorum quenching (QQ), which is the inactivation of the signal molecules by means of enzymatic degradation or modification, inhibits the processes of QS rather than killing the pathogens and is a promising antipathogenic strategy to control the bacterial pathogens. In this study, an AHL lactonase gene (named *aaiK*) was cloned from *Kurthia huakui* LAM0618^T and the AHL lactonase AiiK was expressed by *Escherichia coli*. AiiK exhibits a variable substrate spectrum and efficient degradation of the AHL compounds. The enzyme assays demonstrated that AiiK behaves as an AHL lactonase that can hydrolyze the lactone bond of the AHLs. The total hydrolytic efficiency of AiiK for C₁₀-HSL is 3.9 s⁻¹·mM⁻¹. AiiK can also maintain 20% activity after 12 h incubation at 37 °C and demonstrate great resistance to α-chymotrypsin, trypsin, and protease K. Furthermore, AiiK significantly inhibits the biofilm formation and attenuates extracellular proteolytic activity and pyocyanin production of *Pseudomonas aeruginosa* PAO1, which indicates the potential application of AiiK as a biocontrol agent or an anti-pathogenic drug.

Quorum sensing (QS) is a common and important bacterial communication mechanism that monitors the population density of bacteria and controls the expression of specialized structural gene sets by using specific receptors that can sense the accumulation of signal molecules. QS systems regulate a variety of functions, including biofilm maturation, bioluminescence, sporulation, production of secondary metabolites, and competence for DNA uptake¹. A diverse array of diseases are caused by a range of pathogens that utilize these QS systems, such as the opportunistic pathogen bacteria *Pseudomonas aeruginosa* can develop persistent complicated biofilm associated infections and stimulate the resistance to antibiotics, and the plant pathogen bacteria *Erwinia carotovora* can produce pectolytic enzymes that hydrolyze pectin between individual plant cells, which causes a disease plant pathologists term bacterial soft rot¹.

N-Acyl homoserine lactones (AHLs) act as key QS signal molecules in gram-negative bacteria. A representative AHL molecule comprises a homoserine lactone and an acyl chain containing an even number of carbon chains or a carbonylation at the C-3 position². Quorum quenching (QQ), which is the inactivation of these signaling molecules by enzymatic degradation or modification, interferes with the QS systems and inhibits the processes of QS³. QQ enzymes, including AHL lactonases, AHL acylases, and AHL oxidoreductases, play an important role in QQ^{1,4-7}. AHL lactonases belong to the metallo-β-lactamase superfamily and catalyze the

¹State Key Laboratory of Agricultural Microbiology and College of Life Science and Technology, Huazhong Agricultural University, Wuhan, 430070, China. ²Institute of Agricultural Resources and Regional Planning, CAAS, Beijing, 100081, China. ³College of life science, Xinjiang Normal University, Xinjiang, 830046, China. Correspondence and requests for materials should be addressed to S.Z. (email: shumiaozhao@mail.hzau.edu.cn) or Z.R. (email: ruanzhiyong@caas.cn)




Figure 1. SDS-PAGE analysis of the purified AiiK. M: molecular weight marker; Lane 1: pET32a vector plasmid (control); Lane 2: supernatant of the sonication product; Lane 3: precipitate of the sonication product; Lane 4: AiiK purified via Ni-NTA.

opening of homoserine lactone bonds in AHLs^{8–10}. Unlike AHL acylases and AHL oxidoreductases, AHL lactonases present variable substrate spectra and efficient degradation of AHLs signals^{5,7,11}.

To date, over 20 QQ enzymes have been reported and the sources of which range from bacteria to fungi and even mammals^{12,13}. QQ enzymes have been considered as novel biocontrol agents, particularly in antivirulence and antibacterial. The first reported AHL lactonase, AiiA, was identified in *Bacillus* sp. strain 240B1 and heterologously expressed in *E. carotovora* strain SCG1. AiiA significantly attenuated the pathogenicity of the *E. carotovora* strain SCG1 on many vegetables, such as carrot, potato, cauliflower, and so on⁴. The heterologous expression of AHL lactonase AiiM by *Escherichia coli* quenched the virulence in *Pectobacterium carotovorum* subsp. *carotovorum*¹⁰. MomL, a novel type of AHL lactonase from flounder mucus-derived *Muricauda olearia*, significantly alleviated the production of virulence factors in *P. aeruginosa* PAO1¹⁴. HqiA, a novel AHL-degrading enzyme type has AHL lactonase activity, interfered the swarming motility and production of maceration enzymes of *P. carotovorum*¹⁵. Therefore, the AHL lactonases provide a promising prospect to be applied as biocontrol agents.

In our previous studies, *K. huakuii* LAM0618^T was isolated from biogas slurry¹⁶. Genomic data¹⁷ revealed that *K. huakuii* LAM0618^T contains a putative protein sequence annotated as an “N-acyl homoserine lactonase”. Bioinformatic analysis suggested that it may represent a potential AHL lactonase. In the current work, the putative *aaiK* gene from *K. huakuii* LAM0618^T was cloned and heterologously expressed in *E. coli* based on method of genome-mining. The physicochemical properties of the purified recombinant AiiK protein (AiiK) and its application in QQ on *P. aeruginosa* PAO1 were investigated.

Results

Sequence analysis and identification of AiiK. The *aaiK* gene was cloned from *K. huakuii* LAM0618^T by using the primers *aaiKR* and *aaiKF*, the full-length 837 bp gene, which encoded a 278-amino acid polypeptide belonging to putative metallo- β -lactamase. No signal peptide was detected in the intact amino acid sequence of AiiK by the analysis of SignalP 4.1. The *aaiK* gene was inserted into expression plasmid pET32a and the recombinant plasmid pET32a-*aaiK* was transformed into *E. Coli* BL21 (DE3). The recombinant AiiK was expressed by the inducing of isopropyl- β -D-thiogalactopyranoside (IPTG) and then purified by using a Ni-NTA column. The molecular weight of the purified fusion protein was approximately 50 kDa, as determined by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) analysis (Fig. 1); this value was much larger than the predicted molecular weight of 31.76 kDa because the purified recombinant AiiK contained complex soluble tags (Trx-Tag, His-Tag and S-Tag). A commonly conserved domain (HXHXDH-H-D) in the AHL lactonases has been detected in the amino acid sequence of AiiK. The results of *vivo* bioassays suggested that *E. coli* BL21 (DE3) cells harboring pET32a-*aaiK* has the ability to degrade N-Hexanoyl-L-homoserine lactone (C₆-HSL) with the decreasing of purple pigment violacein’s diameter (Fig. 2a), and the results of *vitro* bioassays elucidated that AiiK can degrade 10 μ M C₆-HSL within 1 h at 25 °C by the disappearance of purple pigment violacein in comparison to the CK (Fig. 2b). The enzyme assays have confirmed that the AiiK was capable of degrading AHLs, including C₆-HSL, N-(3-oxohexanoyl)-L-homoserine lactone (3-Oxo-C₆-HSL), N-decanoyl-L-homoserine lactone (C₁₀-HSL), and N-tetradecanoyl-L-homoserine lactone (C₁₄-HSL) (data not shown). The opened lactone ring of AHL, which was




Figure 2. (a) AHL-degrading activity bioassays *in vivo*. Subcultures of *E. coli* BL21 (DE3) cells harboring pET32a-*aaiK* were mixed with 10 μ M C₆-HSL and incubated at 37 °C for 0 h, 4 h, 6 h, 8 h, respectively. (b) AHL-degrading activity bioassays *in vitro*. The purified AiiK (4 μ g/mL) were mixed with 10 μ M C₆-HSL and incubated at 25 °C for 1 h. CK means controls of PBS with the same amount of C₆-HSL. The residual AHL was detected using strain *C. violaceum* CV026. (c) HPLC analysis of C₁₀-HSL degradation. CK means controls of PBS with the same amount of C₁₀-HSL; hydrolysis assay (AiiK: 4 μ g/mL, C₁₀-HSL: 50 μ M) was carried out at 25 °C for 10 min, and then SDS was used to terminate the reaction; acidified controls assay (AiiK: 4 μ g/mL, C₁₀-HSL: 50 μ M) was performed at 25 °C for 10 min, and SDS was used to terminate the reaction. The mixture was acidified to pH 2 by adding concentrated HCl and incubated at 25 °C overnight without agitation. The elution time of C₁₀-HSL was at 6.57 \pm 0.01 min and C₁₀-HSL was monitored at 210 nm. All experiments were carried out in triplicates.

| AHL | Mean \pm SD | | k_{cat}/K_m (s ⁻¹ ·mM ⁻¹) |
|---------------------------|------------------------------|---------------------------|--|
| | k_{cat} (s ⁻¹) | K_m (mM ⁻¹) | |
| C ₆ -HSL | 0.649 \pm 0.069 | 0.204 \pm 0.031 | 3.2 |
| 3-Oxo-C ₆ -HSL | 0.168 \pm 0.026 | 0.066 \pm 0.020 | 2.7 |
| C ₁₀ -HSL | 1.321 \pm 0.214 | 0.315 \pm 0.065 | 3.9 |
| C ₁₄ -HSL | 0.038 \pm 0.002 | 0.036 \pm 0.004 | 0.92 |

Table 1. Kinetic constants of AiiK for degrading AHLs. Note: Reactions were performed at pH 7.4 and 25 °C. And all data in this study were exhibited as mean \pm the standard deviation (SD).

catalyzed by AHL lactonase, was acidified at a low pH and the re-lactonization of the opened ring will be taken place resulting in an intact AHL^{18–20}. Hydrolysis assay (Fig. 2c) and acidified control assay (Fig. 2c) suggested that the opened lactone ring of C₁₀-HSL catalyzed by AiiK was re-lactonized when acidified at pH 2 over night, which indicated that AiiK was an AHL lactonase (Fig. 2c). And the way that hydrolysis assay and acidified control assay contributes a novel method to identifying AHL lactonase just by utilizing high-performance liquid chromatography (HPLC) analysis instead of using tandem mass spectrometry (MS). These results demonstrated that *aaiK* encodes an AHL lactonase belonging to AHL-degrading enzymes functioning as QQ enzymes. Hence, we named this gene the autoinducer inactivation gene from *K. huakuii* LAM0618^T (*aaiK*), which is the first autoinducer inactivation gene identified from the genus *Kurthia*.

Kinetic properties of AiiK. To evaluate the substrate specificity, kinetic constants of AiiK for degrading different AHLs ranging from the number of carbon atoms to the carbonylation at the C-3 position of the N-acyl chain were performed by using a series of different concentrations of substrates and corresponding calculated velocities. GraphPad Prism 5 software was utilized to calculate the K_m and k_{cat} values by quoting the Michaelis-Menten equation. The k_{cat} values for all substrates ranged from 0.038 s⁻¹ to 1.321 s⁻¹ and the K_m values of recombinant AiiK ranged from 0.036 mM⁻¹ to 0.315 mM⁻¹ (Table 1), demonstrating AiiK can hydrolyze varieties of AHLs.




Figure 3. Characterization of AiiK. (a) Optimum temperature for enzyme activity. (b) Temperature stability of AiiK. (c) Thermostability of AiiK at 37°C and 45°C. (d) Effect of pH on AiiK. All experiments were carried out in triplicates.




Figure 4. (a) Effect of metal ions and EDTA on AiiK activity. (b) Effect of zinc on AiiK activity *in vitro*. All experiments were carried out in triplicates. (PC stands for positive control, containing 4 μg/mL AiiK, 50 μM C₁₀-HSL, and 10 mM PBS (pH 7.4), then the reaction was performed at 45°C for 10 min).

Characterization of AiiK. The effects of temperature, pH, metal ions, and EDTA on AiiK were detected using C₁₀-HSL as substrate. The optimal temperature of AiiK for degrading C₁₀-HSL was 45°C. AiiK maintained relatively high activity at temperatures ranging from 18°C to 55°C and, interestingly, AiiK retained 27% of the maximum activity at 0°C (Fig. 3a). In terms of thermostability, AiiK possessed more than 95% activity when the temperature was lower than 45°C, and the relative activity significantly declined when the temperature exceeded 45°C. However, AiiK still displayed 28% activity after a preincubation at 65°C for 30 min (Fig. 3b). AiiK also exhibited approximately 20% activity after 12 h of preincubation at 37°C and 45°C (Fig. 3c). The optimal pH of AiiK for degrading C₁₀-HSL at 25°C was 7.5 (Fig. 3d). Low C₁₀-HSL-degrading activity was detected when the pH was below 6, and AiiK maintained more than 53% activity when the pH level ranged from 7 to 10 (Fig. 3d). AiiK was stable at pH values ranging from 6 to 8, maintaining over 70% activity after preincubations for 30 min at 25°C (data not shown).

AiiK activity increased with the addition of 1 mM Co²⁺, Mn²⁺, Mg²⁺, and Ni²⁺, with the most notable stimulation up to 152% with the addition of 1 mM Ni²⁺ (Fig. 4a). With the concentration of metal ions reaching at 5 mM, AiiK activity was all reduced. In addition, AiiK activity was extremely abolished in the presence of Cu²⁺ (Fig. 4a), which parallels MomL from *Muricauda olearia* Th120¹⁴. Moreover, AiiK activity decreased by approximately 19%




Figure 5. Resistance of AiiK to proteolysis. All experiments were carried out in triplicates. PC stands for positive control, containing 4 μg/mL AiiK, 50 μM C₁₀-HSL, and 10 mM PBS (pH 7.4), then the reaction was carried out at 45 °C for 10 min.




Figure 6. Effect of AiiK on biofilm formation in *P. aeruginosa* PAO1. **(a)** Biofilm formation by *P. Aeruginosa* PAO1 represented by crystal violet staining and planktonic cells of *P. Aeruginosa* PAO1. **(b)** The crystal violet staining of biofilm formed by *P. aeruginosa* PAO1. Lanes 1–3: 0 μg/mL of AiiK; Lanes 4–6: 5 μg/mL of AiiK; Lanes 7–9: 10 μg/mL of AiiK. **(c)** Fluorescence microscopy for visualization of biofilm formation by *P. aeruginosa* PAO1 stained with acridine orange.

and 45% after incubations with 1 and 5 mM EDTA, respectively (Fig. 4a). The effect of zinc on AiiK showed that a slight increase in activity of AiiK was observed with the increasing of Zn²⁺ concentration, then a maximum activity between 0.25 mM and 0.50 mM Zn²⁺, and a significant decrease was exhibited when the Zn²⁺ concentration exceeded 2.0 mM (Fig. 4b).

The resistance of AiiK to α-chymotrypsin, trypsin, and protease K was detected using C₁₀-HSL as substrate. AiiK maintained 87% and 101% activity after preincubations with α-chymotrypsin at 37 °C for 30 min and 60 min, respectively (Fig. 5). AiiK activity was enhanced up to 109% after preincubations with trypsin and protease K at 37 °C for 30 and 60 min (Fig. 5). It's very interesting that AiiK exhibits higher activity when preincubated with these proteases at 37 °C for 60 min than that of the same condition for 30 min; this activity is better than observed for AiiA_{A196}²¹. Therefore, these results suggested that AiiK exhibited great protease-resistance.

AiiK's application in QQ on *P. aeruginosa* PAO1. To assess the effect of AiiK on biofilm formation in *P. aeruginosa* PAO1, assays were carried out in a 96-well microtiter plate. Biofilm cells were stained with crystal violet to evaluate the total intact biomass in the microtiter plate under the same experimental conditions. It was found that the inhibition rate of biofilm formation was enhanced with the increasing of application amount of AiiK (Fig. 6a and b) and AiiK did not kill *P. aeruginosa* PAO1 from the O.D at 600 nm (Fig. 6a). *P. aeruginosa* PAO1 biofilm formation was visualized by fluorescence microscopy, indicating significant suppression of biofilm formation on special glass slides compared with the control (Fig. 6c).

It was determined whether AiiK affected extracellular proteolytic activity in *P. aeruginosa* PAO1 and the results declared that 10 μg/mL AiiK reduced extracellular proteolytic activity by 47.3% after incubation with *P. aeruginosa* PAO1 at 37 °C for 12 h (Fig. 7a). Moreover, pyocyanin production of *P. aeruginosa* PAO1 was




Figure 7. Effect of AiiK on virulence factor production. (a) The extracellular proteolytic activity of *P. aeruginosa* PAO1. (b) The pyocyanin production of *P. aeruginosa* PAO1. Bottle 1: 0 µg/mL of AiiK; Bottle 2: 5 µg/mL of AiiK; Bottle 3: 10 µg/mL of AiiK. All experiments were carried out in triplicates.

| AiiK concentration (µg/mL) | OD ₅₂₀ | pyocyanin production (µg/mL) |
|----------------------------|-------------------|------------------------------|
| 0 | 0.223 ± 0.013 | 3.807 ± 0.222 |
| 5 | 0.207 ± 0.011 | 3.534 ± 0.188 |
| 10 | 0.064 ± 0.007 | 1.092 ± 0.120 |

Table 2. Effect of AiiK on pyocyanin production in *P. aeruginosa* PAO1.

significantly decreased with the increasing of application amounts of AiiK (Fig. 7b and Table 2). Therefore, these results demonstrated that AiiK obviously inhibited the biofilm formation as well as the production of virulence factors consisting of extracellular proteolytic activity and pyocyanin production in *P. aeruginosa* PAO1, which contributes to the vital role of AiiK in QQ on *P. aeruginosa* PAO1.

Discussion

Homology analysis based on the identified AHL lactonases at the level of amino acid sequence showed that AiiK shares only 26% identity with AiiA, the first enzyme identified capable of AHL degradation, from the strain *Bacillus* sp. 240B1⁴ (Fig. 8). When assessing similarity with other AHL lactonases, it was found AiiK shares 42% identity with AiiB from *Agrobacterium fabrum* str. C58²², 26% identity with AidC from *Chryseobacterium* sp. StRB126²³, 31% identity with AttM from *A. fabrum* str. C58²⁴, 23% identity with MomL from *Muricauda olearia*¹⁴, and 26% identity with Aii20J from *Tenacibaculum* sp. 20J²⁵ (Fig. 8). The amino acid sequences of these AHL lactonases were retrieved from the NCBI or UniProt database. Sequence alignment was assessed using the MEGA 4.0 and GeneDoc. Compared with these AHL lactonases, the AHL-degradation activity of AiiK from *K. huakuii* LAM0618^T involves a conserved domain (HXHXDH-H-D) essential for QQ activity⁴ (Fig. 8). The HXHXD motif is essential to zinc-binding^{14,26,27}; therefore, we studied the effect of zinc on AiiK *in vitro* and the effect of zinc on AiiK was similar to that on AidC²⁸.

Analysis of the kinetic properties and characterization of AiiK revealed that AiiK exhibited a variable substrate spectrum and an efficient degradation of the AHL compounds used in this study, regardless of the number of carbon atoms of the *N*-acyl chain contained and whether there is carbonylation at the C-3 position. AiiK exhibited higher efficiencies at degrading C₆-HSL and C₁₀-HSL than 3-Oxo-C₆-HSL and C₁₄-HSL, demonstrating that AiiK presents a slightly preference for middle-long chain AHLs and is better at degrading unsubstituted AHLs at C-3 position than those with an oxo-substitution. In this study, C₁₀-HSL was the optimal substrate for AiiK at 25 °C and pH 7.4. Furthermore, only 4 µg/mL AiiK was required to completely hydrolyze 30 µM C₁₀-HSL within only 10 min at 25 °C compared to 20 µg/mL Aii20J from *Tenacibaculum* sp. 20J, which could hydrolyze only 10 µM C₁₀-HSL in 20 min at 22 °C²⁵. The optimal temperatures for AiiA_{SS10}²⁹, AiiA_{B546}³⁰, and AidC²³ to degrade AHLs ranged from 20 °C to 30 °C. This range was lower than that for MomL (40 °C)¹⁴, AiiK (45 °C, this study), and AiiM (50 °C)¹⁰. In terms of thermostability, AiiK displayed 28% and 16% activity after preincubations at 65 °C and 75 °C for 30 min, respectively. Meanwhile Aii20J retained approximately 20% activity after a preincubation at 60 °C for 10 min as well as MomL maintaining approximately 18% activity after a preincubation at 80 °C for 30 min^{14,25}. There are few cold-adapted AHL lactonases reported before, AiiK retained cold-adaption performance with 27% of the maximum activity at 0 °C.

As a common opportunistic pathogen, *P. aeruginosa* can cause severe diseases in plants and mammals, particularly in humans^{31,32}. The virulence factors of *P. aeruginosa* associated with QS include pyocyanin production, extracellular protease secretion, and biofilm formation, which cause complicated biofilm associated infections and stimulate drug resistance³³. These processes of QS are under the control of AHLs in *P. aeruginosa* reminds




Figure 8. Multiple sequence alignment of amino acid sequences of AiiK (WP_029500404) and other representative AHL lactonases. The amino acid sequences were retrieved from NCBI or UniProt database. Sequence alignment was assessed by using of the MEGA 4.0 and GeneDoc. AiiK homologue from *Bacillus* sp. KCTC 13219 (KYG91086) and *Solibacillus silvestris* (BAK54000) showed the highest score and second highest score by BLAST searching in NCBI database. Other amino acid sequences of AHL lactonase are the first enzyme AiiA from *Bacillus* sp. strain 240B1 (AAF62398), AiiB (NP 396590) and AttM (AAD43990) from *Agrobacterium fabrum* str. C58, AidC from *Chryseobacterium* sp. strain StRB126 (BAM28988), MomL from *Muricauda olearia* (AIY30473), and Aii20J (AKN24544) from *Tenacibaculum* sp. 20J. These lactonases share a conserved domain (HXHXDH-H-D) essential for AHLs degradation.

us that AHL lactonases can play a vital role in quenching these effects³⁴. AiiK was demonstrated to be an AHL lactonase with a wide substrate spectrum and efficient degradation of the AHL compounds as well as AiiK remained approximately 20% activity after 12 h of incubation at 37 °C and was resistant to proteases, including α -chymotrypsin, trypsin, and protease K at 37 °C, providing the basis for experimental research by using AiiK to inhibit processes of QS in *P. aeruginosa* PAO1. The great protease-resistance and stability at 37 °C of AiiK play a vital role in the application of AiiK on QQ in *P. aeruginosa* PAO1, because the strain *P. aeruginosa* PAO1 could secrete proteases by QS when cultured at 37 °C. Our research confirmed that AiiK did display great abilities of suppressing the biofilm formation, production of pyocyanin, extracellular proteolytic activity in *P. aeruginosa* PAO1 rather than killing *P. aeruginosa* PAO1 while the progress of biofilm formation, indicating this is a promising anti-pathogenic strategy by which to control these pathogenic bacteria and to prevent antibiotic resistance.

| Strain or plasmid | Description | Reference or source |
|---|---|------------------------|
| Strains | | |
| <i>Kurthia huakuii</i> LAM0618 ^T | Wild type | ACCC 06121 |
| <i>Escherichia coli</i> DH5 α | $\lambda^- \phi 80\text{dlacZ}\Delta\text{M15 } \Delta(\text{lacZYA-argF}) \text{ U169 recA1 endA hsdR17} (\text{r}_\text{K}^- \text{ m}_\text{K}^-) \text{ supE44 thi-1 gyrA relA1}$ | Tiagen |
| <i>Escherichia coli</i> BL21 (DE3) | F- <i>ompT hsdS_B (r_B-, m_B-) dcm gal</i> λ (DE3) pLysS Cm ^r | Tiagen |
| <i>Chromobacterium violaceum</i> CV026 | ATCC 31532 derivative, <i>cviI::Tn5xylE</i> Km ^r , Sm ^r | From Dr. Guishan Zhang |
| <i>Pseudomonas aeruginosa</i> PAO1 | Wild type | From Dr. Haijin Xu |
| Plasmids | | |
| pET32a | Cloning vector, Amp ^r | Novagen |
| pET32a- <i>aiiK</i> | pET32a containing <i>aiiK</i> gene | This study |

Table 3. Bacterial strains and plasmids.

Therefore, these abilities of AiiK empower a more valuable approach by quenching virulence factors and biofilm formation for controlling the biofilm-related infections caused by these pathogenic bacteria and expecting to be an anti-pathogenic drug.

In our study, the *aiiK* gene was the first autoinducer inactivation gene identified from the genus *Kurthia*. We firstly used hydrolysis assay and acidified control assay to confirm that AiiK was an AHL lactonase, which contributes to a novel method of identifying the AHL lactonases instead of using tandem MS. AiiK clearly quenched the effects of QS in *P. aeruginosa* PAO1 and did exhibit great characteristics such as good thermostability, cold adaptation, variable substrate spectrum, efficient degradation of AHLs, and great protease-resistance enable a wide application of AiiK, especially applied as a biocontrol agent or an anti-pathogenic drug. However, a detailed study about potential pharmaceutical application is still needed to investigate.

Materials and Methods

Bacterial strains, growth conditions, and chemicals. Selected bacterial strains and plasmids used in this study are listed in Table 3. The strains *K. huakuii* LAM0618^T and *P. aeruginosa* PAO1 were cultured in tryptic soy broth (TSB; Nippon Becton Dickinson, Tokyo, Japan) at 30 °C and 37 °C, respectively. *E. coli* DH5 α and BL21 (DE3) strains were propagated in Luria-Bertani (LB) medium at 37 °C and appropriate antibiotic was added when required (final concentration of 100 $\mu\text{g}/\text{mL}$ ampicillin). Reporter strain *C. violaceum* CV026 was fostered in LB medium containing 50 $\mu\text{g}/\text{mL}$ kanamycin at 30 °C. The biofilm formation of *P. aeruginosa* PAO1 was tested on a M9 minimal media supplemented with 0.4% (w/v) glucose. Pseudomonas broth (PB, 20 g of Bacto-Peptone (Difco), 10 g of K₂SO₄, and 1.4 g of MgCl₂) is a medium to promote the production of pyocyanin in liquid culture. And B-broth medium (5 g of yeast extract (Difco), 1 g of glucose, 10 g of casein hydrolysate 140 (Gibco), 5 g of NaCl and 1 g of K₂HPO₄, pH 7.3) was used for extracellular proteolytic activity assay of *P. aeruginosa* PAO1. C₆-HSL, 3-oxo-C₆-HSL, C₁₀-HSL, and C₁₄-HSL were purchased from Sigma-Aldrich (St. Louis, MO). All of these substrates were dissolved and prepared in chromatographic-grade methanol.

Cloning of the *aiiK* gene from *K. huakuii* LAM0618^T. A bacterial genome extraction kit (Tiagen, China) was used to extract the genomic DNA of *K. huakuii* LAM0618^T, and this genomic DNA was applied as the template for PCR amplification. The *aiiK* coding region of the LAM0618^T genome was amplified using FastPfu DNA polymerase (TransGen Biotech, Beijing, China) with primers *aiiKR* (5'-CCGGAATTCATGTGTCAAATAAAAAAGTTGTAC-3') and *aiiKF* (5'-CCCAAGCTTTTATTCGTAATACCCCTCCGTTGA-3') that containing *EcoRI* and *HindIII* restriction sites (underlined). The PCR was carried out in accordance with the following cycling parameters: 94 °C for 5 min, 94 °C for 30 s, 55 °C for 30 s, and 72 °C for 1 min for 31 cycles, followed by 72 °C for 10 min. Both PCR products and pET-32a vector were digested with endonucleases *EcoRI* and *HindIII*, and then the digested PCR products were linked (at 25 °C for 1 h) onto the *EcoRI-HindIII* sites of pET32a through T4 DNA ligase for contribution to expression plasmid pET32a-*aiiK*. The nucleotide sequence was sequenced by Life Technologies Company (China).

Expression and purification of recombinant AiiK. *E. coli* BL21 (DE3) cells harboring pET32a-*aiiK* were inoculated into 250 mL of fresh LB medium containing ampicillin at 37 °C and 180 rpm until an optical density at 600 nm of 0.6 was reached (approximately 3 h). Afterward, 0.5 mM IPTG was added to the culture medium, and then the cultures were incubated at 25 °C with moderate shaking (120 rpm) for 12 h. After incubation, the cultures were centrifuged at 10,000 rpm at 4 °C to harvest the *E. coli* BL21 (DE3) cells, and the cells were resuspended with lysis buffer (300 mM NaCl, 10 mM imidazole, and 50 mM Tris-HCl, pH 8.0). Lysozyme (final concentration of 1 mg/mL) and Benzoylase nuclease (final concentration of 20 U/mL) (QIAGEN, Germany) were added to the lysis buffer, followed by incubation on ice batch for 45 min with moderate shaking. The suspension mixture was sonicated on ice batch for 15 min and then centrifuged at 10,000 rpm for 20 min at 4 °C to remove the cell debris. Supernatants were loaded onto Ni-NTA (QIAGEN, Germany), resulting in target protein bound to Ni-NTA which was later washed at least twice with wash buffer (30 mM imidazole, 50 mM Tris-HCl, and 300 mM NaCl, pH 8.0). AiiK was eluted using elution buffer (300 mM imidazole, 50 mM Tris-HCl, and 300 mM NaCl, pH 8.0). Subsequently, NaCl and imidazole were removed from the eluted fraction by ultrafiltration (molecular weight cutoff of 10 kDa) with 10 mM phosphate buffer saline (PBS, pH 7.4)³⁵. SDS-PAGE and BAC protein assay kit (Shengye Biocolor, Shanghai) were used to measure the purity and concentrations of purified recombinant AiiK,

respectively. The purified recombinant AiiK protein (AiiK) was filtration sterilization by using 0.22 μm filters and then mixed with glycerol for storage at -20°C for further study.

AHL-degrading activity bioassays and enzyme assays. Reporter strain *C. violaceum* CV026 was endowed with the ability of evaluating AHL-degrading activity by producing the purple pigment violacein induced by short-chain AHLs³⁶. 1 mL culture of strain *C. violaceum* CV026 incubated at 30°C and 180 rpm for 16 h was mixed with 24 mL LB agar (1.6%) medium and poured in the plates. A 5.5 mm sterile paper disk was placed on an agar plate, and the AHL samples were applied on paper disk. Bioassay plates were incubated at 30°C for 16 h, and then the appearance of pigment was detected. For *in vivo* bioassays, *E. coli* BL21 (DE3) cells harboring pET32a-*aaiK* were precultured into 5 mL of fresh LB medium containing ampicillin and 0.5 mM IPTG at 37°C with rotary shaking at 180 rpm for 12 h. The culture was inoculated at a ratio of 2% into a new 5 mL LB medium containing ampicillin and $10\ \mu\text{M}$ C_6 -HSL, then incubated at 37°C for 0 h, 4 h, 6 h, 8 h, respectively. After incubation of different time, the culture supernatants were collected by centrifugation and used for the above-mentioned AHL-degrading activity bioassays. For *in vitro* bioassays, purified AiiK (final concentration of $4\ \mu\text{g}/\text{mL}$) was mixed with $10\ \mu\text{M}$ C_6 -HSL, incubated at 25°C for 1 h, and then used for the AHL-degrading activity bioassays immediately.

For the hydrolysis assay, the reaction mixture (500 μL) contained 5 μL AiiK (final concentration of $4\ \mu\text{g}/\text{mL}$), 50 μM AHLs (C_6 -HSL, 3-Oxo- C_6 -HSL, C_{10} -HSL, or C_{14} -HSL), and 10 mM PBS (pH 7.4). 2% SDS (m/v) was utilized to terminate the reaction after the mixture was incubated at 25°C for 10 min. The remaining AHLs were then extracted three times using isopycnic ethyl acetate. Subsequently, ethyl acetate was evaporated with nitrogen flux at 25°C , and the remaining AHLs were redissolved in 500 μL of methanol for HPLC analysis and quantification. For the control, which was handled and extracted via hydrolysis assay, comprised AiiK replaced with 10 mM PBS plus the same amount of AHLs. Furthermore, the acidified control assay, which contained the same amount of AiiK and AHLs, was performed at 25°C for 10 min and terminated by 2% SDS. The mixture was acidified to pH 2 by adding concentrated HCl and incubated at 25°C overnight without agitation, and then extracted following the hydrolysis assay.

An Agilent Technologies 1260 series HPLC system was applied to analyze and quantify AHLs degradation. The remaining AHLs were separated in an Agilent ZORBAX Eclipse Plus C18 column at 22°C with a constant flow rate of 0.7 mL/min in isocratic elution and then detected at 210 nm. Eution procedure consisted of an isocratic profile of acetonitrile/water (27:73, v/v) for C_6 -HSL, acetonitrile/water (30:70, v/v) for 3-Oxo- C_6 -HSL, acetonitrile/water (60:40, v/v) for C_{10} -HSL, and acetonitrile/water (80:20, v/v) for C_{14} -HSL.

Kinetic assay of AiiK activity. In order to determine the kinetics of AiiK for AHLs degradation, different concentration (25–150 μM , with an interval of 25 μM) of C_6 -HSL, 3-Oxo- C_6 -HSL, C_{10} -HSL, or C_{14} -HSL were added to PBS within AiiK ($4\ \mu\text{g}/\text{mL}$). After this reaction was carried out at 25°C for 10 min, the following procedure was based on the descriptions in enzyme assays section. The remaining AHLs were quantified by calculating the peak areas for a given retention time compared to those AHLs solutions of known concentrations via the analysis of HPLC. Furthermore, all assays were carried out in triplicates and AiiK replaced by 10 mM PBS plus the same amount of AHLs made up the control.

AiiK characterization. Enzymatic properties of AiiK were characterized based on enzyme assays with 50 μM C_{10} -HSL as the substrate. The optimal reaction temperature of AiiK was determined by evaluating activity at temperatures from 0°C to 85°C (0°C , 4°C , 18°C , 25°C , 30°C , 35°C , 40°C , 45°C , 50°C , 55°C , 65°C , 75°C , and 85°C). To evaluate the thermostability of AiiK, the residual activity of AiiK was measured after preincubations at different temperatures (25°C , 35°C , 45°C , 55°C , 65°C , 75°C , and 85°C) for 30 min. The optimal reaction pH of AiiK was estimated after incubations in a variety of pH levels, ranging from 2 to 11 ($\text{Na}_2\text{HPO}_4/\text{citric acid}$ for pH 2.0 to 7.0, Tris-HCl for pH 8.0 to 9.0, and Gly/NaOH for pH 10.0 to 11.0), for 10 min at 25°C . The pH stability of AiiK was performed via detecting the residual activity of AiiK after it was preincubated at different pH levels for 30 min at 25°C .

The effects of metal ions and EDTA on AiiK were determined by preincubation with 1 mM and 5 mM concentrations of various metal ions (Co^{2+} , Cu^{2+} , Fe^{2+} , Mn^{2+} , Ca^{2+} , Mg^{2+} , Ni^{2+} , and Zn^{2+}) and EDTA at 4°C for 30 min. The effect of zinc on AiiK was detected by preincubation with various concentrations of Zn^{2+} (0.01 mM–10 mM) at 4°C for 30 min. Subsequently, the residual activity of AiiK was measured at 45°C for 10 min using the same method as that of the enzyme assays described above.

To study resistance of AiiK to proteolysis, AiiK was preincubated with α -chymotrypsin, trypsin, and protease K (at a ratio of 1:10, w/w; protease/AiiK) in 10 mM PBS (pH 7.4) at 37°C for 30 min and 60 min. Subsequently, the residual activity of AiiK was measured at 45°C for 10 min by using the same method as that of the enzyme assays described above. All experiments were carried out in triplicates and the detailed experiments were performed under the precondition of more than 20% of the C_{10} -HSL had been remained.

Effect of AiiK on biofilm formation in *P. aeruginosa* PAO1. A static microtiter plate assay, modified from the research of Cady *et al.*³⁷ and Rajesh and Ravishankar Rai³⁸, was used to test the effect of AiiK on biofilm formation in *P. aeruginosa* PAO1. Strain *P. aeruginosa* PAO1 was precultured in TSB overnight at 37°C with rotary shaking at 150 rpm. After preincubation, the cultures were centrifuged at 10,000 rpm to harvest the *P. aeruginosa* PAO1 cells, and the cells were rinsed with sterile 10 mM PBS (pH 7.4) for three times. After the cells were resuspended in M9 minimal media with a concentration of 2×10^7 cfu/mL (determined through optical density and plate count assay), the *P. aeruginosa* PAO1 inocula were then premixed with AiiK (0, 5, or $10\ \mu\text{g}/\text{mL}$). For control wells, 10 mM PBS was added into the inocula instead of AiiK. A total of 200 μL of these cells/AiiK mixtures were dispensed into a 96-well microtiter plate and then incubated at 37°C for 24 h without shaking. Planktonic cells from the plate were transferred gently to a new 96-well microtiter plate to measure the optical

density at 600 nm. The cells at the bottom of the 96-well microtiter plate were stained with 20 μ L of 0.2% crystal violet at 25 °C for 15 min to estimate the quantity of biofilm formation. The wells were washed very gently with sterile distilled water three times until the crystal violet was removed. Finally, absorbance at 590 nm was detected after the crystal violet was extracted from the cells by adding 100 μ L of 95% ethanol.

The effects of AiiK on biofilm formation in *P. aeruginosa* PAO1 were visualized and assessed as previously reported with minor modifications³⁸. Briefly, the growth, resuspension and premixing with AiiK (0, 5, or 10 μ g/mL) of *P. aeruginosa* PAO1 were performed via static microtiter plate assay, and control samples were not treated with AiiK. Subsequently, 2 mL of these cells/AiiK mixtures were dispensed into sterile PA bottles containing special sterile glass slides, and biofilm can grow well on the 5 \times 5 mm special glass slides with incubation at 37 °C for 24 h without agitation. After incubation, the slides were dried and stained with 0.1% of acridine orange (Sigma, China) for 2.5 min. The biofilm cell attachment was observed under a fluorescence microscope (excitation at 490 nm and emission at 525 nm), and the inhibition of biofilm formation in AiiK-treated and control samples was compared visually.

Effect of AiiK on extracellular proteolytic activity in *P. aeruginosa* PAO1. The effect of AiiK on extracellular proteolytic activity in *P. aeruginosa* PAO1 was determined as previously described with modifications³⁹, where azocasein was used as substrate. *P. aeruginosa* PAO1, cultured at 37 °C for 12 h with rotary shaker at 150 rpm, was inoculated in B-broth containing different concentrations of AiiK (0, 5, or 10 μ g/mL). The absorbance of the culture was measured at 590 nm. The supernatants were then utilized as crude extract to determine the extracellular proteolytic activity of the culture through centrifugation of the culture at 12,000 rpm. The reaction mixture (500 μ L) for extracellular proteolytic activity assay contained 150 μ L of crude extract and 250 μ L of 2% azocasein dissolved in 10 mM PBS (pH 7.4) and was incubated at 37 °C for 30 min. A total of 1.2 mL of 10% trichloroacetic acid was added to terminate the reaction. The mixture was then incubated at 25 °C for 15 min. After this incubation, 1.2 mL supernatants of the mixture, which were collected by centrifugation at 3,000 g for 5 min, was mixed with 1 mL of 1 M NaOH in a new tube, and the measurement of absorbance was then determined at 440 nm. Furthermore, the effect of AiiK on extracellular proteolytic activity in *P. aeruginosa* PAO1 was measured as OD₄₁₅/OD₅₉₀.

Effect of AiiK on pyocyanin production in *P. aeruginosa* PAO1. The effect of AiiK on pyocyanin production in *P. aeruginosa* PAO1 was detected as previously described with some modifications⁴⁰. *P. aeruginosa* PAO1 was incubated in PB containing different concentrations of AiiK (0, 5, or 10 μ g/mL) at 37 °C for 18 h with rotary shaker at 150 rpm. A total of 5 mL culture supernatants collected by centrifugation at 12,000 rpm were mixed with 3 mL of chloroform, and then the mixture was reextracted by adding 1 mL of 0.2 M HCl. Subsequently, the measurement of absorbance was determined at 520 nm, and the concentration of pyocyanin was estimated by multiplying the optical density at 520 nm (OD₅₂₀) by 17.072.

Data availability statement. The authors confirm that all data are available.

References

- Fetzner, S. Quorum quenching enzymes. *Journal of biotechnology* **201**, 2–14 (2015).
- Yajima, A. Recent progress in the chemistry and chemical biology of microbial signaling molecules: quorum-sensing pheromones and microbial hormones. *Tetrahedron Letters* **55**, 2773–2780 (2014).
- Grandclement, C. *et al.* Quorum quenching: role in nature and applied developments. *FEMS microbiology reviews* **40**, 86–116 (2016).
- Dong, Y. H. *et al.* AiiA, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 3526–3531 (2000).
- Lin, Y. H. *et al.* Acylhomoserine lactone acylase from *Ralstonia* strain XJ12B represents a novel and potent class of quorum-quenching enzymes. *Molecular Microbiology* **47**, 849–860 (2003).
- Yang, F. *et al.* Quorum quenching enzyme activity is widely conserved in the sera of mammalian species. *FEBS letters* **579**, 3713–3717 (2005).
- Draganov, D. I. *et al.* Human paraoxonases (PON1, PON2, and PON3) are lactonases with overlapping and distinct substrate specificities. *Journal of lipid research* **46**, 1239–1247 (2005).
- Kim, M. H. *et al.* The molecular structure and catalytic mechanism of a quorum-quenching *N*-acyl-L-homoserine lactone hydrolase. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 17606–17611 (2005).
- Liu, D. *et al.* Mechanism of the quorum-quenching lactonase (AiiA) from *Bacillus thuringiensis*. 1. Product-bound structures. *Biochemistry* **47**, 7706–7714 (2008).
- Wang, W. Z. *et al.* AiiM, a novel class of *N*-acylhomoserine lactonase from the leaf-associated bacterium *Microbacterium testaceum*. *Applied and environmental microbiology* **76**, 2524–2530 (2010).
- Wang, L. H. *et al.* Specificity and Enzyme Kinetics of the Quorum-quenching *N*-Acyl Homoserine Lactone Lactonase (AHL-lactonase). *Journal of Biological Chemistry* **279**, 13645–13651 (2004).
- Dong, Y. H. *et al.* Identification of Quorum-Quenching *N*-Acyl Homoserine Lactonases from *Bacillus* Species. *Applied and environmental microbiology* **68**, 1754–1759 (2002).
- Czajkowski, R. & Jafra, S. Quenching of acyl-homoserine lactone dependent quorum sensing by enzymatic disruption of signal molecules. *Acta. Biochim. Pol.* **56**, 1–16 (2009).
- Tang, K. *et al.* MomL, a novel marine-derived *N*-acyl homoserine lactonase from *Muricauda olearia*. *Applied and environmental microbiology* **81**, 774–782 (2015).
- Marta, T. *et al.* HqiA, a novel quorum-quenching enzyme which expands the AHL lactonase family. *Sci. Rep.* **7**, 943, <https://doi.org/10.1038/s41598> (2017).
- Ruan, Z. *et al.* *Kurthia huakuii* sp. nov., isolated from biogas slurry, and emended description of the genus *Kurthia*. *International journal of systematic and evolutionary microbiology* **64**, 518–521 (2014).
- Ruan, Z. *et al.* Draft Genome Sequence of *Kurthia huakuii* LAM0618T, an Organic-Pollutant-Degrading Strain Isolated from Biogas Slurry. *Genome announcements* **2** (2014).
- Yates, E. A. *et al.* *N*-Acylhomoserine Lactones Undergo Lactonolysis in a pH-, Temperature-, and Acyl Chain Length-Dependent Manner during Growth of *Yersinia pseudotuberculosis* and *Pseudomonas aeruginosa*. *Infection and Immunity* **70**, 5635–5646 (2002).
- Dong, Y. H. *et al.* Quenching quorum-sensing-dependent bacterial infection by an *N*-acyl homoserine lactonase. *Nature* **411**, 813–817 (2001).
- Wah, S. *et al.* AidP, a novel *N*-Acyl homoserine lactonase gene from Antarctic *Planococcus* sp. *Sci. Rep.* **7**, 42968, <https://doi.org/10.1038/srep42968> (2017).

21. Cao, Y. *et al.* Orally administered thermostable *N*-acyl homoserine lactonase from *Bacillus* sp. strain AI96 attenuates *Aeromonas hydrophila* infection in zebrafish. *Applied and environmental microbiology* **78**, 1899–1908 (2012).
22. Liu, D. *et al.* Structure and Specificity of a Quorum-Quenching Lactonase (AiiB) from *Agrobacterium tumefaciens*. *Biochemistry* **46**, 11789–11799 (2007).
23. Wang, W. Z. *et al.* AidC, a novel *N*-acylhomoserine lactonase from the potato root-associated *cytophaga-flavobacteria-bacteroides* (CFB) group bacterium *Chryseobacterium* sp. strain StRB126. *Applied and environmental microbiology* **78**, 7985–7992 (2012).
24. Matthyse, A. G. *et al.* A region of the *Agrobacterium tumefaciens* chromosome containing genes required for virulence and attachment to host cells. *Biochimica. et. Biophysica. Acta.* **1490**, 208–212 (2000).
25. Mayer, C. *et al.* Aii20, a wide-spectrum thermostable *N*-acylhomoserine lactonase from the marine bacterium *Tenacibaculum* sp. 20J, can quench AHL-mediated acid resistance in *Escherichia coli*. *Applied microbiology and biotechnology* **99**, 9523–9539 (2015).
26. Thomas, P. W. *et al.* The quorum-quenching lactonase from *Bacillus thuringiensis* is a metalloprotein. *Biochemistry* **44**, 7559–7569 (2005).
27. Momb, J. *et al.* Mechanism of the Quorum-Quenching Lactonase (AiiA) from *Bacillus thuringiensis*. 2. Substrate Modeling and Active Site Mutations. *Biochemistry* **47**, 7715–7725 (2008).
28. Mascarenhas, R. *et al.* Structural and Biochemical Characterization of AidC, a Quorum-Quenching Lactonase with Atypical Selectivity. *Biochemistry* **54**, 4342–4353 (2015).
29. Qiu, J. *et al.* Enzymatic characterization and function of *N*-acylhomoserine lactonase SS10. *Acta. Phytopathol. Sinica.* **37**, 629–636 (2007).
30. Chen, R. *et al.* High yield expression of an AHL lactonase from *Bacillus* sp. B546 in *Pichia pastoris* and its application to reduce *Aeromonas hydrophila* mortality in aquaculture. *Microbial Cell Factories* **9**, 39 (2010).
31. Reimmann, C. *et al.* Genetically programmed autoinducer destruction reduces virulence gene expression and swarming motility in *Pseudomonas aeruginosa* PAO1. *Microbiology* **148**, 923–932 (2002).
32. Loukas, A. *et al.* Vaccination with recombinant aspartic hemoglobinase reduces parasite load and blood loss after hookworm infection in dogs. *PLoS medicine* **2**, e295 (2005).
33. Whiteley, M. *et al.* Gene expression in *Pseudomonas aeruginosa* biofilms. *Nature* **413**, 860–864 (2001).
34. Flickinger, S. T. *et al.* Quorum sensing between *Pseudomonas aeruginosa* biofilms accelerates cell growth. *Journal of the American Chemical Society* **133**, 5966–5975 (2011).
35. Guo, X. *et al.* Characterization of a highly thermostable and organic solvent-tolerant copper-containing polyphenol oxidase with dye-decolorizing ability from *Kurthia huakuii* LAM0618^T. *PLoS ONE* **11**, e0164810 (2016).
36. Morohoshi, T. *et al.* *N*-Acylhomoserine lactone regulates violacein production in *Chromobacterium violaceum* type strain ATCC 12472. *FEMS Microbiology Letters* **279**, 124–130 (2008).
37. Cady, N. C. *et al.* Inhibition of biofilm formation, quorum sensing and infection in *Pseudomonas aeruginosa* by natural products-inspired organosulfur compounds. *PLoS ONE* **7**, e38492 (2012).
38. Rajesh, P. S. & Ravishankar Rai, V. Quorum quenching activity in cell-free lysate of endophytic bacteria isolated from *Pterocarpus santalinus* Linn., and its effect on quorum sensing regulated biofilm in *Pseudomonas aeruginosa* PAO1. *Microbiological research* **169**, 561–569 (2014).
39. Ayora, S. & Giötz, F. Genetic and biochemical properties of an extracellular neutral metalloprotease from *Staphylococcus hyicus* subsp. *hyicus*. *Mol. Gen. Genet.* **242**, 421–430 (1994).
40. Essar, D. W. *et al.* Identification and characterization of genes for a second anthranilate synthase in *Pseudomonas aeruginosa*: interchangeability of the two anthranilate synthases and evolutionary implications. *J. Bacteriol.* **172**, 884–900 (1990).

Acknowledgements

This study was funded by National Natural Science Foundation of China (NSFC, No. 31670006), National Nonprofit Institute Research Grant of CAAS (No. 2014-30) and the Special Fund for Agro-scientific Research in the Public Interest (No. 201403019). We thank Haijin Xu of Nankai University for his kindly assistance in providing the strain *P. aeruginosa* PAO1 and Dr. Guishan Zhang for his kindly providing of strain *C. violaceum* CV026.

Author Contributions

S.M.Z. and Z.Y.R. conceived and supervised the study. W.W.D. designed the experiments, performed the experiments, acquired the data, and drafted the manuscript. J.Z., X.G., D.L.K., Q.Z., Y.Q.Z. and X.Y.L. performed statistical analyses and contributed to the manuscript. All authors reviewed the manuscript and provided meaningful intellectual contributions to the study.

Additional Information

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018