

Characterization of Microsatellite Loci in the Lichen Fungus *Lobaria pulmonaria* (Lobariaceae)

Author(s): Silke Werth , Carolina Cornejo , and Christoph Scheidegger

Source: Applications in Plant Sciences, 1(2) 2013.

Published By: Botanical Society of America

URL: <http://www.bioone.org/doi/full/10.3732/apps.1200290>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

CHARACTERIZATION OF MICROSATELLITE LOCI IN THE LICHEN FUNGUS *LOBARIA PULMONARIA* (LOBARIACEAE)¹

SILKE WERTH^{2,3}, CAROLINA CORNEJO², AND CHRISTOPH SCHEIDEGGER²

²Swiss Federal Research Institute WSL, Zuercherstrasse 111, CH-8903 Birmensdorf, Switzerland

- *Premise of the study:* Microsatellite loci were developed for the threatened haploid lichen fungus *Lobaria pulmonaria* to increase the resolution to identify clonal individuals, and to study its population subdivision.
- *Methods and Results:* We developed 14 microsatellite markers from 454 DNA sequencing data of *L. pulmonaria* and tested for cross-amplification with *L. immixta* and *L. macaronesica*. The number of alleles per locus ranged from two to 23. Nei's unbiased gene diversity, averaged over loci, ranged from 0.434 to 0.517 in the three studied populations.
- *Conclusions:* The new markers will increase the genetic resolution in studies that aim at disentangling clones in *L. pulmonaria* and may be useful for closely related species within *Lobaria* sect. *Lobaria*.

Key words: Ascomycetes; Ascomycota; lichen fungi; *Lobaria pulmonaria*; microsatellites; population subdivision.

Lobaria pulmonaria (L.) Hoffm. (Lobariaceae, Peltigerales) is a widely distributed lichen in the northern hemisphere and afro-temperate forests in South Africa. In central Europe, the species has faced a severe decline in the past decades, and the species is therefore of conservation concern. In the past decade, *L. pulmonaria* has become a model species for the population biology and conservation biology of lichens (Scheidegger and Werth, 2009). Eight microsatellite markers have thus far been published for the lichen fungus *L. pulmonaria*, a species that mainly reproduces clonally (Dal Grande et al., 2012; Werth and Scheidegger, 2012). Here, we develop 14 additional microsatellite markers to increase the genetic resolution for detailed studies of population subdivision and the reproductive system of this species, and we test for cross-amplification with two Macaronesian endemics closely related to *L. pulmonaria*—*L. immixta* Vain. and *L. macaronesica* C. Cornejo & Scheid.

METHODS AND RESULTS

We collected thallus fragments from 190 individuals from São Miguel Island, Azores (SM1: 37.85132°N, 25.78249°W), and from El Hierro, Canary Islands (SH2: 27.74317°N, 17.98651999°W; SH3: 27.73292°N, 18.01134°W) (see Appendix 1 for voucher information). Samples were air-dried after collection and stored at -20°C until DNA extraction using the DNeasy Plant Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. The fungus *L. pulmonaria* was grown in axenic culture from ascospores (voucher: GB1-10; 36.69658°N, 5.02322°W), and total genomic DNA isolated from the axenic culture was used for 454 pyrosequencing following standard protocols on a GS FLX instrument (Roche, Schlieren, Switzerland) performed at Microsynth

(Balgach, Switzerland). The data amounted to 233260 reads of an average length of 313.8 bases, in total 73 171 881 bases.

MSATCOMMANDER version 0.8.2 (Faircloth, 2008) was used to find di-, tri-, tetra-, penta-, and hexanucleotide repeats and design primers with Primer3 (Rozen and Skaletsky, 2000) using default values (annealing temperature [T_a] = 60°C, GC content 35–75%, primer length 19 bp) based on the 454 database. A total of 478 contigs contained microsatellite repeats (43 di-, 359 tri-, 37 tetra-, 11 penta-, and 28 hexanucleotides). For 304 of these contigs, flanking regions allowed primer design. Subsequently, 16 loci were tested, each with more than 11 repeats. Fourteen of these loci amplified successfully, and polymorphism was assessed based on 99 thalli of *L. pulmonaria*, 59 of *L. macaronesica*, and 32 of *L. immixta*. Multiplex PCR reactions were carried out in 5 µL reaction volumes using the M13 method (Schuelke, 2000), adding 1 µL of primer mix containing all loci to be labeled with the same fluorescent dye (primer without M13-tail: 0.15 µM, M13-tailed primer: 0.01 µM, dye-labeled M13 primer: $n \times 0.15$ µM, n = number of loci in multiplex), 2.5 µL Jump-Start *Taq* ReadyMix (Sigma-Aldrich, Buchs, Switzerland), 0.5 µL of genomic DNA, and 1 µL of ddH₂O. For primer sequences, see Table 1. PCR amplifications used an initial denaturation at 94°C for 5 min; followed by 30 cycles of 30 s at 94°C, 45 s at the annealing temperature of 60°C, 45 s at 72°C; followed by eight cycles of 30 s at 94°C, 45 s at 53°C, 45 s at 72°C to incorporate the dye-labeled M13 primer (5'-TGTAACGACGGCCAGT-3'); and a final extension at 72°C for 60 min.

Fragment analysis was performed after pooling PCR products labeled with four dyes, using GeneScan-500 LIZ (Life Technologies, Rotkreuz, Switzerland) as an internal size standard on a 3130xl Genetic Analyzer (Life Technologies). Genotyping was performed with GeneMapper version 3.7 (Life Technologies). Polymorphism was determined using our own code (available upon request) in R (R Development Core Team, 2011).

Of the 14 loci assessed, 13 amplified and were polymorphic in *L. pulmonaria*, eight in *L. macaronesica*, and seven in *L. immixta*. None of the loci amplified within a culture of *Dictyochochloropsis reticulata*, the green-algal photobiont of *L. pulmonaria*. The number of alleles per locus ranged from two to 23, and the maximum gene diversity was 0.846 (Table 2). Eight of the new markers and five of the previously published markers had 10 or more alleles in the studied individuals. Hence, the resolution for future studies will be increased. Moreover, with seven and eight loci working for *L. macaronesica* and *L. immixta*, the new markers appear to be useful for population genetic studies of closely related species in *Lobaria* sect. *Lobaria*.

CONCLUSIONS

By increasing the marker resolution, the newly developed polymorphic microsatellite loci will allow us to perform detailed

¹Manuscript received 9 June 2012; revision accepted 27 July 2012.

The authors thank the Swiss National Science foundation (grants 3100AO-105830 and 31003A_127346 to C.S.) for funding. The regional governments of the Canary Islands and Azores kindly provided collecting permits. Dr. Andreas Beck provided an axenic culture of *Dictyochochloropsis reticulata*.

³Author for correspondence: silke.werth@wsl.ch

TABLE 1. Overview of the microsatellite loci designed for the lichen fungus *Lobaria pulmonaria*.

Locus	Primer name	Primer sequences (5'–3') ^a	T _a (°C)	Repeat motif	Dye	Size (bp)	Multiplex	GenBank accession no.
LPu04843	M13-LPu04843F	F: TGTAAACGACGGCCAGTACCTACGAAGCCCAAGGTCG R: CATCGTCAGGCTTGAGCAC	60	(AAG) ₁₆	PET	304–389	1	JX126841
LPu08412	M13-LPu08412F	F: TGTAAACGACGGCCAGTACACAAAGGCTCCGGTAAAG R: ACCAGTCGAAACCCAGGAC	60	(CT) ₁₄	PET	220–236	1	JX126842
LPu17457	M13-LPu17457F	F: TGTAAACGACGGCCAGTGTAGGTCAGGAAGCACCG R: GGAGACACTCTGTAGCCACC	60	(CTGCTT) ₁₂	VIC	182–294	1	JX126845
LPu26427	M13-LPu26427F	F: TGTAAACGACGGCCAGTGCACGGGAGTATATCGGC R: CGCGAAGCGTATGATAAGAGG	59	(CTT) ₁₄	VIC	344–389	1	JX126846
LPu32425	LPu32425F	F: ATACCGAATACCGGCTCC	60	(AAG) ₁₈	NED	109–336	1	JX126847
LPu39713	M13-LPu32425R	R: TGTAAACGACGGCCAGTCTGCTGCTTGTACGAC	60	(GCT) ₁₉	FAM	226–428	1	JX126852
LPu40211	M13-LPu39713F	F: TGTAAACGACGGCCAGTCTTACCAGAGGTGGTTC R: TCCTTCTAATCCGAACCCCTG	57	(CTT) ₁₃	FAM	240–273	1	JX126854
LPu34888	M13-LPu40211F	F: TGTAAACGACGGCCAGTGGATGATGGCAACCG R: AACGGACCTGGGATTTCC	60	(ATC) ₁₄	FAM	315–360	2	JX126849
LPu38061	LPu34888F	F: TCGGACGATGTGGGAATGG	60	(AG) ₁₄	FAM	261–272	2	JX126851
LPu37451	M13-LPu38061R	R: ATTTCTGGAACCCGGACTG F: TGTAAACGACGGCCAGTATGCACAGCAGGTCAAACG	60	(AGT) ₁₃	NED	367–445	2	JX126850
LPu39912	LPu37451F	F: CGCGCCTTTGAACACCTG	60	(CTT) ₁₆	NED	303–388	2	JX126853
LPu13707	M13-LPu37451R	R: TGTAAACGACGGCCAGTCTCCGCTGGCAATAAACCG F: CATCGTCAGGCTTGAGCAC	60	(ACT) ₁₅	PET	165–516	2	JX126843
LPu30668	LPu39912F	F: TGTAAACGACGGCCAGTACCTACGAAGCCCAAGGTCG R: GGTGGAACAGAAAGGTGGC	60	(AG) ₁₃	NED	198–200	2	JX126848
LPu14122	LPu13707F	F: TGTAAACGACGGCCAGTCTCCGCTGGCAATAAACCG R: GGTGGAACAGAAAGGTGGC	60	(ATCAGT) ₁₁	PET	322–364	2	JX126844
	LPu30668F	F: GGTGGAACAGAAAGGTGGC	60					
	LPu14122F	F: GGTGGAACAGAAAGGTGGC	60					
	M13-LPu30668R	R: TGTAAACGACGGCCAGTACCTACGAAGCCCAAGGTCG F: GGTGGAACAGAAAGGTGGC	59					
	LPu14122F	F: GGTGGAACAGAAAGGTGGC	60					
	M13-LPu14122R	R: TGTAAACGACGGCCAGTAAATACGGCCGGGATCAGG F: CACCGCTGGAATAGGTAGC	58					

Note: T_a = annealing temperature.
^aThe M13-sequence is underlined.

TABLE 2. Polymorphism in 14 new and eight previously published microsatellite loci developed for the tree lungwort *Lobaria pulmonaria*, and cross-amplified for its close relatives *L. immixta* and *L. macaronensis*.^a

Locus ^b	Total		<i>L. pulmonaria</i> SH2 (N = 54)		<i>L. pulmonaria</i> SH3 (N = 37)		<i>L. pulmonaria</i> SM1 (N = 8)		<i>L. macaronensis</i> SM1 (N = 56)		<i>L. immixta</i> SM1 (N = 32)	
	N	A	A	H _c	A	H _c	A	H _c	A	H _c	A	H _c
LPu04843	190 [†]	19	9	0.831	10	0.785	2	0.250	5	0.702	1	0.000
LPu08412	190	5	4	0.539	4	0.685	3	0.607	—	—	—	—
LPu17457	190	10	4	0.523	3	0.419	4	0.750	—	—	2	0.272
LPu26427	96	5	1	0.000	1	0.000	2	0.250	2	0.198	2	0.175
LPu32425	190	10	2	0.201	2	0.105	4	0.821	3	0.482	2	0.516
LPu39713	190	7	2	0.331	2	0.450	1	0.000	3	0.395	2	0.121
LPu40211	96	10	6	0.652	3	0.520	3	0.607	2	0.071	3	0.567
LPu34888	190	10	3	0.461	2	0.054	5	0.893	2	0.226	6	0.756
LPu38061	190	4	3	0.542	2	0.054	3	0.667	1	0.000	—	—
LPu37451	190	14	10	0.846	6	0.668	4	0.821	—	—	—	—
LPu39912	190	19	9	0.835	10	0.785	2	0.250	5	0.705	1	0.000
LPu13707	190	23	6	0.440	4	0.683	3	0.464	11	0.615	2	0.389
LPu30668*	190	2	—	—	—	—	1	0.000	2	0.333	2	0.533
LPu14122*	190	6	—	—	—	—	4	0.750	1	0.000	2	0.272
Average		10.9	4.2	0.517	3.5	0.434	2.9	0.509	2.6	0.339	1.8	0.327
<i>LPu03</i>	190	3	2	0.075	1	0.000	1	0.000	1	0.000	—	—
<i>LPu09</i>	190	10	4	0.519	5	0.630	3	0.750	2	0.335	4	0.587
<i>LPu15</i>	190	16	4	0.623	5	0.714	3	0.679	10	0.867	2	0.121
<i>LPu23</i>	190	11	3	0.479	3	0.160	1	0.000	1	0.000	6	0.552
<i>LPu24</i>	190	5	2	0.042	1	0.000	1	0.000	1	0.000	1	0.000
<i>LPu25</i>	190	25	14	0.863	7	0.784	6	0.929	1	0.000	4	0.504
<i>LPu28</i>	190	40	13	0.898	9	0.786	3	0.679	18	0.858	2	0.175
<i>MS4</i>	190	3	2	0.082	2	0.056	1	0.000	—	—	—	—
Average		14.1	5.5	0.448	4.1	0.391	2.4	0.380	4.9	0.294	3.2	0.323

Note: A = number of alleles; H_c = Nei's unbiased gene diversity; N = total number of samples analyzed.

^aPopulations used in the study: SM1 = São Miguel, Azores, 37.85132°N, 25.78249°W; SH2 = El Hierro, Canary Islands, 27.74317°N, 17.98651999°W; and SH3 = El Hierro, Canary Islands, 27.73292°N, 18.01134°W.

^bLoci printed in italics have been published previously.

*Marker was not analyzed for populations SH2 and SH3 in *L. pulmonaria*.

[†]Overall number of samples includes three additional thalli of *L. macaronensis* from site SH2.

studies of the reproductive system of *L. pulmonaria* (and its close relatives), and hence aid our understanding of the population biology of a fascinating lichen symbiosis.

LITERATURE CITED

DAL GRANDE, F., I. WIDMER, H. H. WAGNER, AND C. SCHEIDEGGER. 2012. Vertical and horizontal photobiont transmission within populations of a lichen symbiosis. *Molecular Ecology* 21: 3159–3172.

FAIRCLOTH, B. C. 2008. MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92–94.

R DEVELOPMENT CORE TEAM. 2011. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

ROZEN, S., AND H. J. SKALETSKY. 2000. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.

SCHEIDEGGER, C., AND S. WERTH. 2009. Conservation strategies for lichens: Insights from population biology. *Fungal Biology Reviews* 23: 55–66.

SCHUELKE, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* 18: 233–234.

WERTH, S., AND C. SCHEIDEGGER. 2012. Congruent genetic structure in the lichen-forming fungus *Lobaria pulmonaria* and its green-algal photobiont. *Molecular Plant-Microbe Interactions* 25: 220–230.

APPENDIX 1. Herbarium vouchers of *Lobaria* species used in this study residing in the personal herbarium of Christoph Scheidegger at WSL. All samples analyzed are stored frozen at –20°C.

Site	Species	Voucher	Record
SM1	<i>L. immixta</i>	SM1-01a	14703
SM1	<i>L. macaronensis</i>	SM1-01c	14705
SM1	<i>L. pulmonaria</i>	SM1-01e	14707
SH2	<i>L. pulmonaria</i>	SH2-01a	10746
SH3	<i>L. pulmonaria</i>	SH3-02b	11170
GB1	<i>L. pulmonaria</i>	GB1-10	10164