

SHORT REPORT

Open Access

# Characterization of microsatellite markers developed from *Prosopis rubriflora* and *Prosopis ruscifolia* (Leguminosae - Mimosoideae), legume species that are used as models for genetic diversity studies in Chaquenan areas under anthropization in South America

Fábio M Alves<sup>1,2</sup>, Maria I Zucchi<sup>3</sup>, Ana MG Azevedo-Tozzi<sup>1</sup>, Ângela LB Sartori<sup>4</sup> and Anete P Souza<sup>1,2\*</sup>

## Abstract

**Background:** *Prosopis rubriflora* and *Prosopis ruscifolia* are important species in the Chaquenan regions of Brazil. Because of the restriction and frequency of their physiognomy, they are excellent models for conservation genetics studies. The use of microsatellite markers (Simple Sequence Repeats, SSRs) has become increasingly important in recent years and has proven to be a powerful tool for both ecological and molecular studies.

**Findings:** In this study, we present the development and characterization of 10 new markers for *P. rubriflora* and 13 new markers for *P. ruscifolia*. The genotyping was performed using 40 *P. rubriflora* samples and 48 *P. ruscifolia* samples from the Chaquenan remnants in Brazil. The polymorphism information content (PIC) of the *P. rubriflora* markers ranged from 0.073 to 0.791, and no null alleles or deviation from Hardy-Weinberg equilibrium (HW) were detected. The PIC values for the *P. ruscifolia* markers ranged from 0.289 to 0.883, but a departure from HW and null alleles were detected for certain loci; however, this departure may have resulted from anthropic activities, such as the presence of livestock, which is very common in the remnant areas.

**Conclusions:** In this study, we describe novel SSR polymorphic markers that may be helpful in future genetic studies of *P. rubriflora* and *P. ruscifolia*.

**Keywords:** *Prosopis*, Pantanal, Chaco, Population Genetics, Conservation, Short Tandem Repeats

## Findings

### Background

The genus *Prosopis* L. belongs to the Leguminosae botanical family, which contains 44 species. *Prosopis* L. is predominantly restricted to the neotropics [1]. *Prosopis rubriflora* [2] and *Prosopis ruscifolia* [3] are tree species

known locally as “espinheiro” and “algarroba,” respectively. These species are important both economically and ecologically. For example, the fruits and seeds of *P. ruscifolia* are reported to be good sources of nutrition for humans and animals [4], and the flowers of *P. rubriflora*, which are present throughout the year, provide important food resources, such as pollen and nectar, for the local fauna [5]. *P. rubriflora* has a narrow distribution range and is limited to Paraguay and Brazil, but *P. ruscifolia* is also found in Argentina and Bolivia [6,7].

In Brazil, *P. rubriflora* and *P. ruscifolia* are associated with Chaquenan areas [8] and are limited to the southern portion of the Pantanal [9,10]. Both species are excellent

\* Correspondence: anete@unicamp.br

<sup>1</sup>Departamento de Biologia Vegetal (BV), Instituto de Biologia, Universidade Estadual de Campinas - UNICAMP, Cidade Universitária Zeferino Vaz, CP 6109, CEP 13083-970 Campinas, SP, Brazil

<sup>2</sup>Centro de Biologia Molecular e Engenharia Genética (CBMEG) - Universidade Estadual de Campinas - UNICAMP, Cidade Universitária Zeferino Vaz, CP 6010, CEP 13083-970 Campinas, SP, Brazil

Full list of author information is available at the end of the article

**Table 1 Primers developed for *Prosopis rubriflora* and *Prosopis ruscifolia***

Marker	GenBank register no.	Primer sequences (5'-3')	Motifs	T <sub>a</sub> (°C)	Size <sup>a</sup> (bp)	Crossed amplification
Prb1	KF923365	F: AACTACCGCAGCACTTTTCAGA R: ACTACTTGGAGATGCCGTGGA	(gt)7	62.7	255-267	267 <sup>b</sup>
Prb2	KF923366	F: GAAAGCCGCGCTCCTAAG R: ATTCTTTTGTGTCTTGTCTTCTCG	(gc)4(ac)7	61.0	140-146	126 <sup>b</sup>
Prb3	KF923367	F: TCCAAAGACCGCAAGAAGAT R: AGGCCAAAAAGGACTCAAAAT	(ca)7	61.0	149-159	145 <sup>b</sup>
Prb4	KF923368	F: ATCCGATAAATACACCTTCTGG R: GGTGTATCGTAAAAGCCTGG	(ca)8	61.0	194-230	203 <sup>b</sup>
Prb5	KF923369	F: TTTAAACATTGCACGTGAACCTAT R: TTCACCCCTAAACCCCTT	(ac)9	56.4	149-155	-
Prb6	KF923370	F: TCATCTCTCAAAGAAAACGCACTC R: CCGCAGAGAAGCCCCTACATA	(tg)10	56.4	115-133	125 <sup>b</sup>
Prb7	KF923371	F: GGCTTAGCATCACCTCCAT R: CTTACCCCTTCAGTCCATTTACCA	(ac)8	61.0	219-225	220 <sup>b</sup>
Prb8	KF923372	F: CAACACCAAAACGCGCAGATGAT R: TTCGCCAAACGCCAGCATTAG	(gt)13	61.0	144-164	154 <sup>b</sup>
Prb9	KF923373	F: TTCTTCTCCTTCTTCATCTTCTCC R: ACAACGTTGATCCCAAAACCTAAG	(ac)9	62.7	167-175	190 <sup>b</sup>
Prb10	KF923374	F: TTTTGGTGGATTGATAGAGCC R: GAGTGGGGTCAAGAAAGAACAG	(tca)5	56.4	223	-
Prsc1	KC753210	F: AATGGAGTTTGTGTGTCTGTGG R: ATTACGGATACATCGAGCCTTCTT	(ac)9(ct)5	56.5	279-297	300 <sup>b</sup>
Prsc2	KC753211	F: GCGGAATCCAAACGACAA R: ACAGCAACACCCCTCACTCTCAA	(ac)9	64.7	224-252	250 <sup>b</sup>
Prsc3	KC753212	F: CCACAAGCACACGCACTCAGAC R: CCAGCACTAGACTTCGCCACCAAC	(ca)6	64.7	156-160	122
Prsc4	KC753213	F: CAAAATCCAACAAATAACACACC R: GGCGGATTCCTGGCTCTCT	(caa)2(ga)4	63.9	218-232	230 <sup>b</sup>
Prsc5	KC753214	F: CGCGTTAAGTCTGCCTTGCTTT R: CTCATGGTATTTCCCTTGTCTGCTCC	(gt)8	59.0	220-240	218
Prsc6	KC753215	F: CGAGCGGCGAAAAATGATAAA R: GCTGCTTCCCATAACTCTCTCCT	(gt)8	63.9	184-210	200
Prsc7	KC753216	F: CAGGGATTTAATCTCTTTGGGTAG R: ACAAGCTGGAAAGAGTCGCA	(tg)8...(gtgg)2(gt)5	59.0	122-156	122 <sup>b</sup>
Prsc8	KC753217	F: AGTGACGTGAACACGCTGAGG R: TGCTGATGTGTGTGTTTTGAGAT	(tg)10	62.7	98-120	114 <sup>b</sup>
Prsc9	KC753218	F: TCAGACTCCCGTAACCAG R: CGCACTCGAGCAGCATCT	(tg)9	59.0	112-122	-
Prsc10	KC753219	F: AACGCAACGCGCCGCAACTAT R: ACAAACGCTCGAATACTGGGGG	(ca)7(ct)7	56.5	260-284	-
Prsc11	KC753220	F: CCCGGCAACTCAAATCAACTTCATA	(ac)11	62.7	229-371	244 <sup>b</sup>

**Table 1 Primers developed for *Prosopis rubriflora* and *Prosopis ruscifolia* (Continued)**

		R: GGTCTAATTCTATTGGTGGGCTCTCTGG				
Prsc12	KC753221	F: GGGGTGCATGTTGGGGATTG R: TTTGGCCGGATTAACAGAGCA	(gt)10	59.0	185-223	220 <sup>b</sup>
Prsc13	KC753222	F: CTTACCATCACCGATTTCCCTT R: GCAACGAAGCAGCTGAAGAACAC	(ctt)5	62.7	102	116

T<sub>a</sub> - Optimal annealing temperature defined after gradient tests of the corresponding markers. <sup>a</sup>Range of the fragment sizes from the polymorphic markers and the sequenced size of the monomorphic markers; <sup>b</sup>Polymorphism observed for the transferred markers based on the 5 samples used.

indicators of Chaquenan areas in Brazil; *P. rubriflora* is usually associated with arboreal physiognomy, and *P. ruscifolia* is frequently associated with forest physiognomy. Both species can be used as models for genetic studies of diversity in these areas.

While estimating genetic diversity, the use of molecular markers has been helpful in defining alleles and studying genetic flow, population structure, paternity, inheritability, genetic maps and conservation genetics [11]. Simple sequence repeat markers (SSRs), commonly referred to as microsatellite markers, are desirable tools because they are co-dominant in nature, multi-allelic and widely distributed in the genome; they are also currently cheap, reproducible and relatively easy to analyze [12]. This work reports the development, characterization and transferability of microsatellite markers for *P. rubriflora* and *P. ruscifolia*.

**Construction of a microsatellite-enriched library**

DNA was extracted from *P. rubriflora* and *P. ruscifolia* using the DNeasy® Plant Mini Kit (Qiagen, Hilden, DE) according to the manufacturer's instructions. Microsatellite-enriched libraries for *P. rubriflora* and *P. ruscifolia* were constructed as described by Billote et al. [13]. The genomic DNA was digested with *AfaI* after enrichment

with streptavidin-coated magnetic beads (Streptavidin MagneSphere Paramagnetic Particles, Promega, Madison, WI); biotinylated (CT)<sub>8</sub> and (GT)<sub>8</sub> microsatellite probes were added for the dinucleotide-enriched library. The fragments were amplified by PCR and cloned into the pGEM-T vector (Promega, Madison, WI). XL1-Blue (*Escherichia coli*) competent cells were transformed with the recombinant plasmids and then cultivated on agar medium containing ampicillin (100 mg/ml), X-galactosidase 2% (100 µg/ml) and IPTG (100 mM). The selected clones were added to a Big Dye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA) and sequenced using an ABI 377 sequencer (Applied Biosystems, Foster City, CA). The sequences were aligned and edited using SeqMan Software (DNAStar, Madison, WI), and the adapters and restriction sites were removed using Microsat Software (A. M. Risterucci, CIRAD, personal communication). To identify microsatellite-enriched regions, we used the Simple Sequence Repeat Identification Tool (SSRIT) [14] and defined the following numbers of repeats/motifs: five/dinucleotides, four/trinucleotides and three/tetra- or pentanucleotides. After these steps, primers were designed using the PrimerSelect software (DNAS-tar, Madison, WI).

**Table 2 Markers developed for *Prosopis rubriflora***

Marker	N <sub>a</sub>			H <sub>o</sub>		H <sub>e</sub>		PIC	Null alleles		HW (P-value)	
	T	FSM	FSV	FSM	FSV	FSM	FSV		FSM	FSV	FSM	FSV
Prb1	7	6	7	0.526 <sup>a</sup>	0.474	0.522 <sup>a</sup>	0.737	0.602	0.044	0.145	0.348	0.045
Prb2	3	1	3	0.000	0.150 <sup>a</sup>	0.000	0.145 <sup>a</sup>	0.073	0.001	0.000	-	1.000
Prb3	3	3	2	0.316	0.150	0.428	0.296	0.313	0.103	0.134	0.101	0.069
Prb4	12	11	8	0.650	0.850 <sup>a</sup>	0.799	0.803 <sup>a</sup>	0.766	0.092	0.000	0.457	0.291
Prb5	4	4	4	0.500	0.600	0.583	0.683	0.576	0.039	0.007	0.689	0.254
Prb6	6	4	6	0.500 <sup>a</sup>	0.400	0.458 <sup>a</sup>	0.432	0.413	0.000	0.000	1.000	0.335
Prb7	4	4	4	0.684 <sup>a</sup>	0.650 <sup>a</sup>	0.605 <sup>a</sup>	0.499 <sup>a</sup>	0.473	0.000	0.000	0.355	0.121
Prb8	5	5	3	0.500	0.250	0.524	0.304	0.370	0.000	0.050	0.821	0.468
Prb9	10	10	9	0.684	0.650	0.835	0.819	0.791	0.053	0.062	0.177	0.016

FSM - Fazenda São Manoel, FSV - Fazenda Santa Vergínia, N<sub>a</sub> - Number of alleles, H<sub>o</sub> - Observed heterozygosity, H<sub>e</sub> - Expected heterozygosity, PIC - Polymorphism information content, P-values of Hardy-Weinberg (HW) equilibrium (P-value > 0.0055 after Bonferroni correction), null alleles (null frequency < 0.20). <sup>a</sup>Populations where the values of H<sub>o</sub> were higher than those of H<sub>e</sub>.

**Table 3 Markers developed for *Prosopis ruscifolia***

Marker	N <sub>a</sub>			H <sub>o</sub>		H <sub>e</sub>		PIC	Null alleles		HW (P-value)	
	T	FRC	ECD	FRC	ECD	FRC	ECD		FRC	ECD	FRC	ECD
Prsc1	7	4	7	0.600 <sup>a</sup>	0.783	0.566 <sup>a</sup>	0.828	0.701	0.000	0.020	0.916	0.009
Prsc2	8	7	5	0.480	0.565	0.553	0.761	0.658	0.032	0.114	0.034	0.00 <sup>c</sup>
Prsc3	3	3	3	0.360 <sup>a</sup>	0.348	0.344 <sup>a</sup>	0.456	0.348	0.000	0.048	0.674	0.073
Prsc4	5	4	3	0.040	0.043	0.321	0.275	0.289	0.247 <sup>b</sup>	0.221 <sup>b</sup>	0.000 <sup>c</sup>	0.000 <sup>c</sup>
Prsc5	6	4	5	0.440 <sup>a</sup>	0.273	0.378 <sup>a</sup>	0.654	0.484	0.000	0.218 <sup>b</sup>	1.000	0.000 <sup>c</sup>
Prsc6	8	4	8	0.320	0.696	0.653	0.801	0.703	0.196	0.068	0.001 <sup>c</sup>	0.087
Prsc7	9	6	7	0.440	0.565	0.727	0.779	0.755	0.167	0.081	0.001 <sup>c</sup>	0.000 <sup>c</sup>
Prsc8	7	4	7	0.320	0.174	0.577	0.789	0.656	0.165	0.338 <sup>b</sup>	0.008	0.000 <sup>c</sup>
Prsc9	5	3	3	0.240	0.130	0.280	0.559	0.430	0.040	0.267 <sup>b</sup>	0.484	0.000 <sup>c</sup>
Prsc10	7	5	5	0.286	0.227	0.633	0.758	0.670	0.212 <sup>b</sup>	0.294 <sup>b</sup>	0.000 <sup>c</sup>	0.000 <sup>c</sup>
Prsc11	17	8	12	0.560	0.652	0.845	0.884	0.883	0.157	0.127	0.003	0.000 <sup>c</sup>
Prsc12	5	4	5	0.600	0.591	0.569	0.707	0.589	0.000	0.075	0.592	0.013

FRC - Fazenda Retiro Conceição, ECD - Estação do Carandazal, N<sub>a</sub> - Number of alleles, H<sub>o</sub> - Observed heterozygosity, H<sub>e</sub> - Expected heterozygosity, PIC - Polymorphism information content, P-values of Hardy-Weinberg (HW) equilibrium (P-value > 0.0041 after Bonferroni correction). <sup>a</sup>Populations where the values of H<sub>o</sub> were higher than those of H<sub>e</sub>; <sup>b</sup>Possible null alleles (null frequency < 0.20); <sup>c</sup>Departure from HW equilibrium was observed.

### Fragment amplification

The fragments were amplified using polymerase chain reactions containing 8 ng of template DNA, 2 mM MgCl<sub>2</sub>, 50 mM KCl, 20 mM Tris-HCl (pH 8.4), 0.2 mM dNTPs, 0.19 mg/ml BSA (bovine serum albumin), 0.15 mM of each primer and 1 U of *Taq* DNA polymerase; the reactions were then brought to a final volume of 20 µl with ultrapure water. To define the temperatures for the PCR reactions, we adopted the guidelines described by Mottura *et al.* [15]; for the annealing temperatures, we used a gradient program with temperatures ranging from 65°C to 55°C. The samples were collected in the Chaco remnants of Corumbá and Porto Murinho, Mato Grosso do Sul, Brazil. Twenty *P. rubriflora* samples were collected in each of two Chaco remnant locations: Fazenda São Manoel (FSM) (21°47'44.5"S; 57°39'34.6"W) and Fazenda Santa Vergínia (FSV) (22°06'40.5"S; 57°49'57.6"W). Twenty-three *P. ruscifolia* samples were collected in Estação do Carandazal (ECD) (19°48'33.2"S; 57°10'11.0"W), and 25 samples were collected in Fazenda Retiro Conceição (FRC) (21°42'23.7"S; 57°45'58.2"W). The cross-amplification of the markers was evaluated in 5 *P. rubriflora* samples obtained from FRC (21°41'00.7"S; 57°46'43.8"W) and 5 *P. ruscifolia* samples from Chácara Jacaré (21°41'20.1"S; 57°52'15.5"W) using the same conditions as for the native species. The amplified samples were genotyped by vertical electrophoresis using denaturing polyacrylamide gels (6%), and DNA bands were visualized using silver nitrate [16]; the sizes of the resulting fragments were estimated by comparison to a 10-bp DNA ladder (Invitrogen, Carlsbad, CA). Statistical analyses were performed using Microsatellite Toolkit v.3.1.1 [17] to calculate the expected heterozygosity (H<sub>e</sub>),

observed heterozygosity (H<sub>o</sub>) and polymorphism information content (PIC). The Genepop software v.1.2 [18] was used to estimate adherence to Hardy-Weinberg (HW) equilibrium and possible linkage disequilibrium (LD), and the frequency of null alleles was estimated using FreeNA [19].

### Results and discussion

We designed 32 primer pairs: 13 for *P. rubriflora* and 19 for *P. ruscifolia*. However, only 10 of the *P. rubriflora* primer pairs and 13 of the *P. ruscifolia* primer pairs amplified properly. The nine remaining pairs of primers were discarded because amplification errors were observed in the preliminary tests. Polymorphisms were detected in 9 of the native *P. rubriflora* markers and 12 of the native *P. ruscifolia* markers; only one marker from each species had a monomorphic pattern based on the populations analyzed. Eight markers from *P. rubriflora* successfully cross-amplified and were polymorphic for the tested samples, and 2 markers failed during cross-amplification. Eleven *P. ruscifolia* markers were successfully cross-amplified; 7 were polymorphic, and 2 failed this analysis (Table 1).

The number of *P. rubriflora* alleles in the sampled remnants ranged from 3 to 12; the polymorphism information content (PIC) values of these markers ranged from 0.073 to 0.791, the observed heterozygosity (H<sub>o</sub>) ranged from 0.000 to 0.850, and the expected heterozygosity (H<sub>e</sub>) ranged from 0.000 to 0.835. No evidence of null alleles was observed, and no departure from Hardy-Weinberg equilibrium was observed (Table 2). No significant linkage disequilibrium (LD) was observed for any of the markers of this species after Bonferroni correction (P-value for 5% = 0.001389). The number of *P. ruscifolia*

alleles in both of the remnants ranged from 3 to 17, the PIC values ranged from 0.289 to 0.883, the  $H_o$  values ranged from 0.040 to 0.783, and the  $H_e$  values ranged from 0.275 to 0.884. Possible null alleles were observed for the markers Prsc5, Prsc8 and Prsc9 from one remnant (ECD), and the markers Prsc4 and Prsc10 had possible null alleles in both remnants. A departure from HW was observed for Prsc2, Prsc5, Prsc6, Prsc8, Prsc9 and Prsc11 in one of the remnants (the majority were observed in ECD) and for Prsc4, Prsc7 and Prsc10 in both remnants (Table 3). Significant LD was observed for the loci Prsc5 and Prsc6 after Bonferroni correction ( $P$ -value for 1% = 0.00016).

Higher values of  $H_o$  were observed for the Prb1, Prb2, Prb4, Prb6, Prb7, Prsc1, Prsc3 and Prsc5 markers in this study; these higher values may indicate that an insufficient number of samples was collected or may be related to the reproductive patterns of these populations. The ECD populations are highly disturbed, and the FRC population is currently recovering from a relatively recent suppression (within the last 15 years); these factors may underlie the observed departure from HW and the presence of null alleles. A study with new and conserved populations may produce better results for these markers.

These markers are the first microsatellite markers developed for *Prosopis rubriflora* and *Prosopis ruscifolia*, and together with the set of *P. ruscifolia* markers amplified by Bessega et al. [20], they are expected to be useful tools for studies of the conservation genetics, reproductive biology, phylogeography and taxonomy of these species.

#### Availability of supporting data

The original sequences of the developed markers were submitted to the GenBank database (<http://ncbi.nlm.nih.gov>), and the registered codes are available in Table 1.

The testimony samples were deposited at Herbarium Universidade Estadual de Campinas (UEC – Campinas, SP, BR) and registered according to the following: *P. rubriflora* – 74477 (Fazenda São Manoel – Porto Murтинho, MS), 154715 (Fazenda Santa Vergínia – Porto Murтинho, MS); *P. ruscifolia* – 74469 (Fazenda Retiro Conceição – Porto Murтинho, MS), 37266 (Estação do Carandazal – Corumbá, MS).

#### Competing interests

The authors declare that no competing interests exist regarding this research.

#### Authors' contributions

FMA developed and characterized the markers and drafted the manuscript; MIZ provided support during the statistical analysis; ALBS provided support for the plant collection in the field; AMGAT participated in the experimental design; APS conceived the study, participated in its design and coordination and helped draft the manuscript. All of the authors read and approved the final manuscript.

#### Acknowledgments

This work received financial support from Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP, proc. 2010/51242-6) as a graduate fellowship to FMA, Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) as a Research fellowship to APS, and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES/CNPq (Casadinho/Procad # 552352/2011-0).

#### Author details

<sup>1</sup>Departamento de Biologia Vegetal (BV), Instituto de Biologia, Universidade Estadual de Campinas - UNICAMP, Cidade Universitária Zeferino Vaz, CP 6109, CEP 13083-970 Campinas, SP, Brazil. <sup>2</sup>Centro de Biologia Molecular e Engenharia Genética (CBMEG) - Universidade Estadual de Campinas - UNICAMP, Cidade Universitária Zeferino Vaz, CP 6010, CEP 13083-970 Campinas, SP, Brazil. <sup>3</sup>Pólo Apta Centro Sul, Rod. SP 127 km 30, Piracicaba, SP CP 28, CEP 13400-970, Brazil. <sup>4</sup>Departamento de Biologia, Centro de Ciências Biológicas e da Saúde (CCBS), Universidade Federal de Mato Grosso do Sul – UFMS, Cidade Universitária s/n, Caixa Postal 549, CEP 79070-900 Campo Grande, MS, Brazil.

Received: 9 January 2014 Accepted: 12 June 2014

Published: 18 June 2014

#### References

- Lewis GP, Schrire B, Mackinder B, Lock M: *Legumes of the World*. Kew: Royal Botanic Gardens; 2005.
- Hassler E: *Repert Spec Nov Regni Veg* 1910, **8**:552.
- Grisebach A: *Plantae Lorentzianae. Abh Kön Ges Wiss Göttingen* 1874, **19**:130.
- Freyre M, Astrada E, Blasco C, Baigorria C, Rozycki V, Bernardi C: **Valores nutricionais de frutos de vinal (*Prosopis ruscifolia*): consumo animal e humano. *Ciência y Tecnología Alimentaria* 2003, **4**(1):41–46.**
- Freitas TG, Souza CS, Aoki C, Arakaki LMM, Stefanello TH, Sartori ALB, Sigrist MR: **Flora of Brazilian humid Chaco: composition and reproductive phenology. *Check List* 2013, **9**(5):973–979.**
- Burkart AA: **A monograph of the genus *Prosopis* (Leguminosae Subfam. Mimosoideae). *J Arnold Arbore* 1976, **57**:219–247.**
- Prado DE, Gibbs PE, Pott A, Pott VJ: **The Chaco-Pantanal transition in southern Mato Grosso, Brazil. In *Nature and Dynamics of Forest-Savanna Boundaries*. Edited by Furlley PA, Proctor J, Ratter JA. London: Chapman and Hall; 1992:451–470.**
- Pott A, Pott VJ: *Plantas do Pantanal*. Corumbá: EMBRAPA; 1994.
- Furtado PP, Guimarães JG, Fonzar BC: **Vegetação - As regiões fitoecológicas, sua natureza e seus recursos econômicos. In *Projeto RADAMBRASIL. Folha SE 21. Campo Grande: geologia, geomorfologia, pedologia, vegetação e uso potencial da terra. Volume 28*. Edited by Divisão de Publicação. Rio de Janeiro: MME/SG/Projeto RADAMBRASIL; 1982:281–336.**
- Loureiro RL, Lima JPS, Fonzar BC: **Vegetação - As regiões fitoecológicas, sua natureza e seus recursos econômicos. In *Projeto RADAMBRASIL. Folha SE 21. Corumbá e parte da folha SE.20: geologia, geomorfologia, pedologia, vegetação e uso potencial da terra. Volume 27*. Edited by Divisão de Publicação. Rio de Janeiro: MME/SG/Projeto RADAMBRASIL; 1982:329–366.**
- Conner JK, Hartl DL: *A Primer of Ecological Genetics*. Massachusetts: Sinauer Associates; 2004.
- Ferreira ME, Grattapaglia D: *Introdução ao uso de marcadores moleculares em análise genética*. Brasília: Embrapa Recursos Genéticos e Biotecnologia; 1998.
- Billotte N, Lagoda P, Risterucci AM, Baurens FC: **Microsatellite-enriched libraries: applied methodology for the development of SSR markers in tropical crops. *Fruits* 1999, **54**(4):277–288.**
- Temnykh S, DeClerck G, Lukashova A, Lipovich L: **Computational and experimental analysis of microsatellites in rice (*Oryza sativa* L.): Frequency, length variation, transposon associations, and genetic marker potential. *Genome Res* 2001, **11**:1441–1452.**
- Mottura MC, Finkeldey R, Verga AR, Gailing O: **Development and characterization of microsatellite markers for *Prosopis chilensis* and *Prosopis flexuosa* and cross-species amplification. *Mol Ecol Notes* 2005, **5**:487–489.**
- Creste S, Tulmann-Neto A, Figueira A: **Detection of single repeat polymorphisms in denaturing polyacrylamide sequencing gels by silver staining. *Plant Mol Biol Report* 2001, **19**:229–306.**

17. Park SDE: Trypanotolerance in West African Cattle and the Population Genetic Effects of Selection. *PhD thesis*. University of Dublin, Ireland; 2001.
18. Raymond M, Rousset F: GENEPOP (Version 1.2): Population Genetics Software for Exact Tests and Ecumenicism. *J Hered* 1995, **86**(3):248–249.
19. Chapuis MP, Estoup A: Microsatellite null alleles and estimation of population differentiation. *Mol Biol Evol* 2007, **24**(3):621–631.
20. Bessega CF, Pometti CL, Miller JT, Watts R, Saidman BO, Vilardi JC: New microsatellite loci for *Prosopis alba* and *P. chilensis* (Fabaceae). *Appl Plant Sci* 2013, **1**(5):1200324.

doi:10.1186/1756-0500-7-375

**Cite this article as:** Alves et al.: Characterization of microsatellite markers developed from *Prosopis rubriflora* and *Prosopis ruscifolia* (Leguminosae - Mimosoideae), legume species that are used as models for genetic diversity studies in Chaquenian areas under anthropization in South America. *BMC Research Notes* 2014 **7**:375.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

