

Characterization of the complete mitochondrial genome of the Rock pigeon, *Columba livia* (Columbiformes: Columbidae)

X.Z. Kan, X.F. Li, L.Q. Zhang, L. Chen, C.J. Qian, X.W. Zhang and L. Wang

The Provincial Key Laboratory of the Conservation and Exploitation Research of Biological Resources in Anhui, College of Life Sciences, Anhui Normal University, Wuhu, China

Corresponding author: X.Z. Kan
E-mail: xianzhaokan@gmail.com

Genet. Mol. Res. 9 (2): 1234-1249 (2010)

Received March 4, 2010

Accepted April 3, 2010

Published June 29, 2010

DOI 10.4238/vol9-2gmr853

ABSTRACT. The rock pigeon (*Columba livia*), or Rock dove, is a member of the bird family Columbidae. We mapped the complete mitochondrial genome of the Rock pigeon. The mitochondrial genome of this species is a circular molecule of 17,229 bp in length, encoding a standard set of 13 protein-coding genes, two ribosomal RNA genes, and 22 transfer RNA genes, plus a putative control region, demonstrating a structure very similar to that of other birds. As found in other vertebrates, most of these genes are coded on the H-strand, except for NADH dehydrogenase subunit 6 (*nad6*) and eight tRNA genes (Gln, Ala, Asn, Cys, Tyr, Ser(UCN), Pro, Glu). The AT skew and GC skew of the whole genome, protein-coding genes, tRNA, rRNA, and the control region were calculated for the complete mitochondrial genomes of 30 avian species, representing 29 orders. All protein-coding genes initiated with ATG, except for *cox1* and *nad5*, which began with GTG. One extra nucleotide ‘C’ was present in NADH dehydrogenase subunit 3 (*nad3*). All tRNA gene sequences have the potential to fold into typical

cloverleaf secondary structures. Within the control region, conserved sequences were identified in three domains. Although the conserved blocks, such as ETAS1, ETAS2, CSB1, CSB1-like, and boxes C, D, E, and F, are readily identifiable in the *C. livia* control region, the typical origin of H-strand replication (O_H), CSB2 and CSB3 could not be detected. These results provide basic information for phylogenetic analyses of birds, especially Columbiformes species.

Key words: Mitochondrial genome; Rock pigeon; *Columba livia*; Columbiformes

INTRODUCTION

There are around 10,000 living species of birds, making them the most numerous tetrapod vertebrates. All modern birds are classified within the subclass Neornithes, which has two subdivisions: the Palaeognathae, containing mostly flightless birds, like ostriches, and the very diverse Neognathae, containing all other birds (Livezey and Zusi, 2007). Columbiformes is one of the most easily recognized avian orders worldwide and is traditionally subdivided into two families, the Columbidae and the Raphidae (Pereira et al., 2007). The Columbidae currently include over 300 species of pigeons and doves. The Rock pigeon (*Columba livia*), or Rock dove, is a member of the family Columbidae. This species includes the domestic pigeon, the feral pigeon, and the wild Rock pigeon. Wild Rock pigeons are pale grey, with two black bars on each wing; although domestic and feral pigeons are very variable in color and pattern. The Rock pigeon has a restricted natural resident range in western and southern Europe, north Africa, and south Asia. This species has been domesticated for several thousand years, giving rise to the domestic pigeon. Many domestic birds have escaped or have been released over the years, and have given rise to the feral pigeon. Feral pigeons are found in large numbers in cities and towns all over the world.

Genome-scale approaches have played an important role in the inference of evolutionary relationships among diverse organisms (Rokas et al., 2003). Compared with the nuclear genome, the mitochondrial genome has several intrinsic characteristics (small genome size, fast substitution rate and maternal inheritance mode), which render the mitogenome as one of the most suitable markers for phylogenetic (Wang et al., 2008) and population genetics studies, such as analysis of gene flow, hybridization and introgression (Moore, 1995).

Typical metazoan mitochondrial genomes are double-stranded circular molecules, ranging in size from approximately 15 to 20 kb and containing 37 genes: 13 protein-coding genes, 22 transfer RNA genes (tRNAs) and two ribosomal RNA genes (rRNAs) (Boore, 1999). Additionally, one major control region is believed to control the initiation of replication and transcription of animal mitochondrial DNA (mtDNA) (Shadel and Clayton, 1997). To date, nearly 110 complete mtDNA sequences are available for 28 avian species (Table 1); however, no complete mitochondrial sequence has been published for members of the order Columbiformes, except for one sequence deposited in GenBank (NC_013244). Here, we present the complete mitochondrial genome of the Rock pigeon, *C. livia* (Columbiformes: Columbidae) and give a thorough description of its genome features in comparison to other avian species.

Table 1. List of avian species examined in this study, classified according to Howard and Moore (2003).

Order	Family	Species	Accession No.	Reference
Neognathae				
Anseriformes	Anatidae	<i>Branta canadensis</i>	NC_007011	Unpublished data ^a
Apodiformes	Apodidae	<i>Apus apus</i>	NC_008540	Unpublished data ^b
Caprimulgiformes	Aegothelidae	<i>Aegotheles cristatus</i>	NC_011718	(Pratt et al., 2009)
Charadriformes	Haematopodidae	<i>Haematopus ater</i>	NC_003713	(Paton et al., 2002)
Ciconiiformes	Ciconiidae	<i>Ciconia ciconia</i>	NC_002197	Unpublished data ^c
Columbiformes	Columbidae	<i>Hemiphaga novaeseelandiae</i>	NC_013244	Unpublished data ^d
Coraciiformes	Coraciidae	<i>Eurystomus orientalis</i>	NC_011716	(Pratt et al., 2009)
Cuculiformes	Neomorphidae	<i>Geococcyx californianus</i>	NC_011711	(Pratt et al., 2009)
Falconiformes	Falconidae	<i>Falco peregrinus</i>	NC_000878	(Mindell et al., 1998)
Galliformes	Phasianidae	<i>Gallus gallus</i>	NC_007236	(Nishibori et al., 2005)
Gaviiformes	Gaviidae	<i>Gavia stellata</i>	NC_007007	(Slack et al., 2006)
Gruiformes	Rhynochetidae	<i>Rhynchochets jubatus</i>	NC_010091	(Morgan-Richards et al., 2008)
Passeriformes	Menuridae	<i>Menura novaehollandiae</i>	NC_007883	(Slack et al., 2007)
Pelecaniformes	Phaethontidae	<i>Phaethon rubricauda</i>	NC_007979	Unpublished data ^e
Phoenicopteriformes	Phoenicopteridae	<i>Phoenicopterus ruber</i>	NC_010089	(Morgan-Richards et al., 2008)
Piciformes	Ramphastidae	<i>Pteroglossus azara</i>	NC_008549	(Gibb et al., 2007)
Podicipediformes	Podicipedidae	<i>Tachybaptus novaehollandiae</i>	NC_010095	(Morgan-Richards et al., 2008)
Procellariiformes	Diomedeidae	<i>Thalassarche melanophrys</i>	NC_007172	(Slack et al., 2006)
Psittaciformes	Psittacidae	<i>Melopsittacus undulatus</i>	NC_009134	Unpublished data ^f
Rheiformes	Rheidae	<i>Rhea americana</i>	NC_000846	(Harlid et al., 1998)
Sphenisciformes	Spheniscidae	<i>Eudyptula minor</i>	NC_004538	(Slack et al., 2003)
Strigiformes	Strigidae	<i>Ninox novaeseelandiae</i>	NC_005932	(Harrison et al., 2004)
Struthioniformes	Struthionidae	<i>Struthio camelus</i>	NC_002785	(Haddrath and Baker, 2001)
Tinamiformes	Tinamidae	<i>Tinamus major</i>	NC_002781	(Haddrath and Baker, 2001)
Trochiliformes	Trochilidae	<i>Archilochus colubris</i>	NC_010094	(Morgan-Richards et al., 2008)
Trogoniformes	Trogonidae	<i>Trogon viridis</i>	NC_011714	(Pratt et al., 2009)
Palaeognathae				
Apterygiformes	Apterygidae	<i>Apteryx haastii</i>	NC_002782	(Haddrath and Baker, 2001)
Casuariiformes	Casuariidae	<i>Casuarius casuarius</i>	NC_002778	(Haddrath and Baker, 2001)
Dinornithiformes	Dinornithidae	<i>Dinornis giganteus</i>	NC_002672	(Cooper et al., 2001)

Unpublished data: ^aSnyder JC, Snider AR, Senecal AJ, Disantis EJ, et al.; ^bSlack KE, Delsuc F, McLenaghan PA, Bartosch-Haerlid A, et al.; ^cYamamoto Y; ^dGibb GC, Goldberg J, Trewick SA, Powlesland RG, et al.; ^eYamamoto Y, Kakizawa R and Yamagishi S; ^fGuan X, Samuels DC and Smith EJ.

MATERIAL AND METHODS

Sample collection and DNA extraction

An adult domestic male Rock pigeon was collected in Wuhu, Anhui Province, China. Total genomic DNA was extracted from the muscle tissue using standard phenol/chloroform methods (Sambrook and Russell, 2001).

PCR amplification and sequencing

Natural transfer of DNA from the mitochondria to the nucleus generates nuclear copies of mtDNA (NUMTs) and is an ongoing evolutionary process (Hazkani-Covo et al., 2010). To minimize the possibility of obtaining NUMTs, two long overlapping fragments

(~12 kb in length) were first amplified using the long and accurate-polymerase chain reaction (LA-PCR) kit (Takara, Dalian, China). The first LA-PCR primer set was LA16SF (5'-CCTACGTGATCTGAGTTCAGACCGGAGCAATCCAG-3') of Nishibori et al. (2001) and CytbR252 (5'-GATGCAGATGAAGAAGAATGAGGGCGCCGTTGC-3') designed based on the sequence of the cytochrome b gene (*cob*) from Cabot's tragopan. The second primer set was LACLMTF (5'-GAAGCATCATCTCCCACCTAGAACATTGAAAATTC-3') and LACLMTR (5'-CCCTCCACCAGCAGGATCAAAGAAGGTAGTGTT-3') based on certain sequences of *C. livia* using the above LA-PCR primer set (Table 1). The LA-PCR was conducted at 94°C for 1 min, followed by 35 cycles consisting of 10 s denaturation at 98°C and 13 min annealing and extension at 68°C, with a final extension step of 10 min at 72°C, in a DNA thermal cycler TC-3000 (Techne, Barloworld Scientific Ltd., UK). The amplified fragments with a size of approximately 12 kb that were obtained were used as the templates for amplification of the short overlapping fragments (1.1-1.5 kb in length) with 15 primer sets (Table 2). All 15 sets of primers were designed based on the conserved sequences of the known mitochondrial genome sequences in Galliformes or on the determined sequences of fragments of the mitogenome of *C. livia* in this study, which were aligned using CLUSTAL X v2.0.10 (Larkin et al., 2007). Each primer set amplified an mtDNA fragment containing an overlap of at least 100 bp with the adjacent amplified fragment at both termini. PCR was carried out in a volume of 25 µL containing 5-50 ng DNA template, 6.25 pmol of each primer, 0.2 mM of each dNTP, 2 mM MgCl₂, and 0.625 U *DreamTaq*TM DNA polymerase (Fermentas, Burlington, Canada) or Ex *Taq*

Table 2. Primers used in amplifying and sequencing *Columba livia*.

Fragment	Primer name	Primer sequence (5'-3')	Primer size (bp)
1	CLMT1F	TATCACACCTCCCTACACCG	20
	CLMT1R	TCGTGATGGATACTTCTCGC	20
2	AVMT2F	AACCCATTATAATGATACGG	20
	AVMT2R	TIACTGCTGAGTACCCGTGG	20
3	AVMT3F	GCAAAAGACTTAGTCCTAACCC	21
	AVMT3R	CTTTGCGACAGAGACGGGTT	21
4	AVMT4F	AAAGTCGTAACAAGGTAAGTGAC	22
	AVMT4R	CGCCCCAACCGAAAAATGTC	20
5	AVMT5F	AAGACGAGAAAGACCCGTGG	20
	CLMd5F	CGCAGTACCAATCCTAAC	19
6	CLMT5R	CGGTTTGTTCAAGCGAGATT	21
	AVMT6F	TAAGCACCCCTGGCCATCACC	20
7	CLMT6R	GTTAGAAGGGTGAGGTTTG	20
	AVMT7F	ACACAGACAGGAAAATCCT	20
8	CLMT7R	AAGGTCTACAGAGGCTCCG	19
	AVMT8F	CGCATAAATAACATAAGCTC	21
9	AVMT8R	GAAGCATTAAAGTGGTTTGAT	20
	AVMT9F	AAGCCTTCTCAGCAAAACGA	20
10	AVMT9R	GCTTAGGTTCATGGTCAGGT	20
	AVMT10F	ATGACATGCCCAATTAAACC	21
11	AVMT10R	GATGGCTTGTGTTTCGGTTTC	21
	AVMT11F	CAAGCCTAGCCCCAACACCAAG	21
12	AVMT11R	ATGGGGTTAGTCAGTGTAGGC	21
	ColMT12F	CTCATGGTCAGATCACTTAC	21
13	ColMT12R	GTTCCCTCGTTGGGTTATTAG	20
	AVMT13F	ACTACGAACGGACACACAGCCG	22
14	AVMT13R	GAAGG CCAAA TTGAG CGGAT	20
	AVMT14F	ATGACAAGGACGAGCTGAAG	20
15	AVMT14R	ATTATTTTAGTAGGGGGTGT	20
	AVMT15F	GCCAA CCTTC ATCTC ACCATAA	22
	CLMd15R	AGGGTTGCAGGCGTGAAGT	20
	AVMT15R	CTTGTGCGTGGGTTGTCTCGGG	22

DNA polymerase (TaKaRa, Dalian, China). Amplification was conducted in a DNA thermal cycler TC-3000, as above. PCR cycles were as follows: one cycle of 4 min at 70°C, 4 cycles of 40 s at 94°C, 20 s at 52°C, and 2 min and 10 s at 72°C, followed by 36 cycles of 20 s at 94°C, 20 s at 50–55°C, and 2 min and 10 s at 72°C. The process was completed with a final extension step at 72°C for 10 min. The band with the right size was cut out and purified using an EZ Spin Column DNA Gel Extraction Kit (Bio Basic Inc.) and then cloned with pGEM®-T Easy Vector System II (Promega). Internal primers were applied to fragments 5 and 15 (Table 2). All distinct clones were sequenced on an ABI-PRISM 3730 sequencer using a BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems) and the corresponding primer.

Gene identification and genome analyses

DNA sequences were analyzed using BioEdit 7.0.9.0 (Hall, 1999) and DAMBE 4.1.19 (Xia and Xie, 2001) softwares. Contig assembly was performed with the ContigExpress program (a component of Vector NTI Suite 6.0). The boundaries of protein-coding genes and rRNA genes were initially identified via DOGMA (Wyman et al., 2004) using the default setting, and refined by alignment with mitochondrial genomes of other avian species (Table 1). Composition skewness was calculated according to the formulas (AT skew = [A - T] / [A + T]; GC skew = [G - C] / [G + C]) (Perna and Kocher, 1995). Most tRNA genes were identified using tRNAscan-SE1.21 (Lowe and Eddy, 1997) under the ‘cove only’ search mode, with the vertebrate mitochondrial genetic code and ‘mito/chloroplast’ source. Some tRNA genes that were not found by the tRNA-SE1.21 were identified by proposed secondary structures and anti-codons (Shen et al., 2009). A gene map of the mitochondrial genome of *C. livia* was initially generated with OGDRAW (Lohse et al., 2007) and then modified manually.

RESULTS AND DISCUSSION

Genome organization and base composition

The complete sequence of the mtDNA of *C. livia* is 17,229 bp in length (Figure 1 and Table 3) and was deposited in GenBank with the accession No. (GU908131). The size of avian mtDNA ranges from 16,595 bp (*Ninox novaeseelandiae*) to 18,967 bp (*Thalassarche melanophrys*). Similar to the typical mitochondrial genomes of other vertebrates, the Rock pigeon mtDNA consists of 13 typical protein-coding genes, 22 tRNA genes, 2 rRNA genes (*srRNA* and *lrRNA*; small and large rRNA subunits, respectively) and one putative control region (D-loop) (Table 3). As found in other vertebrates, most of these genes are coded on the H-strand, except for one protein-coding gene (*nad6*) and eight tRNA genes (*tRNA^{Gln}*, *tRNA^{Ala}*, *tRNA^{Asn}*, *tRNA^{Cys}*, *tRNA^{Tyr}*, *tRNA^{Ser(UCN)}*, *tRNA^{Pro}*, *tRNA^{Glu}*).

The overall base composition of the H-strand is as follows: A (30.15%), T (23.98%), G (13.98%), C (31.90%); the A+T content of *C. livia* (54.13%) is similar to those of other birds (ranging from 52.46 to 57.56%, Table 4). As in most vertebrates, the overall base composition is skewed against guanine in the *C. livia* mitochondrial genome, which is due to a strong bias against the use of guanine at the third codon position (San Mauro et al., 2004).

The nucleotide composition of the *C. livia* mitogenome is slightly biased toward A+T nucleotides (54.13%, Table 4), which is a higher percentage than that of *P. azara* (52.46%),




Figure 1. Gene map of the mitochondrial genome of *Columba livia*. Genes encoded on the heavy or light strands are shown outside and inside the circular gene map, respectively. The inner ring displays the GC content. Twenty-two tRNA genes are designated by single-letter amino acid codes. This figure was initially generated with OGDRAW and modified manually.

P. rubricauda (52.75%), *B. canadensis* (52.78%), *N. novaehollandiae* (52.8%), *R. americana* (53.06%), *C. ciconia* (53.67%), and *G. gallus* (53.99%), but lower than in the other 22 avian species, ranging from 54.17 to 57.56%. Within 13 protein-coding genes in the *C. livia* mitochondrial genome, the A+T composition is the highest in the *atp8* gene (55.95%), and the lowest in the *cob* gene (51.53%).

Table 3. Localization and features of genes in the mitochondrial genome of *Columba livia*.

Gene/region	Strand	Position		Size		Codon		Anticodon	Intergenic nucleotides ^b
		From	To	Nucleotide	Amino acids	Start	Stop ^a		
<i>tRNA^{Phe}</i>	H	1	69	69				GAA	0
<i>srRNA</i>	H	70	1,042	973					0
<i>tRNA^{Ile}</i>	H	1,043	1,114	72				TAC	0
<i>lRNA</i>	H	1,115	2,700	1586					0
<i>tRNA^{Leu(UUR)}</i>	H	2,701	2,774	74				TAA	11
<i>nad1</i>	H	2,786	3,751	966	324	ATG	AGA		17
<i>tRNA^{Le}</i>	H	3,769	3,839	71				GAT	5
<i>tRNA^{Gln}</i>	L	3,845	3,915	71				TTG	-1
<i>tRNA^{Met}</i>	H	3,915	3,983	69				CAT	0
<i>nad2</i>	H	3,984	5,023	1040	346	ATG	TA-		0
<i>tRNA^{Trp}</i>	H	5,024	5,094	71				TCA	1
<i>tRNA^{Ala}</i>	L	5,096	5,164	69				TGC	2
<i>tRNA^{Asn}</i>	L	5,167	5,239	73				GTT	2
<i>tRNA^{Cys}</i>	L	5,242	5,308	67				GCA	-1
<i>tRNA^{Tyr}</i>	L	5,308	5,379	72				GTA	1
<i>cox1</i>	H	5,381	6,931	1551	516	GTG	AGG		-9
<i>tRNA^{Ser(UCN)}</i>	L	6,923	6,996	74				TGA	2
<i>tRNA^{Asp}</i>	H	6,999	7,067	69				GTC	2
<i>cox2</i>	H	7,070	7,753	684	227	ATG	TAA		1
<i>tRNA^{Lys}</i>	H	7,755	7,825	71				TTT	1
<i>atp8</i>	H	7,827	7,994	168	54	ATG	TAA		-10
<i>atp6</i>	H	7,985	8,668	684	227	ATG	TAA		-1
<i>cox3</i>	H	8,668	9,451	784	261	ATG	T-		0
<i>tRNA^{Gly}</i>	H	9,452	9,520	69				TCC	0
<i>nad3</i>	H	9,521	9,872	352		ATT	TAA		1
<i>tRNA^{Arg}</i>	H	9,874	9,942	69				TCG	1
<i>nad4L</i>	H	9,944	10,240	297	98	ATG	TAA		-7
<i>nad4</i>	H	10,234	11,611	1378	459	ATG	T-		0
<i>tRNA^{His}</i>	H	11,612	11,680	69				GTG	0
<i>tRNA^{Ser(AGY)}</i>	H	11,681	11,746	66				GCT	-1
<i>tRNA^{Leu(CUN)}</i>	H	11,746	11,817	72				TAG	0
<i>nad5</i>	H	11,818	13,632	1815	604	GTG	AGA		11
<i>cob</i>	H	13,644	14,786	1143	380	ATG	TAA		0
<i>tRNA^{Thr}</i>	H	14,787	14,855	69				TGT	6
<i>tRNA^{Pro}</i>	L	14,862	14,931	70				TGG	45
<i>nad6</i>	L	14,977	15,498	522	173	ATG	TAG		3
<i>tRNA^{Glu}</i>	L	15,502	15,572	71				TTC	0
<i>D-loop</i>	H	15,573	17,229	1657					0

^a“-” Indicates termination codons completed via polyadenylation. ^bNegative values represent overlapping nucleotides.

The AT skew and GC skew were calculated for the complete mitogenomes of 30 avian species, representing 29 orders (Table 4). The AT skew for the *C. livia* mitochondrial genome is slightly positive (0.114), indicating the occurrence of more As than Ts. We found similar results in the other 29 avian species (0.068 to 0.180). In contrast, the GC skew in all 30 avian mitochondrial genomes was strongly negative (-0.346 to 0.438), meaning that there is a heavy bias toward Cs and against Gs. Furthermore, one extra nucleotide ‘C’ is present in *nad3* of *C. livia*, which is consistently observed in many other birds and some turtles and is thought not to be translated (Slack et al., 2003). The function of the extra “C” in *nad3* and its phylogenetic implications are still unclear.

Table 4. Composition and skewness in avian mitochondrial genomes.

Species	Size (bp) ^a	A%	C%	G%	T%	A+T%	AT skew	GC skew
Whole genome								
<i>A. apus</i>	17,037	31.37	31.52	13.19	23.92	55.29	0.135	-0.410
<i>A. colubris</i>	16,356	30.62	31.27	14.22	23.86	54.48	0.124	-0.375
<i>A. cristatus</i>	18,607	30.67	30.54	13.63	25.10	55.77	0.100	-0.383
<i>A. haastii</i>	16,980	30.91	29.48	13.33	26.24	57.15	0.082	-0.377
<i>B. canadensis</i>	16,760	30.18	32.07	15.14	22.60	52.78	0.144	-0.359
<i>C. casuarinus</i>	16,756	30.47	29.73	14.45	25.25	55.72	0.094	-0.346
<i>C. ciconia</i>	17,347	30.54	31.98	14.35	23.13	53.67	0.138	-0.381
<i>C. livia</i>	17,229	30.15	31.90	13.98	23.98	54.13	0.114	-0.391
<i>D. giganteus</i>	17,070	30.67	31.25	13.93	24.15	54.82	0.119	-0.383
<i>E. minor</i>	17,611	30.96	31.84	13.53	23.67	54.63	0.133	-0.404
<i>E. orientalis</i>	17,774	30.26	31.74	13.82	24.18	54.44	0.112	-0.393
<i>F. peregrinus</i>	18,068	32.74	30.87	13.57	22.77	55.51	0.180	-0.389
<i>G. californianus</i>	17,091	32.34	31.70	12.40	23.55	55.89	0.157	-0.438
<i>G. gallus</i>	16,785	30.25	32.49	13.52	23.74	53.99	0.121	-0.412
<i>G. stellata</i>	17,573	30.36	31.43	14.39	23.81	54.17	0.121	-0.372
<i>H. ater</i>	16,791	31.59	31.12	13.67	23.62	55.21	0.144	-0.390
<i>H. novaeseelandiae</i>	17,264	30.97	31.89	13.21	23.92	54.89	0.128	-0.414
<i>M. undulates</i>	18,193	31.13	31.73	12.96	24.19	55.32	0.125	-0.420
<i>M. novae Hollandiae</i>	17,839	30.11	30.36	14.45	25.08	55.19	0.091	-0.355
<i>N. novae Hollandiae</i>	16,223	30.83	33.40	13.78	21.97	52.80	0.168	-0.416
<i>P. azara</i>	18,736	28.44	34.10	13.44	24.02	52.46	0.084	-0.435
<i>P. rubber</i>	17,446	31.54	32.03	13.49	22.93	54.47	0.158	-0.407
<i>P. rubricauda</i>	17,777	29.41	32.95	14.29	23.34	52.75	0.115	-0.395
<i>R. american a</i>	16,714	28.69	32.20	14.74	24.37	53.06	0.081	-0.372
<i>R. jubatus</i>	16,937	30.32	30.36	14.15	25.17	55.49	0.093	-0.364
<i>S. camelus</i>	16,595	30.47	30.40	14.21	24.92	55.39	0.100	-0.363
<i>T. major</i>	16,701	30.48	29.88	13.05	26.56	57.04	0.069	-0.392
<i>T. melanophrys</i>	18,967	31.45	30.53	13.64	24.38	55.83	0.127	-0.382
<i>T. novae Hollandiae</i>	18,002	31.75	32.06	13.12	23.08	54.83	0.158	-0.419
<i>T. viridis</i>	17,751	30.73	28.82	13.62	26.83	57.56	0.068	-0.358
Protein-coding genes								
<i>A. apus</i>	11,388	29.00	32.74	13.03	25.22	54.22	0.070	-0.431
<i>A. colubris</i>	11,394	28.59	32.23	13.83	25.35	53.94	0.060	-0.400
<i>A. cristatus</i>	11,388	28.77	32.04	13.57	25.62	54.39	0.058	-0.405
<i>A. haastii</i>	11,385	29.16	30.41	13.10	27.33	56.49	0.032	-0.400
<i>B. canadensis</i>	11,412	27.97	33.04	15.12	23.86	51.83	0.079	-0.372
<i>C. casuarinus</i>	11,394	28.23	30.65	14.12	27.01	55.24	0.022	-0.369
<i>C. ciconia</i>	11,400	28.42	33.89	13.63	24.05	52.47	0.083	-0.426
<i>C. livia</i>	11,388	27.96	32.98	13.68	25.38	53.34	0.048	-0.414
<i>D. giganteus</i>	11,397	28.28	32.27	13.85	25.60	53.88	0.050	-0.400
<i>E. minor</i>	11,400	29.16	33.83	13.01	24.00	53.16	0.097	-0.444
<i>E. orientalis</i>	11,397	28.20	33.25	13.73	24.80	53.00	0.064	-0.416
<i>F. peregrinus</i>	11,406	29.49	32.52	13.19	24.80	54.29	0.086	-0.423
<i>G. californianus</i>	11,403	30.23	32.29	12.56	24.92	55.15	0.096	-0.440
<i>G. gallus</i>	11,397	28.31	33.58	13.37	24.74	53.05	0.067	0.430
<i>G. stellata</i>	11,400	28.48	33.49	13.78	24.25	52.73	0.080	-0.417
<i>H. ater</i>	11,400	29.75	32.15	13.14	24.96	54.71	0.088	-0.420
<i>H. novaeseelandiae</i>	11,391	28.99	32.67	13.21	25.13	54.12	0.071	-0.424
<i>M. undulates</i>	11,400	29.89	33.60	12.26	24.25	54.14	0.104	-0.465
<i>M. novae Hollandiae</i>	11,394	27.92	32.25	14.40	25.43	53.35	0.047	-0.383
<i>N. novae Hollandiae</i>	11,406	29.03	33.72	13.49	23.73	52.76	0.100	-0.429
<i>R. jubatus</i>	11,379	26.09	36.34	13.60	23.97	50.06	0.042	-0.455
<i>P. azara</i>	11,400	29.18	32.92	13.49	24.41	53.59	0.089	-0.419
<i>P. rubber</i>	11,397	27.12	34.61	14.64	23.64	50.76	0.069	-0.405
<i>P. rubricauda</i>	11,403	26.01	33.78	14.57	25.64	51.65	0.007	-0.397
<i>R. americana</i>	11,397	28.53	31.70	13.64	26.14	54.66	0.044	-0.398
<i>S. camelus</i>	11,400	28.01	31.21	14.24	26.54	54.55	0.027	-0.373
<i>T. major</i>	11,382	28.40	30.50	12.91	28.20	56.60	0.004	-0.405
<i>T. melanophrys</i>	11,406	29.14	31.45	13.56	25.85	54.99	0.060	-0.397
<i>T. novae Hollandiae</i>	11,388	30.30	33.42	12.78	23.50	53.80	0.126	-0.447
<i>T. viridis</i>	11,406	29.07	29.38	13.46	28.09	57.16	0.017	-0.372

Continued on next page

Table 4. Continued.

Species	Size (bp)	A%	C%	G%	T%	A+T%	AT skew	GC skew
tRNA								
<i>A. apus</i>	1,548	29.78	21.58	21.32	27.33	57.11	0.043	-0.006
<i>A. colubris</i>	1,551	30.37	20.95	21.41	27.08	57.45	0.057	0.011
<i>A. cristatus</i>	1,539	29.82	20.86	21.25	28.07	57.89	0.030	0.009
<i>A. haastii</i>	1,551	33.27	24.31	16.57	25.85	59.12	0.126	-0.189
<i>B. canadensis</i>	1,540	30.65	21.56	20.78	27.01	57.66	0.063	-0.018
<i>C. casuarinus</i>	1,539	30.86	21.25	19.56	28.33	59.19	0.043	-0.041
<i>C. ciconia</i>	1,550	29.81	21.35	21.42	27.42	57.23	0.042	0.002
<i>C. livia</i>	1,547	28.70	21.78	22.24	27.28	55.98	0.025	0.010
<i>D. giganteus</i>	1,552	30.22	20.94	21.07	27.77	57.99	0.042	0.003
<i>E. minor</i>	1,548	29.20	21.77	21.96	27.07	56.27	0.038	0.004
<i>E. orientalis</i>	1,550	29.23	21.61	21.74	27.42	56.65	0.032	0.003
<i>F. peregrinus</i>	1,563	29.17	21.63	21.69	27.51	56.69	0.029	0.001
<i>G. californianus</i>	1,542	30.80	21.34	20.75	27.11	57.91	0.064	-0.014
<i>G. gallus</i>	1,543	30.14	21.84	20.67	27.35	57.49	0.049	-0.028
<i>G. stellata</i>	1,554	29.79	21.75	21.43	27.03	56.82	0.049	-0.007
<i>H. ater</i>	1,549	30.08	20.92	20.79	28.21	58.30	0.032	-0.003
<i>H. novaeseelandiae</i>	1,549	29.70	20.98	21.37	27.95	57.65	0.030	0.009
<i>M. undulates</i>	1,541	30.24	21.35	20.44	27.97	58.21	0.039	-0.022
<i>M. novae Hollandiae</i>	1,548	29.72	20.93	21.45	27.91	57.62	0.031	0.012
<i>N. novae Hollandiae</i>	1,557	29.42	22.41	21.71	26.46	55.88	0.053	-0.016
<i>P. azara</i>	1,546	29.50	21.54	21.73	27.23	56.73	0.040	0.004
<i>P. rubber</i>	1,551	29.72	21.79	21.28	27.21	56.93	0.044	-0.012
<i>P. rubricauda</i>	1,543	28.78	22.81	21.97	26.44	55.22	0.042	-0.019
<i>R. american a</i>	1,538	29.91	20.42	21.46	28.22	58.13	0.029	0.025
<i>R. jubatus</i>	1,555	29.65	21.29	21.74	27.33	56.98	0.040	0.010
<i>S. camelus</i>	1,538	30.82	20.22	20.81	28.15	58.97	0.045	0.014
<i>T. major</i>	1,537	30.45	20.04	20.82	28.69	59.14	0.030	0.019
<i>T. melanophrys</i>	1,757	30.17	20.38	21.29	28.17	58.34	0.034	0.022
<i>T. novae Hollandiae</i>	1,532	29.63	21.48	21.67	27.22	56.85	0.042	0.004
<i>T. viridis</i>	1,555	30.42	20.51	20.96	28.10	58.52	0.040	0.011
rRNA								
<i>A. apus</i>	2,572	33.90	27.10	18.47	20.53	54.43	0.246	-0.189
<i>A. colubris</i>	2,553	33.37	26.48	19.31	20.84	54.21	0.231	-0.157
<i>A. cristatus</i>	2,561	32.02	27.22	19.25	21.28	53.30	0.202	-0.172
<i>A. haastii</i>	2,638	34.72	25.21	17.78	22.29	57.01	0.218	-0.173
<i>B. canadensis</i>	2,591	33.08	26.24	20.46	20.22	53.30	0.241	-0.124
<i>C. casuarinus</i>	2,559	34.54	25.21	18.80	21.45	56.00	0.234	-0.146
<i>C. ciconia</i>	2,576	33.27	27.91	18.94	19.88	53.14	0.252	-0.191
<i>C. livia</i>	2,559	32.75	26.26	19.42	21.57	54.32	0.206	-0.150
<i>D. giganteus</i>	2,567	34.75	27.04	17.96	20.26	55.01	0.263	-0.202
<i>E. minor</i>	2,583	32.91	27.84	19.09	20.17	53.08	0.240	-0.186
<i>E. orientalis</i>	2,562	32.90	27.40	19.59	21.10	53.01	0.219	-0.166
<i>F. peregrinus</i>	2,578	33.67	27.97	18.89	18.47	53.14	0.292	-0.194
<i>G. californianus</i>	2,572	33.75	26.56	18.04	21.66	55.40	0.218	-0.191
<i>G. gallus</i>	2,598	33.06	28.37	18.17	20.40	53.46	0.237	-0.219
<i>G. stellata</i>	2,588	33.04	27.47	19.32	20.17	53.21	0.242	-0.174
<i>H. ater</i>	2,552	33.03	26.61	19.47	20.89	53.92	0.225	-0.155
<i>H. novaeseelandiae</i>	2,575	33.94	27.34	18.14	20.58	54.52	0.245	-0.202
<i>M. undulates</i>	2,549	33.58	30.29	18.12	18.01	51.59	0.302	-0.251
<i>M. novae Hollandiae</i>	2,575	32.35	25.48	20.31	21.86	54.21	0.194	-0.113
<i>N. novae Hollandiae</i>	2,557	32.93	29.25	19.63	18.19	51.11	0.288	-0.197
<i>P. azara</i>	2,556	32.08	29.93	18.97	19.01	51.10	0.256	-0.224
<i>P. rubber</i>	2,567	33.27	26.76	19.05	20.92	54.19	0.228	-0.168
<i>P. rubricauda</i>	2,581	33.05	29.99	18.09	18.87	51.92	0.273	-0.248
<i>R. american a</i>	2,546	32.88	26.75	19.17	21.21	54.08	0.216	-0.165
<i>R. jubatus</i>	2,553	31.84	25.85	19.98	22.33	54.17	0.176	-0.128
<i>S. camelus</i>	2,546	34.33	26.28	18.07	21.33	55.66	0.234	-0.185
<i>T. major</i>	2,575	34.06	26.06	18.02	21.86	55.92	0.218	-0.182
<i>T. melanophrys</i>	2,573	33.15	26.16	19.43	21.26	54.41	0.219	-0.148
<i>T. novae Hollandiae</i>	2,563	34.14	28.25	18.14	19.47	53.61	0.274	-0.218
<i>T. viridis</i>	2,600	33.00	25.50	19.00	22.50	55.50	0.189	-0.146

Continued on next page

Table 4. Continued.

Species	Size (bp)	A%	C%	G%	T%	A+T%	AT skew	GC skew
Control region								
<i>A. apus</i>	21,470	34.49	24.56	11.90	29.05	63.54	0.086	-0.347
<i>A. colubris</i>	787	29.35	26.18	15.25	29.10	58.45	0.004	-0.264
<i>A. cristatus</i>	3,054	31.57	25.87	11.72	30.71	62.28	0.014	-0.376
<i>A. haastii</i>	1,351	25.54	25.91	12.66	35.90	61.44	-0.169	-0.344
<i>B. canadensis</i>	1,194	29.65	31.49	13.48	25.38	55.03	0.078	-0.400
<i>C. casuarinus</i>	1,138	28.21	26.98	15.20	29.61	57.82	-0.024	-0.279
<i>C. ciconia</i>	1,779	31.08	22.60	17.93	28.39	59.47	0.045	-0.115
<i>C. livia</i>	1,657	30.84	28.49	13.28	27.40	58.24	0.059	-0.364
<i>D. giganteus</i>	1508	30.11	27.45	14.32	28.12	58.22	0.034	-0.314
<i>E. minor</i>	2,040	31.13	23.04	13.24	32.60	63.73	-0.023	-0.270
<i>E. orientalis</i>	2,190	30.78	25.89	11.60	31.74	62.51	-0.015	-0.381
<i>F. peregrinus</i>	2,460	39.88	22.48	14.02	23.25	63.13	0.263	-0.232
<i>G. californianus</i>	1,169	28.74	28.49	11.89	30.88	59.62	-0.036	-0.411
<i>G. gallus</i>	1,232	26.46	26.62	13.47	33.44	59.90	-0.117	-0.328
<i>G. stellata</i>	1,984	29.89	21.57	15.98	32.56	62.45	-0.043	-0.149
<i>H. ater</i>	1,240	32.02	25.97	14.68	27.34	59.35	0.079	-0.278
<i>H. novaeseelandiae</i>	1,690	30.41	29.88	11.78	27.87	58.28	0.044	-0.434
<i>M. undulates</i>	2,685	26.48	23.35	14.23	35.94	62.42	-0.152	-0.243
<i>M. novaehollandiae</i>	2,280	31.32	24.91	11.36	32.41	63.73	-0.017	-0.374
<i>N. novaehollandiae</i>	662	29.46	30.82	14.80	24.92	54.38	0.083	-0.351
<i>P. azara</i>	3,224	27.57	27.14	12.44	32.85	60.42	-0.087	-0.371
<i>P. rubber</i>	1,878	34.40	30.24	11.24	24.07	58.47	0.177	-0.458
<i>P. rubricauda</i>	2,211	29.76	25.33	11.85	33.06	62.82	-0.053	-0.363
<i>R. american a</i>	1,171	30.06	27.50	14.52	27.92	57.98	0.037	-0.309
<i>R. jubatus</i>	1,414	29.84	21.64	15.49	33.03	62.87	-0.051	-0.166
<i>S. camelus</i>	1,034	28.82	26.50	14.31	30.37	59.19	-0.026	-0.299
<i>T. major</i>	1,102	29.67	25.50	12.07	32.76	62.43	-0.049	-0.357
<i>T. melanophrys</i>	2,629	31.95	26.63	13.35	28.07	60.02	0.065	-0.332
<i>T. novaehollandiae</i>	2,437	28.60	25.44	14.53	31.43	60.03	-0.047	-0.273
<i>T. viridis</i>	2,128	29.56	25.89	11.94	32.61	62.17	-0.049	-0.369

^aThe control region of *A. colubris* and *N. novaehollandiae* are incomplete.

Protein-coding genes

The total length of the 13 protein-coding genes in *C. livia* mtDNA is 11,388 bp, accounting for 66.1% of the complete mitochondrial genome (Table 4). The length of the 13 protein-coding genes found in avian species varies from 11,379 (*Rhynochetos jubatus*) to 11,412 bp (*Branta canadensis*). The 13 protein-coding genes found in the *C. livia* mtDNA are similar in length to most other avian species. The longest protein-coding gene of *C. livia* mtDNA is the *nad5* gene (1815 bp), whereas the shortest is the *atp8* gene (168 bp; Table 5).

Analysis of the base composition at each codon position of the concatenated 13 protein-coding genes of *C. livia* mitochondrial genome shows that each codon position has a different AT/GC bias (Table 6). The first and third codon positions of *C. livia* are biased toward A and C, while the second codon positions are biased toward T and C. The AT composition at the first codon position is 49.1%. The values of the second and third codon positions are 58.5 and 52.5%, respectively (Table 6). As with most avian species (except for *A. haastii*, *T. major* and *T. viridis*), the second codon position has the highest AT composition (Table 6). The *nad6* gene of *C. livia* mitogenome has strong skews of T vs A (-0.587), and G vs C (0.602), while the *nad1* gene has a slight skew of T vs A (-0.016), and a strong skew of C vs G (0.452). All other 11 protein-coding genes of the *C. livia* mitochondrial genome have a slight skew of A vs T (0.005 to 0.151), and a strong skew of C vs G (GC skew = -0.328 to -0.757; Table 5).

Table 5. Base composition for protein-coding genes found in the mitochondrial genome of *Columba livia*.

Gene	Length (bp)	Proportion of nucleotides (%)					AT skew	GC skew
		A	C	G	T	A+T		
<i>nad1</i>	966	25.98	34.27	12.94	26.81	52.80	-0.016	-0.452
<i>nad2</i>	1041	31.70	34.20	10.18	23.92	55.62	0.140	-0.541
<i>cox1</i>	1551	26.11	31.98	16.18	25.73	51.84	0.007	-0.328
<i>cox2</i>	684	30.41	30.85	14.62	24.12	54.53	0.115	-0.357
<i>atp8</i>	168	32.14	38.69	5.36	23.81	55.95	0.149	-0.757
<i>atp6</i>	684	28.65	35.09	9.80	26.46	55.12	0.040	-0.563
<i>cox3</i>	786	26.97	33.21	15.01	24.81	51.78	0.042	-0.377
<i>nad3</i>	351	27.07	33.90	12.25	26.78	53.85	0.005	-0.469
<i>nad4L</i>	297	27.61	33.33	14.48	24.58	52.19	0.058	-0.394
<i>nad4</i>	1380	30.51	35.72	10.72	23.04	53.55	0.139	-0.538
<i>nad5</i>	1815	31.13	34.49	11.40	22.98	54.10	0.151	-0.503
<i>cob</i>	1143	26.95	35.87	12.60	24.58	51.53	0.046	-0.480
<i>nad6</i>	522	10.92	9.39	37.74	41.95	52.87	-0.587	0.602
Mean	876	27.40	32.38	14.10	26.12	53.52		

Table 6. Summary of base composition at each codon position of the concatenated 13 protein-coding genes in avian mitochondrial genomes.

Species	1st codon position					2nd codon position					3rd codon position				
	A%	T%	G%	C%	AT%	A%	T%	G%	C%	AT%	A%	T%	G%	C%	AT%
<i>A. apus</i>	28.5	21.2	21.9	28.4	49.7	18.3	39.7	12.8	29.2	58.0	40.2	14.8	4.3	40.7	55.0
<i>A. colubris</i>	28.0	21.9	22.1	28.0	49.9	18.5	39.8	12.6	29.0	58.3	39.2	14.4	6.8	39.6	53.6
<i>A. cristatus</i>	28.7	20.9	22.1	28.3	49.6	18.3	39.8	12.7	29.2	58.1	39.3	16.2	5.9	38.6	55.5
<i>A. haastii</i>	28.6	22.3	21.4	27.6	50.9	18.3	40.5	12.8	28.4	58.8	40.6	19.2	5.0	35.2	59.8
<i>B. canadensis</i>	27.3	20.3	23.8	28.7	47.6	18.2	40.4	12.8	28.6	58.6	38.4	11.0	8.8	41.8	49.4
<i>C. casuarinus</i>	27.9	21.4	22.6	28.1	49.3	18.2	40.3	12.8	28.6	58.5	38.6	19.3	7.0	35.2	57.9
<i>C. ciconia</i>	28.4	19.6	22.3	29.7	48.0	18.3	39.4	12.8	29.4	57.7	38.5	13.1	5.8	42.6	51.6
<i>C. livia</i>	28.5	20.6	22.2	28.7	49.1	18.4	40.1	12.7	28.9	58.5	37.0	15.5	6.2	41.4	52.5
<i>D. giganteus</i>	28.1	21.0	22.1	28.8	49.1	18.2	40.2	12.8	28.8	58.4	38.6	15.6	6.6	39.2	54.2
<i>E. minor</i>	29.5	20.4	21.4	28.7	49.9	18.4	39.2	12.8	29.6	57.6	39.6	12.4	4.8	43.2	52.0
<i>E. orientalis</i>	28.5	20.3	21.9	29.3	48.8	18.5	39.9	12.7	28.8	58.4	37.6	14.1	6.6	41.6	51.7
<i>F. peregrinus</i>	29.5	20.7	21.4	28.4	50.2	18.4	39.5	12.8	29.4	57.9	40.7	14.3	5.3	39.7	55.0
<i>G. californianus</i>	30.2	20.9	21.0	27.9	51.1	18.5	39.7	12.5	29.3	58.2	42.0	14.2	4.1	39.7	56.2
<i>G. gallus</i>	28.8	20.8	21.7	28.6	49.6	18.1	39.9	12.7	29.3	58.0	38.0	13.5	5.7	42.8	51.5
<i>G. stellata</i>	28.5	20.0	22.1	29.4	48.5	18.6	39.9	12.7	28.9	58.5	38.3	12.9	6.6	42.2	51.2
<i>H. ater</i>	29.4	21.2	21.4	27.9	50.6	18.7	39.9	12.8	28.7	58.6	41.1	13.8	5.3	39.8	54.9
<i>H. novaezeelandiae</i>	28.5	20.8	22.0	28.8	49.3	18.5	40.0	12.8	28.7	58.5	40.0	14.6	4.9	40.6	54.6
<i>M. undulatus</i>	29.4	20.8	20.8	28.9	50.2	18.6	39.4	12.4	29.6	58.0	41.6	12.5	3.6	42.3	54.1
<i>M. novaeahollandiae</i>	27.5	20.5	23.5	28.4	48.0	18.2	40.2	12.9	28.6	58.4	38.0	15.5	6.8	39.7	53.5
<i>N. novaeahollandiae</i>	28.9	19.8	21.5	29.8	48.7	18.3	39.0	12.7	30.1	57.3	40.0	12.4	6.3	41.4	52.4
<i>P. azara</i>	28.3	20.8	21.7	29.2	49.1	18.2	39.8	12.7	29.3	58.0	31.7	11.3	6.4	50.6	43.0
<i>P. rubber</i>	28.5	20.6	22.2	28.7	49.1	18.4	39.9	12.8	28.9	58.3	40.6	12.7	5.5	41.2	53.3
<i>P. rubricauda</i>	27.7	19.8	22.2	30.2	47.5	18.1	39.5	13.0	29.4	57.6	35.6	11.5	8.6	44.2	47.1
<i>R. americana</i>	27.0	21.0	22.5	29.4	48.0	18.0	40.0	12.9	29.0	58.0	33.0	15.8	8.2	42.9	48.8
<i>R. jubatus</i>	28.7	20.8	22.4	28.1	49.5	18.5	39.8	12.6	29.1	58.3	38.4	17.8	5.9	37.9	56.2
<i>S. camelus</i>	27.8	21.3	22.6	28.2	49.1	18.3	40.3	12.7	28.7	58.6	37.9	17.9	7.4	36.7	55.8
<i>T. major</i>	29.5	22.7	20.7	27.0	52.2	18.3	40.1	12.8	28.8	58.4	37.3	21.8	5.2	35.7	59.1
<i>T. melanophrys</i>	28.7	21.2	22.0	28.0	49.9	18.4	40.0	12.9	28.7	58.4	40.3	16.3	5.8	37.6	56.6
<i>T. novaeahollandiae</i>	29.4	20.4	21.2	28.9	49.8	18.6	39.5	12.6	29.3	58.1	42.9	10.5	4.6	42.1	53.4
<i>T. viridis</i>	28.7	22.8	21.9	26.6	51.5	18.7	40.1	12.5	28.7	58.8	39.8	21.4	5.9	32.8	61.2

There are four reading frame overlaps within the mitochondrial genome of *C. livia* (*cox1* and *tRNA^{Ser(UCN)}* share nine nucleotides; *atp8* and *atp6* share 10 nucleotides; *nad4L* and *nad4* share seven nucleotides; *atp6* and *cox3* share one nucleotide). Other overlaps are shown in Table 3.

Twelve genes (with the exception of *nad6*) of the 13 typical protein-coding genes (*nad1-6* and *4L*, *cox1-3*, *atp6* and *atp8*, *cob*) are encoded on the heavy strand. All protein-coding genes initiate with ATG, except for *cox1* and *nad5*, which begins with GTG. Five types of stop codons are used by the coding genes, including TAA for *nad3*, *cox2*, *atp8*, *atp6*, *nad4L*, and *cob*; AGA for *nad1* and *nad5*; TAG for *nad6*; AGG for *cox1*, and incomplete stop codon T- or TA- for *cox3*, *nad2* and *nad4*, respectively. The use of an incomplete stop codon T, a common mechanism for stopping protein translations, was also observed in other avian species.

The pattern of codon usage in the *C. livia* mtDNA was also studied (Table 7). There are 3785 codons for all the 13 protein-coding genes, after stop codons are excluded. The most frequently used amino acid was Leu (17.89%), followed by Thr (9.30%), Ala (7.67%), Ile (7.59%), and Ser (7.43%).

Table 7. Codon usage of 13 protein-coding genes in the mitochondrial genome of *Columba livia*.

Amino acid	Codon	Number	Frequency (%)	Amino acid	Codon	Number	Frequency (%)
Phe	TTT	47	1.24	Tyr	TAT	28	0.74
	TTC	166	4.37		TAC	86	2.27
Leu	TTA	65	1.71	Stop	TAA	9	0.24
	TTG	21	0.55		TAG	1	0.03
	CTT	76	2.00	His	CAT	22	0.58
	CTC	187	4.93		CAC	86	2.27
	CTA	299	7.88	Gln	CAA	79	2.08
	CTG	30	0.79		CAG	13	0.34
	ATT	86	2.27	Asn	AAT	24	0.63
Ile	ATC	202	5.32		AAC	105	2.77
	ATA	132	3.48	Lys	AAA	74	1.95
Met	ATG	40	1.05		AAG	9	0.24
	GTT	40	1.05	Asp	GAT	21	0.55
Val	GTC	53	1.40		GAC	43	1.13
	GTA	56	1.48	Glu	GAA	79	2.08
	GTG	22	0.58		GAG	18	0.47
	TCT	34	0.90	Cys	TGT	7	0.18
	TCC	107	2.82		TGC	19	0.50
Ser	TCA	78	2.05	Trp	TGA	93	2.45
	TCG	8	0.21		TGG	12	0.32
	CCT	45	1.19	Arg	CGT	9	0.24
	CCC	82	2.16		CGC	16	0.42
Pro	CCA	93	2.45		CGA	42	1.11
	CCG	5	0.13		CGG	6	0.16
	ACT	58	1.53	Ser	AGT	13	0.34
Thr	ACC	157	4.14		AGC	42	1.11
	ACA	130	3.42	Stop	AGA	2	0.05
	ACG	8	0.21		AGG	1	0.03
Ala	GCT	50	1.32	Gly	GGT	27	0.71
	GCC	139	3.66		GGC	80	2.11
	GCA	93	2.45		GGA	80	2.11
	GCG	9	0.24		GGG	32	0.84

Non-coding regions

The non-coding regions in the *C. livia* mitochondrial genome include a control region (D-loop) and a few spaces (Table 3). The control region (1657 bp) is located between *tRNA^{Glu}* and *tRNA^{Phe}* genes. The length of the control region of avian species varies between 1034 (*Struthio camelus*) and 3224 bp (*Pteroglossus azara*), ranging in AT content from 55.03 (*B. canadensis*) to 63.73% (*E. minor* and *M. novaehollandiae*; Table 4).

Based on the distribution of the variable nucleotide positions and differential frequencies of the nucleotides, the mitochondrial control region is divided into three domains (Brown et al., 1986; Saccone et al., 1991; Randi and Lucchini, 1998). The nucleotide composition of the *C. livia* control region was A = 30.84, T = 27.40, C = 28.49, and G = 13.28% (Table 4), with a bias against G, which is usual for the mtDNA sense strand of vertebrates (Wolstenholme, 1992). Domain I (ETAS, extended termination-associated sequences) contains part A (nt 1-146 in Figure 2) and part B (nt 147-398 in Figure 2). In part A, ETAS1 and ETAS2 are found at positions 71-126 and 108-146 nt, respectively, and overlapped one another by 19 bp, with 63.3 and 58.7% similarity to the consensus mammalian ETAS1 and ETAS2, respectively (Sbisa et al., 1997). In part B (nt 147-398), a CSB1-like block (5'-TAGAACCTTAATGATACC TTAGACATA-3') has 63.0% similarity to the conserved sequence block (CSB1) in domain III (Figure 2). Four conserved sequence boxes in the central domain II (nt 399-897) were identified as boxes C, D, E, and F (Figure 2). In domain III (nt 898-1657), there is a poly (T) sequence (nt 1123-1131), which is located just a few nucleotides downstream from the putative CSB1 (nt 983-1007; Figure 2). Although the conserved blocks, such as ETAS1, ETAS2, CSB1, CSB1-like, and boxes C, D, E, and F, are readily identifiable in the *C. livia* control region, the typical origin of H-strand replication (O_H), CSB2 and CSB3 could not be detected (Walberg and Clayton, 1981). The bidirectional light- and heavy-strand transcription promoters (LSP/HSP) (L'Abbé et al., 1991) are found in *C. livia* (Figure 2). Furthermore, there are 27 tandem repeats (CAAA) at the end of the *C. livia* control region.




Figure 2. Schematic representation of the organization of the *Columba livia* control region. ETAS = extended termination-associated sequences; F through C boxes = conserved sequence boxes in the central domain; CSB = conserved sequence block; CSB-like = a sequence similar to the CSB; LSP = light-strand transcription promoter; HSP = heavy-strand transcription promoter; R = tandem repeats.

Ribosomal and transfer RNA genes

The Rock pigeon mitochondrial genome has two rRNA subunits (*srRNA* and *lrRNA*), as in other vertebrates. The *srRNA* gene is located between *tRNA^{Phe}* and *tRNA^{Val}* genes, and the *lrRNA* gene is located between *tRNA^{Val}* and *tRNA^{Leu(UUR)}* genes. The lengths of *srRNA* and *lrRNA* are 973 and 1586 bp. The A+T content of the *rRNA* is 54.32%, which is within the range observed for other avian species (Table 4).

Twenty-two tRNA genes were observed in the Rock pigeon mitochondrial genome. The tRNA genes are interspersed in the genome, ranging in size from 66 (*tRNA^{Ser(AGY)}*) to

74 (*tRNA^{Trp}*) nucleotides (Table 3). *tRNA^{Cys}* and *tRNA^{Ser(AGY)}*, which were not found by the tRNAscan-SE, were identified by comparison with *G. gallus* counterparts. All tRNA gene sequences have the potential to fold into typical cloverleaf secondary structures (Figure 3). The DHU and TΨC arms contain two and five nucleotide pairs, respectively.




Figure 3. Inferred secondary structures of 22 tRNAs found in the *Columba livia* mitochondrial genome.

This is the first complete nucleotide sequence described for the mitochondrial genome of the Rock pigeon. We also detailed the genome organization and codon usage of *C. livia* mitochondrial DNA. These results provide basic information for phylogenetic analyses among the birds, and especially Columbiformes species.

ACKNOWLEDGMENTS

Research supported by the National Natural Science Foundation of China (#30870172), the Key Program of Natural Science Foundation of the Anhui Higher Education Institutions (#KJ2008A13ZC), the Natural Science Foundation of Anhui Normal University (#2008zxz14), the Key Lab of Biotic Environment and Ecology Safety in Anhui Province, and the Program for Innovative Research Team in Anhui Normal University.

REFERENCES

- Boore JL (1999). Animal mitochondrial genomes. *Nucleic Acids Res.* 27: 1767-1780.
- Brown GG, Gadaleta G, Pepe G, Saccone C, et al. (1986). Structural conservation and variation in the D-loop-containing region of vertebrate mitochondrial DNA. *J. Mol. Biol.* 192: 503-511.
- Cooper A, Lalueza-Fox C, Anderson S, Rambaut A, et al. (2001). Complete mitochondrial genome sequences of two extinct moas clarify ratite evolution. *Nature* 409: 704-707.
- Gibb GC, Kardailsky O, Kimball RT, Braun EL, et al. (2007). Mitochondrial genomes and avian phylogeny: complex characters and resolvability without explosive radiations. *Mol. Biol. Evol.* 24: 269-280.
- Haddrath O and Baker AJ (2001). Complete mitochondrial DNA genome sequences of extinct birds: ratite phylogenetics and the vicariance biogeography hypothesis. *Proc. Biol. Sci.* 268: 939-945.
- Hall AT (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41: 95-98.
- Harlid A, Janke A and Arnason U (1998). The complete mitochondrial genome of *Rhea americana* and early avian divergences. *J. Mol. Evol.* 46: 669-679.
- Harrison GL, McLennan PA, Phillips MJ, Slack KE, et al. (2004). Four new avian mitochondrial genomes help get to basic evolutionary questions in the late cretaceous. *Mol. Biol. Evol.* 21: 974-983.
- Hazkani-Covo E, Zeller RM and Martin W (2010). Molecular poltergeists: mitochondrial DNA copies (numts) in sequenced nuclear genomes. *PLoS Genet.* 6: e1000834.
- Howard R and Moore A (2003). The Howard and Moore Complete Checklist of the Birds of the World. 3rd edn. Christopher Helm, London.
- L'Abbé D, Duhamel JF, Lang BF and Morais R (1991). The transcription of DNA in chicken mitochondria initiates from one major bidirectional promoter. *J. Biol. Chem.* 266: 10844-10850.
- Larkin MA, Blackshields G, Brown NP, Chenna R, et al. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics* 23: 2947-2948.
- Livezey BC and Zusi RL (2007). Higher-order phylogeny of modern birds (Theropoda, Aves: Neornithes) based on comparative anatomy. II. Analysis and discussion. *Zool. J. Linn. Soc.* 149: 1-95.
- Lohse M, Drechsel O and Bock R (2007). OrganellarGenomeDRAW (OGDRAW): a tool for the easy generation of high-quality custom graphical maps of plastid and mitochondrial genomes. *Curr. Genet.* 52: 267-274.
- Lowe TM and Eddy SR (1997). tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25: 955-964.
- Mindell DP, Sorenson MD and Dimcheff DE (1998). Multiple independent origins of mitochondrial gene order in birds. *Proc. Natl. Acad. Sci. U. S. A.* 95: 10693-10697.
- Moore WS (1995). Inferring phylogenies from mtDNA variation: mitochondrial-gene trees versus nuclear-gene trees. *Evolution* 49: 718-726.
- Morgan-Richards M, Trewick SA, Bartosch-Harlid A, Kardailsky O, et al. (2008). Bird evolution: testing the Metaves clade with six new mitochondrial genomes. *BMC Evol. Biol.* 8: 20.
- Nishibori M, Hayashi T, Tsudzuki M, Yamamoto Y, et al. (2001). Complete sequence of the Japanese quail (*Coturnix japonica*) mitochondrial genome and its genetic relationship with related species. *Anim. Genet.* 32: 380-385.

- Nishibori M, Shimogiri T, Hayashi T and Yasue H (2005). Molecular evidence for hybridization of species in the genus *Gallus* except for *Gallus varius*. *Anim Genet.* 36: 367-375.
- Paton T, Haddrath O and Baker AJ (2002). Complete mitochondrial DNA genome sequences show that modern birds are not descended from transitional shorebirds. *Proc. Biol. Sci.* 269: 839-846.
- Pereira SL, Johnson KP, Clayton DH and Baker AJ (2007). Mitochondrial and nuclear DNA sequences support a Cretaceous origin of Columbiformes and a dispersal-driven radiation in the Paleocene. *Syst. Biol.* 56: 656-672.
- Perna NT and Kocher TD (1995). Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. *J. Mol. Evol.* 41: 353-358.
- Pratt RC, Gibb GC, Morgan-Richards M, Phillips MJ, et al. (2009). Toward resolving deep neoaves phylogeny: data, signal enhancement, and priors. *Mol. Biol. Evol.* 26: 313-326.
- Randi E and Lucchini V (1998). Organization and evolution of the mitochondrial DNA control region in the avian genus *Alectoris*. *J. Mol. Evol.* 47: 449-462.
- Rokas A, Williams BL, King N and Carroll SB (2003). Genome-scale approaches to resolving incongruence in molecular phylogenies. *Nature* 425: 798-804.
- Saccone C, Pesole G and Sbisa E (1991). The main regulatory region of mammalian mitochondrial DNA: structure-function model and evolutionary pattern. *J. Mol. Evol.* 33: 83-91.
- Sambrook J and Russell DW (2001). Molecular Cloning: A Laboratory Manual. 3rd edn. Cold Spring Harbor Laboratory Press, New York.
- San Mauro D, Garcia-Paris M and Zardoya R (2004). Phylogenetic relationships of discoglossid frogs (Amphibia: Anura: Discoglossidae) based on complete mitochondrial genomes and nuclear genes. *Gene* 343: 357-366.
- Sbisa E, Tanzariello F, Reyes A, Pesole G, et al. (1997). Mammalian mitochondrial D-loop region structural analysis: identification of new conserved sequences and their functional and evolutionary implications. *Gene* 205: 125-140.
- Shadel GS and Clayton DA (1997). Mitochondrial DNA maintenance in vertebrates. *Annu. Rev. Biochem.* 66: 409-435.
- Shen X, Tian M, Liu Z, Cheng H, et al. (2009). Complete mitochondrial genome of the sea cucumber *Apostichopus japonicus* (Echinodermata: Holothuroidea): the first representative from the subclass Aspidochirota with the echinoderm ground pattern. *Gene* 439: 79-86.
- Slack KE, Janke A, Penny D and Arnason U (2003). Two new avian mitochondrial genomes (penguin and goose) and a summary of bird and reptile mitogenomic features. *Gene* 302: 43-52.
- Slack KE, Jones CM, Ando T, Harrison GL, et al. (2006). Early penguin fossils, plus mitochondrial genomes, calibrate avian evolution. *Mol. Biol. Evol.* 23: 1144-1155.
- Slack KE, Delsuc F, McLennan PA, Arnason U, et al. (2007). Resolving the root of the avian mitogenomic tree by breaking up long branches. *Mol. Phylogenet. Evol.* 42: 1-13.
- Walberg MW and Clayton DA (1981). Sequence and properties of the human KB cell and mouse L cell D-loop regions of mitochondrial DNA. *Nucleic Acids Res.* 9: 5411-5421.
- Wang C, Chen Q, Lu G, Xu J, et al. (2008). Complete mitochondrial genome of the grass carp (*Ctenopharyngodon idella*, Teleostei): insight into its phylogenetic position within Cyprinidae. *Gene* 424: 96-101.
- Wolstenholme DR (1992). Animal mitochondrial DNA: structure and evolution. *Int. Rev. Cytol.* 141: 173-216.
- Wyman SK, Jansen RK and Boore JL (2004). Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* 20: 3252-3255.
- Xia X and Xie Z (2001). DAMBE: software package for data analysis in molecular biology and evolution. *J. Hered.* 92: 371-373.