

Chasing the heaviest black holes of jetted active galactic nuclei

G. Ghisellini,^{1*} R. Della Ceca,² M. Volonteri,³ G. Ghirlanda,¹ F. Tavecchio,¹
L. Foschini,¹ G. Tagliaferri,¹ F. Haardt,^{4,5} G. Pareschi¹ and J. Grindlay⁶

¹INAF, Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate, Italy

²INAF, Osservatorio Astronomico di Brera, Via Brera 28, I-20100 Milano, Italy

³Astronomy Department, University of Michigan, Ann Arbor, MI 48109, USA

⁴Università dell'Insubria, Dipartimento di Fisica e Matematica, Via Valleggio 11, I-22100 Como, Italy

⁵INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy

⁶Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

Accepted 2010 January 30. Received 2010 January 29; in original form 2009 November 25

ABSTRACT

We investigate the physical properties of 10 blazars at redshift greater than 2 detected in the 3-yr all-sky survey performed by the Burst Alert Telescope (BAT) on board the *Swift* satellite. We find that the jets of these blazars are among the most powerful known. Furthermore, the mass of their central black hole, inferred from the optical–ultraviolet bump, exceeds a few billions of solar masses, with accretion luminosities being a large fraction of the Eddington one. We compare their properties with those of the brightest blazars of the 3-month survey performed by the Large Area Telescope (LAT) on board the *Fermi* satellite. We find that the BAT blazars have more powerful jets, more luminous accretion discs and larger black hole masses than LAT blazars. These findings can be simply understood on the basis of the blazar sequence, which suggests that the most powerful blazars have a spectral energy distribution with a high-energy peak at MeV (or even sub-MeV) energies. This implies that the most extreme blazars can be found more efficiently in hard X-rays, rather than in the high-energy γ -ray band. We then discuss the implications of our findings for future missions, such as the *New Hard X-ray Mission (NHXM)* and especially the *Energetic X-ray Imaging Survey Telescope (EXIST)* mission which, during its planned 2-yr all-sky survey, is expected to detect thousands of blazars, with a few of them at $z \gtrsim 6$.

Key words: radiation mechanisms: non-thermal – BL Lacertae objects: general – quasars: general – gamma-rays: theory – X-rays: general.

1 INTRODUCTION

Ajello et al. (2009, hereafter A09) recently published the list of blazars detected in the all-sky survey by the Burst Alert Telescope (BAT) on board the *Swift* satellite, between 2005 March and 2008 March. BAT is a coded mask designed to detect gamma-ray bursts (GRBs), has a large field of view ($120^\circ \times 90^\circ$, partially coded) and is sensitive in the 15–150 keV energy range. This instrument was specifically designed to detect GRBs, but since GRBs are distributed isotropically in the sky, BAT, as a by-product, performed an all-sky survey with a reasonably uniform sky coverage, at a limiting sensitivity of the order of 1 mCrab in the 15–55 keV range (equivalent to 1.27×10^{-11} erg cm⁻² s⁻¹) in 1 Ms exposure (A09). Taking the period 2005 March to 2008 March, and evaluating the image resulting from the superposition of all obser-

vations in this period, BAT detected 38 blazars (A09), of which 26 are flat spectrum radio quasars (FSRQs) and 12 are BL Lac objects, once the Galactic plane ($|b| < 15^\circ$) is excluded from the analysis. A09 reported an average exposure of 4.3 Ms, and considered the 15–55 keV energy range, to avoid background problems at higher energies. The well-defined sky coverage and source-selection criteria make the list of the found blazars a complete, flux-limited sample that enabled A09 to calculate the luminosity function and the possible cosmic evolutions of FSRQs and BL Lacs, together with their contribution to the hard X-ray background. A09 also stressed the fact that the detected BAT blazars at high redshift are among the most powerful blazars and could be associated with powerful accreting systems.

Within the BAT sample, there are 10 blazars (all FSRQs) at redshift greater than 2, and five at redshift between 3 and 4. For comparison, the Large Area Telescope (LAT) on board the *Fermi* satellite detected about 100 blazars (at high significance in the first 3 months), with a maximum redshift of 2.944 (Abdo et al. 2009).

*E-mail: gabriele.ghisellini@brera.inaf.it




Figure 1. Comparison of the redshift distributions of the bright *Fermi*/LAT and *Swift*/BAT blazars, divided in BL Lacs and FSRQs.

Only five blazars have $z > 2$, with only one with $z > 2.5$. Fig. 1 shows the distribution of redshifts for the *Fermi*/LAT and the *Swift*/BAT blazars. Comparing the two redshift distributions and including BL Lac objects, a Kolmogorov–Smirnov (KS) test gives a probability $P = 0.079$ that they are drawn from the same distribution; it decreases to $P = 0.034$ when excluding BL Lacs. The BAT FSRQs have a tail at high redshifts, not present in the case of LAT FSRQs. This presence of this tail in the redshift distribution of BAT blazars is the first indication that a survey in hard X-rays can be more fruitful in pinpointing the most powerful blazars lying at the highest redshifts. Theoretically, this can be understood on the basis of the blazar sequence (Fossati et al. 1998; Ghisellini et al. 1998; Donato et al. 2001; Ghisellini, Maraschi & Tavecchio 2009b). According to this sequence the most powerful blazars, which are all FSRQs, have the high-energy peak of their spectral energy distribution (SED) in the MeV range. As a consequence, these objects are more luminous in hard X-rays than what they are above 100 MeV, and thus become detectable in the BAT survey even if they are undetected in the LAT one. Indeed, *none* of the 10 BAT blazars at $z > 2$ is present in the 3-month LAT survey of bright blazars, while four are present in the *Fermi*/LAT 11-month survey catalogue of sources with a $>4\sigma$ level detection¹ (PKS 0537–286, PKS 0746+254, PKS 0805+6144 and PKS 0836+710), and one additional source (PKS 2149–306) has a *Fermi*/LAT detection reported by Böck et al. (2010). The *Fermi*/LAT fluxes of these sources are just above the limiting sensitivity for a 5σ detection in 1-yr survey.

The aim of this paper is to study the most powerful and distant blazars present in the BAT survey in order to estimate the power carried by their jets in the form of bulk motion of matter and fields. Furthermore, we will estimate the mass of their central black hole and their accretion disc luminosities. To this aim we will take advantage of the data of the other *Swift* instruments (the X-ray Telescope, XRT, and the Ultraviolet (UV) and Optical Telescope, UVOT). We will then compare the overall properties of these powerful BAT blazars with the same properties of the bright γ -ray blazars detected by the LAT and recently studied in Ghisellini et al. (2010, hereafter G10).

¹See http://fermi.gsfc.nasa.gov/ssc/data/access/lat/1yr_catalog/

Finally, we will discuss the implications of our findings for the future missions, such as *New Hard X-ray Mission (NHXM)* and *Energetic X-ray Imaging Survey Telescope (EXIST)*, highlighting the possible discovery space that these missions can have concerning the search and the study of the largest black hole masses of jetted sources at large redshift.

We use a flat cosmology with $h_0 = \Omega_\Lambda = 0.7$. We adopt the convention $Q = 10^x Q_x$ and use CGS units unless specified otherwise.

2 THE SAMPLE

In A09 all BAT blazars were fitted with a simple power law in the 15–55 keV energy range. The resulting energy spectral indices α_X are shown in Fig. 2 as a function of the rest-frame 15–55 keV luminosity, calculated according to

$$L_X = 4\pi d_L^2 \frac{F_X}{(1+z)^{1-\alpha_X}}, \quad (1)$$

where F_X is the observed X-ray flux in the 15–55 keV energy range as listed in A09.

Blazars with $z > 2$ are marked with black diamonds: they are the most luminous, with $L_X > 2 \times 10^{47}$ erg s^{−1} in the (rest-frame) 15–55 keV band. Our cut in redshift therefore corresponds also to a cut in luminosity, as expected for a flux-limited sample.

Table 1 reports the redshift, the BAT X-ray flux in the 15–55 keV range, the energy spectral index in this energy range and the corresponding K -corrected X-ray luminosity. It should be noted that these 10 blazars are *not* the only $z > 2$ blazars detected by BAT. Indeed, there are four additional blazars in the $|b| < 15^\circ$ region (0212+735, with $z = 2.367$; OA 198 at $z = 2.365$; Swift J1656.3–3302 at $z = 2.4$ and PKS 1830–211 at $z = 2.507$), which are present in the catalogues of Cusumano et al. (2010) and Tueller et al. (2010).




Figure 2. The X-ray energy spectral index α_X as a function of the luminosity in the BAT energy range 15–55 keV. In this plane BL Lacs (triangles) and FSRQs (circles and diamonds) are clearly separated. We also mark (black diamonds) the FSRQs with $z > 2$. The labelled sources are the BL Lac IES 0120+340 ($z = 0.272$) and the three sources classified as FSRQs by A09 to the left-hand side of the divide. Note that of these three objects, B0007+107 (i.e. Mkn 1501) is a Seyfert 1 galaxy at $z = 0.09$ with a superluminal jet (Brunthaler et al. 2000) and S5 2116+81 ($z = 0.084$) is a radio galaxy (Stickel, Kühr & Fried 1993). The grey stripes indicate the X-ray luminosity dividing BL Lacs from FSRQs.

Table 1. The 10 BAT blazars in A09 with $z > 2$. The flux F_X is in units of 10^{-11} erg cm $^{-2}$ s $^{-1}$ and is calculated in the 15–55 keV energy range. The luminosity L_X is calculated according to equation (1) and is in units of erg s $^{-1}$.

Name	Alias	z	F_X	α_X	log L_X
0014+813	S5	3.366	1.88 ± 0.21	0.93 ± 0.55	48.03
0222+185	RBS 315	2.69	1.42 ± 0.22	0.48 ± 0.45	47.63
0537–286	PKS	3.10	1.27 ± 0.20	0.56 ± 0.30	47.75
074625+2549	SDSS	2.98	1.49 ± 0.25	0.08 ± 0.38	47.50
0805+6144	GB6	3.033	0.96 ± 0.19	0.58 ± 0.62	47.62
0836+710	4C71.07	2.17	2.85 ± 0.18	0.47 ± 0.14	47.73
1210+330	B2	2.50	0.90 ± 0.17	0.40 ± 0.30	47.32
2126–158	PKS	3.268	1.55 ± 0.27	0.72 ± 0.68	47.99
2149–306	PKS	2.35	3.72 ± 0.26	0.52 ± 0.21	47.95
225155+2217	MG3	3.668	1.00 ± 0.19	0.51 ± 0.33	47.77

We chose the A09 catalogue mainly because of their derivation of the blazar luminosity function performed with the data in that paper.

Several of the sources listed in Table 1 have already been discussed and modelled in the literature: S5 0014+813 was discussed by Ghisellini et al. (2009a) because of its apparently huge black hole mass (40 billion solar masses, but see Ghisellini et al. 2009a for the possibility that this is overestimated because of the assumption of an isotropically emitting accretion disc, and see below for further discussion); RBS 315 was discussed in Tavecchio et al. (2007, presenting also *Suzaku* data), PKS 0537–286 is discussed in Sambruna et al. (2007) and Bottacini et al. (2010, presenting also *INTEGRAL* data); J074625+2549 is discussed in Sambruna et al. (2006) and Watanabe et al. (2009, presenting also *Suzaku* data); 4C 71.07 (i.e. 0836+710) is discussed in Foschini et al. (2006) and Sambruna et al. (2007); PKS 2149–306 is discussed in Sambruna et al. (2007) and Bianchin et al. (2009, presenting also *XMM-Newton* and *INTEGRAL* data); J225155+2217 is discussed in Bassani et al. (2007, presenting also *INTEGRAL* data) and Maraschi et al. (2008).

To the best of our knowledge, the overall SED of the remaining three blazars, GB6 J0805+6144, B2 1210+330 and PKS 2126–158 have not yet been discussed and modelled.

2.1 The BAT blazars’ divide

Fig. 2 shows that BL Lacs and FSRQs are clearly separated in the α_X – L_X plane: BL Lacs have steeper slopes and lower luminosities than FSRQs. This can be readily explained recalling that, in BL Lacs, the X-ray flux often belongs to the tail of the synchrotron hump, while in more powerful FSRQs the X-ray flux always belongs to the very hard portion of the high-energy hump, believed to be produced by inverse Compton process. The separation occurs at $\alpha_X \sim 1$ and $L_X \sim 10^{45}$ erg s $^{-1}$. This ‘divide’ is related to a similar divide present in the α_γ – L_γ plane (Ghisellini et al. 2009b), using the 3-month data of the *Fermi*/LAT survey. However, there is one important difference: in the α_γ – L_γ plane BL Lacs are the *hard* sources, while FSRQs are the softer ones. In the γ -ray energy range the flux originates from the same (inverse Compton) component, and low-luminosity BL Lacs peak at energies close to the end of the LAT energy range, or even at higher energies, and therefore have γ -ray slopes flatter than FSRQs, which instead peaks at energies smaller than 100 MeV. Fig. 2 is therefore another manifestation of the blazar sequence because it agrees with the ideas that low-power blazars have both the synchrotron and the Compton peaks at larger frequencies than FSRQs. In this sense we have ‘blue’ BL Lacs and ‘red’ FSRQs. We propose here that the existence of a

dividing X-ray luminosity is the same as discussed in Ghisellini et al. (2009b), namely a change in the accretion regime of the underlying accretion disc. This interpretation assumes that there is a connection between the non-thermal beamed luminosity and the thermal luminosity produced by accretion, and that the range of black hole masses of the blazars illustrated in Fig. 2 is relatively small. Within this context, the most luminous objects are the ones accreting close to the Eddington rate. The dividing X-ray luminosity is produced by objects whose accretion disc is accreting at a rate a factor of ~ 300 lower. If these sources have the same mass as the most powerful blazars, then they accrete at the 0.3 per cent of the Eddington rate, and at these rates the accretion could change regime, becoming radiatively inefficient (see the discussion in Ghisellini et al. 2009b). In turn, the rather abrupt decrease of ionizing photons occurring at and below the critical accretion rate implies an equally abrupt decrease of the broad-line luminosity, making the blazar appear as a lineless BL Lac.

3 Swift DATA AND ANALYSIS

We collected the public *Swift* data of the 10 sources listed in Table 1. For all but one source, B2 1210+330, we could find at least one observation in the *Swift* archive. We obtained a *Swift* Target of Opportunity for this source: the observations were performed on 2009 October 26 and 27.

3.1 XRT data

The XRT data were processed with the standard procedures (XRTPIPELINE v.0.12.2). We considered photon counting mode data with the standard 0–12 grade selection. Source events were extracted in a circular region of aperture ~ 47 arcsec, the background was estimated in an equal-size circular region far from the source. Response matrices were created through the *xrtmkarf* task. The channels with energies below 0.3 keV and above 10 keV were excluded from the fit and the spectra were rebinned in energy so as to have at least 30 counts per bin. For spectra with very few counts, the Cash statistics was applied. When a sufficiently long exposure is available, we present the data corresponding to those observations. In a few cases we summed the data of different observations, to have a more accurate fit, after having checked that the fits of the single observations gave consistent results.

Each spectrum was analysed through *XSPEC* with an absorbed power law using the Galactic column density ($N_{\text{H}}^{\text{Gal}}$ from Kalberla et al. 2005). The computed errors represent the 90 per cent

Table 2. Results of the X-ray analysis.

Source	Observation date (dd/mm/yyyy)	t_{exp} (s)	$N_{\text{H}}^{\text{Gal}}$ (10^{20} cm^{-2})	Γ_1	Γ_2	E_{break} (keV)	$F_{0.3-10}^{\text{unabs}}$ (10^{-12} CGS)	χ^2 or C_{stat}	d.o.f.
0014+81	Sum ^a	8311	13.6	1.32 ± 0.1			5.4	22.6	19
0222+185	28/07/2006	4150	9.3	1.20 ± 0.08			14 ± 8	344	378
0537–286	08/12/2005	14 737	2.2	1.20 ± 0.08			4 ± 0.1	333	407
074625+2549	05/11/2005	24 999	4.6	1.22 ± 0.06			4.6 ± 0.2	467	543
0805+614	Sum ^b	8121	4.7	1.25 ± 0.13			3.14	10.8	10
0836+710	13/04/2007	7367	2.9	1.42 ± 0.05			17.5 ± 0.5	526	532
1210+330	26/10/2009	3782	1.16	1.9 ± 0.37			0.57 ± 0.15	31	41
2126–158	Sum ^c	34 782	5.0	0.6 ± 0.3	1.5 ± 0.05	0.95 ± 0.2	10.3	204	161
2149–306	10/12/2005	3336	1.6	1.45 ± 0.07			16.2 ± 8	367	394
225155+2217	22/05/2007	12 396	5.0	1.28 ± 0.08			4.66	19	25

^aSum of the observations 11/01/2007, 12/01/2007 and 14/01/2007.

^bSum of the observations 23/01/2009, 17/10/2006 and 21/01/2009.

^cSum of the observations 07/04/2007, 21/11/2008 and 06/04/2007.

Table 3. Summary of *Swift*/UVOT observed magnitudes. Lower limits are at 3σ level. Values of A_V from Schlegel, Finkbeiner & Davis (1998).

Source	A_V	v	b	u	$uvw1$	$uvm2$	$uvw2$
0014+81	0.70	16.43 ± 0.05	17.57 ± 0.06	18.2 ± 0.1	> 19.4	> 19.6	> 20.0
0222+185	0.916	18.62 ± 0.25	19.28 ± 0.13	19.35 ± 0.18	> 20.65	–	> 20.85
0537–286	0.125	19.39 ± 0.1	19.94 ± 0.07	21.05 ± 0.19	> 22.25	> 22.31	> 22.82
074625+2549	0.12	19.42 ± 0.14	19.84 ± 0.14	20.09 ± 0.17	> 21.6	> 21.89	> 22.37
0805+6144	0.189	> 19.3	> 20.4	> 20.1	> 20.5	> 20.3	> 20.4
0836+710	0.101	17.17 ± 0.05	17.29 ± 0.03	16.46 ± 0.02	17.34 ± 0.03	17.88 ± 0.05	17.21 ± 0.04
1210+330	0.041	19.01 ± 0.25	18.97 ± 0.11	18.78 ± 0.12	19.38 ± 0.14	19.00 ± 0.11	20.13 ± 0.17
2126–158	0.264	17.03 ± 0.06	18.20 ± 0.06	19.5 ± 0.2	> 19.8	> 20.3	> 20.6
2149–306	0.083	17.66 ± 0.08	17.88 ± 0.06	17.33 ± 0.05	18.31 ± 0.08	20.44 ± 0.24	20.21 ± 0.13
225155+2217	0.272	–	–	> 20.7	> 22.0	> 22.2	> 21.2

confidence interval on the spectral parameters. In Table 2 we list the results.

3.2 UVOT data

UVOT (Roming et al. 2005) source counts were extracted from a circular region of size 5 arcsec centred on the source position, while the background was extracted from a larger circular nearby source-free region. Data were integrated with the uvotimsum task and then analysed by using the uvotsource task.

The observed magnitudes have been dereddened according to the formulae by Cardelli, Clayton & Mathis (1989) and converted into fluxes by using standard formulae and zero-points from Poole et al. (2008). No further absorption was applied. Table 3 lists the result.

4 LYMAN α AND LYMAN-EDGE ABSORPTION

Being at redshifts between 2 and ~ 3.7 , the optical–UV flux of the blazars in our sample could be affected by absorption of neutral hydrogen in intervening Lyman α absorption systems. Single-ionized helium is not an issue at these redshifts for UVOT wavelengths. Not knowing the real attenuation along individual lines of sight, we estimate the *probable* attenuation using the effective opacity $\tau_{\text{eff}} \equiv -\ln(e^{-\tau})$, where the average is taken over all possible lines of sight. We compute the mean attenuation for six wavelengths approximately centred in the six UVOT filters. Fig. 3 shows the corresponding τ_{eff} as a function of redshift. We adopt the column density distribution described in Haardt & Madau (in preparation), which is based on the mean-free-path measurements of Prochaska, Worseck & O’Meara (2009). Full details of our calculation will be




Figure 3. The optical depth due to neutral hydrogen and helium as a function of redshifts. Vertical lines (and labels) indicate the redshifts of the blazar in our sample. The different curves are for the six filters of UVOT, as labelled.

described in Haardt et al. (in preparation), together with a more refined treatment of the mean attenuation and its variance around the mean. Such procedure is very crude, as the attenuation variance along an individual line of sight is large for a fixed observed wavelength. We must note however that (i) the variance is largely reduced when the actual filter width is taken into account (Madau 1995) and (ii) the variance of the attenuation is mainly driven by the few Lyman-limit systems present along an individual line of sight.

Our absorption model results in a mean number of thick systems which is <1 for $z \lesssim 4$, so we do not expect excessive offset of the attenuation along individual line of sight with respect to the mean value.

When presenting the SED of our sources, we will show both the fluxes and upper limits dereddened for the extinction due to our Galaxy and the fluxes (and upper limits) obtained by de-absorbing them with the τ_{eff} shown in Fig. 3.

5 THE SPECTRAL ENERGY DISTRIBUTION

Although the SED of several blazars of our sample have already been presented in the literature, we show them all in Figs 4–6, adding to the archival data the BAT data from the A09 catalogue. The SEDs show the following general properties.

(i) We show as a grey stripe the detection limits of the LAT instrument: the lower bound corresponds to the pre-flight differential sensitivity limit for 1-yr survey at the 5σ level,² while the upper bound is simply a factor of 4 higher. The latter should mimic the approximate sensitivity limit for the 3-month LAT survey at 10σ confidence level. These curves depend on the assumed spectral index of the source, and have been obtained assuming $\alpha_\gamma = 1$, approximately valid for the expected spectrum of FSRQs. All sources in the present sample have not been detected in the first 3-month LAT survey. Their γ -ray flux should then lie below the upper bound of the grey stripe. This, together with the BAT data, constrains the high-energy peak of the SED to occur in the MeV band. Two blazars, 0537–286 and 0836+710, were detected by Energetic Gamma Ray Experiment Telescope (EGRET) (Thompson et al. 1993; Nandikotkur et al. 2007) with a flux level reported in Figs 4 and 5. This indicates a large-variability amplitude.

(ii) The high-energy peak is dominating the electromagnetic output with luminosities exceeding $10^{48} \text{ erg s}^{-1}$ in all cases but S5 0014+813 (in which it is the accretion disc luminosity that dominates the power output).

(iii) The 0.3–10 keV *Swift*/XRT data indicate a very hard X-ray spectrum, harder than the BAT one, but, in general, the extrapolation from the XRT agrees with the BAT flux. The BAT spectrum is a 3-yr average, while the XRT spectrum is more often a one-epoch spectrum, and rarely an average of the available observations.

(iv) In the optical–UV band the spectrum is almost always steep (exceptions are 1210+330 and 225155+2217), and sometimes shows a peak. This component can be interpreted as the emission from the accretion disc (see G10 and Ghisellini et al. 2009a, and Section 4.2).

(v) When there are multiple X-ray observations, these show variability, with an apparent tendency for a ‘harder when brighter’ behaviour of the X-ray flux (see 0222+185; 0836+710; 2126–158; 2149–306 and 225155+2217). The variability amplitude thus is greater at higher energies, being very modest below a few keV.

5.1 The model

To model the SED we have used a relatively simple leptonic, one-zone synchrotron and inverse Compton model. This model, fully discussed in Ghisellini & Tavecchio (2009), has the following main characteristics.

²See http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation/Cicerone/Cicerone_LAT_IRFs/LAT_sensitivity.html




Figure 4. SEDs of S5 0014+813, 0222+185 (i.e. RBS 315) and PKS 0537–286 together with the fitting models, with parameters listed in Table 4. De-absorbed UVOT, XRT and BAT data are indicated by darker symbols (red in the electronic version), while archival data [from NASA/IPAC Extragalactic Database (NED)] are in light grey. Diamonds (and lower arrows, cyan in the electronic version) indicate UVOT data not de-absorbed by intervening Lyman α clouds. The magenta square symbols for S5 0014+813 are *IRAS* and Two Micron All Sky Survey (2MASS) data points. The short-dashed line is the emission from the IR torus, the accretion disc and its X-ray corona. The long-dashed and the dot-dashed grey lines are the synchrotron self-Compton and the external Compton components, respectively. The thick solid (blue) line is the sum of all components.

We assume that in a spherical region of radius R , located at a distance R_{diss} from the central black hole, relativistic electrons are injected at a rate $Q(\gamma)$ ($\text{cm}^{-3} \text{ s}^{-1}$) for a finite time equal to the light crossing time R/c . For the shape of $Q(\gamma)$ we adopt a smoothly broken power law with a break at γ_b :

$$Q(\gamma) = Q_0 \frac{(\gamma/\gamma_b)^{-s_1}}{1 + (\gamma/\gamma_b)^{-s_1+s_2}}. \quad (2)$$

The emitting region is moving with a velocity βc corresponding to a bulk Lorentz factor Γ . We observe the source at the viewing angle




Figure 5. SED of SDSS 074625+2549, PKS 0805–6144, PKS 0836+710 and B2 1210+330. Symbols and lines are as in Fig. 4.

θ_v and the Doppler factor is $\delta = 1/[\Gamma(1 - \beta \cos \theta_v)]$. The magnetic field B is tangled and uniform throughout the emitting region. We take into account several sources of radiation externally to the jet: (i) the broad-line photons, assumed to re-emit 10 per cent of the accretion luminosity from a shell-like distribution of clouds located




Figure 6. SED of PKS 2126–158, PKS 2149–306 and MG3 J225155+2217. Symbols and lines are as in Fig. 4.

at a distance $R_{\text{BLR}} = 10^{17} L_{\text{d},45}^{1/2}$ cm; (ii) the infrared (IR) emission from a dusty torus, located at a distance $R_{\text{IR}} = 2.5 \times 10^{18} L_{\text{d},45}^{1/2}$ cm; (iii) the direct emission from the accretion disc, including its X-ray corona; (iv) the starlight contribution from the inner region of the host galaxy and (v) the cosmic background radiation. All these contributions are evaluated in the blob comoving frame, where we calculate the corresponding inverse Compton radiation from all these contributions, and then transform into the observer frame.

We calculate the energy distribution $N(\gamma)$ (cm^{-3}) of the emitting particles at the particular time R/c , when the injection process ends. Our numerical code solves the continuity equation which includes injection, radiative cooling and e^\pm pair production and reprocessing. Ours is not a time-dependent code: we give a ‘snapshot’ of the predicted SED at the time R/c , when the particle distribution $N(\gamma)$ and consequently the produced flux are at their maximum.

Since we are dealing with very powerful sources, the radiative cooling time of the particles is short, shorter than R/c even for the

least energetic particles. This implies that, at lower energies, the $N(\gamma)$ distribution is proportional to γ^{-2} , while, above γ_b , $N(\gamma) \propto \gamma^{-(s_2+1)}$. The electrons emitting most of the observed radiation have energies γ_{peak} which is close to γ_b (but these two energies are not exactly equal, due to the curved injected spectrum).

We model at the same time the thermal disc (and IR torus) radiation and the non-thermal jet emission. The link between these two components is given by the amount of radiation energy density (as seen in the comoving frame of the emitting blob) coming directly from the accretion disc or reprocessed by the broad-line region (BLR) and the IR torus. This radiation energy density depends mainly on R_{diss} , but not on the adopted accretion rate or black hole mass (they are in any case chosen to reproduce the observed thermal disc luminosity).

To calculate the flux produced by the accretion disc, we adopt a standard Shakura & Sunyaev (1973) disc (see Ghisellini & Tavecchio 2009). This allows us to fit also the thermal radiation seen in the optical–UV range, and to estimate the accretion rate and the black hole mass, as discussed below.

5.2 Estimate of the black hole mass

For all our sources, we interpret the near-IR (NIR), optical and UV emission as due to the accretion disc. For simplicity, we assume that the disc is an optically thin, geometrically thick, Shakura & Sunyaev (1973) disc, emitting a blackbody at each radius. The maximum temperature (and hence the νF_ν peak of the disc luminosity) occurs at ~ 5 Schwarzschild radii and scales as $T_{\text{max}} \propto (L_d/L_{\text{Edd}})^{1/4} M^{-1/4}$. The total optical–UV flux gives $L_d = \eta \dot{M} c^2$. Once we specify the efficiency η , we can derive both the black hole mass and the accretion rate. Assuming a Schwarzschild black hole, we set $\eta = 0.08$.

In Fig. 7 we show a zoom of the SED of PKS 2149–306 to illustrate the uncertainties on the estimated value of the mass of the central black hole. The three SEDs shown correspond to $M = 3 \times 10^9$, 5×10^9 and 10^{10} solar masses. The maximum temperature of the accretion disc is a (albeit weak) function of the black hole mass: the overall disc emission becomes bluer for smaller black




Figure 7. Zoom of the SED of PKS 2149–306 to show how the theoretical SED changes by changing the black hole mass. Three values are shown: $M = 3 \times 10^9$, 5×10^9 and 10^{10} solar masses. Dotted, short- and long-dashed lines show the contribution of the accretion disc and its X-ray corona, while solid lines show the sum of the thermal and the beamed non-thermal components.

hole masses. From Fig. 7 one can see that the $M = 3 \times 10^9 M_\odot$ case gives a poor fit, while the $M = 10^{10} M_\odot$ case tends to overproduce the flux at lower optical frequencies. We can conclude that, within our assumptions and when the data show the peak of the thermal emission, the resulting estimate of the black hole mass is rather accurate with uncertainties that are significantly smaller than a factor of 2.

Three caveats are in order. The first is that we are assuming a standard Shakura & Sunyaev (1973) disc, namely a disc geometrically thin and optically thick, emitting blackbody radiation at each annulus according to a standard temperature profile (e.g. Frank, King & Raine 2002). For very large accretion rates, close to Eddington, the disc structure might be modified, and in the inner region a funnel may develop, as in the so-called ‘thick’ and ‘slim’ discs (see e.g. Abramowicz et al. 1988; Madau 1988; Szuszkiewicz, Malkan & Abramowicz 1996). We have discussed this possibility for S5 0014+813 (Ghisellini et al. 2009a). If a funnel is present, the emitted disc radiation is not isotropic any longer, and face-on observers (as for blazars) would see an amplified radiation, and this would lead to overestimate the black hole mass. We will discuss again this point later.

The second caveat concerns the assumption of a Schwarzschild, non-rotating black hole. The efficiency η for a Kerr hole and corotating accretion disc is larger, as are the temperatures of the innermost radii. But also the gravitational light bending and redshift are larger, and the calculation of the observed spectrum is not as straightforward as in the Schwarzschild case (see e.g. Li et al. 2005). Comparing the spectra calculated in the Schwarzschild and maximal Kerr cases, for equal mass and accretion rate, and assuming that the disc emits blackbody spectra at all radii, we find that the overall disc emission in the Kerr case is bluer and stronger above the emission peak, and very similar below (smaller frequencies are emitted at larger radii, where the emitted flux is insensitive to the black hole rotation). Decreasing the accretion rate would decrease the emission peak (because the maximum temperature would be smaller), but also the overall flux, both below and beyond the emission peak. Since we end up having a deficit in the low-frequency part of the spectrum, we must change also the mass, in the direction of an increase of it, because this allows us to have a smaller maximum temperature. Therefore, to recover the original SED in the case of a Kerr hole, we must decrease the accretion rate and increase the mass. So assuming Kerr holes would not help in decreasing the derived masses.

The third caveat is that the assumption of a blackbody spectrum may be too simplistic: for some regions of the disc a modified blackbody may be a better choice. Since a modified blackbody is a less efficient radiator than a pure blackbody, the resulting spectrum, for the same accretion rate and black hole mass, will be less powerful and bluer (because the disc will be hotter). If this occurs, then our derived black hole masses should be considered as lower limits.

6 RESULTS

Table 4 lists the parameters used to fit the SED of the 10 blazars. We find that they are all distributed in a narrow range, as expected, because we are dealing with extremely powerful objects, characterized by similar (albeit extreme) properties.

The distance from the black hole, R_{diss} , where dissipation occurs and most of the observed radiation is produced, ranges from 500 to 1.4×10^3 Schwarzschild radii. Three sources (0746+253, 1210+330 and marginally 2149–306) have $R_{\text{diss}} > R_{\text{BLR}}$, while for

Table 4. List of parameters used to construct the theoretical SED. Column 1: name; column 2: redshift; column 3: dissipation radius in units of 10^{15} cm and the values in parentheses are in units of Schwarzschild radii; column 4: black hole mass in solar masses; column 5: size of the BLR in units of 10^{15} cm; column 6: power injected in the blob calculated in the comoving frame, in units of 10^{45} erg s^{-1} ; column 7: accretion disc luminosity in units of 10^{45} erg s^{-1} and the values in parentheses are in units of L_{Edd} ; column 8: magnetic field in gauss; column 9: bulk Lorentz factor at R_{diss} ; columns 10 and 11: break and maximum random Lorentz factors of the injected electrons; columns 12 and 13: slopes of the injected electron distribution $[Q(\gamma)]$ below and above γ_b . The total X-ray corona luminosity is assumed to be in the range of 10–30 per cent of L_d . Its spectral shape is assumed to be always proportional to $\nu^{-1} \exp(-h\nu/150 \text{ keV})$. The viewing angle θ_v is 3° for all sources.

Name	z	R_{diss}	M	R_{BLR}	P'_i	L_d	B	Γ	γ_b	γ_{max}	s_1	s_2
1	2	3	4	5	6	7	8	9	10	11	12	13
0014+813	3.366	9.6e3 (800)	4e10	5.5e3	0.12	3e3 (0.5)	0.33	16	150	2e3	-1	3
0222+185	2.69	660 (550)	4e9	1.45e3	0.08	210 (0.35)	2.94	13.5	40	2e3	0	3.2
0537-286	3.104	420 (700)	2e9	735	0.13	54 (0.18)	1.92	15	50	2e3	-1	3
0746+254	2.979	1.65e3 (1.1e3)	5e9	866	0.24	75 (0.1)	0.1	15	200	5e3	0.75	2.6
0805+614	3.033	270 (600)	1.5e9	581	0.15	34 (0.15)	2.54	14	60	3e3	-0.5	3
0836+710	2.172	540 (600)	3e9	1.5e3	0.22	225 (0.5)	3.28	14	90	2e3	-1	3.6
1210+330	2.502	420 (1.4e3)	1e9	866	0.08	75 (0.5)	0.73	16	80	3e3	-0.5	3.5
2126-158	3.268	1.8e3 (600)	1e10	2.7e3	0.13	750 (0.5)	2.61	14.1	40	2e3	0	3.3
2149-306	2.345	1.2e3 (800)	5e9	1.24e3	0.18	150 (0.2)	1.12	15	60	3e3	0	3.3
225155+2217	3.668	300 (1e3)	1e9	1.06e3	0.15	112 (0.75)	2.78	15	70	2e3	0	4
Average	3	1.8e3 (1e3)	6e9	1.3e3	0.15	180 (0.2)	0.82	15	100	2e3	0	3
(LAT $z > 2$)	2.2	630 (700)	3e9	948	0.1	90 (0.2)	1.65	15	300	4e3	0	2.8

all the others the location of the dissipation region lies within the BLR.

The injected power in the form of relativistic electrons is very similar, around 10^{44} erg s^{-1} for all sources, and since the cooling is severe, all of it is transformed in radiation. The accretion disc luminosity is at the level of 0.1–0.4 L_{Edd} , even if it spans a large range in absolute units, between 3×10^{46} and 2×10^{48} erg s^{-1} , due to the range of the black hole masses. The bulk Lorentz factor is between 13 and 18.

6.1 Comparison with bright *Fermi*/LAT blazars

It is instructive to compare the jet power and accretion disc luminosities of the BAT blazars of our sample with all the bright blazars detected by the LAT 3-month survey, as analysed in G10. The jet carries power in the form of bulk motion of particles (electrons and protons) and magnetic field. Furthermore, we can calculate the power spent by the jet to produce the radiation we see. The different components of the jet power are

$$P_i = \pi r_{\text{diss}}^2 \Gamma^2 \beta c U'_i, \quad (3)$$

where U'_i is the energy density of the i component, as measured in the comoving frame. We define the total jet power as the sum of the electron (P_e), proton (P_p), found assuming one proton per electron, and magnetic field (P_B) contributions, while we call P_r the power spent in producing radiation. Table 5 lists the jet powers for the 10 BAT blazars.

P_r versus L_d : Fig. 8 shows P_r as a function of the accretion luminosity L_d for our BAT blazars, together with all the LAT blazars in G10. These include BL Lac objects for which only an upper limit to their disc luminosity could be found (shown by arrows). Black filled diamonds correspond to the BAT blazars. The power $P_r = \pi r_{\text{diss}}^2 \Gamma^2 \beta c U'_{\text{rad}}$ can be rewritten as [using $U'_{\text{rad}} = L'/(4\pi r_{\text{diss}}^2 c)$]

$$P_r = L' \frac{\Gamma^2}{4} = L \frac{\Gamma^2}{4\delta^4} \sim L \frac{1}{4\delta^2}, \quad (4)$$

where L is the total observed non-thermal luminosity (L' is in the comoving frame) and U'_{rad} is the radiation energy density produced

Table 5. Logarithm of the jet power in the form of radiation, Poynting flux and bulk motion of electrons and protons (assuming one proton per emitting electron). Powers are in erg s^{-1} .

Name	$\log P_r$	$\log P_B$	$\log P_e$	$\log P_p$
0048-071	46.42	46.98	45.73	47.49
0222+185	46.11	46.42	45.27	47.89
0537-286	46.43	45.74	45.52	48.01
0746+253	46.53	44.36	46.28	47.85
0805+614	46.42	45.34	45.64	47.99
0836+710	46.60	46.36	45.54	48.00
1210+330	46.30	44.95	45.27	47.79
2126-158	46.38	47.27	45.12	47.84
2149-306	46.59	46.18	45.25	48.02
225155+2217	46.48	45.77	45.74	48.19
Average	46.43	45.96	45.54	47.92
(LAT $z > 2$)	46.3	46.0	44.8	47.2

by the jet (i.e. excluding the external components). The last equality assumes $\theta_v \sim 1/\Gamma$. This quantity is almost model independent since it depends only on the adopted δ , which can be estimated also by other means, namely superluminal motions. Therefore Fig. 8 shows two quantities that are (almost) model independent. The BAT blazars are the most extreme, lying at the upper end of the P_r – L_d distribution. All follow the trend defined by the LAT FSRQs, with the exception of S5 0014+813, which has accretion disc luminosity larger than that expected from its jet luminosity.

P_{jet} versus L_d : Fig. 9 shows the jet power $P_{\text{jet}} = P_p + P_e + P_B$ as a function of the disc luminosity. Again, the BAT blazars have the most powerful jets and discs, and S5 0014+813 appears to be an outlier with respect to the general trend defined by the ensemble of LAT and BAT blazars.

P_{jet} versus P_r : Fig. 10 shows the jet power as a function of P_r , the power in radiation. It shows the efficiency of the jet in converting its bulk power into radiation. Low-power BL Lacs are the most efficient in converting P_{jet} into P_r , while powerful blazars are less




Figure 8. The power spent by the jet to produce the radiation we see, P_r , as a function of the accretion disc luminosity L_d . BAT blazars (black diamonds) are compared with the blazars in the *Fermi* 3-month catalogue of bright sources detected above 100 MeV. The latter have different symbols according to their γ -luminosity, as labelled.




Figure 9. The jet power P_{jet} as a function of the accretion disc luminosity L_d . Symbols are as in Fig. 8.

efficient. The BAT blazars follows this trend, and in this plot S5 0014+813 is not an outlier.

This suggests that S5 0014+813 indeed has an overluminous accretion disc with respect to its jet, and this favours the hypothesis, discussed in Ghisellini et al. (2009a) that the thermal radiation from the inner parts of the disc is geometrically collimated by the presence of a funnel, possibly as a consequence of the large accretion rate in Eddington units (likely to be even larger than that calculated here and in Ghisellini et al. 2009a because we assumed a standard disc).

For the other BAT blazars, instead, there is no need to invoke a non-standard accretion disc, since they follow the jet power – disc luminosity relation defined by less extreme blazars, selected from the *Fermi*/LAT survey. The latter have luminosities around 0.1 of the Eddington one or even less, and should indeed have standard discs.




Figure 10. The jet power P_{jet} as a function of the power spent by the jet to produce the radiation we see, P_r . Symbols are as in Fig. 8.




Figure 11. Comparison of black hole mass distributions. The top panel shows all *Fermi*/LAT FSRQs (i.e. excluding BL Lacs), and the nine *Swift*/BAT FSRQs with $z > 1.8$. The bottom panel shows the mass distribution of the 10 BAT blazars at $z > 2$.

This suggests that the black hole masses estimated for the other BAT blazars are robust, and not a consequence of our assumption of a standard accretion disc.

It is then interesting to compare the black hole masses derived for our BAT blazars with those of the 3-month survey of LAT, and then to the subsample (of nine objects) of LAT blazars at $z > 1.8$. This is done in Fig. 11: high-redshift BAT blazars have a black hole mass distribution extending to larger values than the high- z LAT blazars. The numbers are too small for a statistical comparison, and a KS test returns a large probability that the two distributions are drawn from the same one. In any case, we can safely conclude that the hard X-ray selection is a very efficient way to pinpoint the most powerful extreme of the blazar population, possibly more than selecting them through their *Fermi*/LAT γ -ray flux. We can wonder if this merely depends on the sensitivities of the current BAT and LAT instruments, or if indeed the most powerful blazar jets are more luminous in hard X-rays than in the 0.1–10 GeV band. In Fig. 12 we try to answer to this question by showing three theoretical SEDs




Figure 12. Average SEDs for the sources in our samples, for all FSRQs detected by *Fermi* and for the subsample of the eight *Fermi* blazars at redshift $z > 1.8$. The top and the bottom panels show the fluxes and luminosities, respectively. The shown frequencies are calculated in the rest frame of the source for the luminosity plot, and are the observed ones for the flux plot. In Table 4 we list the parameters used to construct the shown SEDs for FSRQs, while, for the LAT BL Lacs, we report the same SED as shown in G10.

corresponding to (i) the average SED of BL Lacs detected by the 3-month LAT survey (see the used parameters in G10); (ii) the average SED of the $z > 2$ FSRQs detected by the same LAT survey and (iii) the average SED of our $z > 2$ BAT FSRQs. For the latter two SEDs we used the parameters listed at the end of Table 4. We show both the flux (top panel) and the luminosity (bottom) SEDs. The shown frequencies are calculated in the rest frame of the source for the luminosity plot, and are the observed ones for the flux plot. Fig. 12 shows the following.

- (i) The LAT BL Lacs and high-redshift FSRQs have the same average flux in hard X-rays.
- (ii) The LAT BL Lacs are on average brighter than high- z FSRQs in γ -rays.
- (iii) Comparing high-redshift LAT and BAT FSRQs, we see that the latter are fainter in γ -rays and in fact none of our 10 BAT blazars has been detected in the 3-month LAT survey.
- (iv) In the luminosity plot, LAT BL Lacs are the least luminous (most of them are at $z < 0.5$, see Fig. 1).
- (v) Comparing high- z BAT and LAT FSRQs, we see that BAT blazars are slightly more luminous (in bolometric terms) even if they are less powerful γ -ray sources. In the entire X-ray band they are more powerful than LAT blazars.

We have checked if some of the high-redshift LAT and BAT blazars are present on the Sloan Digital Sky Survey (SDSS), with their black hole mass estimated through the full width at half-maximum of the emission lines and the ionizing continuum luminosity. In the compilation of Shen et al. (2008) we have found RGB J0920+446 ($z = 2.190$) with different mass estimates, ranging from $M = 2 \times 10^9 M_\odot$ and $5 \times 10^{10} M_\odot$, to be compared with our estimate of $M = 6 \times 10^9 M_\odot$ (Ghisellini, Tavecchio & Ghirlanda 2009c); 4C +38.41 (i.e. 1633+382, $z = 1.814$) with a mass in the range $(1.6\text{--}8) \times 10^9 M_\odot$

(our estimate: $M = 5 \times 10^9 M_\odot$; Ghisellini et al. 2009c) and J074625.87+2549 with $M = 6 \times 10^9 M_\odot$ (our estimate: $M = 7 \times 10^9 M_\odot$, see Table 4). Also PKS 1502+106 ($z = 1.839$) is present in the SDSS (DR7) catalogue, but with no mass estimate yet.

All this phenomenology can be understood in simple terms on the basis of the blazar sequence: the most powerful blazars are FSRQs whose synchrotron peak is located in the submillimetre band, and whose high-energy peak is located in the MeV band. So the 0.1–10 GeV luminosity, being in a band relatively far from the peak, is not very large. On the contrary, instruments sensitive in the hard X-ray band can catch these sources at their emission peak, making this band the optimal one in finding the upper end of the blazar luminosity distribution.

7 DISCUSSION

One of our major result is that *all* the 10 BAT blazars studied here have a black hole heavier than 10^9 solar masses. Among the BAT blazars they are the most luminous (all of them have $L_X > 2 \times 10^{47}$ erg s $^{-1}$) and the most distant (all of them have $z > 2$). Since these objects are at high redshifts, our finding has important implications on the number density of heavy black holes, especially if we consider that for each blazar pointing at us, there must be hundreds of similar sources (having black holes of similar masses) pointing elsewhere. Using a mix of simple theoretical and observational considerations we will find in the following what we consider a conservative ‘minimal mass function’ for heavy black holes associated to radio-loud objects. But before doing this we first discuss the implications of our results for future hard X-ray missions.

7.1 Implications for future missions

NHXM: The *NHXM*³ is a project for a satellite hosting four mirror modules able to concentrate X-ray photons in the 0.2–80 keV range (Pareschi et al. 2009). This is achieved in part by a long (10 m) focal length and partly by a multilayer coating. A 2–30 keV polarimeter is foreseen at the focus of one of the mirror modules. The angular resolution is better than 20 arcsec at 30 keV. The sensitivity above 10 keV is two to three orders of magnitude better than anything already flown. On the other hand, the long focal length limits the field of view to $\sim 12 \times 12$ arcmin².

The main contribution of this mission to the issue of finding large black hole masses at high redshift is to observe pre-selected candidates in the hard X-ray region of the spectrum, and consequently to assess the non-thermal nature of this emission. By establishing a large jet power, it will hint to a corresponding large accretion rate and thus a large black hole mass. In this context, we stress that the blazars we have analysed in this paper are all strong (> 0.1 Jy) radio sources and very hard in the ‘classical’ 2–10 keV band. These properties can then be taken as selection criteria to construct a useful sample of radio-loud sources to be observed by *NHXM*.

EXIST: The *EXIST*⁴ is a proposed *Medium Class Mission* to conduct the most sensitive full-sky survey for black holes on all scales (from the stellar to the supermassive ones). To this purpose *EXIST* has been specifically designed to have on board three complementary instruments:

- (i) a large-area (4.5 m²), wide-field (70° × 90°) hard X-ray (5–600 keV) imaging (2-arcmin resolution and ~ 20 -arcsec localization

³See <http://www.brera.inaf.it/NHXM2/>

⁴See <http://exist.gsfc.nasa.gov/>

for 5σ sources) coded mask telescope [High Energy Telescope (HET)];

(ii) a Soft X-ray Imager (SXI; focusing telescope with CCD, ~ 2 -arcsec localizations) operating over the 0.1–10 keV energy range with *XMM-Newton*-like (1-telescope) area;

(iii) an optical/NIR telescope (IRT) with 1.1 m diameter with instruments covering the wavelength range 3000–22 000 Å with $R \sim 3000$ spectrograph and 4×4 -arcmin² imaging array with objective prism capability.

The HET has approximately a 20 times better sensitivity than BAT, and extends the energy range to lower and higher energies: it can then see powerful and distant blazars *at or very close to their peak of emission*. Furthermore, the operation mode (similar to *Fermi*), patrolling the entire sky every three hours, is ideal to discover rare objects. The SXI can give crucial information about the level and shape of the 0.2–10 keV spectrum as well as to locate the X-ray sources on sky with an accuracy of few arcsec. Finally, the IRT can take the IR and optical spectra (and/or photometry) of these rare sources, and can catch powerful blazars *where their accretion disc peaks*. Given the expected flux levels of IR–optical and X-ray fluxes of high-redshift powerful blazars, these ‘follow-up’ observations will be very inexpensive. *EXIST* has been designed to break the ‘multiwavelength investigation bottleneck’ by having on board a suite of instruments on unprecedented wavelength coverage that operate simultaneously.

To illustrate these concepts, Fig. 13 shows the SED of PKS 2149–306, the theoretical fitting model, and the same model for a source located at $z = 8$ having the same total luminosity. We




Figure 13. The SEDs and model for PKS 2149–306 at its actual redshift ($z = 2.345$) and what the model SED would appear if the source were at $z = 8$. We also show the limiting sensitivities for the three instruments foreseen to be on board the *EXIST* mission: the high-energy-coded mask (HET, sensitive in the 5–600 keV range), the soft X-ray telescope (SXI, operating in the 0.1–10 keV energy range) and the IR–optical telescope (IRT). We indicate the exposure time needed to reach the shown sensitivities. It is remarkable that powerful blazars like PKS 2149–306, if it existed even at $z = 8$, could easily be detected by hard X-ray telescopes like *EXIST*/HET. The presence of SXI and especially the IR–optical telescope would also allow us to find the redshift and to very easily provide a complete spectrum of the accretion disc at its emission peak, thus yielding a robust estimate of the accretion rate and the black hole mass.




Figure 14. X-ray luminosity as a function of redshift for a simulated sample of blazars detectable by the 2-yr all-sky survey of *EXIST*. The inset shows the entire distribution, while the main figure zooms the region of high luminosity. This expected sample of blazars detectable by *EXIST* has been produced by extrapolating the cosmological evolution properties of the BAT blazars as derived by A09.

superimpose also the expected sensitivities of the three *EXIST* instruments (for the indicated exposure times). If blazars like PKS 2149–306 exist at high redshifts, *EXIST* will be able to find them and characterize their general physical properties in a rather complete way. These include the jet power, the accretion disc luminosity and the black hole mass, which can be estimated in the same way as done in this paper.

Of course the choice of $z = 8$ (for Fig. 13) does not imply that blazars of this kind do exist at this redshift. To evaluate the number of blazars detectable by *EXIST* in the all-sky survey we used the X-ray luminosity function and the cosmological evolution model of blazars as recently derived by A09 from the analysis of the BAT data. In particular, we used the modified pure luminosity evolution model with best-fitting parameters as reported in fig. 9 of A09,⁵ renormalizing the luminosity function to the number of blazars detected by BAT (38 sources). A power-law spectrum with photon index equal to 1.5 has been assumed to convert the (15–55 keV) A09 luminosity function to the 10–40 keV energy range and to compute *K*-corrections.

Assuming a flux limit in the 10–40 keV energy band of $\sim 8 \times 10^{-13}$ erg cm⁻² s⁻¹ for the 2-yr *EXIST* survey data, we expect ~ 19000 blazars in the all-sky survey. Their distribution in the luminosity–redshift plane is shown in Fig. 14, while their redshift distribution (compared with the redshift distribution of the BAT blazars) is shown in Fig. 15 (solid line). Table 6 lists the expected number of blazars as a function of z , together with the number of high X-ray luminosity blazars possibly hosting a heavy black hole (columns 2 and 3, respectively). This makes the great discovery potential of *EXIST* clear; more than 1500 blazars are expected at redshift greater than 4 and about a dozen at redshift greater than 7.

⁵We used $\gamma_1 = -0.87$, $\gamma_2 = 2.73$, $L_* = 1.8 \times 10^{44}$ erg s⁻¹, $k = 4.8$, $\gamma = -0.36$, see A09 for details.




Figure 15. The redshift distribution of the blazars detectable by the 2-yr all-sky survey of *EXIST* compared with the redshift distribution of the BAT blazars (dashed line). The dotted histogram corresponds to the ‘minimal’ LF as discussed in the text.

Table 6. Number of blazars detectable by *EXIST* above a given redshift, according to the luminosity function derived in A09 and the limiting sensitivity of *EXIST* for the 2-yr all-sky survey. The third column gives the number of objects, above a given z and with X-ray luminosities above $L_{\text{thr}} = 2 \times 10^{47} \text{ erg s}^{-1}$. The fourth and fifth columns give the total number of blazars, and those exceeding L_{thr} , for the ‘minimal’ LF discussed in the text.

z	$N_{\text{A09}}(>z)$	$L_X > L_{\text{thr}}$	$N_{\text{min}}(>z)$	$L_X > L_{\text{thr}}$
0	1.9e4	223	1.83e4	121
1	1.6e4	223	1.53e4	121
2	1.0e4	222	9.7e3	120
3	4.9e3	199	4.3e3	102
4	1.6e3	154	1.0e3	57
5	335	76	41	5
6	58	24	1	0
7	11	9	0	0
8	3	3	0	0

The expected number of high X-ray luminosity blazars reported in column 3 of Table 6 clearly depends on the extrapolation of the A09 cosmological evolution model at redshift greater than ~ 4 , a redshift range where no BAT data are available. To derive a ‘strict’ lower limit to the number of high X-ray luminosity blazars detectable by *EXIST* we have used a cosmological evolution model that is equal to the A09 one up to $z \sim 4.3$ (where they measure the peak of the density of high X-ray luminosity blazars) but having an exponential cut-off beyond $z = 4.3$ [$L_z = L_{z=4.3} \times \exp(z - 4.3)$ for $z > 4.3$]. This luminosity function (LF) is consistent with the lower limits on the number of blazars known between $z = 4$ and 5 (see Fig. 16). In the following we will call this evolution model ‘minimal LF’. The expected number of blazars computed using this ‘minimal LF’ are reported in Table 6 (columns 4 and 5), while their redshift distribution is shown in Fig. 15 (dotted line). As expected the only differences are at redshift greater than 4, where for this rather extreme ‘minimal LF’ we expect to detect about 60 high X-ray luminosity blazars at $z > 4$, five at $z > 5$ and none at $z > 6$. The real number of high X-ray luminosity




Figure 16. The number density of black holes with $M > 10^9 M_{\odot}$ as a function of redshift. The grey stripe is based purely on connecting black hole mass to halo mass (upper envelope: Pizzella et al. 2005; bottom: Baes et al. 2003) and estimating the number density of black holes via the number density of their host dark matter haloes using the Press & Schechter formalism (see text). In the lower part of the figure we report, as grey stripes (red and green in the electronic version), the mass function $\Phi(z, M > 10^9 M_{\odot})$ for blazars as derived from the luminosity function of A09, considering $L_X > 2 \times 10^{47} \text{ erg s}^{-1}$. These stripes correspond to two ‘extreme’ cases of cosmological evolution for the blazars population beyond redshift 4: the upper (red) stripe corresponds to the cosmological evolution model of A09 extrapolated up to $z \sim 9$, while the lower stripe corresponds to its ‘high- z cut-off version’ as discussed in the text. The filled square in the $3 < z < 4$ bin is taken directly from fig. 10 of A09. The filled pentagons and arrows are the lower limits derived from the existence of a few blazars in the 2 redshift bins. In the upper part of the figure we show the same points/stripes upshifted by the factor $2\Gamma^2 = 450$ to account for misaligned sources. In this way the upper stripe of the mass function $\Phi(z, M > 10^9 M_{\odot})$ for radio-loud sources is in conflict, at high z , with the estimates derived by massive haloes, while the lower (green) stripe derived through the ‘minimal LF’ is consistent.

blazars detectable by *EXIST* are probably in-between the numbers reported in Table 6 (columns 3 and 5).

7.2 Black hole–dark halo connection at high redshifts

It is well known that growing a black hole of a billion solar masses within a billion years from the big bang is a challenge for hierarchical models of structure formation (e.g. Haiman 2004; Shapiro 2005; Volonteri & Rees 2005, 2006; Tanaka & Haiman 2009). Assuming accretion at the Eddington rate, a black hole mass increases in time as

$$M(t) = M(0) \exp\left(\frac{1 - \epsilon}{\epsilon} \frac{t}{t_{\text{Edd}}}\right), \quad (5)$$

where $t_{\text{Edd}} = 0.45 \text{ Gyr}$ and ϵ is the radiative efficiency. For a ‘standard’ radiative efficiency $\epsilon \approx 0.1$, and a seed mass $M(0) = 10^2 - 10^4 M_{\odot}$, it takes at least 0.7–0.9 Gyr to grow up to $\sim 10^{10} M_{\odot}$. The cosmic time at $z = 4$ is 1.5 Gyr, but it is only 0.9 Gyr at $z = 6$, and 0.7 Gyr at $z = 7$. We expect therefore that black holes of billion solar masses at higher and higher redshift become increasingly rare.

We will provide an estimate of the number density of black holes with $M > 10^9 M_{\odot}$ as a function of redshift that is independent of the formation and growth efficiency of black holes. Empirical correlations have been found between the black hole mass (M)

and the central stellar velocity dispersion (σ) of the host (Gültekin et al. 2009, and references therein), and between the central stellar velocity dispersion and the asymptotic circular velocity (V_c) of galaxies (Ferrarese 2002; Baes et al. 2003; Pizzella et al. 2005):

$$\sigma = 200 \text{ km s}^{-1} \left(\frac{V_c}{320 \text{ km s}^{-1}} \right)^{1.35} \quad (6)$$

and

$$\sigma = 200 \text{ km s}^{-1} \left(\frac{V_c}{339 \text{ km s}^{-1}} \right)^{1.04} \quad (7)$$

as suggested by Pizzella et al. (2005) and Baes et al. (2003), respectively.

The latter is a measure of the total mass of the dark matter halo of the host galaxies. A halo of mass M_h collapsing at redshift z has a circular velocity

$$V_c = 142 \text{ km s}^{-1} \left(\frac{M_h}{10^{12} M_\odot} \right)^{1/3} \left(\frac{\Omega_m}{\Omega_m^z} \frac{\Delta_c}{18\pi^2} \right)^{1/6} (1+z)^{1/2}, \quad (8)$$

where Δ_c is the overdensity at virialization relative to the critical density. For a *WMAP5* (the five years data release of the Wilkinson Microwave Anisotropy Probe) cosmology we adopt here the fitting formula (Bryan & Norman 1998) $\Delta_c = 18\pi^2 + 82d - 39d^2$, where $d \equiv \Omega_m^z - 1$ is evaluated at the collapse redshift, so that $\Omega_m^z = \Omega_m(1+z)^3 / [\Omega_m(1+z)^3 + \Omega_\Lambda + \Omega_k(1+z)^2]$.

We will further assume that the black hole– σ scaling is

$$M = 10^9 M_\odot \left(\frac{\sigma}{356 \text{ km s}^{-1}} \right)^4 \quad (9)$$

and that these scaling relations observed in the local universe hold at all redshifts. Therefore we derive the relationship between black hole and dark matter halo mass (see also Bandara, Crampton & Simard 2009):

$$M_h = 4.1 \times 10^{13} M_\odot \left(\frac{M}{10^9 M_\odot} \right)^{0.56} \left(\frac{\Omega_m}{\Omega_m^z} \frac{\Delta_c}{18\pi^2} \right)^{-1/2} (1+z)^{-3/2} \quad (10)$$

and

$$M_h = 7.1 \times 10^{13} M_\odot \left(\frac{M}{10^9 M_\odot} \right)^{0.72} \left(\frac{\Omega_m}{\Omega_m^z} \frac{\Delta_c}{18\pi^2} \right)^{-1/2} (1+z)^{-3/2}. \quad (11)$$

The number density of black holes with $M > 10^9 M_\odot$ therefore corresponds to the number density of haloes with mass $M_h > M_{\text{thr}}$, if M_{thr} is the mass of a halo that hosts a black hole of billion solar masses. We estimate the number density of dark matter haloes using the Press & Schechter formalism (Sheth & Tormen 1999). In Fig. 16 we show the results as the grey stripe, encompassing the values found by the two methods above. This estimate ignores any issue related to black hole formation and growth time-scale. The grey stripe can then be considered as a rough upper limit to the number density of heavy black holes as a function of redshift.

7.3 Large black hole masses at high redshift

These model predictions can be compared with the volume density of high-redshift blazars hosting a black hole of mass larger than $10^9 M_\odot$. The latter can be found using the cosmological evolution model of A09 along its high- z cut-off (i.e. ‘minimal’) version (see Section 7.1), assuming, as we have found in this paper that all blazars with $L_X > 2 \times 10^{47} \text{ erg s}^{-1}$ have a black hole of mass

$M > 10^9 M_\odot$. We cannot exclude that blazars with lower X-ray luminosity also host massive black holes, so the ‘observational’ points, strictly speaking, are the lower limits.

The lower part of Fig. 16 shows two stripes (red and green, in the electronic version) corresponding to two mass functions $\Phi(z, M > 10^9 M_\odot)$ of blazars, both calculated for X-ray luminosities larger than $2 \times 10^{47} \text{ erg s}^{-1}$. The flatter (red) stripe corresponds to the cosmological evolution model of A09 extrapolated up to $z \sim 9$, while the steeper (green) stripe corresponds to its ‘high- z cut-off version’ (‘minimal’ LF, see Section 7.1 for details).

The lower limits shown with a (blue) pentagon and arrow in the 4–5 redshift bin corresponds to the existence of at least four blazars for which we have estimated a black hole mass larger than $10^9 M_\odot$. They are RXJ 1028.6–0844 ($z = 4.276$; Yuan et al. 2005); GB 1508+5714 ($z = 4.3$; Hook et al. 1995); PMN J0525–3343 ($z = 4.41$; Worsley et al. 2004a) and GB 1428+4217 ($z = 4.72$; Worsley et al. 2004b). There are other three blazars with $4 < z < 5$ discussed in Yuan et al. (2006), but they have X-ray luminosities smaller than $10^{47} \text{ erg s}^{-1}$. The other lower limit shown by the (blue) pentagon and arrow in the 5–6 redshift bin corresponds to the existence of at least one blazar, Q0906+6930 at $z = 5.47$, with an estimated black hole mass of $2 \times 10^9 M_\odot$ (Romani 2006).

All these points concerns sources pointing at us. The real density of heavy black holes must account for the much larger population of misaligned sources. We have then multiplied the mass function $\Phi(z, M > 10^9 M_\odot)$ of blazars and the other lower limits by $2\Gamma^2 = 450$, i.e. we have assumed an average Γ factor of 15, appropriate for the BAT blazars analysed here. Fig. 16 shows the resulting points.

The mass function of heavy black holes of all jetted sources is now close or even greater (if we extend the cosmological evolution model of A09 beyond $z \sim 4$) than the upper limit defined by ‘halo–black holes’ (grey stripe) at the largest redshifts. The mass function derived by the ‘minimal’ LF is instead consistent.

To summarize, the BAT blazar survey allowed us to meaningfully construct the hard X-ray LF of blazars. We have shown that its high-luminosity end can be translated into the mass function of black holes with more than one billion solar masses. Up to $z = 4$, where we do see blazars, the cosmological evolution model, as derived by A09, is secure. Beyond $z = 4$ it depends strongly on the assumed evolution. We have then constructed the minimal evolution consistent with the existing data and the (few) existing lower limits. As Fig. 16 shows, the true mass function of heavy black hole in jetted sources should be bracketed by the two shown mass functions derived from the A09 and the ‘minimal’ LF. The true mass density should then lie in-between the two possible choices. The implications of this finding are far-reaching, and we plan to investigate them in a forthcoming paper (Volonteri et al. in preparation).

8 CONCLUSIONS

We summarize here our main conclusions.

(i) In the α_X – L_X plane the ensemble of blazars detected by BAT separates quite clearly in BL Lac objects and FSRQs: the former have steeper spectra and lower luminosities. This is a manifestation of the blazar sequence since low-power blazars are characterized by a population of emitting electrons with a large energy break, implying larger synchrotron and inverse Compton frequency peaks. In the hard X-ray range we often see, in BL Lacs, the steep synchrotron tail of emission, while in more powerful blazars (i.e. FSRQs), we see the hard inverse Compton component.

(ii) In the same plane there is an indication of a ‘divide’, namely BL Lacs and FSRQs separate in luminosity, at a few $\times 10^{45} \text{ erg s}^{-1}$.

This behaviour mirrors what occurs for *Fermi*/LAT blazars: the dividing luminosity indicates when the accretion disc changes mode of accretion, becoming radiatively inefficient for luminosities of the order of 0.3–1 per cent of the Eddington one.

(iii) The 10 BAT blazars at $z > 2$ are among the most powerful known. Not only their beamed jet bolometric luminosity, but also their jet power and the accretion luminosities are among the largest ones.

(iv) The black hole masses are also very large to account for the observed disc luminosities and spectra. They are all greater than one billion solar masses and a few approach 10 billions.

(v) The ‘record holder’ S5 0014+813, having a black hole mass of 40 billion solar masses, is an outlier with respect to the jet power – disc luminosity correlation defined by the *Fermi*/LAT FSRQs and obeyed by the high-redshift BAT blazars. This leads us to favour the hypothesis that its accretion disc is non-standard, having, in its inner regions, a funnel collimating the radiation around the jet axis. An anisotropy factor of ~ 10 is enough to make this object consistent with the jet power – disc luminosity correlation. Consequently, its mass could be smaller (by a similar factor of 10).

(vi) For each blazar pointing at us and detected through its beamed non-thermal emission, there should be other $\sim \Gamma^2$ at the same redshift with similar properties, including the mass of the black hole. This puts a lower limit to the density of heavy black holes in the $3 < z < 4$ redshift range.

(vii) Hard X-ray surveys can catch powerful and distant blazars where their high-energy SED peaks. This implies that future X-ray missions such as *EXIST* and *NHXM* will be the most effective way to find and study the most extreme radio-loud objects. In the *Fermi*/LAT 0.1–100 GeV energy range these objects, having steep spectra, are less conspicuous and can be missed by *Fermi*. According to the *EXIST* sensitivity in hard X-rays, powerful FSRQs can be easily detected even at redshift 8, if they exist. According to the X-ray luminosity function derived by A09, *EXIST* should detect 500–1500 blazars at $z > 4$ and 20–60 at $z > 6$, allowing us to derive the mass function of radio-loud active galactic nuclei up to very large redshifts.

ACKNOWLEDGMENTS

We thank the anonymous referee for his/her comments. We thank Andrea Merloni and Marco Ajello for discussions. This work was partly financially supported by a 2007 COFIN-MIUR and an ASI I/088/06/0 grants. This research made use of the NED which is operated by the Jet Propulsion Laboratory, Caltech, under contract with NASA, and the *Swift* public data made available by the HEASARC archive system. We also thank Neil Gehrels and the *Swift* team for quickly approving and performing the requested too observations of 1210+330.

REFERENCES

Abdo A. A. et al., 2009, *ApJ*, 700, 597
 Abramowicz M. A., Czerny B., Lasota J. P., Szuszkiewicz E., 1988, *ApJ*, 332, 646
 Ajello M. et al., 2009, *ApJ*, 699, 603 (A09)
 Baes M., Buyle P., Hau G. K. T., Dejonghe H., 2003, *MNRAS*, 341, L44
 Bandara K., Crampton D., Simard L., 2009, *ApJ*, 704, 1135
 Bassani L. et al., 2007, *ApJ*, 669, L1
 Bianchin V. et al., 2009, *A&A*, 496, 423
 Böck M., Kadler M., Tosti G., Burnett T., Ohia R., Müller C., Wilms J., 2010, *Proc. 2009 Fermi Symp.*, preprint (arXiv:0912.4192)
 Bottacini E. et al., 2010, *A&A*, 509, A69
 Brunthaler A. et al., 2000, *A&A*, 357, L45
 Bryan G. L., Norman M. L., 1998, *ApJ*, 495, 80

Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
 Cusumano G. et al., 2010, *A&A*, 510, A48
 Donato D., Ghisellini G., Tagliaferri G., Fossati G., 2001, *A&A*, 375, 739
 Ferrarese L., 2002, *ApJ*, 578, 90
 Foschini L. et al., 2006, *A&A*, 453, 829
 Fossati G., Maraschi L., Celotti A., Comastri A., Ghisellini G., 1998, *MNRAS*, 299, 433
 Frank J., King A., Raine D. J., 2002, *Accretion Power in Astrophysics*. Cambridge Univ. Press, Cambridge
 Ghisellini G., Tavecchio F., 2009, *MNRAS*, 397, 985
 Ghisellini G., Celotti A., Fossati G., Maraschi L., Comastri A., 1998, *MNRAS*, 301, 451
 Ghisellini G., Foschini L., Volonteri M., Ghirlanda G., Haardt F., Burlon D., Tavecchio F., 2009a, *MNRAS*, 399, L24
 Ghisellini G., Maraschi L., Tavecchio F., 2009b, *MNRAS*, 396, L105
 Ghisellini G., Tavecchio F., Ghirlanda G., 2009c, *MNRAS*, 399, 2041
 Ghisellini G., Tavecchio F., Foschini L., Ghirlanda G., Maraschi L., Celotti A., 2010, *MNRAS*, 402, 497
 Gültekin K. et al., 2009, *ApJ*, 698, 198
 Haiman Z., 2004, *ApJ*, 613, 36
 Hook I. M., McMahon R. G., Patnaik A. R., Browne I. W. A., Wilkinson P. N., Iwrin M. J., Hazard C., 1995, *MNRAS*, 273, L63
 Kalberla P. M. W., Burton W. B., Hartmann D., Arnal E. M., Bajaja E., Morras R., Pöppel W. G. L., 2005, *A&A*, 440, 775
 Li L.-X., Zimmerman E. R., Narayan R., McClintock J. E., 2005, *ApJS*, 157, 335
 Madau P., 1988, *ApJ*, 327, 116
 Madau P., 1995, *ApJ*, 441, 18
 Maraschi L. et al., 2008, *MNRAS*, 391, 1981
 Nandikotkur G., Jahoda K. M., Hartman R. C., Mukherjee R., Sreekumar P., Böttcher M., Sambruna R. M., Swank J. H., 2007, *ApJ*, 657, 706
 Pareschi G. et al., 2009, in Pareschi G. et al., eds, *Proc. SPIE Vol. 7437, Optics for EUV, X-Ray, and Gamma-Ray Astronomy IV*. SPIE, Bellingham, p. 743704
 Pizzella A., Corsini E. M., Dalla Bontà E., Sarzi M., Coccatto L., Bertola F., 2005, *ApJ*, 631, 785
 Poole T. S. et al., 2008, *MNRAS*, 383, 627
 Prochaska J. X., Worseck G., O’Meara J. M., 2009, *ApJ*, 705, L113
 Romani R. W., 2006, *AJ*, 132, 1959
 Roming P. W. A. et al., 2005, *Space Sci. Rev.*, 120, 95
 Sambruna R. M. et al., 2006, *ApJ*, 646, 23
 Sambruna R. M., Tavecchio F., Ghisellini G., Donato D., Holland S. T., Markwardt C. B., Tueller J., Mushotzky R. F., 2007, *ApJ*, 669, 884
 Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
 Shakura N. I., Sunyaev R. A., 1973, *A&A*, 24, 337
 Shapiro S. L., 2005, *ApJ*, 620, 59
 Shen Y., Greene J. E., Strauss M. A., Richards G. T., Schneider D. P., 2008, *ApJ*, 680, 169
 Sheth R. K., Tormen G., 1999, *MNRAS*, 308, 119
 Stickel M., Kühr H., Fried J. W., 1993, *A&AS*, 97, 483
 Szuszkiewicz E., Malkan M. A., Abramowicz M. A., 1996, *ApJ*, 458, 474
 Tavecchio F., Maraschi L., Ghisellini G., Kataoka J., Foschini L., Sambruna R. M., Tagliaferri G., 2007, *ApJ*, 665, 980
 Thompson D. J. et al., 1993, *ApJ*, 415, L13
 Tueller J. et al., 2010, *ApJS*, 186, 378
 Volonteri M., Rees M. J., 2005, *ApJ*, 633, 624
 Volonteri M., Rees M. J., 2006, *ApJ*, 650, 669
 Watanabe S. et al., 2009, *ApJ*, 694, 294
 Worsley M. A., Fabian A. C., Turner A. K., Celotti A., Iwasawa K., 2004a, *MNRAS*, 350, 207
 Worsley M. A., Fabian A. C., Celotti A., Iwasawa K., 2004b, *MNRAS*, 350, L67
 Yuan W., Fabian A. C., Celotti A., McMahon R. G., 2005, *MNRAS*, 358, 432
 Yuan W., Fabian A. C., Worsley M. A., McMahon R. G., 2006, *MNRAS*, 368, 985

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.