

## Checklist of the species in the subfamily Vespinae (Insecta: Hymenoptera: Vespidae)

James M. Carpenter<sup>1</sup> and Jun-ichi Kojima<sup>2</sup>

<sup>1</sup>Department of Entomology, American Museum of Natural History,  
Central Park West at 79<sup>th</sup> Street, New York, NY 10024, U. S. A.

<sup>2</sup>Natural History Laboratory, Faculty of Science, Ibaraki University, Mito 310, Japan

**Abstract** A checklist of the species in the subfamily Vespinae is presented, including synonyms and distributional summaries. Sixty-seven extant species in four genera are treated as valid, with an additional ten fossil species listed. No subspecies are recognized. Lamarck (1801) did not designate *Vespa crabro* Linnaeus as the type species of the genus *Vespa*. However, Latreille's (1810) designation of the same species is the first valid one, therefore existing generic nomenclature need not be disturbed. The following nomenclatural change is made: *Vespa variabilis* du Buysson, 1905, is a junior primary homonym of *Vespa variabilis* Fabricius, 1781. It is replaced by *Vespa variabilis fumida* van der Vecht, 1959, which is elevated to species rank, NEW STATUS. The lectotype of *Vespa annulata* Smith, 1858, is designated.

**Key words** Hymenoptera, Vespidae, Vespinae, *Vespa*, checklist

### Introduction

In recent years, several checklists of hornets and yellowjackets (Vespinae) have been published (Edwards, 1980; Matsuura and Yamane, 1984, 1990; Archer, 1989). Their usefulness as taxonomic tools is quite limited however. Matsuura and Yamane's and Archer's lists include summaries of distributional data, but otherwise all four lists consist of no more than names, authors and the year of publication. The original publications are not cited at all, the minimal requirement for utility in taxonomic research. Users of these lists must canvass the literature on their own to access the data contained in the original publications. In addition, all four lists perpetuate some nomenclatural errors.

The present checklist provides original references, and moreover includes information on types, namely, sex, type locality, and type depository. Preparation of this list also revealed some problems in the nomenclature of Vespinae, which are rectified here.

The status given each taxon generally follows Archer (1989, 1991a), but without recognition of any subspecies. Archer has studied the entire subfamily worldwide, and he (1991b, 1992, 1994b, 1995, 1997) is in the process of synonymizing all subspecies in the genus *Vespa* (Zalat, 1992, has already done so for some subspecies of *orientalis*). We concur: subspecies have no place in a phylogenetic system, rather, diagnosable taxa are to be treated as species (Nelson and Platnick, 1981; Nixon and Wheeler, 1990). We accept Archer's work as a basis for extending the synonymy to the remaining subspecies here, namely sinking all subspecies under the nominate species. There is an exception to Archer's classification. We follow Yamane and Tano (1985) in treating *Vespula arisana* as a valid species, rather than Archer's (1982) treatment as a subspecies of *orbata*, which would lead to synonymy here. Yamane and Tano (1985) illustrated differences in the male genitalia between *arisana* and *orbata*. Their sample was not large (one male specimen of *arisana*), but that is irrelevant until intergrades are shown to exist.

We have followed Archer in other synonymies for which there is variance in recent usage. First, Matsuura and Yamane (1984, 1990) and Yamane and Tano (1985) recognized *Vespa schrenckii* as a valid species, in contrast to Archer's (1981) treatment as a subspecies of *V. rufa*. Yamane et al. (1980: 34, footnote) suggested that two species occurred in northeastern Asia, and Matsuura and Yamane (1984: 305 and 1990: 250) argued that because *rufa* and *schrenckii* coexist, two species are indicated. Overlap while remaining distinct might indeed indicate phylogenetic species, however Matsuura and Yamane's discussion indicated that instead the forms do not remain distinct, rather the color characters by which they are distinguished intergrade. The forms are therefore treated as synonyms.

Second, Eck (1993) continued to recognize *Dolichovespula loekenae* as a species (which she described), but Pekkarinen (1995) has shown that it intergrades with *D. norvegicoides* and *pacifica*. Archer (1989) treated *loekenae* as a subspecies of *pacifica*, while Matsuura and Yamane (1984, 1990) and Pekkarinen (1995) treated both *loekenae* and *pacifica* as subspecies of *norvegicoides*. (Note also that Pekkarinen (1995: 66) confirmed a diagnostic difference, the elevation between gena and clypeus, which Archer (1989) had used to distinguish the two species, *norvegicoides* and *pacifica* (including *loekenae*).) Pekkarinen (1995: 68) cited allopatry as a reason for regarding them as subspecies. This is similar to the argument by Starr (1992) and Martin (1993) for recognition of *Vespa wilemani* as a species, merely on the basis that that is geographically segregated from *V. vivax*. Extrinsic features such as distribution are scarcely characters, diagnostic in the phylogenetic sense (Nixon and Wheeler, 1990; Carpenter, 1997).

It may be noted that the synonymy of subspecies leads to recognition of five Holarctic species (*Dolichovespula adulterina* and *D. norwegica*, and *Vespa austriaca*, *V. rufa* and *V. vulgaris*), aside from two other adventive species (*Vespa crabro* and *Vespa germanica*). The Palearctic and Nearctic representatives of the Holarctic species were first described as separate species, but their identity was recognized by Bequaert (1932). He treated some of these species as varieties of Holarctic species, but recognized none for *austriaca* and *vulgaris*. Ever since, these latter two taxa have been treated as Holarctic species. Miller (1958, 1961) raised the nearctic varieties of the other species to species rank. Subsequent study has confirmed this status for *Vespa acadica*, *V. atropilosa*, *V. consobrina* and *V. vidua*, and *Dolichovespula norvegicoides* (Wagner, 1978; Archer, 1989). But *D. albida* and *D. arctica*, and *V. intermedia* have come to be treated as subspecies, of *D. norwegica* and *D. adulterina*, and *V. rufa*, respectively (Eck 1981, 1984; Archer, 1981, 1989; Pekkarinen, 1995). There are no fixed differences, even of color, which might serve to diagnose phylogenetic species (Archer, 1981; Matsuura and Yamane, 1984 and 1990; Pekkarinen, 1995). The Nearctic and Palearctic taxa are therefore synonymized here. Note also that the distinction between *D. maculata* and *D. media* merits further study (Bequaert, 1932: 113; Eck, 1988), although there appear to be fixed differences in color.

The arrangement of the checklist is alphabetical. Each species is followed by the original citation, and then synonyms, which are listed chronologically. Each publication is cited in abbreviated form, to facilitate ready reference. Where two dates are listed, the first is the true date of publication, so far as can be determined, with the date following in parentheses being the date printed on the publication. The original locality is quoted for each description, and the type depository is then listed in parentheses. Museums or institutions where the types are kept are as follows: the code word designating the museum or institution in the present checklist is the city containing the collection.

Beijing: Institute Zoology Academia Sinica

Bergen: Zoologisk Museum, Universitetet I Bergen  
 Berlin: Zoologisches Museum der Humboldt–Universität  
 Budapest: Hungarian Natural History Museum  
 Calcutta: Zoological Survey of India  
 Cambridge: Museum of Comparative Zoology, Harvard University  
 Dresden: Staatliches Museum für Tierkunde  
 Edinburgh: Royal Scottish Museum  
 Fukuoka: Department of Entomology, Faculty of Agriculture, Kyushu University  
 Gainesville: American Entomological Institute  
 Genève: Muséum d'Histoire Naturelle  
 Genova: Museo Civico di Storia Naturale  
 Hangzhou: Zhejian Museum of Natural History  
 Kagoshima: Faculty of Science, Kagoshima University  
 Kobenhavn: Universitets Zoologisk Museum  
 Leiden: Nationaal Natuurhistorisch Museum  
 Linnean Society, London  
 London: The Natural History Museum  
 Lund: Museum of Zoology, Lund University  
 Marburg: Philipp-Universität Marburg, Zoologisches Institut  
 München: Zoologisches Sammlung des Bayerischen Staates  
 Ottawa: Biosystematics Research Institute  
 Oxford: Hope Entomological Collection, University Museum  
 Paris: Laboratoire d'Entomologie, Muséum National d'Histoire Naturelle  
 Philadelphia: Department of Entomology, Philadelphia Academy of Natural Sciences  
 Sapporo: Systematic Entomology, Faculty of Agriculture, Hokkaido University  
 Shinjuku, Tokyo: National Science Museum  
 Seoul: Korea University  
 St. Petersburg: Zoological Institute, Russian Academy of Sciences  
 Stockholm: Naturhistoriska Riksmuseet (Entomologiska)  
 Taichung: Taiwan Agricultural Research Institute  
 Torino: Instituto e Museo di Zoologia, Università di Torino  
 Tsukuba: National Institute of Agro-environmental Studies  
 Uppsala: Entomology Department, Zoological Museum, Uppsala University  
 Washington: U. S. National Museum of Natural History  
 Wien: Zoologische Sammlung, Naturhistorisches Museum

In the final paragraph for each species, under Distribution:, distributional data are summarized by country, where the states in the U. S. A. and provinces in Canada are abbreviated as follows.

AK: Alaska	HI: Hawaii	MI: Michigan
AL: Alabama	IA: Iowa	MN: Minnesota
AR: Arkansas	ID: Idaho	MO: Missouri
AZ: Arizona	IL: Illinois	MS: Mississippi
CA: California	IN: Indiana	MT: Montana
CO: Colorado	KS: Kansas	NC: North Carolina
CT: Connecticut	KY: Kentucky	ND: North Dakota
DC: Washington, DC	LA: Louisiana	NE: Nebraska
DE: Delaware	MA: Massachusetts	NH: New Hampshire
FL: Florida	MD: Maryland	NJ: New Jersey
GA: Georgia	ME: Maine	NM: New Mexico

NV: Nevada	TX: Texas	B. C.: British Columbia
NY: New York	UT: Utah	Man.: Manitoba
OH: Ohio	VA: Virginia	N. B.: New Brunswick
OK: Oklahoma	VT: Vermont	N. W. T.: North West Territories
OR: Oregon	WA: Washington	Ont.: Ontario
PA: Pennsylvania	WI: Wisconsin	P. E. I.: Prince Edward Island
RI: Rhode Island	WV: West Virginia	Que.: Quebec
SC: South Carolina	WY: Wyoming	Sask.: Saskatchewan
SD: South Dakota		
TN: Tennessee	Alta.: Alberta	

Varvio-Aho et al. (1984) published phenetic and cladistic analyses of an allozyme data set for eight species of European Vespinae, and Schmitz and Moritz (1990) published a phenogram based on an mtDNA restriction fragment size polymorphism data set for six species of European Vespinae. Carpenter (1987a, 1992) showed that the phylogenetic conclusions of neither study were actually supported by the data sets, which were demonstrated to be quite ambiguous.

Phylogenetic relationships among the genera are: *Vespa* + (*Provespa* + (*Dolichovespula* + *Vespula*)) (Carpenter, 1987b). Carpenter (1987b) synonymized all subgenera, and also discussed support for species groups in *Dolichovespula* and *Vespula*. Archer (1994a) provided a cladistic analysis of the species within the genus *Vespa*.

Before proceeding further, we must take up the troubling matter of the type species of the genus *Vespa*.

### What is the type species of the genus *Vespa*?

For two centuries, the type species of *Vespa* has usually been taken to be *Vespa crabro*, one of 17 originally included species, and the first listed (Linnaeus, 1758). The initial subdivision of the genus is largely due to Latreille (1802), who separated such genera as *Polistes*, *Eumenes*, *Synagris* and *Odynerus* (all Vespidae in the present sense). Among presently recognized species, that work left three of the originally included species in *Vespa: crabro*, *rufa* and *vulgaris*. Latreille (1802: 364) cited only *vulgaris* as "Exemple" of *Vespa*. But he later (1810: 438) cited *crabro* as "Type." This type designation was generally followed. Thus, in the first subdivision of *Vespa* in the restricted sense (= Vespinae in the present sense), Thomson (1869) included only one species, *crabro*, in the subgenus *Vespa*. He included eight species in his subgenus *Vespula*, among them *rufa* and *vulgaris*.

However, in this century, most workers have followed Bequaert (1930), who considered Lamarck's (1801: 271) earlier citation of *crabro* to have designated the type species of *Vespa*. A few previous workers (Bingham, 1897; Ashmead, 1902) had accepted Latreille's (1802) citation as designating *vulgaris* as the type species. This disagreement presented no particular difficulties, until Thomson's subgenus came to be recognized as a genus (Ashmead, 1902) and was subdivided in turn (Rohwer, 1916), with those subdivisions themselves being raised to generic status (Blüthgen, 1938). Bischoff (1931b) accepted Latreille's citation of *vulgaris* as the valid type designation, with the consequence that *Vespula* is a synonym of *Vespa* (although Bischoff did not draw that conclusion, erroneously refusing to accept Ashmead's (1902) designation of *austriaca* as type species of *Vespula*). It then followed that *Macrovespa* Dalla Torre, 1904, is the available name for the genus including *crabro*, while *Vespa* applies to Thomson's *Vespula*.

Bequaert (1932: 75–76) decried Bischoff's classification, stating:

"The divergence of opinion in this case seems to show once more the utter futility of hoping for stability if one follows an intricate system of rules, recommendations and opinions, which invariably leads to mere casuistry."

Bequaert rejected Bischoff's classification, and in this he has been followed ever since by vespid taxonomists.

However, the "intricate system of rules, recommendations and opinions" to which Bequaert referred was Opinion 79 of the International Commission on Zoological Nomenclature (1924), which Bischoff had followed. That Opinion specifically stated that Lamarck's (1801) work "is not to be accepted as designation of type species" (and see Direction 32, International Commission on Zoological Nomenclature, 1956). Bequaert (1932) thus contravened the Commission, and thereby, the present International Code of Zoological Nomenclature (Article 78). And this fact has not occasioned comment from subsequent vespid taxonomists, most of whom have followed Bequaert, including the authors of the most recent generic catalog (van der Vecht and Carpenter, 1990).

Bequaert (1930: 61) characterized the consequences of accepting Latreille's (1802) citation of *vulgaris* as "disturbing." Indeed. But Bequaert's course will not do for those who follow the Rules. Lamarck's citation must be set aside.

An earlier subdivision of *Vespa* than that of Latreille (1802) exists, but has been ignored, namely that of Christ (1791). Christ (1791: 205) listed *Vespa* as the name for "Die Wespen." That author then proposed (1791: 212) a division of *Vespa* into two genera, which he called *Vespa crabro*, for "Hornisse," and *Vespa*, for "Gemeine Wespen." On the same page, he stated "Die Hornisse. V. Crabro." He then proceeded to describe hornets, and stated (p. 213) "Das Wort *Crabro* leget Linne der Hornisse bei," leaving no doubt he was referring to the species *Vespa crabro* Linnnaeus. It was the only species listed in this section. The designation "Die Hornisse. V. Crabro" was repeated on p. 214, under the heading "Beischreibung der Arten." Christ included 16 taxa within "*Vespa crabro*," all with trinomina. Christ intended his *Vespa crabro* in the near-universal sense followed since, as meaning Hornisse (hornets in English, frelon in French). Thus, Christ's repeated citation of *Vespa crabro* as Hornisse appears a clearer "type designation" of *Vespa* in the sense of Thomson and auct. (including Latreille), than the oft-disputed "examples" cited by the systematists of a decade later than Christ. That unambiguous distinction was followed by Thomson when he subdivided *Vespa*. Interpreting this distinction as a type designation would allow existing nomenclature to remain undisturbed.

We prefer however an alternative course. Arguments about whether the citations by Christ (1791), Lamarck (1801), and Latreille (1802) are valid are obviated by rejecting all of them. The first valid designation of the type species of *Vespa* in the scope of the Code is that by Latreille (1810: 438), as suggested by Blüthgen (1938: 289) and discussed below. Generic nomenclature is not disturbed, as *crabro* remains the type species of *Vespa*.

Christ's (1791) "*Vespa crabro*" was a genus in his terminology. The Code is of course quite clear that nomenclature above the species level is uninominal (Article 4). Christ's newly proposed names for taxa within his *Vespa crabro* have nevertheless universally been considered as available, as species level trinomina (see, e.g., *Vespa crabro germana*). Interpreting Christ's trinomina as subspecies is evidently unexceptionable. But however Christ intended his *Vespa crabro*, his citation could thus be interpreted as meaning no more than the designation of a type of "Hornisse" – not the type of the genus *Vespa*.

The Code is also quite clear that mere mentioning of a species as an example of a genus is not a designation (Article 67(c)). And, as alluded to above, Opinion 79 specifically stated that

Lamarck (1801) cited “a known species, or rarely several” as examples. Opinion 79 therefore excluded Lamarck’s listings as valid type designation.

In Latreille (1802), the description of each genus (or subdivision within a genus) is followed by citation of one or a few species as “Exemple.” It is clear that such mention of a species as an example is not a valid designation of a type species under the Code. Therefore, Latreille’s citation of *vulgaris* is an example, and not designation of a type species.

It should be mentioned that Bequaert (1930) also stated that Latreille (1804: 181) designated *crabro* as type species of *Vespa*. This is an error; on the cited page, Latreille in fact mentioned no species under *Vespa*.

In contrast to all of the foregoing works, in 1810 Latreille cited species in the genera he listed under the heading “Table des genres avec l’indication de l’espèce qui leur sert de type.” For *Vespa* he cited only *crabro*. And Opinions 11 (1910) and 136 (1939) of the International Commission on Zoological Nomenclature have specifically ruled this particular list as validly designating type species, where no previous valid designation existed, and where Latreille cited only one (originally included) species. Those conditions are fulfilled in this case. *Vespa crabro* is therefore (and thankfully) really the type species of the genus *Vespa*.

There do not seem to be any grounds for disputing this conclusion, but if for some reason other workers were to disagree, an appeal should be made to the Commission for use of plenary powers, either explicitly to rule Christ’s (1791) citation as a binding type designation, to modify Opinion 79 to accept Lamarck’s (1801) citation as a type designation, or to set aside Latreille’s (1802) citation in favor of Latreille (1810). The present generic nomenclature has been stable for more than sixty years as regards the interpretation of *Vespa*, and with minor exceptions, for more than a century (and indeed longer, as shown here). However accomplished, this stability should be maintained.

### Designations of lectotypes

In order to avoid future taxonomic confusion, we here designate the lectotype of *Vespa annulata* Smith, 1859, which is in Oxford and labelled as the lectotype by C. O’Toole; that designation has never been published. The specimen is labelled “SAR” (circular label), “*Vespa annulata* Smith”, “Lectotype *Vespa annulata* Smith, 1859, Design. C. O’Toole, 1993”. Another female in Oxford with the same locality label is the paralectotype.

In his 1957 and 1959 work, van der Vecht designated the lectotypes of 13 taxa previously described in *Vespa* (*Vespa barthelemyi* du Buysson, 1905, is now placed in *Provespa*). Besides those properly designated as the lectotypes, he referred to type specimens of the following taxa as “the type”; *Vespa parallela* var. *biroi* du Buysson, 1905 (in Paris; 1957: 13), *Vespa rubricans* Pérez, 1910 (in Paris; 1957: 24), and *Vespa mongolica* var. *divergens* Pérez, 1910 (in Paris; 1957: 38). Du Buysson (1905: 513) listed Budapest and London as the depository of the types of *V. parallela* var. *biroi*, but judging from the fact that the label data given by van der Vecht agree with those in the original description, a specimen from the type series may have kept by du Buysson and deposited in Paris. Thus, van der Vecht’s statement inadvertently referring it as “the type” is considered as the designation of the lectotype (under Article 74(b) in the Code).

Pérez (1910: 10) provided a range of the body length for *Vespa rubricans*, suggesting that his description was based on more than one specimen. However, he did not give any statement allowing the inference of the number of specimens he examined when he described

*Vespa mongolica* var. *divergens*. Subsequently, van der Vecht statements are also considered as the designations of the lectotypes of both taxa.

Moreover, van der Vecht's two articles include ambiguous statements with respect to certain lectotype designations. He (1957: 41) mentioned "*Vespa velutina celebensis* Pérez . . . Bua Kraëng . . . (type in coll. Pérez, MP)". Judging from the description by Pérez (1910: 12), the taxon was based on a single female; thus the specimen is the holotype.

In the list of synonyms of *Vespa multimaculata multimaculata*, van der Vecht (1957: 44) stated "*V. multimaculata*; "Brunei, Java"; type in MP) (Brunei is in North Borneo! v.d.V.)", and in the list of the specimens he examined, "Brunei (type in coll. Pérez, MP)". However, he (1957: 44) mentioned, in the discussion of synonymy, "the type in the collection Pérez (MP) bears a label "Brunei, Borneo". Pérez (1910: 14), providing a range of the body length, may have described the species based on more than one specimen; van der Vecht's mention of "the type" is here regarded as the designation of the lectotype.

Van der Vecht (1959: 214) gave the label data of the type of *Vespa alduini* Guérin-Méneville, 1831, and said "(type, MCG)". The description (figure) of *Vespa alduini* was very probably based on a single specimen, and that is the holotype.

Finally, concerning *Vespa oberthuri* du Buysson, 1902, van der Vecht (1959: 218) stated "*Vespa crabro oberthuri* . . . 3♀ Se-Tchouen, Sio-Lou, leg. R. Oberthür (type and paratypes, MP); 2♀ Ta-Tsien-Lou (paratypes, MP, ML)". All of these five females are apparently syntypes. Under the Code, van der Vecht's statement cannot be regarded as the designation of the lectotype. We have not seen the specimens that van der Vecht examined, and therefore leave the designation of the lectotype of this taxon for future study.

### Checklist of Vespinae (Vespariae Latreille, 1802)

#### Genus *Dolichovespula* Rohwer

*Dolichovespula* Rohwer, 1916, Conn. State Geol. Nat. Hist. Surv. Bull. 22: 642, subgenus of *Vespula* Thomson.

Type species: *Vespa maculata* Linnaeus, 1763, by original designation.

*Pseudovespula* Bischoff, 1931 (1930), Sitz.-Ber. Ges. Naturf. Freunde Berl.: 346, subgenus of *Pseudovespa* Schmiedeknecht.

Type species: "*Ps. adulterina* Buyss." [= *Vespa norwegica* var. *adulterina* du Buysson, 1905], by original designation.

*Boreovespula* Blüthgen, 1943, Stettin. Ent. Ztg. 104: 149, subgenus of *Dolichovespula* Rohwer.

Type species: "*Dolichovespula norwegica* (F.)" [= *Vespa norwegica* Fabricius, 1781], by original designation.

*Metavespula* Blüthgen, 1943, Stettin. Ent. Ztg. 104: 149, subgenus of *Dolichovespula* Rohwer.

Type species: "*Dolichovespula silvestris* (Scop.)" [!= *Vespa sylvestris* Scopoli, 1763], by original designation.

#### *adulterina* (du Buysson)

*Vespa borealis* Lewis, 1897, Trans. Am. Ent. Soc. 24: 171 (key), 174, female, male –

"United States, New York, Mt. Washington, British Columbia; Amherst, Mass." (Philadelphia?). Junior primary homonym of *Vespa borealis* Kirby, 1837, *Vespa*

*borealis* Zetterstedt, 1838, *Vespa borealis* Smith, 1843, and of *Vespa crabro* var. *borealis* Radoszkowski, 1863.

*Vespa norwegica* var. *adulterina* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73 (3): 499 (key), 600, 628, female – “Europe: Piémont . . . ; Vosges, Remiremont . . . ; Suisse, Engadine, Rosegethal . . . Amérique; Orégon, Corvallis” “Mongolie . . . Ile Sakhaline. Village de Lasevo, près de la station de Viterbsk, Russie” (Paris, München, Washington, St. Petersburg).

*Vespula arctica* Rohwer, 1916, Conn. State Geol. Nat. Hist. Surv. Bull. 22: 642 (in subgenus *Dolichovespula*). Replacement name for *Vespa borealis* Lewis.

*Vespa norwegica saxonica* natio *colchica* Birula, 1930, Ann. Mus. Zool. Acad. Sci. U. R. S. S. 31 (2): 314, male – [Georgia] “Adsharien (Batum-Gebiet, Alpenwiesen, s. g. Jailag, bei Sarytschair auf der Höhe von etwa 2300 m ü. d. M . . . )” (St. Petersburg). Unavailable under Articles 45e and 45f(iii) of the International Code of Zoological Nomenclature.

*Dolichovespula adulterina montivaga* Yamane, 1982, Trans. Shikoku Ent. Soc. 16: 109, female, male – “Mt. Kariba (1500 m alt.), Hokkaido, Japan” (holotype female Kagoshima); also from Honshu.

Distribution: Norway; Sweden; Finland; Belgium; France; Switzerland; Italy; Poland; Russia: Kola Peninsula, Caucasus to southwestern Siberia, Primor’ye, Sakhalin; Turkey; Georgia; Mongolia; Taiwan; Japan; Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, NV, ID, MT, WY, UT, CO, AZ, NM, ND, MN, WI, IL, MI, IN, KY, TN, OH, NY, PA, VT, NH, ME, MA, CT, RI, NJ, DE, MD, WV, VA, NC, GA.

#### *alpicola* Eck

*Dolichovespula alpicola* “Wagner” Eck, 1984, Ent. Abhand. Staat. Mus. Tierk. Dresden 48 (4): 40 (key), fig. 3L, female – “westliches Nordamerika”; 1987, Ent. Abhand. Staat. Mus. Tierk. Dresden 50 (10): 191 (lectotype female from “WYO Albany County, Medicine Bow Nat. For.”, Dresden).

Distribution: Canada: Yukon Terr., B. C., Alta.; U. S. A.: AK, WA, OR, MT, WY, CO, UT, NM, AZ.

#### *arenaria* (Fabricius)

*Vespa arenaria* Fabricius, 1775, Syst. Ent.: 365 – “in America arenosis” (London).

*Vespa borealis* Kirby, 1837, Fauna Bor.-Am. 4, Ins.: 264 – [Canada] “Lat. 65°” (type depository unknown).

*Vespa Diabolica* de Saussure, 1854, Ét. Fam. Vespi. 2: 138, female – [U. S. A.] “L’Amérique du Nord, Philadelphie” (Torino).

*Vespa fernaldi* Lewis, 1897, Trans. Am. Ent. Soc. 24: 171 (key), 173, female – [U. S. A.] “Colorado” (Philadelphia?).

Distribution: Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, NV, ID, MT, WY, UT, CO, AZ, NM, NE, ND, SD, MN, NE, IA, WI, IL, MI, IN, KY, TN, OH, NY, PA, VT, NH, ME, MA, CT, RI, NJ, DE, MD, WV, VA, NC, GA.

#### *asiatica* Archer

*Vespa sylvestris* var. *intermedia* Birula, 1930, Ann. Mus. Zool. Acad. Sc. U. R. S. S. 31 (2): 330, 331, female, male – [Uzbekistan] “Zeravschan-Tal und der östlichen

Buchara" (lectotype female from "Ost-Buchara . . . im Gebirgsrücken Peter des Grossen . . . Engpass Kamtschirák" in St. Petersburg). Junior primary homonym of *Vespa rufa* var. *intermedia* du Buysson, 1905. – Archer, 1981, Ent. Scand. 12: 189 (designation of lectotype).

*Dolichovespula asiatica* Archer, 1981, Ent. Scand. 12: 187, 189. Replacement name for *Vespula sylvestris* var. *intermedia* Birula.

Distribution: Kazakhstan; Uzbekistan; Turkmenistan; Afghanistan; Pakistan; India: Kashmir; Mongolia; China: Xinjiang, Gansu.

#### *baileyi* Archer

*Dolichovespula baileyi* Archer, 1987, Ent. Mon. Mag. 123: 30, female – "CHINA, Szechwan, Lin Yang, 2500 m" (Edinburgh); also from another locality in Sichuan.

Distribution: China: Sichuan.

#### *flora* Archer

*Dolichovespula flora* Archer, 1987, Ent. Mon. Mag. 123: 27, female – "BURMA, Adung Valley" (London); also from China.

Distribution: Myanmar; China: Sichuan, Yunnan.

#### *kuami* Kim and Yoon

*Dolichovespula kuami* Kim and Yoon, 1996, Korean J. Syst. Zool. 12(3): 199, figs. 1, 2, female – "Korea: Chungcheongbuk-do Province, Sokrisan" (Seoul); also from Kyeonggi-do Prov.

Distribution: South Korea.

#### *lama* (du Buysson)

*Vespa lama* du Buysson, 1903, Bull. Soc. Ent. France 1903: 175, female – [India] "Sikkim" (holotype Paris).

*Dolichovespula nyalamensis* Lee, 1986, Sinozool. 10 (4): 195, 197, figs. 1(4), 2(2), female – {China} "Nelam, Xizang (Tibet), 3300 m" (Beijing).

Distribution: India: Uttar Pradesh, Sikkim; China: Xizang.

#### *maculata* (Linnaeus)

*Vespa maculata* Linnaeus, 1763, Amoenit. Acad. 6: 412 – [U. S. A.] "Pennsylvania" (lectotype female Stockholm). – Day, 1979, Biol. J. Linn. Soc. 12: 66 (designation of lectotype).

*Vespa maculata americana* Christ, 1791, Naturgesch. Ins.: 239 – "mitternachtlichen Amerika" (destroyed). Junior primary homonym of *Vespa americana* Fabricius, 1775.

Distribution: Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, NV, ID, MT, WY, UT, CO, AZ, NM, ND, SD, MN, NE, IA, KS, MO, OK, AR, TX, LA, AR, MO, WI, IL, MI, IN, OH, NY, PA, VT, NH, ME, MA, CT, RI, NJ, DE, MD, KY, WV, VA, TN, NC, SC, MS, AL, GA, FL.

#### *media* (Retzius)

*Vespa Crabro medius* Retzius, 1783, C. Degeer Gen. Spec. Ins.: 63 (type depository unknown).

- Vespa Geerii* Lepeletier, 1836, Hist. Nat. Ins. Hym. 1: 510, female, male – “La Suède et les environs de Paris . . . Versailles . . . dans la forêt de Saint-Germain-en-Laye” (type depository unknown).
- Vespa crassa* Herrich-Schäffer, 1841, Fauna Ins. Germ. 179: 33, fig. 179.4, male (destroyed).
- Vespa similis* Schenck, 1853, Jahrb. Ver. Naturk. Nassau 9 (1): 15 (key), 24, female, male – [Germany] “Weilburg” (Marburg?).
- Vespa rufoscutellata* Schenck, 1853, Jahrb. Ver. Naturk. Nassau 9 (1): 15 (key), 28, male – [Germany] “Wiesbaden” (Marburg?).
- Vespa flavicincta* Schenck, 1853, Jahrb. Ver. Naturk. Nassau 9 (1): 16 (key), 27, female – [Germany] “Weilburg” (Marburg?). Junior primary homonym of *Vespa flavicincta* Gravenhorst, 1807.
- Vespa Geeri*; Dalla Torre, 1904, Gen. Ins. 19: 65. Incorrect spelling of *Vespa geerii* Lepeletier.
- Vespa media* var. *lineolata* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 17, female – [France] “Millau” (Paris).
- Vespula media media* var. *lugubris* Birula, 1927, Ann. Mus. Zool. Acad. Sci. U. R. S. S. 28: 80, female – “Nord- und Mittelteilen Ost-Europas . . . Norden” (St. Petersburg). Unavailable under Articles 45e and 45f(iii) of the International Code of Zoological Nomenclature.
- Dolichovespula media* var. *conjugens* Paul, 1943, Stettin. Ent. Ztg. 104: 142 – [Poland] “Stettin: Eckerberger Wald” (type depository unknown).
- Dolichovespula media sugare* Ishikawa, 1969, Bull. Nat. Sci. Mus. Tokyo 12 (2): 180, figs. 1–3, female, male – “Japan (Honshu) . . . Tsuta, Towada, Aomori Pref.” (holotype female Shinjuku, Tokyo); also from numerous other localities on Honshu.
- Dolichovespula borealis* Lee, 1986, Sinozool. 10 (4): 195, 196, figs. 1(2), 2(3), female – [China] “Hengdaohezi, Heilongjiang province” (Beijing). Junior secondary homonym of *Vespa borealis* Kirby, 1837, *Vespa borealis* Zetterstedt, 1838, and *Vespa borealis* Lewis, 1897.

Distribution: Norway; Sweden; Finland; Denmark; France; Spain; Germany; Austria; Italy; Poland; Bulgaria; across Russia to the Sea of Okhotsk, Lake Baikal, southwestern Siberia, Sakhalin; Mongolia; China: Heilongjiang, Liaoning; Korea; Japan. Occasionally U. K.: England.

#### *norvegicoides* (Sladen)

-  *Vespa norvegicoides* Sladen, 1918, Ottawa Nat. 32: 71, male, female – [Canada] “N.S. to B.C.” (lectotype female from “Amherst, Nova Scotia” in Ottawa). – Miller, 1961, Can. Ent., Suppl. 22: 36, 38 (designation of lectotype).

Distribution: Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, ID, MT, WY, ND, MN, WI, MI, KY, TN, NY, PA, VT, NH, ME, MA, CT, RI, NJ, MD, WV, VA, NC, SC.

#### *norwegica* (Fabricius)

-  *Vespa norwegica* Fabricius, 1781, Spec. Ins. 1: 460 – “in Norwegia” (Kobenhavn).  
*Vespa britannica* Leach, 1814, Zool. Misc. 1: 112, pl. 50, female, male – “in Britannia; in Anglia rarius, in Caledonia vulgatissime” (London).

- Vespa marginata* Kirby, 1837, Fauna Bor.-Am. 4, Ins.: 265, pl. VI fig. 2, female – [Canada] “Taken in the Route from New York, and again in Lat. 65°” (type depository unknown). Junior primary homonym of *Vespa marginata* Gmelin, 1790.
- Vespa borealis* Zetterstedt, 1838, Ins. Lappon. Descr.: 454, female – “Lapponia” (Lund?). Junior primary homonym of *Vespa borealis* Kirby, 1837.
- Vespa norvegica* Olivier, 1792, Encycl. Méthod. Ins. 6: 680; Smith, 1858, Cat. Brit. Fossor. Hym., Form. Vesp.: 220. Unjustified emendation of *Vespa norwegica* Fabricius.
- ? *Vespa peruana* de Saussure, 1867, Reise Novara., Zool. 2, Hym.: 18, female – “Peru altior; Quito” [error] (Wien).
- Vespa albida* Sladen, 1918, Ottawa Nat. 32: 71, male, female – [U. S. A.] “Alaska” (lectotype male Ottawa). – Miller, 1961, Can. Ent., Suppl. 22: 28, 29 (designation of lectotype).
- Vespa saxonica* var. *arctica* Friese, 1919, Mém. Acad. Sci. Russ. (8) 28 (15): 1, female, male – [Russia] “Chanema-Fluss . . . Sobj-Tal . . . Longot-johan-Tal . . . Berezov” (St. Petersburg). Junior secondary homonym of *Vespula arctica* Rohwer, 1916.
- Dolichovespula norwegica* var. *Zetterstedti* Blüthgen, 1937, Mitt. Ent. Ges. Halle 15: 37. Replacement name for *Vespa saxonica* var. *arctica* Friese.

Distribution: Norway; Sweden; Finland; Denmark; Ireland; U. K.: Scotland, Wales, England; Belgium; France; Spain; Germany; Austria; Poland; Slovakia; Hungary; Bulgaria; across Russia to Sakhalin; Turkey; China: Xinjiang; Mongolia; Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, ME.

#### *omissa* (Bischoff)

- Vespa tripunctata* Schenck, 1861, Jahrb. Ver. Naturk. Nassau 16: 17 (key), 25, female – [Germany] “Bamberg” (München). Junior primary homonym of *Vespa tripunctata* Fabricius, 1787.

- Vespula norvegica saxonica* morpha *ingrica* Birula, 1930, Ann. Mus. Zool. Acad. Sci. U. R. S. S. 31 (2): 309, female – [Russia] “Leningrader Gouvernement” (lectotype from “Leningrad, Kaipolova”, St. Petersburg). Unavailable under Articles 45e and 45f(iii) of the International Code of Zoological Nomenclature. – Blüthgen, 1961. Abh. Dtsch. Akad. Wissenschaft. Berlin, Kl. Chem. geol. Biol. 2: 42 (designation of lectotype).

- Vespa omissa* Bischoff, 1931, Mitt. Deutsch. Ent. Ges. 2 (1): 6, female – [Austria] “Thüringen” (Berlin); also from Tirol.

- Vespula ingrica* (“Birula”) Bequaert, 1932, Ent. Am. n. ser. 12 (2): 89, footnote. Available under Articles 10c, 23j and 50c of the International Code of Zoological Nomenclature.

Distribution: Norway; Sweden; Finland; Denmark; Belgium; France; Spain; Germany; Switzerland; Austria; Italy; Poland; Czech Republic; Slovakia; Russia; Bulgaria; Turkey; Iran.

#### *pacifica* (Birula)

- Vespula norvegica pacifica* Birula, 1930, Ann. Mus. Zool. Acad. Sci. U. R. S. S. 31 (2): 295, 317, female, male – [Russia] “Im Küstengebiete des Pazifischen Ozeans und auf den naheliegenden Inseln” (lectotype female from “Insel Sachalin”, St. Petersburg). – Eck, 1984, Ent. Abh. Staat. Mus. Tierk. Dresden 47 (11): 195 (designation of lectotype).

*Dolichovespula loekenae* Eck, 1980, Reichenbachia 18 (30): 213, female, male – [Norway] “Espeland, HOY: Alversund” (Bergen); also from Sweden; Finland.

Distribution: Norway; Sweden; Finland; Russia: northwestern region, Ob' River to east of Lake Baikal, Primor'ye, Sakhalin; Korea; Japan.

**panda** Archer

*Dolichovespula* [!] *panda* Archer, 1981 (1980), Entomon 5 (4): 341, fig. 1, female [male = worker] – “CHINA, Szechuan, 14.4 km south-west Tarsienlu, 2600–4000 m” (Washington); also from other localities in Sichuan.

Distribution: China: Sichuan.

**saxonica** (Fabricius)

*Vespa saxonica* Fabricius, 1793, Ent. Syst. 2: 256 – “in Germania” (type depository unknown).

*Vespa bavarica* von Schrank, 1802, Fauna Boica 2 (2): 350, female – [Germany] “Waldungen” (type depository unknown).

*Vespa tridens* Schenck, 1853, Jahrb. Ver. Naturk. Nassau 9 (1): 18 (key), 38, female, male – [Germany] “Weilburg . . . in der Gegend von Wiesbaden” (Marburg?).

*Vespula norvegica saxonica* morpha *monticola* Birula, 1930, Ann. Mus. Zool. Acad. Sci. U. R. S. S. 31 (2): 313 – [Russia] “Gebirgen des Kaukasus” (St. Petersburg). Unavailable under Articles 45e and 45f(iii) of the International Code of Zoological Nomenclature.

*Dolichovespula saxonica nipponica* Yamane, 1975, Kontyû 43 (3): 349, female – [Japan] “HOKKAIDO. Otoineppu” (Sapporo); also from elsewhere in Hokkaido, and Honshu.

*Dolichovespula saxonica kamtschatkensis* Eck, 1983, Ent. Abhand. Staat. Mus. Tierk. Dresden 46 (8): 171, female, male – [Russia] “Petropawlowsk, Flußtal Poganka, Kamtschatka” (holotype female St. Petersburg); also from elsewhere in Kamchatka.

*Dolichovespula saxonica nigrescens* Eck, 1983, Ent. Abhand. Staat. Mus. Tierk. Dresden 46 (8): 172, female – [Russia] “Primorje, Umgeb. Anisimowka” (Dresden); also from elsewhere in Russia; Mongolia; Korea.

Distribution: Norway; Sweden; Finland; Denmark; Belgium; France; Spain; Germany; Switzerland; Austria; Poland; Slovakia; Bulgaria; across Russia to Kamchatka Peninsula, Kurile Is., Sakhalin; Turkey; Mongolia; China: Xinjiang, Gansu, Sichuan, Liaoning; Korea; Japan.

**stigma** Lee

*Dolichovespula stigma* Lee, 1986, Sinozool. 10 (4): 195, 198, figs. 1(3), 2(1), female – [China] “Mt. Emei, Sichuan Province, 1800–2100 m” (Beijing); also from 1800–1900 m.

*Dolichovespula sinensis* Archer, 1987, Ent. Mon. Mag. 123: 29, female – “CHINA, Szechwan, Suifu” (Washington); also from Xinjiang.

Distribution: China: Sichuan, Xinjiang.

**sylvestris** (Scopoli)

*Vespa sylvestris* Scopoli, 1763, Ent. Carniol.: 309 – [Slovenia] “Carnioliae” (destroyed?).

*Vespa parietum* Harris, 1776, Exposit. Eng. Ins.: 128, pl. XXXVII fig. 4, male – “English” (destroyed?). Junior primary homonym of *Vespa parietum* Linnaeus, 1758.

*Vespa holsatica* Fabricius, 1793, Ent. Syst. 2: 257 – [Germany] “Kiliae” (Kobenhavn).

*Vespa frontalis* Latreille, 1802, Ann. Mus. Hist. Nat. 1: 290, pl. XXI fig. 7 – [France] “Chartreaux de Paris” (type depository unknown).

*Vespa campanaria* Fowler, 1833, Ann. Mag. Nat. Hist. 6: 536, 538, fig. 69 (type depository unknown).

Note: Edwards (1980: 362) attributed *campanaria* to “(Shaw, 1803).”

*Vespa anglica* “Leach” Smith, 1843, Zoologist 1: 168, as a synonym of *Vespa holsatica* Fabricius, 1793. Available under Art. 11(e) of the International Code of Zoological Nomenclature.

*Vespa pilosella* Costa, 1858, Ric. Ent. Monti Partenii: 20, 28, female, male – [Italy] “Monti Partenii” (type depository unknown). Junior primary homonym of *Vespa pilosella* Gravenhorst, 1807.

*Vespa silvestris*; Dalla Torre, 1894, Cat. Hym. 9: 156. Incorrect spelling of *Vespa sylvestris* Scopoli.

*Vespa silvestris* [!] var. *sumptuosa* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73 (3): 493 (key), 603, female – “Perse centrale” (lectotype Wien). – Archer, 1981, Ent. Scand. 12: 191 (designation of lectotype, inadvertently referring it as “the holotype”).

*Dolichovespula xinjiangensis* Lee, 1986, Sinozool. 10 (4): 195, 199, figs. 1(1), 2(4), female, male – [China] “Xinyuan, Xinjiang Uygur Aut. Reg., 850–1200 m” (holotype female Beijing); also from Altay.

Note: Edwards (1980: 362) listed “*arbustorum* (Blanchard, 1845)” as a synonym of *sylvestris*. However, Blanchard (1845: 62) merely referred to the common name “*Guêpe des arbustes*” given by Réaumur. That name is of course not available.

Distribution: Norway; Sweden; Finland; Denmark; Ireland; U. K.: Scotland, Wales, England; Germany; Belgium; France; Spain; Morocco; Austria; Slovenia; Italy; Poland; Slovakia; Bulgaria; across Russia to Siberia, Primor’ye; Turkey; Armenia; Iran; Afghanistan; Pakistan; India: Kashmir; Mongolia; China: Xinjiang, Gansu, Fujian.

#### *xanthicincta* Archer

*Dolichovespula pacifica xanthicincta* Archer, 1981 (1980), Entomon 5 (4): 341, 343, female [male = female] – “CHINA-TIBET BORDER, near Tang Gu, 4300 m” (Washington); also from Xizang; Myanmar.

Distribution: China: Xizang; Bhutan; Myanmar.

#### Genus *Provespa* Ashmead

*Provespa* Ashmead, 1903, Ent. News 14: 182, genus.

Type species: *Vespa dorylloides* de Saussure, 1854 [= *Vespa anomala* de Saussure, 1854], by original designation.

#### *anomala* (de Saussure)

*Vespa anomala* de Saussure, 1854, Ét. Fam. Vesp. 2: 112, pl. XIV figs. 2, 2a, female – [Indonesia] “l’ile de Java” (Torino, London).

*Vespa dorylloides* de Saussure, 1854, Ét. Fam. Vesp. 2: 256, Errata. Unnecessary replacement name for *Vespa anomala* de Saussure.

*Vespa dorylloides*; du Buysson, 1905 (1904), Ann. Soc. Ent. France 73 (3): 497, 616. Incorrect spelling of *Vespa dorylloides* de Saussure.

Distribution: India: Uttar Pradesh, Sikkim; Myanmar; Thailand; Malaysia: Peninsular Malaysia, Sarawak; Singapore; Indonesia: Sumatra, Bangka, Batu Is., Java, Borneo (Kalimantan).

*barthelemyi* (du Buysson)

*Vespa Barthelemyi* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 492 (key), 618, pl. V figs. 3, 4, female, male – “Cambodge, Pnom-penh” (lectotype female Paris); also from India; Vietnam; Myanmar. – van der Vecht, 1957, Zool. Verh., Leiden 34: 10 (designation of lectotype).

Distribution: India: Sikkim, Assam; Bhutan; China: Yunnan, Guangxi; Myanmar; Thailand; Laos; Cambodia; Vietnam; Malaysia: Peninsular Malaysia.

*nocturna* van der Vecht

*Provespa nocturna* van der Vecht, 1935, Ent. Meded. Ned. Indië 1: 41, fig. 1, female, male – [Indonesia] “Sumatra, Palembangsche Bovenlanden” (holotype female Leiden); also from Borneo, ? Sulawesi.

Distribution: Vietnam; Malaysia: Peninsular Malaysia, Sarawak; Indonesia: Sumatra, Bangka, Borneo (Kalimantan).

Genus *Vespa* Linnaeus

*Vespa* Linnaeus, 1758, Syst. Nat., ed. 10, 1: 343, 572, genus (17 species).

Type species: “*Vespa crabro*, Fab.” [= *Vespa crabro* Linnaeus, 1758], by subsequent designation of Latreille, 1810, Consid. Gén. Crust. Arachn. Ins.: 438.

*Macrovespa* Dalla Torre, 1904, Gen. Ins. 19: 64, group of genus *Vespa* Linnaeus.

Type species: *Vespa crabro* Linnaeus, 1758, by subsequent designation of Bequaert, 1930, Bull. Brooklyn Ent. Soc. 25: 64.

*Nyctovespa* van der Vecht, 1959, Zool. Meded., Leiden 36 (13): 210, subgenus of *Vespa* Linnaeus.

Type species: *Vespa binghami* du Buysson, 1905, by original designation.

*affinis* (Linnaeus)

*Apis affinis* Linnaeus, 1764, Mus. Ludov. Ulricae : 417 – “in Calidis regionibus” (holotype female Uppsala).

*Vespa affinis* Fabricius, 1787, Mant. Ins. 1: 287 – “in China” (Kobenhavn). Junior secondary homonym of *Vespa affinis* (Linnaeus, 1764).

*Vespa unifasciata* Olivier, 1792, Encycl. Méthod. Ins. 6: 677 – “Indes orientales” (type depository unknown). Junior primary homonym of *Vespa unifasciata* Gmelin, 1790.

*Vespa Alduini* Guérin-Méneville, 1831, in Duperrey, Voyage Coquille, Zool., Ins. Hym. Atlas: pl. 9 fig. 6; 1838, Texte 2 (2): 264 – [Indonesia] “l’île de Bourou, l’une des Moluques” (holotype female Genova).

*Vespa bimaculata* Guérin-Méneville, 1838, in Duperrey, Voyage Coquille, Zool., Ins. Hym. Texte 2 (2): 264. Unnecessary replacement name for *Vespa alduini* Guérin-Méneville. Junior primary homonym of *Vespa bimaculata* Geoffroy, 1785, and of *Vespa bimaculata* Olivier, 1792.

*Vespa Nigripennis* de Saussure, 1854, Ét. Fam. Vesp. 2: 156 – “Les Philippines” (London). Junior primary homonym of *Vespa nigripennis* Degeer, 1773.

*Vespa cincta* var. *picea* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 488 (key), 537, female – “Nouvelle-Guinée, Tupuseleia . . . ; Ile Yule . . . ; N<sup>le</sup>-Guinée, Kapa-Kapa” (lectotype Genova). – van der Vecht, 1959, Zool. Meded., Leiden 36: 214 (designation of lectotype).

- Vespa Indosinensis* Pérez, 1910, Act. Soc. Linn. Bordeaux 64: 8, female – “Cochinchine; Annam (M. Boutan); Malacca; Célèbes septentrionalis; India” (lectotype Paris). – van der Vecht, 1957, Zool. Verh., Leiden 34: 28 (designation of lectotype).
- Vespa formosana* Sonan, 1927, Trans. Nat. Hist. Soc. Formosa 17 (89): 125, female, male – [Taiwan] “Taihoku, Kagi, Takezaki, Nanto, Daimokko, Maruyama” (Taichung).
- Vespa affinis continentalis* Bequaert, 1936, Treubia 15 (4): 346 (key), 350, female – “INDIA: Mangalore” (Washington); also from another locality in India; Myanmar; Sri Lanka; Thailand.
- Vespa affinis* var. *hainanensis* Bequaert, 1936, Treubia 15 (4): 347 (key), 349, female – “CHINA: Hainan Island . . . (2 miles south of Nodoa)” (Cambridge).
- Vespa affinis nigriventris* van der Vecht, 1957, Zool. Verh., Leiden 34: 28 (key), 29, female – [Philippines] “Palawan . . . Puerto Princesa” (Gainesville); also from another locality on Palawan.
- Vespa affinis rufonigrans* van der Vecht, 1957, Zool. Verh., Leiden 34: 28 (key), 29, female, male – [Indonesia] “Palu, North West Celebes” (holotype female Leiden); also from elsewhere on Sulawesi, nearby islands, Borneo, northern Moluccas, Misool, Salawati, Waigeo, New Guinea.
- Vespa affinis archboldi* van der Vecht, 1957, Zool. Verh., Leiden 34: 28 (key), 32, female – [Indonesia] “Hollandia” (Leiden); also from elsewhere in northern New Guinea.
- Vespa affinis moluccana* van der Vecht, 1957, Zool. Verh., Leiden 34: 28 (key), 32, female, male – [Indonesia] “Saparua I. near Amboina” (holotype female Leiden); also from southern Moluccas, Kai, Aru, New Guinea.
- Vespa affinis alticincta* van der Vecht, 1957, Zool. Verh., Leiden 34 28 (key), 33, female, male – “New Britain” (holotype female London).

Distribution: India: Bihar, West Bengal, Maharashtra, Karnataka, Tamil Nad, Kerala, Sikkim, Assam; Sri Lanka; Myanmar; China: Hainan, Xianggang, Guangdon, Fujian, Guangxi, Hunan, Hubei, Shanghai; Taiwan; Japan in the Ryukyu Is. south of Miyako I.; Thailand; Laos; Vietnam; Malaysia: Peninsular Malaysia, Sabah, Sarawak; Singapore; Indonesia: Sumatra, Nias, Bangka, Java, Borneo (Kalimantan), Sulawesi, Talaud Is., Buton, Salayar, Sunda Is., Timor, Moluccas (Morotai, Ternate, Tidore, Halmahera, Bacan, Buru, Ambon, Saparua, Haruku, Seram), Kai, Misool, Salawati, Waigeo, Sorong, Yapen, New Guinea (Irian Jaya); Papua New Guinea, including New Britain and New Ireland; Philippines: Palawan, Dumaran, Labuan, Mindanao, Samar, Luzon.

#### *analis* Fabricius

- Vespa analis* Fabricius, 1775, Syst. Ent.: 363 – “in Cap. B. S.” [error; probably Java] (London).
- Vespa Crabro sphinx* Christ, 1791, Naturgesch. Ins.: 217, pl. 18 fig. 5, female (destroyed).
- Vespa tyrranica* Smith, 1857, Cat. Hym. Brit. Mus. 5, Vesps.: 119, female – “Singapore” (lectotype London). – van der Vecht, 1959, Zool. Meded., Leiden 36: 215 (designation of lectotype).
- Vespa japonica* Smith, 1868, Ent. Mon. Mag. 4: 279, male – “Hakodadi, in Japan” (lectotype London). Junior primary homonym of *Vespa japonica* Radoszkowski, 1857, and of *Vespa japonica* de Saussure, 1858. – Kojima, 1997, Nat. Hist. Bull. Ibaraki Univ. 1: 20 (designation of lectotype).
- Vespa parallelula* André, 1884, Ann. Soc. Ent. France (6) 4, Bull.: lxi, female – [Russia] “Wladivostock, sur l’Amour, Sibérie orientale” (Paris?).

*Vespa insularis* Dalla Torre, 1894, Cat. Hym. 9: 147. Replacement name for *Vespa japonica* Smith.

*Vespa tridentata* Cameron, 1903, Entomologist 36: 278, female – “Japan” (lectotype London). – Kojima, 1997, Nat. Hist. Bull. Ibaraki Univ. 1: 20 (designation of lectotype).

*Vespa nigra* du Buysson, 1903, Bull. Soc. Ent. France 1903: 175, male – [China] “Yunnan, Tsé-kou” (Paris).

*Vespa parallela* var. *Biroi* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 487 (key), 513, female – “Singapore” (lectotype Paris). – van der Vecht, 1957, Zool. Verh., Leiden 34: 14 (designation of lectotype).

*Vespa analis* var. *tenebrosa* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 516, female – [Indonesia] “Java, Goban . . . Gondang” (lectotype Paris). – van der Vecht, 1957, Zool. Verh., Leiden 34: 14 (designation of lectotype).

*Vespa analis* var. (or subsp.) *barbouri* Bequaert, 1939, Trans. Am. Ent. Soc. 65: 38 (key), 40, female – [India] “Teesta Valley, Sikkim” (Cambridge).

*Vespa analis* var. (or subsp.) *kuangsiiana* Bequaert, 1939, Trans. Am. Ent. Soc. 65: 38 (key), 42, female – “Kwangsi, China” (Cambridge).

*Vespa analis eisa* Yamane, 1987, Kontyū 55 (4): 628 (key), 631, female – [Japan] “Yona, Okinawa-jima” (Kagoshima); also from elsewhere on Okinawa, and central Ryukyus.

*Vespa analis nagatomii* Yamane, 1987, Kontyū 55 (4): 628 (key), 632, female – [Japan] “Koza-dake, Iriomote-jima” (Kagoshima); also three other localties on Iriomote.

Distribution: India: Kashmir, Uttar Pradesh, West Bengal, Tamil Nadu, Meghalaya, Sikkim, Assam; Nepal; China: Sichuan, Yunnan, Hainan, Guangxi, Fujian, Hubei, Zhejiang, Heilongjiang; Myanmar; Thailand; Laos; Malaysia: Peninsular Malaysia; Singapore; Indonesia: Sumatra, Bangka, Sunda Is. (Sebesi, Sangijiang, Sebuku, Sertung, Madura, Lombok, Legundi, Rakata, Krakatau, Panaitan, Deli), Java, Bawean, Bali; Taiwan; Korea; Japan, including Ryukyu Is.; Russia: Amur, Primor’ye.

#### *auraria* Smith

*Vespa auraria* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 46, pl. VIII fig. 8, female – “Northern India” (London, Oxford?).

*Vespa velutina pruthii* van der Vecht, 1959, Zool. Meded., Leiden. 36 (13): 228, female – [India] “Kashmir, Srinagar” (Calcutta); also from Pakistan.

Distribution: Afghanistan; Pakistan; India: Kashmir, Himachal Pradesh, Punjab, Uttar Pradesh, West Bengal, Meghalaya, Sikkim, Assam; Nepal; Myanmar; Thailand; Laos; China: Yunnan; Malaysia: Peninsular Malaysia. Introduced Yemen.

#### *basalis* Smith

*Vespa basalis* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 46, female – “Nepaul” (London).

*Vespa oblitterata* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 47, female – “Northern India” (London).

*Vespa basilis*; Dover, 1929, Bull. Raffles Mus. 2: 48. Incorrect spelling of *Vespa basalis* Smith.

Distribution: Pakistan; India: Himachal Pradesh, West Bengal, Meghalaya, Sikkim, Assam; Nepal; China: Sichuan, Yunnan, Fujian; Taiwan; Myanmar; Thailand; Laos; Vietnam; Indonesia: Sumatra.

Note: The record from Korea in Kim et al. (1994) was based on misidentification (Kim, personal communication).

***bellicosa*** de Saussure

*Vespa Bellicosa* de Saussure, 1854, Ét. Fam. Vespa. 2.: 146, pl. XIV fig. 10, female – “Java” [error; Sumatra or Borneo] (lectotype Torino). – van der Vecht, 1959, Zool. Meded., Leiden 36: 216 (designation of lectotype).

Distribution: Malaysia: Sarawak; Indonesia: Sumatra, Bengkalis, Borneo (Kalimantan).

***bicolor*** Fabricius

*Vespa bicolor* Fabricius, 1787, Mant. Ins. 1: 288 – “in China” (Kobenhavn).

*Vespa lutea* Coquebert, 1804, Illustr. Iconogr. Insect. 3: 94, pl. XXI fig. 10 – “Massiliae lecta, in navi ex India” (type depository unknown).

*Vespa auraria* var. *citriventris* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 552, female – [India] “Sikkim, de Padamtsin à Lingtou” (lectotype Paris); also from West Bengal; and China. – van der Vecht, 1959, Zool. Meded., Leiden 36: 218 (designation of lectotype).

Distribution: India: Uttar Pradesh, West Bengal, Sikkim, Assam, Meghalaya; Bhutan; Nepal; China: Sichuan, Yunnan, Guangxi, Jiangxi, Fujian; Xianggang; Myanmar; Thailand; Laos; Cambodia; Vietnam.

***binghami*** du Buysson

*Vespa Binghami* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 488 (key), 523, female – [Myanmar] “Tenasserim, Taungoo Hills” (lectotype Paris); also from China; and “Indes orientales.” – van der Vecht, 1959, Zool. Meded., Leiden 36: 210 (designation of lectotype).

*Vespa suprunenkoi* Birula, 1925 (1924), Arch. Naturgesch. (Abt. A) 90 (12): 89 (key), 92, female – [Russia] “Insel Sachalin” (St. Petersburg); also from Korea.

Distribution: India: West Bengal, Assam, Meghalaya; Bhutan; Myanmar; Thailand; Laos; Vietnam; China: Xizang, Sichuan, Yunnan, Jiangsu, Shanghai; Russia: Primor’ye, Sakhalin; Korea.

***crabro*** Linnaeus

*Vespa Crabro* Linnaeus, 1758, Syst. Nat., ed. 10, 1: 572 – “in Europae” (holotype female, Linnaean Society, London).

*Vespa vexator* Harris, 1776, Exposit. Eng. Ins.: 127, pl. XXXVII figs. 2–3, female – “English” (destroyed).

*Vespa Crabro major* Retzius, 1783, C. Degeer Gen. Spec. Ins.: 63 (type depository unknown).

? *Vespa pratensis* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 437 – [France] “in Agro Parisiensi” (type depository unknown).

*Vespa Crabro germana* Christ, 1791, Naturgesch Ins.: 215, pl. 18 fig. 3 (destroyed).

*Vespa Crabroniformis* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 40, female, male – “North China” (London).

*Vespa crabro* var. *borealis* Radoszkowski, 1863, Horae Soc. Ent. Ross. 2: 128, pl. XVI fig. 15, female, male – [Russia] “Pargolova i Osinovo Roshchi” (type depository unknown). Junior primary homonym of *Vespa borealis* Kirby, 1837, *Vespa borealis* Zetterstedt, 1840, and of *Vespa borealis* Smith, 1843.

- Vespa crabro* var. *anglica* Gribodo, 1892 (1891), Bull. Soc. Ent. Ital. 23: 242, female – “Inghilterra” (Genova). Junior primary homonym of *Vespa anglica* Smith, 1843.
- Vespa oberthuri* du Buysson, 1902, Bull. Soc. Ent. France 1902: 140, female – “Chine: Se-tchouen, Siao-Lou” (Paris, Leiden); also from Ta-tsien-lou.
- Vespa flavo-fasciata* Cameron, 1903, Entomologist 36: 280, female – “Japan . . . Nügata (Shinanogawa)” (lectotype London). – Kojima, 1997, Nat. Hist. Bull. Ibaraki Univ. 1: 21 (designation of lectotype).
- Vespa crabro* var. *tartarea* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73 (3): 492 (key), 506, female – “Japon: Yokohama . . . Columbia” (Budapest, coll. Martinez Escalera).
- Vespa Crabro* var. *Altaica* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 5, female – “Altai” (Paris).
- Vespa Crabro* var. *Caspica* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 6, female – [Azerbaijan] “Talysch et Lenkoran, région Caspienne” (Paris).
- Vespa crabro vulgata* Birula, 1925 (1924), Arch. Naturgesch. (Abt. A) 90 (12): 100, female, male – “West-Europa” (St. Petersburg).
- Vespa crabro meridionalis* Birula, 1925 (1924), Arch. Naturgesch. (Abt. A) 90 (12): 100, female – “Transkaukasien, Lagodechi, . . . Gouv. Tiflis, . . . Lenkoran . . . Kutais, . . . Transkaspien, Krassnovodsk” (St. Petersburg).
- Vespa crabro chinensis* Birula, 1925 (1924), Arch. Naturgesch. (Abt. A) 90 (12): 101, female, male – “China, auf dem Wege zwischen Peking und She-che . . . Tien-zsin, . . . Tal Lu-li-cho, 65 Werst südlich von Peking, . . . Prov. Sze-tschan von Tshan-Tshou zu Tsha-mynj” (St. Petersburg). Junior primary homonym of *Vespa chinensis* Fabricius, 1793.
- Vespa crabro* var. *birulai* Bequaert, 1931, Konowia 10 (2): 105. Replacement name for *Vespa crabro chinensis* Birula.
- Vespa crabro* var. *gribodoi* Bequaert, 1931, Konowia 10 (2): 105. Replacement name for *Vespa crabro* var. *anglica* Gribodo.

Distribution: Norway; Sweden; Finland; Denmark; U. K.: Scotland, Wales, England; Belgium; France; Spain; Germany; Switzerland; Austria; Italy; Poland; Hungary; Albania; Ukraine; Turkey; Georgia; Azerbaijan; Iran; across Russia to Siberia, Sakhalin; Mongolia; China: Sichuan, Yunnan, Hubei, Beijing, Jilin; Taiwan; Korea; Japan. Introduced into Canada (Ont.); U. S. A. (ND, SD, WI, IN, OH, NY, PA, MA, CT, RI, NJ, DE, MD, DC, KY, WV, VA, TN, NC, SC, AL, GA).

Note: Bequaert (1931: 101) regarded du Buysson’s (1905) record for Algeria as either an accidental introduction, or error in labelling.

#### *ducalis* Smith

- Vespa ducalis* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 39, female, male – [China] “Tein-tung, near Nigo-po-foo” (London).
- Vespa ducalis* var. *pulchra* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 519, female – “Corée, Séoul . . . Japon . . . : Yokohama” (lectotype from “Yokohama” Paris). – van der Vecht, 1959, Zool. Meded., Leiden 36: 224 (designation of lectotype).
- Vespa matsumurai* Sonan, 1935, Trans. Nat. Hist. Soc. 25 (145): 370, figs. 1–3, female – [Japan] “Tokyo” (Taichung).
- Vespa esakii* Sonan, 1935, Trans. Nat. Hist. Soc. Formosa 25 (145): 371, figs. 4–6, female – [Japan] “Izuhara in Tsushima Island” (Fukuoka).

*Vespa tropica loochooensis* Bequaert, 1936, Treubia 15 (4): 336 (key), 343, female – [Japan] “Iriomote Island, in the Loo-CHOO (or Riu-Kiu) ARCHIPELAGO” (Cambridge).

*Vespa tropica pseudosoror* van der Vecht, 1959, Zool. Meded. 36 (13): 224, female – [Vietnam] “Indo-China: Annam . . . Lién-Chiéen près Tourane, 1000 m” (Paris); also from elsewhere in Vietnam; and Taiwan.

Distribution: India: Sikkim, Meghalaya; Nepal; Mynmar; Thailand; Vietnam; China: Gansu, Sichuan, Hainan, Guangdong, Fujian, Hunan, Shanghai, Jiangsu, Liaoning; Taiwan; Russia: Primor’ye; Korea; Japan, including Ryukyu Is.

#### *dybowskii* André

*Vespa Dybowskii* “Rad. in litt.” André, 1884, Spec. Hym. Europe Algér. 2: 582, female – [Russia] “Sibérie” (Paris?).

*Vespa Dubowskii*; Dalla Torre, 1904, Gen. Ins. 19: 65. Incorrect spelling of *Vespa dybowskii* André.

*Vespa Walkeri*; du Buysson, 1905 (1904), Ann. Soc. Ent. France 73 (3): 490 (key), 539, female, male – “Chine: Hai-ning” (London, Paris).

*Vespa dybowskii mutata*; Ma, 1937, Ent. Phytopath. 5 (2): 30, male (in subgenus *Vespa*) – [China] “Hangchow” (Hangzhou, destroyed?).

Distribution: Myanmar; Thailand; China: Xizang, Zhejiang; Russia: Primor’ye; Korea; Japan.

#### *fervida* Smith

*Vespa fervida* Smith, 1858, J. Proc. Linn. Soc. Lond. 3 (9): 23, female – [Indonesia] “Celebes” (London).

Distribution: Indonesia: Sulawesi, Salayar, Buton.

#### *fumida* van der Vecht, NEW STATUS

*Vespa variabilis* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 490 (key), 522, 627, female – “Chine: Mou-pin” (lectotype Paris); also from elsewhere in China; and Indonesia [error]. Junior primary homonym of *Vespa variabilis* Fabricius, 1781. – van der Vecht, 1959, Zool. Meded., Leiden 36: 228 (designation of lectotype).

*Vespa variabilis fumida* van der Vecht, 1959, Zool. Meded. 36 (13): 228, female – “India . . . ‘British Bootan’, Padong” (Paris); also from Darjeeling.

Distribution: India: West Bengal, Sikkim, Arunachal Pradesh, Assam; Bhutan; Nepal; Myanmar; China: Sichuan, Yunnan, Fujian, Hubei.

#### *luctuosa* de Saussure

*Vespa luctuosa* de Saussure, 1854, Ét. Fam. Vesp. 2.: 143, male – “Les Philippines” (London).

*Vespa bellicosa* var. *Semperi* du Buysson, 1905 (1904), Ann. Soc. Ent. France 63: 542, female – “Comigu” [Philippines: ?Camiguin] (lectotype Wien); also from Mindanao, Lubang, Luzon. – van der Vecht, 1957, Zool. Verh., Leiden 34: 48 (designation of lectotype).

*Vespa luctuosa luzonensis* Bequaert, 1934, Bull. Mus. R. Hist. Nat. Belg. 10 (28): 3 (key), 5, female – “Luzon, Philippines” (Berlin).

*Vespa luctuosa negrosensis* Kojima and Tano, 1985, Philipp. Ent. 6 (4): 436, female – [Philippines] “Mambucal, Negros Is.” (Tsukuba).

Distribution: Philippines: Luzon, Lubang, Mindoro, Panay, Negros, Cebu, Biliran, Samar, Leyte, Panaon, Mindanao.

***mandarinia*** Smith

*Vespa Mandarinia* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 38, pl. VIII fig. 1, female – [China] “Tein-tung, near Ning-po-foo, amongst the mountains” (London).

*Vespa magnifica* Smith, 1852, Trans. Ent. Soc. Lond. (N. S.) 2 (2): 45, female – “Nepaul” (London, Oxford).

*Vespa japonica* Radoszkowski, 1857, in Motschulsky, Ét. Ent. 6: 410 – “Japon” (St. Petersburg).

*Vespa bellona* Smith, 1871, Proc. Zool. Soc. Lond.: 248, pl. XVIII fig. 6, female – [China] “Yunan” (London).

*Vespa magnifica* var. *latilineata* Cameron, 1903, Entomologist 36: 278, female – “Japan . . . Hitoyoshi” (London).

*Vespa mandarina*; Dalla Torre, 1894, Cat. Hym. 9: 149. Incorrect spelling of *Vespa mandarinia* Smith.

*Vespa magnifica* var. *nobilis* Sonan, 1929, Trans. Nat. Hist. Soc. Formosa. 19 (101): 137 (key), 140, female – [Taiwan] “Musha” (Taichung); also from other localities on Taiwan.

*Vespa magnifica sonani* Matsumura, 1930, Illust. Thousand Ins. Japan 2: 1, pl. 1 figs. 2, 5 female – “Taiwan, Sina” (lectotype Sapporo). – Kojima, 1997, Nat. Hist. Bull. Ibaraki Univ. 1: 22 (designation of lectotype).

Distribution: India: Himachal Pradesh, Uttar Pradesh, West Bengal, Sikkim, Assam, Meghalaya; Sri Lanka; Nepal; Bhutan; Myanmar; Thailand; Laos; Malaysia: Peninsular Malaysia; China: Sichuan, Yunnan, Xianggang, Jiangxi, Fujian, Hubei, Shanghai, Jiangsu; Taiwan; Russia: Primor’ye; Korea; Japan, including Yaku-shima I. and Tanega-shima I. in Ryukyu Is.

***mocsaryana*** du Buysson

*Vespa Mocsaryana* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 490 (key), 537, pl. VII fig. 3, female, male – “Presqu’ile de Malacca: Péراك . . . ; Sikkim . . . ; Assam: Khasia Hills . . . ; Tenasserim: Thagatā” (Budapest, Wien, London, Genova).

Distribution: India: Meghalaya, Sikkim, Assam; China: Sichuan, Fujian, Anhui; Myanmar; Thailand; Laos; Vietnam; Malaysia: Peninsular Malaysia; Indonesia: Sumatra.

***multimaculata*** Pérez

*Vespa annulata* Smith, 1858, J. Proc. Linn. Soc. Lond., Zool. 2 (6): 116, female – [Malaysia] “Borneo (Sarawak). Malacca.” (lectotype Oxford). Junior primary homonym of *Vespa annulata* Rossi, 1790.

*Vespa multimaculata* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 14, female – “Brunei (Java)” [Borneo] (lectotype Paris). – van der Vecht, 1957, Zool. Verh., Leiden 34: 44 (designation of lectotype).

*Vespa luctuosa* var. *malayana* Bequaert, 1934, Bull. Mus. R. Hist. Nat. Belg. 10 (28): 3 (key), 4. Replacement name for *Vespa annulata* Smith.

*Vespa multimaculata pendleburyi* van der Vecht, 1957, Zool. Verh., Leiden 34: 43, 44, female, male – “Catchment Area near Jitra, Kedah, N. Malaya” (holotype female Leiden); also from elsewhere in Peninsular Malaysia; Thailand; Singapore; Indonesia: Sumatra.

Distribution: Thailand; Malaysia: Peninsular Malaysia, Sarawak; Singapore; Indonesia: Sumatra, Borneo (Kalimantan); Brunei.

*orientalis* Linnaeus

*Vespa orientalis* Linnaeus, 1771, Mant. Plant., ed. 2: 540 – “Oriente” (holotype female, Linnaean Society, London).

*Vespa turcica* Drury, 1773, Illustr. Nat. Hist. 2: 74, pl. XXXIX fig. 1, index – [Turkey] “Smyrna” (destroyed?).

*Vespa quadripunctata* Forskål, 1775, Descr. Anim.: 84 – [Egypt] “Kahirae” (Kobenhavn?).

*Vespa Crabro fusca* Christ, 1791, Naturgesch. Ins.: 216 – [Turkey] “Smirna” (destroyed).

*Vespa AEgyptiaca* Vallot, 1802, Concord. Syst.: 170 – “Égypte” (type depository unknown).

*Vespa Nilotica* Vallot, 1802, Concord. Syst.: 170 – “Égypte” (type depository unknown).

*Vespa Jurinei* de Saussure, 1854, Ét. Fam. Vesp. 2: 133 – “L’Albanie” (lectotype female London). – Guiglia, 1971, Faune Europe Bassin Méditerran. 6: 94 (designation of lectotype).

*Vespa orientalis* var. *AEGYPTIACA* André, 1884, Spec. Hym. Europe Algér. 2: 584 – [Egypt] “Le Caire, Alexandrie” (Paris). Junior primary homonym of *Vespa aegyptiaca* Vallot, 1802.

*Vespa indica*; Wroughton, 1889, J. Bombay Nat. Hist. Soc. 4: 35. Lapsus for *Vespa orientalis* Linnaeus.

*Vespa orientalis* var. *Zavattarii* Guiglia and Capra, 1933, Mem. Soc. Ent. Ital. 12: 168, female – “Fezzan . . . Ubari [Libya] . . . Oued Tizzi (Algeria)” (Genova).

*Vespa orientalis* var. *somalica* Giordani Soika, 1934, Boll. Soc. Ent. Ital. 66 (8): 184, female – “Somalia italiana . . . di Carim” (lectotype Venezia); also from Bender Cassim; 1973, Boll. Mus. Civ. Stor. Nat. Venezia 24: 11 (designation of lectotype).

*Vespa orientalis arabica* Giordani Soika, 1957, Brit. Mus. (Nat. Hist.) Exp. S. W. Arabia 1 (31): 482, female, male – “WESTERN ADEN PROTECTORATE: Al Milah, 1600 ft.” (holotype female London); also from elsewhere in Yemen; and Oman.

Distribution: Italy, including Sicily; Malta; Albania; Romania; Greece, including Crete; Bulgaria; Turkey; Cyprus; Syria; Lebanon; Iraq; Israel; Jordan; Saudi Arabia; Bahrein; Yemen; Oman; U. A. E.; Algeria; Libya; Egypt; Ethiopia; Somalia; Russia; Turkmenistan; Uzbekistan; Tajikistan; Iran; Afghanistan; Pakistan; India: Kashmir, Punjab, Haryana, Uttar Pradesh, Bihar, Delhi, West Bengal, Maharashtra, Andhra Pradesh; Nepal; China: Xinjiang. Introduced Madagascar.

*philippinensis* de Saussure

*Vespa Philippinensis* de Saussure, 1854, Ét. Fam. Vesp. 2.: 148, male – “Les Philippines” (London).

Distribution: Philippines: Luzon, Negros, Samar, Leyte, Mindanao.

*simillima* Smith

*Vespa simillima* Smith, 1868, Ent. Mon. Mag. 4: 280, female – “Japan . . . Hakodadi” (London).

*Vespa mongolica* André, 1884, Ann. Soc. Ent. France (6) 4, Bull.: lix, male, female – [Russia] “Wladivostock, sur l’Amour, dans la Sibérie orientale” (Paris?).

*Vespa xanthoptera* Cameron, 1903, Entomologist 36: 278, male – “Japan . . . Michzusawa” (London).

*Vespa micado* Cameron, 1903, Entomologist 36: 279, female – “Japan . . . Nagasaki” (lectotype London). – Kojima, 1997, Nat. Hist. Bull. Ibaraki Univ. 1: 20 (designation of lectotype).

*Vespa Mongolica* var. *Sexpunctata* Pérez, 1905, Bull. Mus. Hist. Nat. Paris 1905: 24 (*sexpunctata*), 79, female – [Japan] “Yokohama” (holotype Paris).

*Vespa Mongolica* var. *flavata* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 17, female – “Chine” (Paris).

Distribution: China: Liaoning; Russia: Primor’ye, Sakhalin, Kurile Is.; Korea; Japan, including Ryukyu Is. (Yaku-shima I.). Introduced into Canada (B. C.).

#### *soror* du Buysson

*Vespa ducalis* var. *soror* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 490 (key), 519, female – “Chine . . . Kiang-si” (lectotype Paris); also from elsewhere in China; India; Vietnam. – van der vecht, 1957, Zool. Verh., Leiden 34: 16 (designation of lectotype).

Distribution: China: Yunnan, Hainan, Fujian; Thailand; Laos; Vietnam.

#### *tropica* (Linnaeus)

*Sphex tropica* Linnaeus, 1758, Syst. Nat., ed. 10, 1: 571 – “in Indiis” (holotype female Uppsala).

*Vespa cincta* Fabricius, 1775, Syst. Ent.: 362 – [India] “ad littora Malabarica” (Kobenhavn). Junior primary homonym of *Vespa cincta* Drury, 1773, and of *Vespa cincta* Degeer, 1773.

*Vespa Crabro tenebrionis* Christ, 1791, Naturgesch. Ins.: 216, pl. 18 fig. 4 (destroyed).

*Vespa deusta* Lepeletier, 1836, Hist. Nat. Ins. Hym. 1: 506, female – “Patrie inconnue” (lectotype Torino). – van der vecht, 1959, Zool. Meded., Leiden 36: 226 (designation of lectotype).

*Vespa unicolor* Smith, 1863, J. Proc Linn. Soc. Lond., Zool. 7: 44, female – [Indonesia] “Bouru” (Oxford). Junior primary homonym of *Vespa unicolor* Lichtenstein, 1796.

*Vespa cinta*; Wroughton, 1889, J. Bombay Nat. Hist. Soc. 4: 35. Incorrect spelling of *Vespa cincta* Fabricius.

*Vespa eulemoides* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 491 (key), 530, female – [India] “Ile Andaman: port de Blair” (Genova).

*Vespa affinis* var. *trisignata* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 8, female – [Indonesia] “Timor” (Paris).

*Vespa rubricans* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 10, female – “Lindi (Afrique orientale allemande)” (lectotype Paris). – van der Vecht, 1957, Zool. Verh., Leiden 34: 24 (designation of lectotype).

*Vespa tropica* var. *anthracina* Bequaert, 1936, Treubia 15 (4): 335 (key), 341, female, male – “PHILIPPINE ISLANDS: Sibuwan” (holotype female Washington); also from other localities in the Philippines.

*Vespa tropica* var. *haematodes* Bequaert, 1936, Treubia 15 (4): 336 (key), 338, female, male – “INDIA: Kooloo” (holotype female Cambridge); also from six other localities in India; Sri Lanka; Myanmar; China.

*Vespa tropica leefmansi* van der Vecht, 1957, Zool. Verh., Leiden 34: 19, female, male – [Indonesia] “Solok, Padang, Sumatra” (holotype female Leiden); also from Vietnam; Thailand; Malaysia; and other islands in Indonesia.

*Vespa tropica trimeres* van der Vecht, 1957, Zool. Verh., Leiden 34: 19 (key), 22, female – [Indonesia] “Palu, West Celebes” (Leiden).

*Vespa tropica cebuana* Kojima and Reyes, 1984, Kontyû 52 (2): 260, female – [Philippines] “Cebu City” (Tsukuba); also from Talisay.

Distribution: Afghanistan; Pakistan; India: Himachal Pradesh, Bihar, West Bengal, Maharashtra, Karnataka, Tamil Nadu, Kerala, Sikkim, Assam, Meghalaya, Andaman and Nicobar Is.; Sri Lanka; Nepal; Bhutan; China: Yunnan, Xianggang, Fujian; Myanmar; Thailand; Laos; Cambodia; Vietnam; Malaysia: Peninsular Malaysia, Sarawak; Indonesia: Anambas Is., Sumatra, Nias, Batu, Enggano, Bangka, Belitung, Sunda Is. (Sangijiang, Sebesi, Sebuku, Legundi, Krakatau, Panaitan), Java, Bawean, Madura, Kangean Is., Bali, Lombok, Komodo, Sumbawa, Sumba, Flores, Wetar, Timor, Borneo (Kalimantan), Sulawesi, Salayar, Sula Is., Moluccas (Morotai, Halmahera, Ternate, Tidore, Bacan, Obi, Buru, Ambon, Haruku, Seram), Banda, Waigeo, Aru, New Guinea (Irian Jaya); Papua New Guinea, including New Britain; Philippines: Luzon, Lubang, Mindoro, Samar, Leyte, Sibuyan, Panay, Negros, Cebu, Palawan, Jolo, Mindanao, Basilan.

#### *velutina* Lepeletier

*Vespa velutina* “De Haan” Lepeletier, 1836, Hist. Nat. Ins. Hym. 1: 507, female – [Indonesia] “Inde: Java” (coll. Serville).

*Vespa crabro* var. *immaculata* Morawitz, 1889, Horae Soc. Ent. Ross. 23: 161, female – [China] “Kansu. Nanpin.” (St. Petersburg). Junior primary homonym of *Vespa immaculata* Gmelin, 1790.

*Vespa fruhstorferi* Stadelmann, 1894, Sitz.-Ber. Ges. Naturf. Freunde Berl. 1894 (3): 89, female – [Indonesia] “in den alpinen Theilen von West-Java auf dem Gunung-Gede . . . in einer Höhe von 8000 Fuss” (Berlin?).

*Vespa velutina* var. *ardens* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 550, female, male – [Indonesia] “Lombok, Sapit (2000’ . . . ); I. Sumbawa, près Java” (Wien, Budapest, München, Paris).

*Vespa auraria* var. *nigrithorax* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 553, female, male – [India] “Dardjiling” (lectotype female Paris); also from Sikkim; and Bhutan; China; Indonesia: Sumatra, Sulawesi. – van der Vecht, 1957, Zool. Verh., Leiden 34: 37 (designation of lectotype).

*Vespa velutina* var. *Celebensis* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 12, female – [Indonesia] “Bua-Kraeng (Sud-Célèbes)” (holotype Paris).

*Vespa velutina* var. *Megei* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 13, female – “Kouei-Tchéou (Chine centrale)” (Paris).

*Vespa Mongolica* var. *divergens* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 16, female – [Malaysia] “Perak” (lectotype Paris). – van der Vecht, 1957, Zool. Verh., Leiden 34: 38 (designation of lectotype).

*Vespa flavitarsus* Sonan, 1929, Trans. Nat. Hist. Soc. Formosa 19 (101): 137 (key), 142, female – [Taiwan] “Taiko-san (Kagi)” (coll. M. Kato); also from other localities on Taiwan.

*Vespa auraria flavitarsis*; Ma, 1937, Ent. Phytopath. 5 (2): 31. Incorrect spelling of *Vespa flavitarsus* Sonan.

*Vespa velutina sumbana* van der Vecht, 1957, Zool. Verh., Leiden 34: 34 (key), 40, female, male – [Indonesia] “Kananggar, E. Sumba, 700 m” (holotype female Leiden); also from elsewhere on Sumba.

*Vespa velutina variana* van der Vecht, 1957, Zool. Verh., Leiden 34: 35 (key), 37, female – “North Siam . . . Doi Hua Mot, 3000” (Leiden); also from elsewhere in Thailand; Laos; Vietnam.

*Vespa velutina karnyi* van der Vecht, 1957, Zool. Verh., Leiden 34: 35 (key), 38, female, male – [Indonesia] “Bandar Buru, Deli” (holotype female Leiden); also from elsewhere in Sumatra.

*Vespa velutina timorensis* van der Vecht, 1957, Zool. Verh., Leiden 34: 35 (key), 40, female – [Indonesia] “Mollo, 1350 m, Timor” (Basel); also from another locality on Timor.

*Vespa velutina floresiana* van der Vecht, 1957, Zool. Verh., Leiden 34: 35 (key), 40, female – [Indonesia] “Labuan Bajo, Flores” (Leiden); also from elsewhere on Flores, and Komodo.

Distribution: India: West Bengal, Sikkim, Assam; Bhutan; China: Sichuan, Jiangxi, Fujian, Hubei, Zhejiang, Xianggang; Taiwan; Myanmar; Thailand; Laos; Vietnam; Malaysia: Peninsular Malaysia; Indonesia: Sumatra, Java, Bali, Lombok, Sumbawa, Komodo, Sumba, Flores, Timor, Sulawesi.

#### *vivax* Smith

*Vespa vivax* Smith, 1870, in Horne and Smith, Trans. Zool. Soc. Lond. 7 (3): 190, pl. XXI fig. 9, female – “Binsur, Kumaon, North-west Provinces of India” (London, Oxford?).

*Vespa velutina* var. *mediozonalis* Pérez, 1910, Actes Soc. Linn. Bordeaux 64: 14, female – [India] “Sikkim” (Paris).

*Vespa wilemani* Meade-Waldo, 1911, Ann. Mag. Nat. Hist. (8) 7: 104, female – “Formosa” (London).

Distribution: India: Kashmir, Himachal Pradesh, Uttar Pradesh, Sikkim; Nepal; Myanmar; Thailand; China: Xizang, Sichuan, Yunnan; Taiwan.

#### Genus *Vespula* Thomson

*Vespula* Thomson, 1869, Opusc. Ent. 1: 79, subgenus of *Vespa* (eight species).

Type species: *Vespa austriaca* Panzer, 1799, by subsequent designation of Ashmead, 1902, Can. Ent. 34: 164.

*Pseudovespa* Schmiedeknecht, 1881, Ent. Nachr. 7: 314, subgenus of *Vespa* Linnaeus.

Type species: *Vespa austriaca* Panzer, 1799, by monotypy.

*Paravespula* Blüthgen, 1938 (1937), Konowia 16: 271, subgenus of *Dolichovespula* Rohwer.

Type species: *Vespa vulgaris* Linnaeus, 1758, by original designation.

*Allovespula* Blüthgen, 1943, Stettin. Ent. Ztg. 104: 149, subgenus of *Paravespula* Blüthgen.

Type species: “*Paravespula rufa* (Linné)” [= *Vespa rufa* Linnaeus, 1758], by monotypy.

*Rugovespula* Archer, 1982, Kontyû 50 (2): 261, 264, subgenus of *Vespula* Thomson.

Type species: *Vespa koreensis* Radoszkowski, 1887, by original designation.

*acadica* (Sladen)

*Vespa rufa* var. *americana* du Buysson, 1905 (1904), Ann. Soc. Ent. France 63: 499 (key), 592, male – “Canada, Lévis près Québec” (type depository unknown). Junior primary homonym of *Vespa americana* Fabricius, 1775, and of *Vespa maculata americana* Christ, 1791.

*Vespa acadica* Sladen, 1918, Ottawa Nat. 32: 72, male, female – [Canada] “Ottawa” (lectotype female Ottawa) also from Nova Scotia, N.B., B.C. – Miller, 1961, Can. Ent., Suppl. 22: 15, 17 (designation of lectotype).

*Vespa rufa* var. *sladeni* Bequaert, 1932, Ent. Am. (N. S.) 12 (2): 98, 102, female, male – [U. S. A.] “WASHINGTON STATE” (holotype female Cambridge); also from OR, CA, WY, UT, CO; Canada: BC.

Distribution: Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, NV, ID, MT, WY, UT, CO, AZ, NM, SD, ND, MN, NE, WI, MI, NY, PA, VT, NH, ME, MA, CT, RI, NJ, MD, WV, VA, NC.

*arisana* Sonan

*Vespa arisana* Sonan, 1929, Trans. Nat. Hist. Soc. Formosa. 19 (101): 137 (key), 147, female – [Taiwan] “Funkiko (Arisan 4,600 ft.)” (Taichung); also from elsewhere on Taiwan.

Distribution: Taiwan.

*atropilosa* (Sladen)

*Vespa atropilosa* Sladen, 1918, Ottawa Nat. 22: 72, female – [Canada] “Lethbridge, Alta.” (lectotype Ottawa); also from B.C. – Miller, 1961, Can. Ent., Suppl. 22: 17, 19 (designation of lectotype).

Distribution: Canada: B. C., Alta.; U. S. A.: WA, OR, CA, NV, ID, MT, WY, UT, CO, AZ, NM, ND, NE; Mexico: Baja California Norte.

*austriaca* (Panzer)

*Vespa austriaca* Panzer, 1799, Faunae Ins. German. 63: 2, fig. 2, male – “Viennae Austr.” (destroyed).

*Vespa borealis* Smith, 1843, Zoologist 1: 170, fig., female – “Yorkshire . . . the north of Scotland” (London). Junior primary homonym of *Vespa borealis* Kirby, 1837, and of *Vespa borealis* Zetterstedt, 1840.

*Vespa arborea* Smith, 1849, Zoologist 7, App.: lx. Replacement name for *Vespa borealis* Smith.

*Vespa biloba* Schilling, 1850, Jahresber. Schles. Ges. Vaterländ. Kultur 28: 77 – [Poland] “Schlesien” (type depository unknown).

*Vespa Infernalis* de Saussure, 1854, Ét. Fam. Vesp. 2: 139, female – [U. S. A.] “L’Amériquée du Nord. Philadelphie” (Torino).

*Vespa tripunctata* Packard, 1870, Trans. Chicago Acad. Sci. 2: 26, pl. II fig. 11, female – [U. S. A.] “Alaska” (type depository unknown). Junior primary homonym of *Vespa tripunctata* Fabricius, 1787, and of *Vespa tripunctata* Schenck, 1861.

*Vespa arborea* “Ross” Dalla Torre, 1894, Cat. Hym. 9: 138, as a synonym of *Vespa austriaca* Panzer. Junior primary homonym of *Vespa arborea* Smith, 1849.

Distribution: Norway; Sweden; Finland; Denmark; Ireland; U. K.: Scotland, Wales, England; Belgium; Luxembourg; France; Spain; Germany; Switzerland; Austria; Italy; Poland; Czech Republic; Bulgaria; Turkey; Georgia; across Russia to Kamchatka, Amur;

Pakistan; India: Kashmir; Mongolia; China: Xinjiang, Hebei, Liaoning; Korea; Japan; Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, ID, MT, UT, CO, NM, AZ, NM, ND, WI, MI, NY, VT, NH, ME, MA, CT, RI, NJ.

*consobrina* (de Saussure)

*Vespa Consobrina* de Saussure, 1854, Ét. Fam. Vespa. 2: 141, female – [Canada] “L’île de Terre-Neuve” (Paris).

*Vespa scelestā* McFarland, 1888, Trans. Am. Ent. Soc. 15: 297 (key), 298, female, male – [U. S. A.] “Montana” (lectotype female Philadelphia); also from PA, VA, NH, CO, ME, WA, MA. – Cresson, 1928, Mem. Am. Ent. Soc. 5: 57 (designation of lectotype).

*Vespa sulcata*; Howard, 1901, Insect Book: pl. VI fig. 18. Lapsus for *Vespa scelestā* McFarland.

Distribution: Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: WA, OR, CA, ID, MT, WY, UT, CO, ND, MN, WI, IL, MI, IN, OH, NY, PA, VT, NH, ME, MA, CT, RI, NJ, DE, MD, KY, WV, VA, TN, NC, GA.

*flaviceps* (Smith)

*Vespa japonica* de Saussure, 1858, Rev. Mag. Zool. (2) 10: 261, female – “Japon” (Genève). Junior primary homonym of *Vespa japonica* Radoszkowski, 1857.

*Vespa flaviceps* Smith, 1870, in Horne and Smith, Trans. Zool. Soc. Lond. 7 (3): 174, 191, pl. XXI figs. 10, 11, female – “Binsur, Kumaon, North-west Provinces of India” (London).

“*Vespa Lewisii*, Sauss. Guêpes Soc. (MS.)” Smith, 1873, Trans. Ent. Soc. Lond. 1873 (2): 198. Nomen nudum.

*Vespa “lewisii*, Saus. MS.” Cameron, 1903, Entomologist 36: 280, as a synonym of *Vespa japonica* de Saussure, 1858. Available under Art. 11(e) of the International Code of Zoological Nomenclature.

*Vespa Saussurei* Schulz, 1906, Spolia Hym.: 231. Replacement name for *Vespa japonica* de Saussure.

*Vespa karenkona* Sonan, 1929, Trans. Nat. Hist. Soc. Formosa. 19 (101): 137 (key), 148, female – [Taiwan] “Karenkô (3000–6500 ft.)” (Taichung).

*Vespa 4-maculata* Sonan, 1929, Trans. Nat. Hist. Soc. Formosa. 19 (101): 137 (key), 148, female – [Taiwan] “Arisan” (coll. M. Maki).

*Vespa vulgaris* var. *flavitor* Stolfa, 1934, Bull. Soc. Venez. Stor. Nat. 1: 49, female – “India: Bagarkote 8000 piedi. Kumaon W. Himalaia . . . Dal, sopra Dharmasala, 5500 piedi Punjab . . . Pindi Point, Murree 7242 piedi. Punjab . . . Rive del fiume Deo Gad e Foresta S. O. di Pinath, 7 miglia da Kausani Distr. Almora” (type depository unknown).

*Vespula japonica pionganensis* Giordani Soika, 1976, Ann. Hist.-Nat. Mus. Natl. Hung. 68: 287, 290, female (in subgenus *Vespula*) – [Korea] “Za-mo san, 60 km NE from Pyongyang” (Budapest); also from other localities in Korea.

*Vespa vulgaris* var. *flavitor*; Das and Gupta, 1984 (1983), Orient. Ins. 17: 449. Incorrect spelling of *Vespa vulgaris* var. *flavitor* Stolfa.

*Vespula gracilia* Lee, 1986, Sinozool. 10 (4): 201, 203, 206, fig. 2, female – [China] “Zhoushan, Zhejiang Province” (Beijing); also from Fujian.

Distribution: India: Kashmir, Himachal Pradesh, Punjab, Uttar Pradesh, Sikkim, Assam, Manipur, Meghalaya; Nepal; Myanmar; Thailand; China: Xizang, Yunnan, Shanxi, Sichuan, Heilongjiang, Jilin, Jiangsu, Beijing, Fujian, Zhejiang; Taiwan; Russia: Primor'ye; Korea; Japan, including Amami Is. in Ryukyu Is.

*flavopilosa* Jacobson

*Vespa flavopilosa* Jacobson, 1978, in Jacobson et al., Ann. Ent. Soc. Am. 71: 299, 301 (key), 303, figs. 1a, 2c, ra, 5c, d, j, 8, 9, 17, 22–29, female, male – [U. S. A.] “Ithaca, Tompkins Co., NY” (holotype female Washington); also from elsewhere in NY, and NC, ND, MN, MO, IA, WI, IL, IN, MI, OH, KY, TN, GA, VA, WV, DC, MD, DE, PA, NJ, CT, MA, VT, NH, ME; Canada: Manitoba, Ontario, Nova Scotia.

Distribution: Canada: Man., Ont., Que., Nova Scotia; U. S. A.: ND, SD, MN, NE, IA, MO, WI, IL, MI, IN, OH, NY, PA, VT, NH, ME, MA, CT, NJ, DE, MD, DC, KY, WV, VA, TN, NC, SC, GA.

*germanica* (Fabricius)

?*Vespa maculata* Scopoli, 1763, Ent. Carniol.: 312 – [Slovenia] “Carnioliae” (destroyed?). Junior primary homonym of *Vespa maculata* Linnaeus, 1763.

?*Vespa macularis* Olivier, 1792, Encycl. Méth. Ins. 6: 695. Unjustified emendation of *Vespa maculata* Scopoli.

*Vespa germanica* Fabricius, 1793, Ent. Syst. 2: 256 – [Germany] “Kiliae” (type depository unknown).

Distribution: Norway; Sweden; Denmark; Ireland; U. K.: Scotland, Wales, England; Netherlands; Belgium; Luxembourg; France; Spain including Canary Islands; Portugal: Azores; Morocco; Algeria; Tunisia; Germany; Switzerland; Austria; Malta; Poland; Czech Republic; Hungary; Slovenia; Croatia; Albania; Greece; Belarus; Ukraine; across Russia to Primor'ye; Romania; Bulgaria; Turkey; Georgia; Armenia; Iran; Syria; Israel; Kazakhstan; Uzbekistan; Turkmenistan; Tajikistan; Afghanistan; Pakistan; India: Kashmir; Mongolia; China: Gansu, Qinghai, Hebei, Beijing, Liaoning, Neimenggu, Heilongjiang; Taiwan; Korea. Introduced into Iceland; New Zealand; Australia (New South Wales., Australian Capital Territory, Western Australia, Tasmania); Ascension Island; South Africa; Chile; Argentina (Neuquén, Rio Negro, Chubut); Canada (Ont., Que.); U. S. A. (WA, CA, WI, MI, IN, OH, NY, PA, VT, MA, CT, NJ, DE, MD).

*inxpectata* Eck

*Vespa inexpectata* Eck, 1994, Ent. Abhand. Staat. Mus. Tierk. Dresden 56 (6): 125, female, figs. 1–6, 7 – “W. slope, Popocatepetl, Mx. MEX . . . 10000” (Ithaca); also from two other localities in México, and Michoacan.

Distribution: Mexico: México, Michoacan.

*kingdonwardi* Archer

*Vespa kingdonwardi* Archer, 1981, Kontyû 49 (1): 54, 57 (key), 60, fig. 4C–D, female, male – [China] “TIBET, 28.25 N 97.55 E” (holotype female London); also from other localities in Xizang; Myanmar.

*Vespa hirsuta* Lee, 1986, Sinozool. 10 (4): 201, 204, 206, fig. 4, female – [China] “Markham County, Xizang (Tibet) Aut. Reg., 3300 m” (Beijing).

Distribution: China: Xizang; Nepal; Myanmar.

*koreensis* (Radoszkowski)

*Vespa Koreensis* Radoszkowski, 1887, Horae Soc. Ent. Ross. 21: 432, male, female – “Korée” (lectotype female London). – Archer, 1982, Kontyû 50: 263, 267 (designation of lectotype).

*Vespa germanica* var. *stizoides* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 494 (key), 615, female – “Nord de l’Inde” (holotype London).

?*Vespa orbata* var. *aurulenta* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73 (3): 579, female – “Chine, Mou-pin . . . ; Amur” (Paris, Wien).

*Pseudovespa birulai* Gussakovskii, 1932, Ark. Zool. 24A (10): 52, female – [Russia] “Ussuri-Geb.: Tigrovaja . . . ; Sutshan . . . Sedanka . . . Kreis Chabarowsk . . . Wladiwostok” (Stockholm).

*Vespula koreensis salebrosa* Archer, 1982, Kontyû 50 (2): 261, 267, female – “CHINA: Szechuen, Suifu” (London); also from other localities in China.

*Vespula hainanensis* Lee, 1986, Sinozool. 10 (4): 201, 202, 206, fig. 1, female – [China] “Hainan Island, Guangdong Province” (Beijing).

Distribution: India: Sikkim, Meghalaya; Thailand; Laos; China: Sichuan, Yunnan, Hainan, Jiangxi, Fujian, Hunan, Anhui, Zhejiang, Hebei, Beijing, Liaoning, Heilongjiang; Korea; Russia: Amur, Primor’ye.

*maculifrons* (du Buysson)

*Vespa maculifrons* Harris, 1853, in Hitchcock, Rep. Geol. Mineral. Bot. Zool. Mass., ed. 2: 589. Nomen nudum.

*Vespa maculifrons* “H.” du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 608 – [U. S. A.] “Delaware, Wilmington” (London), as a synonym of *Vespa communis* de Saussure, 1857. Available under Art. 11(e) of the International Code of Zoological Nomenclature.

*Vespa communis* var. *flavida* Sladen, 1918, Ottawa Nat. 32: 71, female – “Common in eastern Canada . . . More southern range.” (Ottawa).

Distribution: Canada: Man., Ont., Que., N. B., Nova Scotia; U. S. A.: MT, WY, CO, NM, ND, SD, MN, NE, IA, KS, MO, OK, AR, TX, LA, WI, IL, MI, IN, OH, NY, PA, VT, NH, ME, MA, CT, NJ, DE, MD, KY, WV, VA, TN, NC, SC, MS, AL, GA, FL; Mexico: Tamaulipas, Hidalgo, Nuevo León.

*nursei* Archer

*Vespula nursei* Archer, 1981, Kontyû 49 (1): 54, 57, fig. 4A–B, female, male – “INDIA, Kashmir, 8–9,000 ft” (holotype female London); also from other localities in Kashmir, Punjab, Himachal Pradesh; China; Philippines.

Distribution: India: Kashmir, Himachal Pradesh, Punjab; China: Fujian, Hebei; Philippines: Luzon.

*orbata* (du Buysson)

*Vespa orbata* du Buysson, 1902, Bull. Soc. Ent. France 1902: 140, female – [India] “Dardjiling” (lectotype Paris); also from Sikkim; China. – Archer, 1982, Kontyû 50: 262, 268 (designation of lectotype).

*Vespa minuta* Dover, 1925 (1924), J. Asiat. Soc. Bengal (N. S.) 20: 304, fig. 9, female – [Myanmar] “Tenasserim” (holotype London). Junior primary homonym of *Vespa minuta* Linnaeus, 1767, and of *Vespa minuta* Lichtenstein, 1796.

Distribution: India: Uttar Pradesh, West Bengal, Assam; Nepal; Myanmar.

*pensylvanica* (de Saussure)

*Vespa pensylvanica* de Saussure, 1857, Stettin. Ent. Ztg. 18: 116, 117, female – “America septentr., in Canada, et in montibus mexicanis” (lectotype from New Mexico, Genève). – Jacobson et al., 1978, Ann. Ent. Soc. Am. 71: 300 (designation of lectotype).

*Vespa occidentalis* Cresson, 1874, Trans. Am. Ent. Soc. 5: 100, female, “male” [worker] – [U. S. A.] “Nevada” (lectotype female Philadelphia); also from NM. Junior primary homonym of *Vespa occidentalis* Olivier, 1792. – Cresson, 1916, Mem. Am. Ent. Soc. 1: 106 (designation of lectotype).

*Vespa pennsylvanica*; Dalla Torre, 1894, Cat. Hym. 9: 152. Incorrect spelling of *Vespa pensylvanica* de Saussure.

*Vespa pensylaniva*; Bequaert, 1932, Ent. Am. (N. S.) 12 (2): 95. Incorrect spelling of *Vespa pensylvanica* de Saussure.

Distribution: Canada: B. C., Alta., Sask., Man.; U. S. A.: WA, OR, CA, NV, ID, MT, WY, CO, UT, AZ, NM, ND, SD, MN, NE, TX, WI; Mexico: Baja California Norte, México, Michoacan. Introduced into U. S. A. (HI).

*rufa* (Linnaeus)

*Vespa rufa* Linnaeus, 1758, Syst. Nat., Ed. 10, 1: 572 – “in Europa” (lectotype female, Linnaean Society, London). – Day, 1979, Biol. J. Linn. Soc. 12: 71 (designation of lectotype).

*Vespa Schrenckii* Radoszkowski, 1861, Horae Soc. Ent. Ross. 1: 84, figs. 4–5 – [Russia] “Amur” (type depository unknown).

*Vespa Sibirica* André, 1884, Spec. Hym. Europe Algér. 2: 599, female, male – [Russia] “Sibérie” (Paris?).

*Vespa schrenki*; Dalla Torre, 1904, Gen. Ins. 19: 66. Incorrect spelling of *Vespa schrenckii* Radoszkowski.

*Vespa rufa* var. *intermedia* du Buysson, 1905 (1904), Ann. Soc. Ent. France 73: 499 (key), 591, female, male – “Mandchourie, région du lac Hanka . . . ; Amérique du Nord: Hudson’s Bay” (Paris, London).

*Vespula rufa grahami* Archer, 1981, Kontyû 50 (1): 54, 62, female, male – “CHINA, Szechuen, 30 miles north of Tatsienku” (Washington); also from elsewhere in Sichuan.

*Vespula obscura* Lee, 1986, Sinozool. 10 (4): 201, 205, 206, fig. 5, female – [China] “Zogang County, Xizang (Tibet) Aut. Reg., 3800 m” (Beijing); also from another locality in Xizang.

?*Vespula yichunensis* Lee, 1986, Sinozool. 10 (4): 202, 203, 206, fig. 3, female – [China] “Yichun City, Heilongjiang Province” (Beijing).

Note: Edwards (1980: 363) listed “*rubra* (Laboulbène, 1848)” as a misspelling of *rufa*.

We have not been able to trace this.

Distribution: Norway; Sweden; Finland; Denmark; Ireland; U. K.: Scotland, Wales, England; Netherlands; Belgium; Luxembourg; France; Spain; Italy; Germany; Switzerland; Austria; Poland; Czech Republic; Croatia; Bulgaria; Albania; Ukraine; across Russia to Kamchatka, Primor’ye, Sakhalin, Kurile Is.; Turkey; Georgia; Uzbekistan; Mongolia; China: Xinjiang, Xizang, Sichuan, Yunnan, Liaoning, Heilongjiang; Nepal; Taiwan; Korea; Japan; Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Que., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, VT.

*shidai* Ishikawa, Yamane and Wagner

*Vespula shidai* Ishikawa, Yamane and Wagner, 1980, in Yamane et al., Ins. Matsum. (n. ser.) 19: 1, 44, fig. 135A; female, male – “Sapporo, Hokkaidō, Japan” (holotype male Sapporo); also from other localities in Japan; and Russia.

*Vespula shidai amamiana* Yamane, 1987, Kontyū 55 (4): 634 (key), 635, female, male – [Japan] “Naze, Amami-ōshima” (holotype female Sapporo); also from elsewhere on Amami, and Kakeroma.

Distribution: China: Liaoning, Heilongjiang; Korea; Russia: Primor’ye, Kurile Is.; Japan, including Amami Is. in Ryukyu Is.

*squamosa* (Drury)

*Vespa squamosus* Drury, 1773, Illustr. Nat. Hist. 2: Index to vol. I [pl. XLIII fig. 7 in Vol. 1, female – “New York”] (destroyed?).

*Vespa lineata* Fabricius, 1775, Syst. Ent.: 365 – “in America” (type depository unknown).

*Sphex conchacea* Christ, 1791, Naturgesch. Ins.: 259, pl. 25 fig. 5 – [U. S. A.] “Neuiork” (destroyed).

*Vespa cuneata* Fabricius, 1804, Syst. Piez.: 258 – [U. S. A.] “in Carolina” (Kobenhavn).

*Vespa cruciata* Lepeletier, 1836, Hist. Nat. Ins. Hym. 1: 514. Unjustified emendation of *Vespa cuneata* Fabricius.

✓ *Vespa bistriata* McFarland, 1888, Trans. Am. Ent. Soc. 15: 297 (key), 298, female – “North America” (Philadelphia). Junior primary homonym of *Vespa bistriata* Fabricius, 1804.

*Vespa macfarlandi* Lewis, 1897, Trans. Am. Ent. Soc. 24: 172 (key), 180. Replacement name for *Vespa bistriata* McFarland.

*Vespula squamosa* var. (or subsp.) *michoacana* Bequaert, 1941, Ent. News 52: 249, female – “Tancitaro, 6600 ft., State of Michoacan, Mexico” (Cambridge).

Distribution: Canada: Ont.; U. S. A.: NE, IA, KS, MO, OK, AR, TX, LA, WI, IL, MI, IN, OH, NY, PA, NJ, DE, MD, KY, WV, VA, TN, NC, SC, MS, AL, GA, FL; Mexico: Guerrero, Oaxaca, Tamaulipas, Vera Cruz, Michoacan, Chiapas, Zacatecas; Guatemala; Sau Luis Potosi.

*structor* (Smith)

*Vespa structor* Smith, 1870, in Horne and Smith, Trans. Zool. Soc. Lond. 7 (3): 191, pl. XXI fig. 12, female – “Binsur, Kumaon, North-west Provinces of India” (London).

*Vespa structrix* Schulz, 1906, Spolia Hym.: 231. Unjustified emendation of *Vespa structor* Smith.

Distribution: India: Kashmir, Himachal Pradesh, Uttar Pradesh, West Bengal, Sikkim, Assam; Nepal; China: Xizang, Sichuan; Myanmar.

*sulphurea* (de Saussure)

✓ *Vespa sulphurea* de Saussure, 1854, Ét. Fam. Vesp. 2: 137, female – [U. S. A.] “La Californie” (London).

Distribution: U. S. A.: OR, CA, NV, AZ; Mexico: Baja California Norte.

*vidua* (de Saussure)

✓ *Vespa vidua* de Saussure, 1854, Ét. Fam. Vesp. 2: 136, female – [U. S. A.] “La Caroline” (Paris).

Distribution: Canada: Ont., Que, N. B., Nova Scotia; U. S. A.: ND, SD, MN, IA, WI, IL, MI, IN, OH, NY, PA, VT, NH, ME, MA, CT, RI, NJ, DE, KY, WV, MD, VA, TN, NC, SC, GA.

*vulgaris* (Linnaeus)

*Vespa vulgaris* Linnaeus, 1758, Syst. Nat., ed. 10, 1: 572 – “in Europa” (lectotype female, Linnaean Society, London). – Day, 1979, Biol. J. Linn. Soc. 12: 77 (designation of lectotype).

*Vespa sexcincta* Panzer, 1799, Fauna Ins. German. 63: 1, fig. 1, male – “Viennae Austr.” (destroyed).

*Vespa communis* de Saussure, 1857, Stettin. Ent. Ztg. 18: 116, 117, female – “America septentr.” (Genève?). Junior primary homonym of *Vespa communis* von Schrank, 1785.

*Vespa alasensis* Packard, 1870, Trans. Chicago Acad. Sci. 2: 27, pl. II fig. 10, female – [U. S. A.] “Alaska” (type depository unknown).

*Vespa westwoodii* Shipp, 1893, Psyche 6: 450 – “N. Amer. Bor.” (holotype female Oxford).

*Vespa westwoodi*; Dalla Torre, 1904, Gen. Ins. 19: 66. Incorrect spelling of *Vespa westwoodii* Shipp.

*Vespa vulgaris* var. *pseudogermanica* Stolfa, 1932, Mem. Soc. Ent. Ital. 11 (1): 26 (key), female – [Italy] “Lago superiore di Weissenfels (Tarvisio) . . . Senoseccchia” (type depository unknown).

*Vespa mixta* “Schenck” Edwards, 1980, Soc. Wasps: 363. Nomen nudum.

Note: Edwards (1980: 363) listed “*flava major* (Merret, 1767)” and “*major* (Schwenkfurt, 1603)” as synonyms of *vulgaris*. Merret’s work actually dates to 1667, and does not contain any such name, rather the words “*flavam*” and “*majorem*” occur in his description on p. 196. But of course any names dated from before 1758 are unavailable anyway (Art. 3).

Distribution: Finland; Norway; Sweden; Denmark; Ireland; U. K.: Scotland, Wales, England; Netherlands; Belgium; Luxembourg; France; Spain; Portugal; Germany; Switzerland; Austria; Italy; Poland; Czech Republic; Hungary; Slovenia; Albania; Greece; Belarus; Romania; Moldova; Bulgaria; Russia from the Kola Peninsula to Primor’ye, Sakhalin; Turkey; Georgia; Kazakhstan; Iran; India: Kashmir; Mongolia; China: Sichuan, Yunnan, Hebei, Beijing, Liaoning, Heilongjiang; Korea; Japan; Canada: Yukon Terr., N. W. T., B. C., Alta., Sask., Man., Ont., Newfoundland, N. B., P. E. I., Nova Scotia; U. S. A.: AK, WA, OR, CA, NV, ID, MT, WY, UT, CO, AZ, NM, ND, SD, MN, NE, IA, KS, WI, IL, MI, IN, OH, NY, PA, VT, NH, ME, MA, CT, RI, NJ, DE, MD, KY, WV, VA, TN, NC, GA; Mexico: Durango, Oaxaca. Introduced into Iceland; New Zealand; Australia (Victoria); U. S. A. (HI).

## Fossil Taxa

### Genus *Paleovespa* Cockerell

*Paleovespa* Cockerell, 1906, Bull. Mus. Comp. Zool. 50 (2): 54, genus (3 species).

Type species *Paleovespa florissantia* Cockerell, 1906, by original designation.

*baltica* Cockerell, 1910 (1909), Schrift. Phys.-Ökon. Ges. Königsberg 50 (1): 5.

Oligocene of Germany.

*florissantia* Cockerell, 1906, Bull. Mus. Comp. Zool. 50 (2): 54.

Miocene of U. S. A.: CO.

*gillettei* Cockerell, 1906, Bull. Mus. Comp. Zool. 50 (2): 55.

Miocene of U. S. A.: CO.

*selecta* Cockerell, 1923, Ent. News 34: 270.

Miocene of U. S. A.: CO.

*scudderri* Cockerell, 1906, Bull. Mus. Comp. Zool. 50 (2): 55.

Miocene of U. S. A.: CO.

*wilsoni* Cockerell, 1915, Proc. Acad. Nat. Sci. Philad. 1914: 640.

Miocene of U. S. A.: CO.

#### Genus *Vespa* Linnaeus

*bilineata* Statz, 1936, Decheniana 93: 281, pl. X fig. 31.

Oligocene of Germany.

*cordifera* Statz, 1936, Decheniana 93: 280, pl. X fig. 30.

Oligocene of Germany.

*crabroniformis* Heer, 1867, Neue Denkschr. Allg. Schweiz. Ges. Gesammt. Naturwiss. 22: 6,

pl. III fig. 15a. Junior primary homonym of *Vespa crabroniformis* Smith, 1852.

Miocene of Croatia.

*nigra* Statz, 1936, Decheniana 93: 282, pl. XI fig. 32. Junior primary homonym of *Vespa*

*nigra* Geoffroy, 1785.

Oligocene of Germany.

#### Unrecognized Taxa

Dalla Torre (1904), in his list of unrecognized species of *Vespa*, attributed several species to Weber (1801), but that author was listing species of Lichtenstein (1796). A proposal to suppress Lichtenstein's names, with a few exceptions, is now pending before the International Commission on Zoological Nomenclature (Kerzhner, 1994). That proposal would eliminate all of the Lichtenstein names listed here.

Dalla Torre (1904) also attributed various names to Fourcroy (1785), but those names are properly attributed to Geoffroy, in Fourcroy. He also attributed two names to Olivier (1792), but that work proposed no new names in *Vespa*. The names should be attributed to Reich (1795), who published a list of names for the Piezata recorded by Olivier (1792).

*Vespa aethale* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 202  
(type depository unknown).

*Vespa bicincta* Villers, 1789, C. Linn. Ent. 3: 280 – “Europa. In Galliae Aust.” (destroyed).  
Non Fabricius, 1781.

*Vespa 2-pustulata* Villers, 1789, C. Linn. Ent. 3: 280 – “Europa” (destroyed).

*Vespa capitata* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 440 – [France] “in Agro Parisiensi”  
(type depository unknown).

*Vespa cayana* Reich, 1795, Mag. Thierr. 1 (3): 131 – [Guyane] “Cayenne” (type depository  
unknown).

- Vespa chrysoptera* Villers, 1789, C. Linn. Ent. 3: 284 – “Europa. In Gallia Australiori” (destroyed).
- Vespa chrysopteraarmata* Villers, 1789, C. Linn. Ent. 3: 284 – “Europa. In Gallia Callidiori” (destroyed).
- Vespa communis* von Schrank, 1785, Neues Mag. Liebhab. Ent. 2 (4): 328 – [Germany] “Berchtesgaden” (type depository unknown).
- Vespa condigna* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201, male, female – [India] “Coromandel” (type depository unknown).
- Vespa conifera* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 202 – “India orientali” (type depository unknown).
- Vespa consimilis* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 – “Sina” (type depository unknown).
- Vespa crassipes* Christ, 1791, Naturgesch. Ins.: 246, pl. 23 fig. 10 (destroyed).
- Vespa cribriformis* Christ, 1791, Naturgesch. Ins.: 237, pl. 22 fig. 4 (destroyed).
- Vespa curvipes* Christ, 1791, Naturgesch. Ins.: 246, pl. 23 fig. 9 (destroyed).
- Vespa daedalea* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “America” (type depository unknown).
- Vespa dentata* Olivier, 1792, Encycl. Méthod. Ins. 6: 674 – “l’Amérique méridionale” (type depository unknown).
- Vespa diadema* Christ, 1791, Naturgesch. Ins.: 223, pl. 19 fig. 7 (destroyed).
- Vespa discincta* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “Sina” (type depository unknown).
- Vespa erythrocephala* Gmelin, 1790, Linné Syst. Nat., ed. 13, 1 (5): 2760 (Mus. Lesk.; destroyed).
- Vespa exotica* Gmelin, 1790, Linné Syst. Nat., ed. 13, 1 (5): 2760 – “extra Europam” (Mus. Lesk.; destroyed).
- Vespa ferruginata* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 – [India] “Coromandel” (type depository unknown).
- Vespa ferruginea* Olivier, 1792, Encycl. Méthod. Ins. 6: 683 – “l’Amérique méridionale” (type depository unknown). Non Gmelin, 1790.
- Vespa florisequa maior* Christ, 1791, Naturgesch. Ins.: 244, pl. 23 fig. 4 (destroyed).
- Vespa florisequa minor* Christ, 1791, Naturgesch. Ins.: 244, pl. 23 fig. 5 (destroyed).
- Vespa fulva* Gmelin, 1790, Linné Syst. Nat., ed. 13, 1 (5): 2757 – “imperio Russico propre Zaritzyn” (type depository unknown).
- Vespa fulvipes* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 439 – [France] “in Agro Parisiens” (type depository unknown).
- Vespa fusus maior* Christ, 1791, Naturgesch. Ins.: 242, pl. 23 fig. 1 (destroyed).
- Vespa fusus minor* Christ, 1791, Naturgesch. Ins.: 243, pl. 23 fig. 2 (destroyed).

- Vespa geniculata* Gravenhorst, 1807, Vergl. Übers.: 276 – Europe (type depository unknown).
- Vespa guttata* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 379 – [France] “in Agro Parisiensi” (type depository unknown).
- Vespa quadrum* Olivier, 1792, Encycl. Méthod. Ins. 6: 694.
- Vespa hirsutissima* Villers, 1789, C. Linn. Ent. 3: 284 – “Europa. In Gallia Aust.” (destroyed).
- Vespa horticola* Müller, 1766, Mélang. Phil. Math. Soc. R. Turin 1762–1765, Misc. Taurin. 3: 196 – [Italy] “Taurinensium” (type depository unknown).
- Vespa horticula*; Dalla Torre, 1894, Cat. Hym. 9: 147. Incorrect spelling of *Vespa horticola* Müller.
- Vespa intersecta* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 439 – [France] “in Agro Parisiensi” (type depository unknown).
- Vespa italicica* Christ, 1791, Naturgesch. Ins.: 221 – [Italy] “Florenz” (destroyed).
- Vespa ligata* Müller, 1764, Faun. Ins. Fridrichsdal.: 74 – “Agri Fridrichsdalensis . . . In parietibus ligneis” (type depository unknown).
- Vespa longicornis* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 433 – [France] “in Agro Parisiensi” (type depository unknown).
- Vespa macrocephala* Christ, 1791, Naturgesch. Ins.: 245, pl. 23 fig. 7 (destroyed).
- Vespa menstrua* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “Sina” (type depository unknown).
- Vespa minima* Poda, 1761, Ins. Mus. Graec.: 109.
- Vespa minuta* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “America” (type depository unknown). Non Linnaeus, 1767.
- Vespa mucronata* von Schrank, 1802, Fauna Boica 2 (2): 355 – [Germany] “um Gern.” (type depository unknown).
- Vespa nigra* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 438 – [France] “in Agro Parisiensi” (type depository unknown).
- Vespa ochropygos* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 202 – “India orientali” (type depository unknown).
- Vespa ochrosticta* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 – [India] “Coromandel” (type depository unknown). Non *Vespa ochrosticta* Lichtenstein, 1796: 203, a possible synonym of *Agelaia fulvofasciata* (Degeer).
- Vespa ocreata* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 202 – “India orientali” (type depository unknown)
- Vespa ochreata* “Weber”; Dalla Torre, 1904, Gen. Ins. 19: 67. Incorrect spelling of *Vespa ocreata* Lichtenstein.
- Vespa ordinata* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 435 – [France] “in Agro Parisiensi” (type depository unknown).
- Vespa seriata* Olivier, 1792, Encycl. Méthod. Ins. 6: 693.
- Vespa Pacicephala* Scopoli, 1772, Ann. Hist.-Nat. 5: 122 – [Hungary] “*Cremnitzium*” (destroyed?).

- Vespa petiolata* von Schrank, 1802, Fauna Boica 2 (2): 358 – [Germany] “auf die Doldenblüthen” (type depository unknown).
- Vespa picipes* Reich, 1795, Mag. Thierr. 1 (3): 132 – [Guyane] “Cayenne” (type depository unknown).
- Vespa pilosella* Gravenhorst, 1807, Vergl. Übers.: 275 – Europe (type depository unknown).
- Vespa pteropoda* Scopoli, 1772, Ann. Hist. Nat. 5: 123 – [Hungary] “Cremnitzium” (destroyed?).
- Vespa quadripunctata* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 436 – [France] “in Agro Parisiensi” (type depository unknown). Non Forskal, 1775.
- Vespa quadriguttata* Olivier, 1792, Encycl. Méthod. Ins. 6: 693.
- Vespa quinquefasciata* Müller, 1766, Mélang. Phil. Math. Soc. R. Turin 1762–1765, Misc. Taurin. 3: 196 – [Italy] “Taurinensium” (type depository unknown).
- Vespa 5-fasciata* von Schrank, 1802, Fauna Boica 2 (2): 357 – [Germany] “Sie besucht die Doldenblüthen” (type depository unknown). Non Müller, 1766.
- Vespa scutellata* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “India orientali” (type depository unknown).
- Vespa sericea* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “Sina” (type depository unknown). Non Olivier, 1792.
- Vespa sessilis* Olivier, 1792, Encycl. Méthod. Ins. 6: 683 – “l’Amérique méridionale” (type depository unknown)
- Vespa 6.maculata* Müller, 1766, Mélang. Phil. Math. Soc. R. Turin 1762–1765, Misc. Taurin. 3: 196 – [Italy] “Taurinensium” (type depository unknown).
- Vespa 6-pustulata* Villers, 1789, C. Linn. Ent. 3: 281 – “Europa. In Gallia Aust.” (destroyed).
- Vespa sinuata* Geoffroy, 1785, in Fourcroy, Ent. Paris 2: 438 – [France] “in Agro Parisiensi” (type depository unknown).
- Vespa sphaerogaster* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 – [India] “Coromandel” (type depository unknown).
- Vespa taenia* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 – [India] “Coromandel” (type depository unknown).
- Vespa Tahitensis* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 (type depository unknown).
- Vespa tenebricosa* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “Sina” (type depository unknown).
- Vespa tesserazonia* von Schrank, 1802, Fauna Boica 2 (2): 357 – [Germany] “auf die Doldenblüthen” (type depository unknown).
- Vespa trizonata* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 202 – “India orientali” (type depository unknown).
- Vespa unicolor* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 203 – “Sina” (type depository unknown).

*Vespa usta* Lichtenstein, 1796, Cat. Mus. Zool. Dit. Hamburgi . . . Sect. Tertia Ins.: 201 – [India] “Coromandel” (type depository unknown).

*Vespa velox* Christ, 1791, Naturgesch. Ins.: 245, pl. 23 fig. 6 (destroyed).

### Acknowledgments

We thank Arnold Menke, Bob Jacobson and Michael Archer for many useful suggestions, Sôichi Yamane for comments, and Chris O’Toole for an assistance in examining the types in Oxford. The present study was partly supported by a grant from the Nederlandse Organisatie voor Wetenschappelijk Onderzoek and Japan Society for the Promotion of Science to JK, which allowed him to stay in the Nationaal Natuurhistorisch Museum in Leiden for a year when he could access much of the old literatures cited in this study. Our thanks are also due to Xuexin Chen for information on the type of *Vespa dybowskii mutata*.

### References

- Archer, M. E. 1981. The Euro-Asian species of the *Vespula rufa* group (Hymenoptera, Vespidae), with descriptions of two new species and one new subspecies. *Kontyû* 49: 54–64.
- Archer, M. E. 1982. A revision of the subgenus *Rugovespula* nov. of the genus *Vespula* (Hymenoptera, Vespidae). *Kontyû* 50: 261–269.
- Archer, M. E. 1989. A key to the world species of the Vespinae (Hymenoptera). *Res. Monogr. College of Ripon & York St. John*, 2. York.
- Archer, M. E. 1991a. The number of species that can be recognized within the genus *Vespa* (Hym., Vespinae). *Ent. Mon. Mag.* 127: 161–164.
- Archer, M. E. 1991b. Taxonomy and bionomics of the *Vespa tropica* group (Hym., Vespinae). *Ent. Mon. Mag.* 127: 225–232.
- Archer, M. E. 1992. The taxonomy of *Vespa crabro* L. and *V. dybowskii* André (Hym., Vespinae). *Ent. Mon. Mag.* 128: 157–163.
- Archer, M. E. 1994a. A phylogenetic study of the species of the genus *Vespa* (Hymenoptera: Vespinae). *Ent. Scand.* 24: 469–478.
- Archer, M. E. 1994b. Taxonomy, distribution and nesting biology of the *Vespa bicolor* group (Hym., Vespinae). *Ent. Mon. Mag.* 130: 149–158.
- Archer, M. E. 1995. Taxonomy, distribution and nesting biology of the *Vespa mandarinia* group (Hym., Vespinae). *Ent. Mon. Mag.* 131: 47–53.
- Archer, M. E. 1997. Taxonomy, distribution and nesting biology of *Vespa affinis* (L.) and *Vespa mocsaryana* du Buysson (Hym., Vespinae). *Ent. Mon. Mag.* 133: 27–38.
- Ashmead, W. H. 1902. Classification of the fossorial, predaceous and parasitic wasps, or the superfamily Vespoidea. Paper no. 6. Family XXVIII. Vespidae. *Can. Ent.* 34: 163–166.
- Bequaert, J. 1930. On the generic and subgeneric divisions of the Vespinae (Hymenoptera). *Bull. Brooklyn Ent. Soc.* 25: 59–70.
- Bequaert, J. 1931. The color forms of the common hornet, *Vespa crabro* Linnaeus. *Konowia* 10 (2): 101–109.
- Bequaert, J. 1932. A tentative synopsis of the hornets and yellow-jackets (Vespinae; Hymenoptera) of America. *Ent. Am. (N. S.)* 12 (2): 71–138.
- Bingham, C. T. 1897. Fauna of British India, including Ceylon and Burma. Hymenoptera I. Wasps and Bees. Taylor and Francis, London.
- Bischoff, H. 1931 (1930). Zur Kenntnis der Gattung *Pseudovespa*. *Sitz.-Ber. Ges. Naturf. Freunde Berl.*: 329–346.
- Blanchard, E. 1845. *Histoire des Insectes* 1. Didot Frères, Paris.
- Blüthgen, P. 1938 (1937). Systematisches Verzeichniss der Faltenwespen Mitteleuropas, Skandanaviens und Englands. *Konowia* 16: 270–295.
- Buysson, R. du. 1905 (1904). Monographie des guêpes ou *Vespa*. *Ann. Soc. Ent. France* 73 (3): 485–556.
- Carpenter, J. M. 1987a. On “The evolutionary genetics of social wasps” and the phylogeny of the Vespinae (Hymenoptera, Vespidae). *Ins. Soc.* 34: 58–64.
- Carpenter, J. M. 1987b. Phylogenetic relationships and classification of the Vespinae (Hymenoptera: Vespidae). *Syst. Ent.* 12: 413–431.
- Carpenter, J. M. 1992. Distances, assumptions and social wasps. *Cladistics* 8: 155–160.
- Carpenter, J. M. 1997. Phylogenetic relationships among European *Polistes* and the evolution of social parasitism (Hymenoptera: Vespidae; Polistinae). *Mém. Mus. Paris* (in press).

- Christ, J. L. 1791. Naturgeschichte, Classification und Nomenclatur der Insekten vom Bienen, Wespen und Ameisengeschlecht. Frankfurt am Main.
- Dalla Torre, K. von. 1904. Vespidae. Genera Insectorum 19: 1–108.
- Eck, R. 1981. Zur Verbreitung und Variabilität von *Dolichovespula norwegica* (Hymenoptera, Vespidae). Ent. Abhand. Staat. Mus. Tierk. Dresden 44: 133–152.
- Eck, R. 1984. Bestimmungsschlüssel für die Arten der Gattung *Dolichovespula* Rohwer, 1916. Ent. Abhand. Staat. Mus. Tierk. Dresden 48: 35–44.
- Eck, R. 1988. Probleme der Statusbestimmung im Artbereich Morphologisch-zoogeographische Untersuchung an *Dolichovespula media* und *Dolichovespula maculata* (Insecta, Hymenoptera, Vespidae). Ent. Abhand. Staat. Mus. Tierk. Dresden 51: 93–141.
- Eck, R. 1993. Die Verbreitung der *Dolichovespula sylvestris* im Vergleich mit den übrigen europäischen Arten der Gattung (Hym.: Vespidae). Mitt. Deutsch. Ges. Allg. Angewand. Ent. 8: 919–921.
- Edwards, R. 1980. Social Wasps. Rentokil Limited, East Grinstead.
- Fourcroy, A. F. de 1785. Entomologia Parisiensis; Sive Catalogus Insectorum quae in Agro Parisiensi reperiuntur. Paris.
- International Commission on Zoological Nomenclature. 1910. Opinion 11. Smithson. Publ. 1938: 17–18.
- International Commission on Zoological Nomenclature. 1924. Opinion 79. Smithson. Misc. Coll. 73 (2): 275–276.
- International Commission on Zoological Nomenclature. 1939. Opinion 136. Opinions I. C. Z. N. 2: 15–19.
- International Commission on Zoological Nomenclature. 1956. Direction 32. Opinions and Declarations I. C. Z. N. 1C: 307–328.
- Kerzhner, I. M. 1994. Case 2862. A. A. H. Lichtenstein's (1796, 1797) *Catalogus musei zoologici . . . Section tertia. Continens Insecta* and D. H. Schneider's (1800) *Verzeichniss einer Parthei Insekten . . .*: proposed suppression, with conservation of some Lichtenstein (1796) names (Insecta and Arachnida). Bull. Zool. Nomencl. 51 (2): 108–115.
- Kim, J. K., T.-Y. Moon and I. B. Yoon. 1994. Systematics of vespine wasps from Korea, I. genus *Vespa* Linnaeus (Vespidae, Hymenoptera). Korean J. Entomol. 24: 107–115.
- Lamarck, J. B. P. A. de Monet. 1801. Système des Animaux sans Vertèbres. Paris.
- Latreille, P. A. 1802. Histoire Naturelle, Générale et Particulière des Crustacés et des Insectes, III. Sonnini's suites à Buffon, Paris.
- Latreille, P. A. 1804. Nouveau Dictionnaire d'Histoire Naturelle, XXIV. Paris.
- Latreille, P. A. 1810. Considérations Générales sur l'Ordre Naturel des Animaux Composant les Classes des Crustacés, des Arachnides, et des Insectes. F. Schoell, Paris.
- Lichtenstein, A. A. H. 1796. Catalogus Musei Zoologici Ditissimi Hamburgi, d III. Februar 1796. Auctionis lege distrahendi. Sectio tertia. Continens Insecta. Hamburg.
- Linnaeus, C. 1758. Systema Naturae, ed. 10, 1. Laurentii Salvii, Holmiae.
- Martin, S. J. 1993. Two nests of the hornet *Vespa wilemani* (Hymenoptera, Vespidae). Jpn. J. Ent. 61 (4): 679–682.
- Matsuura, M. and Sk. Yamane. 1984. Comparative Ethology of the Vespine Wasps. Hokkaido Univ. Press, Sapporo. [in Japanese]
- Matsuura, M. and Sk. Yamane. 1990. Biology of the Vespine Wasps. Springer-Verlag, Berlin [translation of the 1984 book "Comparative Ethology of the Vespine Wasps," published by Hokkaido University Press, Sapporo, in Japanese].
- Merret, C. 1667. Pinax Rerum Naturalium Britannicarum, continens Vegetabilia, Animalia, et Fossilia, in hac Insula reperta inchoatus, ed. 2. T. Roycroft, Londini.
- Miller, C. D. F. 1958. Distributional and nomenclatorial problems in some forms of *Vespula* in North America (Hymenoptera: Vespidae). Proc. Tenth Intl. Congr. Ent., Montreal, 1956, 1: 257–264.
- Miller, C. D. F. 1961. Taxonomy and distribution of Nearctic *Vespula*. Can. Ent. 93, Suppl. 22: 1–52.
- Nelson, G. and N. I. Platnick 1981. Systematics and Biogeography: Cladistics and Vicariance. Columbia Univ. Press, New York.
- Nixon, K. C. and Q. D. Wheeler 1990. An amplification of the phylogenetic species concept. Cladistics 6: 211–223.
- Olivier, A. G. 1792. Catalogue des insectes envoyés de Cayenne, a la Société d'Histoire Naturelle de Paris, par M. Le Blond. Actes Soc. Hist. Nat. Paris 1: 120–125.
- Pekkarinen, A. 1995. Geographic variation and taxonomy of the species of *Dolichovespula* in the boreal zone of the Holarctic region (Hymenoptera, Vespinae). Acta Zool. Fenn. 199: 61–70.
- Reich, G. C. 1795. Kurze Beschreibung verschiedener neuern, oder noch wenig bekannten Thiere, welche Herr Le Blond der naturforschenden Gesellschaft zu Paris aus Cayenne als Geschenk überschickt hat. Magazin des Thierreichs 1 (3): 128–132.
- Rohwer, S. A. 1916. Vespidae. In: Guide to the Insects of Connecticut, Part III. The Hymenoptera, or Wasp-like Insects of Connecticut. Conn. State Geol. Nat. Hist. Surv. Bull. 22: 640–644.

- Schmitz, J. and R. F. A. Moritz. 1990. Mitochondrial DNA variation in social wasps (Hymenoptera: Vespidae). *Experientia* 46: 1068–1072.
- Starr, C. K. 1992. The social wasps (Hymenoptera: Vespidae) of Taiwan. *Bull. Natl. Mus. Nat. Sci.* 3: 93–138.
- Thomson, C. G. 1869. *Opuscula Entomologica* 1: 78–82.
- Wagner, R. E. 1978. The genus *Dolichovespula* and an addition to its known species of North America (Hymenoptera: Vespidae). *Pan-Pacif. Ent.* 54: 131–142.
- Weber, F. 1801. *Observationes entomologicae, continentes novorum quae condidit generum characteres, et nuper detectarum specierum descriptiones.* Kiel.
- Varvio-Aho, S. L., P. Pamilo and A. Pekkarinen. 1984. Evolutionary genetics of social wasps (Hymenoptera, Vespidae, *Vespula*). *Ins. Soc.* 31: 375–386.
- Vecht, J. van der, and J. M. Carpenter. 1990. A catalog of the genera of the Vespidae (Hymenoptera). *Zool. Verhand.*, Leiden 260: 3–62.
- Yamane, Sk. and T. Tano. 1985. Supplements to the *Vespula* fauna of Taiwan (Hymenoptera, Vespidae). *Kontyû* 53: 420–425.
- Yamane, Sk., R. E. Wagner and Sô. Yamane. 1980. A tentative revision of the subgenus *Paravespula* of Eastern Asia (Hymenoptera: Vespidae). *Ins. Matsum. n. ser.* 19: 1–46.
- Zalat, S. M. 1992. The Vespidae (social wasps) of Egypt. *J. Egypt Ger. Soc. Zool. (D, Invert. Parasitol.)* 9: 247–258.

Received 11 April 1997;  
revised 1 September 1997;  
accepted 5 September 1997.

### Index of Vespinae

Species- and genus-group names are indexed. For the species-group names genus names are given in parentheses; when they are regarded as valid in the present study, the generic names are given in boldface; when they are synonyms and/or unrecognized taxa, the generic names of the original descriptions are not boldfaced. For the fossile taxa, "FT" is given, and for unrecognized taxa, "UT".

- acadica* (*Vespula*), 75
- adulterina* (*Dolichovespula*), 57
- aegyptiaca* (*Vespa*), 71
- aethale* (*Vespa*) UT, 82
- affinis* (*Vespa*), 64
- affinis* (*Vespa*), 64
- alascensis* (*Vespa*), 81
- albida* (*Vespa*), 61
- alduini* (*Vespa*), 57, 64
- Allovespula*, 74
- alpicola* (*Dolichovespula*), 58
- altaica* (*Vespa*), 68
- alticincta* (*Vespa*), 65
- amamiana* (*Vespula*), 80
- americana* (*Vespa*), 59, 74
- analis* (*Vespa*), 65
- anglica* (*Vespa*), 63, 68
- annulata* (*Vespa*), 56, 70
- anomala* (*Provespa*), 63
- anthracina* (*Vespa*), 72
- arabica* (*Vespa*), 71
- arborea* (*Vespa*), 75
- arbustum*, 63
- archboldi* (*Vespa*), 65
- arctica* (*Vespa*), 61
- arctica* (*Vespula*), 58
- ardens* (*Vespa*), 73
- arenaria* (*Dolichovespula*), 58
- arisana* (*Vespula*), 75
- asiatica* (*Dolichovespula*), 58
- atropilosa* (*Vespula*), 75
- auraria* (*Vespa*), 66
- aurulenta* (*Vespa*), 78
- austriaca* (*Vespula*), 75
- baileyi* (*Dolichovespula*), 59
- baltica* (*Paleovespa*) FT, 81
- barbouri* (*Vespa*), 66
- barthemyi* (*Provespa*), 64
- basalis* (*Vespa*), 66
- basilis* (*Vespa*), 66
- bavarica* (*Vespa*), 62
- bellicosa* (*Vespa*), 67
- bellona* (*Vespa*), 70
- bicincta* (*Vespa*) UT, 82
- bicolor* (*Vespa*), 67
- bilineata* (*Vespa*) FT, 82
- biloba* (*Vespa*), 75
- bimaculata* (*Vespa*), 64
- binghami* (*Vespa*), 67
- bipustulata* (*Vespa*) UT, 82
- biroi* (*Vespa*), 56, 66
- birulai* (*Pseudovespa*), 78
- birulai* (*Vespa*), 68
- bistriata* (*Vespa*), 80
- borealis* (*Dolichovespula*), 60
- borealis* (*Vespa*), 57, 58, 61, 67, 75
- Boreovespula*, 57
- britannica* (*Vespa*), 60
- campanaria* (*Vespa*), 63
- capitata* (*Vespa*) UT, 82
- caspica* (*Vespa*), 68
- cayana* (*Vespa*) UT, 82
- cebuana* (*Vespa*), 73
- celebensis* (*Vespa*), 57, 73
- chinensis* (*Vespa*), 68
- chrysoptera* (*Vespa*) UT, 83
- chrysopteraarmata* (*Vespa*) UT, 83
- cincta* (*Vespa*), 72
- cinta* (*Vespa*), 72
- citriventris* (*Vespa*), 67
- colchica* (*Vespa*), 58
- communis* (*Vespa*), 81
- communis* (*Vespa*) UT, 83
- conchacea* (*Sphex*), 80
- condigna* (*Vespa*) UT, 83
- conifera* (*Vespa*) UT, 83
- conjugens* (*Dolichovespula*), 60
- consimilis* (*Vespa*) UT, 83
- consobrina* (*Vespula*), 76
- continentalis* (*Vespa*), 65
- cordifera* (*Vespa*) FT, 82
- crabro* (*Vespa*), 67
- crabroniformis* (*Vespa*), 67
- crabroniformis* (*Vespa*) FT, 82
- crassa* (*Vespa*), 60
- crassipes* (*Vespa*) UT, 83
- cribriformis* (*Vespa*) UT, 83
- cruciata* (*Vespa*), 80
- cuneata* (*Vespa*), 80
- curvipes* (*Vespa*) UT, 83
- daedalea* (*Vespa*) UT, 83
- dentata* (*Vespa*) UT, 83

- deusta* (*Vespa*), 72  
*diabolica* (*Vespa*), 58  
*diadema* (*Vespa*) UT, 83  
*discincta* (*Vespa*) UT, 83  
*divergens* (*Vespa*), 57, 73  
*Dolichovespula*, 57  
*dorylloides* (*Vespa*), 63  
*doryloides* (*Vespa*), 63  
*dubowskii* (*Vespa*), 69  
*ducalis* (*Vespa*), 68  
*dybowskii* (*Vespa*), 69
- eisa* (*Vespa*), 66  
*erythrocephala* (*Vespa*) UT, 83  
*esakii* (*Vespa*), 68  
*eulemoides* (*Vespa*), 72  
*exotica* (*Vespa*) UT, 83
- fernaldi* (*Vespa*), 58  
*ferruginata* (*Vespa*) UT, 83  
*ferruginea* (*Vespa*) UT, 83  
*fervida* (*Vespa*), 69  
*flava*, 81  
*flavata* (*Vespa*), 72  
*flaviceps* (*Vespula*), 76  
*flavicincta* (*Vespa*), 60  
*flavida* (*Vespa*), 78  
*flavitor* (*Vespa*), 76  
*flavior* (*Vespa*), 76  
*flavitorsis* (*Vespa*), 73  
*flavitorsus* (*Vespa*), 73  
*flavofasciata* (*Vespa*), 68  
*flavopilosa* (*Vespula*), 77  
*flora* (*Dolichovespula*), 59  
*floresiana* (*Vespa*), 74  
*florisequa* (*Vespa*) UT, 83  
*florissantia* (*Paleovespa*) FT, 81  
*formosana* (*Vespa*), 65  
*frontalis* (*Vespa*), 63  
*fruhstorferi* (*Vespa*), 73  
*fulva* (*Vespa*) UT, 83  
*fulvipes* (*Vespa*) UT, 83  
*fumida* (*Vespa*), 69  
*fusca* (*Vespa*), 71  
*fusus* (*Vespa*), 83
- geeri* (*Vespa*), 60  
*geerii* (*Vespa*), 60  
*geniculata* (*Vespa*) UT, 84  
*germania* (*Vespa*), 67  
*germanica* (*Vespula*), 77  
*gillettei* (*Paleovespa*) FT, 82  
*gracilia* (*Vespula*), 76  
*grahami* (*Vespula*), 79  
*gribodoi* (*Vespa*), 68  
*guttata* (*Vespa*) UT, 84
- haematodes* (*Vespa*), 72  
*hainanensis* (*Vespa*), 65  
*hainanensis* (*Vespula*), 78  
*hirsuta* (*Vespula*), 77  
*hirsutissima* (*Vespa*) UT, 84  
*holsatica* (*Vespa*), 62  
*horticola* (*Vespa*) UT, 84  
*horticula* (*Vespa*) UT, 84
- immaculata* (*Vespa*), 73  
*indica* (*Vespa*), 71  
*indosinensis* (*Vespa*), 65  
*inxspectata* (*Vespula*), 77  
*infernalis* (*Vespa*), 75  
*ingrica* (*Vespula*), 61  
*insularis* (*Vespa*), 66  
*intermedia* (*Vespa*), 79  
*intermedia* (*Vespula*), 58  
*intersecta* (*Vespa*) UT, 84  
*italica* (*Vespa*) UT, 84
- japonica* (*Vespa*), 65, 70, 76  
*jurinei* (*Vespa*), 71
- kamtschatkensis* (*Dolichovespula*), 62  
*karenkona* (*Vespa*), 76  
*karnyi* (*Vespa*), 74  
*kingdonwardi* (*Vespula*), 77  
*koreensis* (*Vespula*), 78  
*kuami* (*Dolichovespula*), 59  
*kuangsiana* (*Vespa*), 66
- lama* (*Dolichovespula*), 59  
*latilineata* (*Vespa*), 70  
*leefmansi* (*Vespa*), 72  
*lewisii* (*Vespa*), 76  
*ligata* (*Vespa*) UT, 84  
*lineata* (*Vespa*), 80  
*lineolata* (*Vespa*), 60  
*loekena* (*Dolichovespula*), 62  
*longicornis* (*Vespa*) UT, 84  
*loochooensis* (*Vespa*), 69  
*luctuosa* (*Vespa*), 69  
*lugubris* (*Vespula*), 60  
*lutea* (*Vespa*), 67  
*luzonensis* (*Vespa*), 69
- macfarlandi* (*Vespa*), 80  
*macrocephala* (*Vespa*) UT, 84  
*Macrovespa*, 64  
*macularis* (*Vespa*), 77  
*maculata* (*Dolichovespula*), 59  
*maculata* (*Vespa*), 77  
*maculifrons* (*Vespula*), 78  
*magnifica* (*Vespa*), 70  
*maior* (*Vespa*) UT, 83  
*major*, 81

- major* (*Vespa*), 67  
*malayana* (*Vespa*), 70  
*mandarina* (*Vespa*), 70  
*mandarinia* (*Vespa*), 70  
*marginata* (*Vespa*), 61  
*matsumurai* (*Vespa*), 68  
*media* (*Dolichovespula*), 59  
*mediozonalis* (*Vespa*), 74  
*megei* (*Vespa*), 73  
*menstrua* (*Vespa*) UT, 84  
*meridionalis* (*Vespa*), 68  
*Metavespula*, 57  
*micado* (*Vespa*), 72  
*michoacana* (*Vespula*), 80  
*minima* (*Vespa*) UT, 84  
*minor* (*Vespa*) UT, 83  
*minuta* (*Vespa*), 78  
*minuta* (*Vespa*) UT, 84  
*mixta* (*Vespa*), 81  
*mocsaryana* (*Vespa*), 70  
*moluccana* (*Vespa*), 65  
*mongolica* (*Vespa*), 71  
*monticola* (*Vespula*), 62  
*montivaga* (*Dolichovespula*), 58  
*mucronata* (*Vespa*) UT, 84  
*multimaculata* (*Vespa*), 57, 70  
*mutata* (*Vespa*), 69
- nagatomii* (*Vespa*), 66  
*negrosensis* (*Vespa*), 69  
*nigra* (*Vespa*) FT, 82  
*nigra* (*Vespa*) UT, 84  
*nigrans* (*Vespa*), 66  
*nigrescens* (*Dolichovespula*), 62  
*nigripennis* (*Vespa*), 64  
*nigrihorax* (*Vespa*), 73  
*nigriventris* (*Vespa*), 65  
*nilotica* (*Vespa*), 71  
*nipponica* (*Dolichovespula*), 62  
*nobilis* (*Vespa*), 70  
*nocturna* (*Provespa*), 64  
*norvegica* (*Vespa*), 61  
*norvegicoides* (*Dolichovespula*), 60  
*norwegica* (*Dolichovespula*), 60  
*nursei* (*Vespula*), 78  
*nyalamensis* (*Dolichovespula*), 59  
*Nyctovespa*, 64
- oberthuri* (*Vespa*), 57, 68  
*obliterata* (*Vespa*), 66  
*obscura* (*Vespula*), 79  
*occidentalis* (*Vespa*), 79  
*ochreata* (*Vespa*) UT, 84  
*ochropygos* (*Vespa*) UT, 84  
*ochrosticta* (*Vespa*) UT, 84  
*ocreata* (*Vespa*) UT, 84  
*omissa* (*Dolichovespula*), 61
- orbata* (*Vespula*), 78  
*ordinata* (*Vespa*) UT, 84  
*orientalis* (*Vespa*), 71
- pacicephala* (*Vespa*) UT, 84  
*pacifica* (*Dolichovespula*), 61  
*Paleovespa*, 81  
*panda* (*Dolichovespula*), 62  
*parallela* (*Vespa*), 65  
*Paravespula*, 74  
*parietum* (*Vespa*), 62  
*pendleburyi* (*Vespa*), 70  
*pennsylvanica* (*Vespa*), 79  
*pensylvanica* (*Vespa*), 79  
*pensylvanica* (*Vespula*), 79  
*peruana* (*Vespa*), 61  
*petiolata* (*Vespa*) UT, 85  
*philippinensis* (*Vespa*), 71  
*picea* (*Vespa*), 64  
*picipes* (*Vespa*) UT, 85  
*pilosella* (*Vespa*), 63  
*pilosella* (*Vespa*) UT, 85  
*pionganensis* (*Vespula*), 76  
*pratensis* (*Vespa*), 67  
*Provespa*, 63  
*pruthii* (*Vespa*), 66  
*pseudogermanica* (*Vespa*), 81  
*pseudosoror* (*Vespa*), 69  
*Pseudovespa*, 74  
*Pseudovespula*, 57  
*pteropoda* (*Vespa*) UT, 85  
*pulchra* (*Vespa*), 68
- quadriguttata* (*Vespa*) UT, 85  
*quadrimaculata* (*Vespa*), 76  
*quadripunctata* (*Vespa*), 71  
*quadripunctata* (*Vespa*) UT, 85  
*quadrum* (*Vespa*) UT, 84  
*quinquefasciata* (*Vespa*) UT, 85
- relecta* (*Paleovespa*) FT, 82  
*rubra*, 79  
*rubricans* (*Vespa*), 56, 72  
*rufa* (*Vespula*), 79  
*rufonigrans* (*Vespa*), 65  
*rufoscutellata* (*Vespa*), 60  
*Rugovespula*, 74
- salebrosa* (*Vespula*), 78  
*saussurei* (*Vespa*), 76  
*saxonica* (*Dolichovespula*), 62  
*scelestia* (*Vespa*), 76  
*schrenckii* (*Vespa*), 79  
*schrenki* (*Vespa*), 79  
*scudderii* (*Paleovespa*) FT, 82  
*scutellata* (*Vespa*) UT, 85  
*semperi* (*Vespa*), 69

- seriata* (*Vespa*) UT, 84  
*sericea* (*Vespa*) UT, 85  
*sessilis* (*Vespa*) UT, 85  
*sexmaculata* (*Vespa*) UT, 85  
*sexcincta* (*Vespa*), 81  
*sexpunctata* (*Vespa*), 72  
*sexpustulata* (*Vespa*) UT, 85  
*shidai* (*Vespula*), 80  
*sibirica* (*Vespa*), 79  
*silvestris* (*Vespa*), 63  
*similis* (*Vespa*), 60  
*simillima* (*Vespa*), 71  
*sinensis* (*Dolichovespula*), 62  
*sinuata* (*Vespa*) UT, 85  
*sladeni* (*Vespula*), 75  
*somalica* (*Vespa*), 71  
*sonani* (*Vespa*), 70  
*soror* (*Vespa*), 72  
*sphaerogaster* (*Vespa*) UT, 85  
*sphinx* (*Vespa*), 65  
*squamosa* (*Vespula*), 80  
*stigma* (*Dolichovespula*), 62  
*stizoides* (*Vespa*), 78  
*structor* (*Vespula*), 80  
*structrix* (*Vespa*), 80  
*sugare* (*Dolichovespula*), 60  
*sulcata* (*Vespa*), 76  
*sulphurea* (*Vespula*), 80  
*sumbana* (*Vespa*), 73  
*sumptuosa* (*Vespa*), 63  
*suprunenkoi* (*Vespa*), 67  
*sylvestris* (*Dolichovespula*), 62
- taenia* (*Vespa*) UT, 85  
*tahitensis* (*Vespa*) UT, 85  
*tartarea* (*Vespa*), 68  
*tenebricosa* (*Vespa*) UT, 85  
*tenebrionis* (*Vespa*), 72  
*tenebrosa* (*Vespa*), 66  
*tesserazonia* (*Vespa*) UT, 85  
*timorensis* (*Vespa*), 74
- tridens* (*Vespa*), 62  
*tridentata* (*Vespa*), 66  
*trimeres* (*Vespa*), 73  
*tripunctata* (*Vespa*), 61, 75  
*trisignata* (*Vespa*), 72  
*trizonata* (*Vespa*) UT, 85  
*tropica* (*Vespa*), 72  
*turcica* (*Vespa*), 71  
*tyrannica* (*Vespa*), 65
- unicolor* (*Vespa*), 72  
*unicolor* (*Vespa*) UT, 85  
*unifasciata* (*Vespa*), 64  
*usta* (*Vespa*) UT, 86
- variabilis* (*Vespa*), 69  
*variana* (*Vespa*), 74  
*velox* (*Vespa*) UT, 86  
*velutina* (*Vespa*), 73  
*Vespa*, 64, 82  
*Vespula*, 74  
*vexator* (*Vespa*), 67  
*vidua* (*Vespula*), 80  
*vivax* (*Vespa*), 74  
*vulgaris* (*Vespula*), 81  
*vulgata* (*Vespa*), 68
- walkeri* (*Vespa*), 69  
*westwoodi* (*Vespa*), 81  
*westwoodii* (*Vespa*), 81  
*wilemani* (*Vespa*), 74  
*wilsoni* (*Paleovespa*) FT, 82
- xanthicincta* (*Dolichovespula*), 63  
*xanthoptera* (*Vespa*), 71  
*xinjiangensis* (*Dolichovespula*), 63
- yichunensis* (*Vespula*), 79
- zavattarii* (*Vespa*), 71  
*zetterstedti* (*Dolichovespula*), 61