

Research Article

Chemical Analysis and Evaluation of Antioxidant, Antimicrobial, and Photoprotective Activities of *Schinopsis brasiliensis* Engl. (Anacardiaceae)

Sarah Raquel Gomes de Lima-Saraiva,^{1,2} Fernanda Granja da Silva Oliveira,¹
Raimundo Gonçalves de Oliveira Junior,¹ Camila de Souza Araújo,¹
Ana Paula de Oliveira,¹ Alessandra Gomes Marques Pacheco,¹
Larissa Araújo Rolim,¹ Elba Lúcia Cavalcanti de Amorim,²
Francine Celise Siqueira César,³ and Jackson Roberto Guedes da Silva Almeida¹

¹Center for Studies and Research of Medicinal Plants, Federal University of Vale do São Francisco, 56304-205 Petrolina, PE, Brazil

²Federal University of Pernambuco, 50670-901, Recife, PE, Brazil

³University of São Paulo, 14040-900, Ribeirão Preto, SP, Brazil

Correspondence should be addressed to Jackson Roberto Guedes da Silva Almeida; jackson.guedes@univasf.edu.br

Received 13 April 2017; Revised 16 August 2017; Accepted 5 September 2017; Published 15 October 2017

Academic Editor: Cristina Garcia-Viguera

Copyright © 2017 Sarah Raquel Gomes de Lima-Saraiva et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Schinopsis brasiliensis Engl. is a native plant of Caatinga which has high concentrations of compounds capable of absorbing ultraviolet light, suggesting its potential application for the development of sunscreen preparations. After its identification and collection, this vegetable drug was submitted to a physicochemical analysis through the preparation of ethanolic extract. The phytochemical screening and analysis of extracts were carried out by high-performance liquid chromatography (HPLC) evaluation. The antioxidant activity of the extract was evaluated by 2,2-diphenyl-1-picrylhydrazyl (DPPH) method and β -carotene bleaching test. Inhibitory hemolytic activity and morphological deformation of erythrocytes induced by H_2O_2 were also demonstrated and the antimicrobial activity was analyzed by the minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) method. For the *in vitro* determination of the sun protection factor (SPF), the spectrophotometric method was used. From the analyses carried out with this species, this plant showed significant results for the antioxidant and antimicrobial activities, as well as sunscreen action. Important flavonoids were identified. These data are an important step for the development of new photoprotective cosmetic with Caatinga species, revealing importance and representing another incentive for the preservation of the species involved and analyzed in the study.

1. Introduction

The Caatinga is an exclusive Brazilian biome that covers approximately 844 km² of the country, which corresponds to 11% of the national territory. This biome is rich in biodiversity since vegetation has adapted to the semiarid climate conditions and currently more than 1500 endemic species have been registered [1]. Due to its geographical location and its proximity to the Equator, the region receives about 3,000 hours of solar rays incidence per year and its

perpendicular incidence allows the region to have marked evapotranspiration indices among other meteorological phenomena. Caatinga also registers high averages of temperature around 26°C [2].

The high solar incidence in this biome interferes directly with the morphology of the vegetation and especially with its chemical composition, since some substances such as flavonoids play a special role in protection against ultraviolet radiation [3].

The sun emits ultraviolet radiation, which is divided according to the wavelengths: UVC (100–290 nm), UVB (290–320 nm), and UVA (320–400 nm). UVC radiation is absorbed by the ozone layer in the stratosphere and represents a low percentage of sunlight on the earth's surface. On the other hand, UVB and UVA radiations are biologically important because they may cause several changes in the skin, such as irregular pigmentation, photoaging, and premalignant and even malignant lesions [4].

The use of sunscreens is an alternative to protect the skin against the incidence of UV radiation. Currently, the development of products with raw materials from natural sources has been reported as a trend due to its good acceptance and the potential activity of natural products [5, 6]; for example, the extracts from the flowers of *Bromelia laciniosa* showed high sun protection factor [7], as well as extracts from the leaves of *Neoglaziovia variegata* [8].

The choice of new photoprotective compounds mainly involves their capacity of ultraviolet light absorption by the chromophores present in the molecules. Generally, structures containing aromatic rings can absorb UV radiation [9]. Thus, there are several natural products with structures containing aromatic rings, such as polyphenols [10].

Several native plant species of Caatinga have high concentrations of these phenolic compounds, such as flavonoids, which are capable of absorbing ultraviolet light and whose absorption spectrum occurs with two peaks, one between 240 and 280 nm and another between 300 and 550 nm. Thus, these plants species may be used for the developments of sunscreens in photoprotective preparations [11]. Therefore, *Schinopsis brasiliensis* Engl. species native to the Caatinga belonging to the family Anacardiaceae, while it was not used popularly as a photoprotective, was selected to study the composition of secondary metabolites potentially applicable for this purpose. Then, the aim of the study was to evaluate the photoprotective effect, antioxidant and antibacterial activity, and chemical analysis of the ethanol extract from the barks of this species.

2. Materials and Methods

2.1. Plant Material. The plant material was collected in the city of Petrolina in May 2013. Voucher number 2362 of the species was coded and deposited at Herbário Vale do São Francisco (HVASF). The botanical material was identified by comparing the samples collected with a voucher of the species deposited at the HVASF.

After collecting the material, the barks of *Schinopsis brasiliensis* Engl. were dried in circulating air oven at 40°C for 72 h and pulverized in a mill.

2.2. Plant Drug Physicochemical Quality Control. Among the characteristics necessary to guarantee the quality of the vegetal material, the recommendations of the Brazilian Pharmacopoeia were followed: foreign material, loss on drying, particle size, total and sulfated ashes, foam index, intumescence index, and hemolytic assay [12].

Additionally, assays were adapted for the determination of pH in aqueous solution, following another study found in the literature [13].

2.3. Extraction. The powdered vegetable material was transferred to a flask and subjected to maceration with 95% ethanol, with three extractions at intervals of 72 h, at room temperature, protected from light. The extractive solution was filtered and concentrated under vacuum, and thus the crude ethanolic extract (Sb-EtOH) was obtained [14].

2.4. Determination of Total Phenolic Compounds and Tannins. To determine the total phenolic content, the Folin-Ciocalteu method was used and the methodology for precipitation by casein followed by Folin-Ciocalteu was used for tannins. The content of tannins corresponds to the difference between the total and residual phenolic contents [15]. To quantify total phenolic content, aliquots of the sample (1.0 mg/mL) were mixed with 5.0 mL of Folin-Ciocalteu reagent (10%, v/v), 10.0 mL of Na₂CO₃ (7.5%, w/v), and 84.0 mL of distilled water. The solution was allowed to stand in the dark for 30 minutes. The absorbance was measured at 760 nm.

To quantify the tannins, aliquots of 10.0 mL of the sample (1.0 mg/mL) were shaken for three hours with 1.0 g of casein. After filtration, the volume was completed to 25.0 mL with distilled water. 2.0 mL aliquots of this filtrate were quantified by the Folin-Ciocalteu method. The correlation equation constructed with tannic acid was $y = 0.0677x + 0.0125$ ($R^2 = 0.9981$). The total phenolic and tannin contents were expressed in milligrams of tannic acid per g of extract (mg EAT/g).

2.5. Determination of Flavonoids. The flavonoid content was determined using the colorimetric method based on the formation of the flavonoid-aluminum complex [16]. 2.0 mL aliquots of the sample (1.0 mg/mL) were mixed with 0.6 mL of glacial acetic acid, 10.0 mL of pyridine (20.0%, v/v), 2.5 mL of aluminum chloride (5.0%, w/v), and 10.9 mL of distilled water.

The solution was allowed to stand in the dark for 30 minutes. At 420 nm, the absorbance of the solution was measured. The calibration curve was $y = 0.0290x + 0.0022$ ($R^2 = 0.9999$). The results were expressed in mg of rutin equivalent per g of extract (mg RE/g).

2.6. HPLC Analysis. For the qualitative and quantitative determination of secondary metabolites in the extract, a liquid chromatograph (Prominence LC-20AT, Shimadzu®, Japan) with a diode array detector (DAD, SPD-M20A) coupled to an LC Solution ChemStation data-processing station was used.

The mobile phase used was composed of 2 solvents: solvent A (ultrapure water and acetic acid, 0.5%) and solvent B (acetonitrile, 100%) with 1 mL/min flow rate and gradient according to Table 1.

The stationary phase was a TSKgel Super-ODS (Supelco®) column C18 maintained at 30°C. The volume of the samples injected was 20 µL, being monitored at 270 and 340 nm. For quantitative determination of the analytical standards apigenin, catechin, epicatechin, and gallic acid (Sigma-Aldrich®, analytical grade with purity higher than 99%) in the samples, analytical curves were obtained at the concentrations of 50,

TABLE 1: Mobile phase gradient used for determination of secondary metabolites.

Time (min)	Solvent A (%)	Solvent B (%)
0.00	100	0
40.00	20	80
50.00	100	0

100, 125, 150, and 200 $\mu\text{g}/\text{mL}$, and the sample was analyzed at 200 $\mu\text{g}/\text{mL}$ in triplicate.

2.7. DPPH Assay. The free radical scavenging activity was measured using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay [17]. After preparation of the samples (1 mg/mL), concentrations of 1, 3, 9, 27, 81, and 243 $\mu\text{g}/\text{mL}$ were then prepared. 2.5 mL of the samples and 1 mL of DPPH (50 $\mu\text{g}/\text{mL}$) were added in glass cuvettes. The solutions were allowed to stand for 30 minutes protected from light, and the samples were measured at 518 nm. Ascorbic acid, BHA, and BHT were used as standards. As a negative control, the DPPH solution was used. The antioxidant activity was calculated from a calibration curve obtained by the percentage of antioxidant activity and expressed as the efficient concentration (EC_{50}), that is, the concentration of the sample needed to reduce the absorbance of the 50% negative control.

2.8. Inhibition of Autooxidation of β -Carotene Assay. The β -carotene bleaching method is based on the loss of the yellow color of β -carotene due to its reaction with radicals formed by linoleic acid oxidation in an emulsion [18]. β -Carotene (2 mg) was dissolved in 10 mL chloroform and linoleic acid (40 mg) and Tween 40 (400 mg) were added to 2 mL of this solution. Chloroform was evaporated under vacuum at 40°C and 100 mL of distilled water was added; then the emulsion was vigorously shaken during two minutes. The emulsion (3.0 mL) was added to a tube containing 0.12 mL of solutions of 1 mg/mL of reference compounds and sample extracts. The reaction was followed by spectrophotometry at $\lambda = 470$ nm and the test emulsion was incubated in a water bath at 50°C for 120 min, when the absorbance was measured again. The results obtained were compared with the reference antioxidants ascorbic acid, BHA, and BHT and expressed as percentage of antioxidant activity (%AA) using the following formula: % antioxidant activity = $[1 - (A_0 - A_t)/(A_0^0 - A_t^0)] \times 100$, where A_0 is the initial absorbance and A_t is the final absorbance measured for the test sample and A_0^0 is the initial absorbance and A_t^0 is the final absorbance measured for the negative control (blank). Tests were carried out in triplicate.

2.9. Inhibitory Hemolytic Activity and Morphological Deformation of Erythrocytes Induced by H_2O_2 . Blood was collected and centrifuged at 1500 RPM for 5 minutes; erythrocytes were separated from the plasma and were washed three times using 10 volumes of 20 mM phosphate-buffered saline (PBS), pH 7.4 [19], obtaining a 10% suspension of erythrocytes. To evaluate the antioxidant potential, gallic acid and ethanolic extract samples were used at the concentrations of 5, 10, 15, 20, and 25 $\mu\text{g}/\text{mL}$ of total phenol equivalents expressed

as gallic acid equivalents. Then, the reaction systems were packed and incubated in an oven at 37°C for 30 minutes. Samples were centrifuged at 2000 RPM for 10 minutes, 500 μL of the supernatant was taken, and 2000 μL of PBS buffer was added. The readings were performed in a spectrophotometer at 410 nm. The percentage of hemolysis was calculated from the ratio between the absorbances of the samples and the negative control.

To evaluate the erythrocyte deformation, a reaction medium containing 50 μL of 10% erythrocytes was used in PBS, 50 μL of standard solution and extract (10 μL) gallic acid equivalents, and 100 μL of 200 mM H_2O_2 solution followed by incubation at 37°C for 30 minutes. Samples were centrifuged at 1500 RPM for 10 minutes. The sample was then fixed with 2% glutaraldehyde and centrifuged again. The slide with the material was prepared and then attached to carbon tape in aluminum stub. Scanning electron microscopy (SEM) analysis was performed after metallization with gold powder for 30 seconds.

2.10. Evaluation of the Antibacterial Activity. The antimicrobial activity was performed by microdilution method [20]. Solutions of 25 mg/mL of the extract were prepared using 20% (v/v) DMSO and 200 μL of this dilution was transferred to the microplate containing 200 μL of Mueller-Hinton broth. Serial dilutions were then performed, resulting in concentrations of 25, 12.5, 6.25, 3.12, 1.56, 0.78, 0.39, and 0.195 mg/mL. Gentamicin was used as a reference at an initial concentration of 1.6 mg/mL, which was diluted to concentrations of 0.8, 0.4, 0.2, 0.1, 0.05, 0.025, and 0.0125 mg/mL. Control of broth sterility and bacterial growth was also performed.

The inoculum containing 5×10^5 CFU/mL (0.5 McFarland scale) was added to each well and the bacterial strains used in this study came from the National Institute of Quality Control in Health (INCQS/FIOCRUZ, Brazil). The microorganisms used were *Bacillus cereus* (ATCC 11778), *Enterococcus faecalis* (ATCC 19433), *Escherichia coli* (ATCC 25922), *Klebsiella pneumoniae* (ATCC 13883), *Pseudomonas aeruginosa* (Clinical Isolate), *Salmonella enterica* (ATCC 10708), *Serratia marcescens* (ATCC 13880), *Shigella flexneri* (ATCC 12022), and *Staphylococcus aureus* (ATCC 25923). The microplates were incubated under aerobic conditions for 18–24 hours at 37°C, when 10 μL of 2% 2,3,5-triphenyltetrazolium chloride (CTT) was added to each well for detection of the color change of CTT from colorless to red, reflecting active bacterial metabolism. To determine the MBC (minimum bactericidal concentration), 10 μL aliquots were withdrawn from each well containing the extracts and transferred to Petri dishes containing Mueller-Hinton agar. The plates were incubated for 24 hours at 37°C. The appearance of bacterial colonies for a given concentration indicates that it was not able to kill 99.9% or more of bacterial inoculum used. The assays were performed in triplicate.

2.11. Determination of the In Vitro Sun Protection Factor (SPF). The photoprotective activity of the extract was measured by determination of the maximum absorption wavelength (λ_{max}) and *in vitro* sun protection factor (SPF). For this, the samples were diluted in absolute ethanol, obtaining

concentrations of 5, 25, 50, and 100 mg/L. Samples were analyzed in triplicate. Subsequently, a spectrophotometer scan was performed at wavelengths between 260 and 400 nm, with intervals of 5 nm. The readings were made using quartz cuvettes, and the ethanol was used as blank [21]. Calculation of SPF was obtained according to the equation proposed by Mansur, and the values of $EE \times I$ are constant [22].

$$SPF_{\text{spectrophotometric}} = CF \times \sum_{290}^{320} EE(\lambda) \times I(\lambda) \times Abs(\lambda), \quad (1)$$

where CF is correction factor (=10); $EE(\lambda)$ is spectrum of the erythematous effect; $I(\lambda)$ is solar intensity spectrum; $Abs(\lambda)$ is absorbance.

2.12. Statistical Analysis. All determinations were performed in triplicate. Values were expressed as means \pm standard deviation and data were considered significantly different at $p < 0.05$. GraphPad Prism® software 5.0 was used, using the one-way ANOVA test, followed by Tukey's test.

3. Results and Discussion

The physicochemical quality control of a plant drug is one of the most important parameters before using the plant material for any purpose, as the quality of a natural plant drug should ensure effectiveness and avoid risks to consumers.

In the physicochemical control, the foreign drug materials are classified into three types: (a) parts of the organism from which the drug derives; (b) any organisms, portions, or products of organisms; and (c) impurities of mineral or organic nature, not inherent in the drug. Within this classification, the plant materials were classified as natural, mineral, or organic impurities [12]. From 25 g of the vegetal drug collected by scaling, 0.3 g of foreign material was separated (or 1.2%). This result is within the limits specified by the Brazilian Pharmacopoeia, since the percentage of foreign elements should not exceed 2% m/m.

The results obtained by the granulometric distribution histogram of the powder of the *Schinopsis brasiliensis* barks demonstrate that the particles of the plant material were found predominantly retained in the 500, 425, and 300 μm mesh, representing, respectively, 42, 12, and 15% of all material (Figure 1).

Following the pharmacopoeia criteria, barks of *Schinopsis brasiliensis* were classified as moderately thick powder [12]. In this context, it should be considered that very fine pulverization is not always viable; there is a risk of the formation of a compact mass, hindering the diffusion of the solvent. Moderately thick powder has advantages for the extraction process, since the adhesion of the particles is difficult, providing better penetration of the solvent, and the use is recommended for most drugs [23].

For the determination of water in plant drugs, the loss on drying method is technically simple and is not applicable when the drug contains volatile substances. The water content is given when a product reaches the thermodynamic equilibrium with air at a given temperature and relative humidity. The amount of water lost by drying was 8.45%. In this context,




FIGURE 1: Granulometric distribution histogram of the barks of *Schinopsis brasiliensis*.

the water content is considered safe once according to the pharmacopoeia acceptable limits range from 8 to 14% [12].

To quantify the inorganic content of the plant drug, total ash/incineration was performed. Thus, the drug calcined at high temperature has all its organic matter transformed into CO_2 , leaving only mineral compounds in the form of ash [12]. Sulphated ash comprises the nonvolatile residue on incineration in the presence of sulfuric acid. In general, the test aims to determine the content of inorganic impurities contained in organic substances. The averages of the percentage of total ashes were 10.56%. For the content of sulfated ash, the average was 14.50%. The maximum values for total and sulfated ashes vary widely according to the monographs of the drugs described in the Brazilian Pharmacopoeia [12]; for example, in the monograph *Eugeniae folium* the maximum limits described for total ashes and sulfated ashes are 11.0% and 14.0%, respectively. According to the literature, these quality parameters may indicate the presence of nonvolatile inorganic residues such as sand and stone, found naturally since they are samples of plant materials [24].

The erythrocytes are the first point of attack by free radicals due to the polyunsaturated lipids that constitute their cell membrane; these cells are anucleated and thus devoid of repair systems [25]. H_2O_2 has a high permeation potential and, with the addition of metal ions, like Fe^{2+} , it promotes the formation of the hydroxyl radical (OH^*), which damages the DNA, RNA, proteins, lipids, and nucleus and mitochondrial membranes [25]. Then, the result of the treatment with the extracts of *Schinopsis brasiliensis* Engl. for the test of inhibitory hemolytic activity and morphological deformation of erythrocytes induced by H_2O_2 was positive, since it protected the hemolysis and the deformation of the erythrocytes.

The Sb-EtOH presented a percentage of 43.84% inhibition of erythrocyte hemolysis induced by H_2O_2 in the concentration of 15 $\mu\text{g}/\text{mL}$ GAE. This result is similar to the value found




FIGURE 2: Morphology of erythrocytes after experimental procedure. (a) Sb-EtOH + H₂O₂ (10 μm); (b) Sb-EtOH + H₂O₂ (2 μm); (c) NaCl 0.9% + H₂O₂ (10 μm); (d) NaCl 0.9% + H₂O₂ (2 μm). Figures are obtained by scanning electron microscopy.

TABLE 2: Percentage of maximum inhibition of erythrocyte hemolysis induced by H₂O₂.

Sample	% maximum inhibition ± MSE
Sb-EtOH	43.84 ± 0.02
Gallic acid	50.68 ± 5.48

MSE: Mean Standard Error.

for the reference substance (gallic acid) at a concentration of 5 μg/mL GAE, as shown in Table 2.

The results in Table 2 show the direct correlation between the total phenol content and the potential for hemolytic inhibition. According to the literature, phenolic compounds at high concentrations have prooxidant potentials and, therefore, the results for high concentrations of the crude extract did not inhibit hemolysis as much, as the increase in total

phenol content induced hemolysis rather than preventing it [25–27].

Regarding the analysis of the erythrocyte morphology, the images obtained by scanning electron microscopy (SEM) show that, in general, the preservation of the erythrocyte morphology by Sb-EtOH did not occur when submitted to attack by the oxidizing agent H₂O₂ (Figure 2). The results obtained may be related to phenolic compounds present in the sample [27].

The results of the total phenol content, flavonoids, tannins, and antioxidant activity of crude extract and standards are presented in the Table 3.

Significant contents of total phenolic compounds, flavonoids, and tannins were found in the extract, with 624.6 ± 0.42 (mg EAT/g), 132.4 ± 1.76 (mg RE/g), and 255.8 ± 2.06 (mg EAT/g), respectively. These results confirm studies previously published with this species [28]. High

TABLE 3: The content of total phenols (FT), flavonoid (F), tannins (T), and antioxidant activity were determined by DPPH (EC_{50}) and β -carotene (% AA).

Sample	Results			Antioxidant activity	
	TP (mgGAE/g)	F (mgRE/g)	T (mgTAE/g)	DPPH (EC_{50})	β -Carotene (% AA)
Sb-EtOH	624.6 \pm 0.42	132.4 \pm 1.76	255.8 \pm 2.06	1.46 \pm 0.07	60.81 \pm 0.67
Ascorbic acid	—	—	—	0.64 \pm 0.22	7.50 \pm 2.12
BHA	—	—	—	1.22 \pm 0.43	97.49 \pm 0.76
BHT	—	—	—	8.16 \pm 7.06	99.10 \pm 1.14

GAE: gallic acid equivalent; RE: rutin equivalent; TAE: tannic acid equivalent. The IC_{50} values were obtained by linear regression with 95% confidence interval. IC_{50} is defined by the concentration sufficient to obtain 50% of the maximum estimated effect at 100%. Values are expressed as mean \pm SD ($n = 3$).




FIGURE 3: Identification of phenolic compounds in the crude ethanolic extract of the barks of *Schinopsis brasiliensis* using HPLC-DAD (270 and 340 nm). (a) Catechin; (b) epicatechin; (c) apigenin; and (d) gallic acid.

levels of phenolic compounds were also found in samples of *Schinopsis brasiliensis* [29], in which 455.81 \pm 50.41 mg TAE/g were tannins and 11.29 \pm 0.94 mg RE/g were flavonoids. The vegetal part used in this study was the barks, while in the literature consulted the leaves were used, which may justify the difference between the results found in the studies.

The Sb-EtOH showed better antioxidant activity than BHT by free radical scavenging method, with a value of IC_{50} of 1.46 \pm 0.07. DPPH reacts with an antioxidant compound, which can donate hydrogen and reduce DPPH, promoting change in color from violet to light yellow; therefore, the lower the IC_{50} , the better the antioxidant activity [17].

In the β -carotene bleaching method, the antioxidant potential of the substance is measured by the ability to sequester the free radical generated during the peroxidation of linoleic acid. The easier its oxidation is, the more it will compete with β -carotene in the reaction with radicals, protecting it. The results obtained with the β -carotene autooxidation inhibition test indicate that the crude ethanolic extract had 60.81%, lower than BHT and BHA but higher than ascorbic acid, since this method is used to evaluate the antioxidant capacity of nonpolar substances. Antioxidant

activity is closely linked to the presence of phenolic compounds, as they have in their structure hydroxyls that can act as electron donor agents. Another structural determinant is the antioxidant capacity of flavonoid attributed to hydroxyls in C4 and C3, which would act to increase the antioxidant potential [30]; these factors contribute to the delocalization of electrons in the aromatic rings, thus allowing the stability of the molecule.

Corroborating the results found, it was possible to identify in the Sb-EtOH the four analytical standards used by HPLC-DAD analysis, by means of the similarity between retention times and absorption spectrum in ultraviolet, as well as to quantify them by the compound majority: (a) catechin in 4 min, (b) epicatechin in 3 min, (c) apigenin in 26 min, and (d) gallic acid in 27 min (Figure 3). The chemical structures of these compounds are shown in Figure 4.

The (+)-catechin and (–)-epicatechin are flavanols, which have a hydroxyl group at the 3-position. The properties of the catechins have been widely cited in the literature and present potent sequestering activity of the peroxy radical about ten times higher than *L*-ascorbate (vitamin C) and β -carotene [31].




FIGURE 4: Structures of the substances detected in Sb-EtOH.

The apigenin is a flavone that has a carbonyl group at the 4-position and a double bond between the 2,3-positions. Gallic acid has several proven properties, such as anti-inflammatory activity and antimutagenic, antitumor, and antioxidant activity [32, 33].

The antimicrobial activity of the Sb-EtOH was evaluated by determining a small amount of substances necessary to inhibit the growth of microorganisms (MIC) and the minimal bactericidal concentration (MBC). The Sb-EtOH inhibited bacterial growth in all tested microorganisms at concentrations ranging from 12.5 to 0.39 mg/mL (Table 4).

Thus, the extract under study showed antibacterial activity, mainly related to inhibition of growth. In the literature, there are studies that prove this activity, such as the work done

with the dry methanolic extract of *Schinopsis brasiliensis* leaf in which it showed good activity against both Gram-positive bacteria and Gram-negative bacteria [25]. The research of new antibiotics in medicinal plants is an important scientific issue due to the increasing incidence of multiple resistances of pathogenic microorganisms to drugs that are currently in clinical use [34].

According to the scanning spectrum of Sb-EtOH (Figure 5), it can be observed that the extract analyzed showed a characteristic spectrum of absorption in UVB, since there was absorbance between the wavelengths of 290 and 320 nm. It was found that an increase in absorbance occurred as concentration increases. Thus, Sb-EtOH has a possible photoprotective potential.

TABLE 4: Antimicrobial activity of the extract of *Schinopsis brasiliensis*.

Bacteria	Sb-EtOH (mg/mL)	
	MIC	MBC
<i>Bacillus cereus</i>	12.5	—
<i>Escherichia coli</i>	12.5	12.5
<i>Enterococcus faecalis</i>	12.5	12.5
<i>Klebsiella pneumoniae</i>	12.5	12.5
<i>Pseudomonas aeruginosa</i>	12.5	—
<i>Serratia marcescens</i>	6.25	12.5
<i>Shigella flexneri</i>	3.12	12.5
<i>Staphylococcus aureus</i>	3.12	12.5
<i>Salmonella enterica</i>	0.39	12.5




FIGURE 5: Spectrophotometric absorption profile of *Schinopsis brasiliensis* extract (260–400 nm).

Flavones such as apigenin have been described in the literature as being chemical protectors that absorb light at shorter wavelengths than those visible to the human eye, protecting plant cells from the damage caused by photooxidation. This class of flavonoids acts protecting cells against excess UVB (280–320 nm) radiation, as they accumulate in the epidermal layers of leaves and stems and absorb light intensely in this region of UVB [35]. In addition, it has been shown that increasing plant exposure to UVB light results in the increased synthesis of flavones and flavonols.

The *in vitro* SPF calculation was determined by the spectrophotometric method [18] using the UVB region. The SPF values of Sb-EtOH [5, 25, 50, and 100 mg/L] are, respectively, 0.21 ± 0.01 , 1.35 ± 0.05 , 2.98 ± 0.27 , and 6.27 ± 0.69 . The results are shown in Figure 6.

According to Resolution RDC 30/12 [36] published by ANVISA (National Health Surveillance Agency), the minimum value of sun protection factor for products used by the Brazilian population is 6 (SPF 6). Thus, the Sb-EtOH fulfills this prerequisite of minimum values for the values of




FIGURE 6: *In vitro* Sun protection factor (SPF) of Sb-EtOH (260–400 nm).

photoprotection. This result can be attributed to the presence of flavonoids in the sample.

Although the test was performed *in vitro*, this method correlates well with the *in vivo* tests because it refers to the absorbance of the substance with the erythemalogenic effect of radiation and light intensity at specific wavelengths between 290 and 320 nm (UVB region).

4. Conclusions

The physicochemical analysis showed that the vegetal drug of *S. brasiliensis* is within the limits specified in the Brazilian Pharmacopoeia. The chemical analysis showed that the ethanolic extract studied contains significant amount of phenolic compounds, specifically flavonoids and tannins. Important flavonoids were identified by HPLC-DAD analysis and this study indicated that this plant extract has potent antioxidant and antimicrobial activities, as well as sunscreen action, important data for the future development of a new photoprotective formulation.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

- [1] Brazilian Ministry of Environment, "Plano de Divulgação do Bioma Caatinga," Brasília, Distrito Federal, Brazil, 2011.
- [2] J. Suassuna, SEMI-ÁRIDO: proposta de convivência com a seca, Fundação Joaquim Nabuco, Brazil, 2002.
- [3] K. R. Markham, K. G. Ryan, S. J. Bloor, and K. A. Mitchell, "An increase in the luteolin: Apigenin ratio in *Marchantia polymorpha* on UV-B enhancement," *Phytochemistry*, vol. 48, no. 5, pp. 791–794, 1998.
- [4] J. D'Orazio, S. Jarrett, A. Amaro-Ortiz, and T. Scott, "UV radiation and the skin," *International Journal of Molecular Sciences*, vol. 14, no. 6, pp. 12222–12248, 2013.

- [5] H. C. Polonini, N. R. Raposo, and M. A. F. Brandão, "Natural Sunscreens for Primary Prevention of Skin Cancer," *Revista de Atenção Primária a Saúde*, vol. 14, no. 2, pp. 216–223, 2011.
- [6] F. C. Siqueira César, F. Carnevale Neto, G. S. Porto, and P. M. Campos, "Patent analysis: a look at the innovative nature of plant-based cosmetics," *Química Nova*, vol. 40, no. 7, pp. 840–847, 2017.
- [7] R. G. de Oliveira-Júnior, C. A. A. Ferraz, G. R. Souza et al., "Phytochemical analysis and evaluation of antioxidant and photoprotective activities of extracts from flowers of *Bromelia laciniosa* (Bromeliaceae)," *Biotechnology and Biotechnological Equipment*, vol. 31, no. 3, pp. 600–605, 2017.
- [8] G. d. Raimundo, R. S. Grasielly, L. G. Amanda et al., "Photoprotective, antibacterial activity and determination of phenolic compounds of *Neoglaziovia variegata* (Bromeliaceae) by high performance liquid chromatography-diode array detector (HPLC-DAD) analysis," *African Journal of Pharmacy and Pharmacology*, vol. 9, no. 22, pp. 576–584, 2015.
- [9] J. Flor, M. R. Davolos, and M. A. Correa, "Protetores solares," *Química Nova*, vol. 30, no. 1, pp. 153–158, 2007.
- [10] C. A. Rice-Evans, N. J. Miller, and G. Paganga, "Structure-antioxidant activity relationships of flavonoids and phenolic acids," *Free Radical Biology and Medicine*, vol. 20, no. 7, pp. 933–956, 1996.
- [11] M. F. Bobin, M. Raymond, and M. C. Martini, "Propriedades de absorção UVA/UVB de produtos naturais," *Cosmetics Toiletries*, vol. 7, pp. 44–50, 1995.
- [12] Brazilian Health Regulatory Agency, *Brazilian Pharmacopoeia*, ANVISA, Brasília, Brazil, 5th edition, 2010.
- [13] F. N. Fonseca, *Technological Development of a Phytoproduct from *Justicia pectoralis* – *Chambá*: Obtainment of Standardized Dry Extract (HPLC-DAD) and Pharmacological Evaluation*, Federal University of Ceará, 2009.
- [14] S. R. G. de Lima-Saraiva, H. C. C. Saraiva, J. C. Silva et al., "Antinociceptive properties and acute toxicity of ethanol extract of *Bromelia laciniosa* Mart. ex Schult. f. (Bromeliaceae)," *Tropical Journal of Pharmaceutical Research*, vol. 13, no. 10, pp. 1659–1666, 2014.
- [15] E. L. C. Amorim, J. E. Nascimento, J. M. Monteiro, T. J. S. Peixoto Sobrinho, T. A. S. Araújo, and U. P. Albuquerque, "A simple and accurate procedure for the determination of tannin and flavonoid levels and some applications in ethnobotany and ethnopharmacology," *Functional Ecosystems and Communities*, vol. 2, no. 1, pp. 88–94, 2008.
- [16] T. J. D. S. P. Sobrinho, C. H. T. P. Da Silva, J. E. Do Nascimento, J. M. Monteiro, U. P. De Albuquerque, and E. L. C. De Amorim, "Validation of spectrophotometric methodology for quantify flavonoid content in *Bauhinia cheilantha* (Bongard) Steudel," *Brazilian Journal of Pharmaceutical Sciences*, vol. 44, no. 4, pp. 683–689, 2008.
- [17] W. Brand-Williams, M. E. Cuvelier, and C. Berset, "Use of a free radical method to evaluate antioxidant activity," *Food Science and Technology Lebensmittel-Wissenschaft Technologie*, vol. 28, no. 1, pp. 25–30, 1995.
- [18] W. Aidi Wannes, B. Mhamdi, J. Sriti et al., "Antioxidant activities of the essential oils and methanol extracts from myrtle (*Myrtus communis* var. *italica* L.) leaf, stem and flower," *Food and Chemical Toxicology*, vol. 48, no. 5, pp. 1362–1370, 2010.
- [19] T. K. Girish, P. Vasudevaraju, and U. J. S. Prasada Rao, "Protection of DNA and erythrocytes from free radical induced oxidative damage by black gram (*Vigna mungo* L.) husk extract," *Food and Chemical Toxicology*, vol. 50, no. 5, pp. 1690–1696, 2012.
- [20] T. G. Santos, R. A. Rebelo, E. M. Dalmarco et al., "Chemical composition and antimicrobial activity of leaf essential oil from *Piper malacophyllum* (C. Presl.) c. DC.," *Química Nova*, vol. 35, no. 3, pp. 477–481, 2012.
- [21] I. M. P. Violante, I. M. Souza, C. L. Venturini, A. F. S. Ramalho, R. A. N. Santos, and M. Ferrari, "Avaliação *in vitro* da atividade fotoprotetora de extratos vegetais do cerrado de Mato Grosso," *Brazilian Journal of Pharmacognosy*, vol. 19, no. 2a, pp. 452–457, 2009.
- [22] J. S. Mansur, M. V. R. Breder, M. C. A. Mansur, and R. D. Azulay, "Determinação do fator de proteção solar por espectrofotometria," *Anais Brasileiros de Dermatologia*, vol. 61, pp. 121–124, 1986.
- [23] N. Sharapin, *Fundamentos de tecnologia de produtos fitoterápicos*, CAB e CYTED Colômbia, 2000.
- [24] C. M. Sim, E. P. Schenkel, G. Gosmann et al., *Farmacognosia: da planta ao medicamento*, UFRGS, Florianópolis, Brazil, 2007.
- [25] C. Q. Alves, J. M. David, J. P. David, M. V. Bahia, and R. M. Aguiar, "Métodos para determinação de atividade antioxidante *in vitro* em substratos orgânicos," *Química Nova*, vol. 33, no. 10, pp. 2202–2210, 2010.
- [26] A. L. Pereira, "Efeito dos flavonoides na oxidação da hemoglobina e peroxidação lipídica em eritrócitos normais e deficientes em glicose-6-fosfato desidrogenase," Campinas State University, 1998.
- [27] A. L. B. S. Barreiros, J. M. David, and J. P. David, "Oxidative stress: relations between the formation of reactive species and the organism's defense," *Química Nova*, vol. 29, no. 1, pp. 113–123, 2006.
- [28] C. C. S. Santos, S. S. Araújo, A. L. L. M. Santos et al., "Evaluation of the toxicity and molluscicidal and larvicidal activities of *Schinopsis brasiliensis* Engl. stem bark extract and its fractions," *Brazilian Journal of Pharmacognosy*, vol. 24, no. 3, pp. 298–303, 2014.
- [29] A. M. Saraiva, R. H. A. Castro, R. P. Cordeiro et al., "In vitro evaluation of antioxidant, antimicrobial and toxicity properties of extracts of *Schinopsis brasiliensis* Engl. (Anacardiaceae)," *African Journal of Pharmacy and Pharmacology*, vol. 5, no. 14, pp. 1724–1731, 2011.
- [30] E. J. Lien, S. Ren, H. H. Bui, and R. Wang, "Quantitative structure- activity relationship analysis of phenolic antioxidants," *Free Radical Biology & Medicine*, vol. 26, no. 3-4, pp. 285–94, 1999.
- [31] Y. Yilmaz, "Novel uses of catechins in foods," *Trends in Food Science and Technology*, vol. 17, no. 2, pp. 64–71, 2006.
- [32] P. Grundhöfer, R. Niemetz, G. Schilling, and G. G. Gross, "Biosynthesis and subcellular distribution of hydrolyzable tannins," *Phytochemistry*, vol. 57, no. 6, pp. 915–927, 2001.
- [33] Y. Y. Soong and P. J. Barlow, "Quantification of gallic acid and ellagic acid from longan (*Dimocarpus longan* Lour.) seed and mango (*Mangifera indica* L.) kernel and their effects on antioxidant activity," *Food Chemistry*, vol. 97, no. 3, pp. 524–530, 2006.
- [34] T. C. C. Leite, A. R. De Sena, T. R. Dos Santos Silva, A. K. A. Dos Santos, A. P. T. Uetanabaro, and A. Branco, "Antimicrobial activity of *Marctia* DC species (Melastomataceae) and analysis of its flavonoids by reverse phase-high performance liquid chromatography coupled-diode array detector," *Pharmacognosy Magazine*, vol. 8, no. 31, pp. 209–214, 2012.
- [35] M. M. M. Ferreira, A. H. C. Oliveira, and N. S. Santos, "Flavones and flavonols: new insights into their chemical structure and

biological function,” *Revista Agroambiente*, vol. 2, no. 2, pp. 57–60, 2008.

- [36] Brazilian Health Regulatory Agency, “Approves the Mercosur Technical Regulation on Sunscreens in Cosmetics and other measures”, Resolution RDC no. 30, June 2012. http://portal.anvisa.gov.br/documents/10181/3134554/RDC_30_2012...pdf/c600a91b-f20f-40f9-9c0c-28724725bb21.



Hindawi

Submit your manuscripts at
<https://www.hindawi.com>

