

## Chemical Composition and Evaluation of Antibacterial and Antioxidant Activities of the Essential oil of *Croton urucurana* Baillon (Euphorbiaceae) Stem Bark

Euclésio Simionatto,<sup>\*a</sup> Vanderléa F. L. Bonani,<sup>a</sup> Ademir Farias Morel,<sup>c</sup> Nilva Ré Poppi,<sup>b</sup> Jorge Luiz Raposo Júnior,<sup>b</sup> Caroline Z. Stuker,<sup>c</sup> Gisele M. Peruzzo,<sup>a</sup> Marize T. L. P. Peres<sup>a</sup> and Sônia C. Hess<sup>a</sup>

<sup>a</sup>Departamento de Hidráulica e Transportes, Universidade Federal de Mato Grosso do Sul, 79070-900 Campo Grande-MS, Brazil

<sup>b</sup>Departamento de Química, Universidade Federal de Mato Grosso do Sul, 79070-900 Campo Grande-MS, Brazil

<sup>c</sup>Departamento de Química, Universidade Federal de Santa Maria, 97105-900 Santa Maria-RS, Brazil

O óleo essencial obtido a partir das cascas do caule de *Croton urucurana* Baillon (Euphorbiaceae) foi analisado por CG e CG-EM. 83 compostos foram identificados, sendo borneol (14,7%), acetato de bornila (5,2%), 1-isopropil-7-metil-4-metileno-1,3,4,5,6,8-hexaidro-2H-naftalen-4a-ol (14,7%), sesquicineol (10,5%) e epóxido de  $\gamma$ -gurjuneno (5,4%) os principais componentes. Para o óleo essencial bruto, foi determinada em 3,21 mg mL<sup>-1</sup> a EC<sub>50</sub> no ensaio de atividade antioxidante frente ao radical livre DPPH. A fração do óleo essencial bruto que apresentou atividade antioxidante foi purificada por CCD sobre sílica gel. Análises de CG e CG-EM revelaram que  $\alpha$ -bisabolol (38,3%),  $\alpha$ -eudesmol (9,3%) e guaiol (8,2%) são os principais componentes da fração antioxidante. A EC<sub>50</sub> medida no teste frente ao DPPH foi de 1,05 mg mL<sup>-1</sup> para a fração bioativa. Foi determinada a atividade antimicrobiana do óleo essencial bruto frente a sete bactérias Gram-positivas e Gram-negativas e três fungos. Os valores de CIM medidos variaram de 1,25 a 10,00 mg mL<sup>-1</sup>.

The essential oil obtained from the stem bark of *Croton urucurana* Baillon (Euphorbiaceae) was analysed by GC and GC-MS. 83 compounds were identified and borneol (14.7%), bornyl acetate (5.2%), 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4a-ol (14.7%), sesquiceneole (10.5%) and  $\gamma$ -gurjunene epoxide (5.4%) were the main components. The EC<sub>50</sub> value of the crude essential oil in the DPPH free radical scavenging assay was 3.21 mg mL<sup>-1</sup>. The fraction of the crude essential oil that presented antioxidant activity was purified by prep-TLC on silica gel. GC and GC-MS analysis revealed that  $\alpha$ -bisabolol (38.3%),  $\alpha$ -eudesmol (9.3%) and guaiol (8.2%) were the main components of the antioxidant fraction. The EC<sub>50</sub> value measured for the bioactive oil fraction in the DPPH assay was 1.05 mg mL<sup>-1</sup>. The antimicrobial activity of the crude essential oil was assayed against seven Gram-positive and Gram-negative bacteria and three yeasts. Measured MIC values ranged from 1.25 to 10.00 mg mL<sup>-1</sup>.

**Keywords:** *Croton urucurana*, essential oil, antioxidant, DPPH assay, antimicrobial activity

### Introduction

*Croton* (Euphorbiaceae) is one of the largest genera of flowering plants, with nearly 1,300 species of herbs, shrubs, and trees that are ecologically prominent and often important elements of secondary vegetation in the tropics and subtropics worldwide.<sup>1</sup> Extracts (or infusions) of the

different parts of the plant of several species of the genus *Croton* as *C. lechleri*, *C. palanostigma*, *C. draconoides* and *C. urucurana* are traditional remedies well known for their healing powers and are extensively used by indigenous cultures of the Amazon River for the treatment of infected wounds and to hasten wound healing.<sup>2-8</sup> Its stem bark, when slashed, releases a blood-red sap and, for this reason, *C. urucurana* and other species of the genus *Croton* are known as Dragon's Blood (or Sangra d'água in Brazil).

\*e-mail: eusimionatto@yahoo.com.br

Previous works reported that the essential oils from northeastern Brazilian *Croton* species, *C. zehntneri*, *C. nepetaefolius* and *C. argyrophyloides*, exhibited good antioxidant activities.<sup>9</sup> *Croton urucurana* Baillon (Euphorbiaceae) is a tree commonly found in Paraguay, northern Argentina, southern Brazil, and Uruguay. Three different products from this species are used primarily in folk medicine: the red sap, the stem bark, and the gum exudate.<sup>10</sup>

Many pharmacological effects of *Croton urucurana* have been described in literature, which include wound and ulcer healing, antidiarrhoeic, anticancer, intestinal anti-inflammatory, antioxidant and antirheumatic properties.<sup>11-14,28</sup> Peres *et al.*<sup>15</sup> reported the antibacterial activity against *Staphylococcus aureus* and *Salmonella typhimurium* of the aqueous-EtOH extract, some fractions of the methanolic extract, catechin and acetyl aleuritic acid obtained from *C. urucurana*.

Previous phytochemical analysis of *Croton urucurana* have identified the presence of acetyl aleuritic acid, catechin, gallic acid, gallo catechin, sonderianin,  $\beta$ -sitosterol and its glucoside, stigmasterol, campesterol and two novel clerodane diterpenes.<sup>13,15,16</sup> Fucoarabinogalactan, a polysaccharide, has been isolated from the gum exudate of *C. urucurana*.<sup>17</sup> In this study, we report the chemical composition, antioxidant<sup>29</sup> and antimicrobial activities of the essential oil obtained from the stem bark of *Croton urucurana*.

## Experimental

### Plant Material

The stem bark of *Croton urucurana* was collected in February and March 2006, from biological reserve of Federal University of Mato Grosso do Sul, at the town of Campo Grande, Mato Grosso do Sul state, Brazil. Voucher specimens (16859) have been deposited at the Herbarium of Federal University of Mato Grosso do Sul.

### Essential oil isolation

The stem bark was subjected to hydrodistillation for 4 h using a modified Clevenger-type apparatus, followed by exhaustive extraction of the distillate with hexane. After removal of the solvent, the yield of the crude oil was 0.05%. The physical properties for oil were:  $[\alpha]_D^{25}$ : 0.91;  $[\alpha]_D^{25}$ : 1.56;  $[\alpha]_D^{25}$ : -12.1 (in  $\text{CHCl}_3$ ,  $c = 0.025$ ).

### Gas Chromatography/Mass Spectrometry analysis

Analysis of the oil was performed using a Varian GC-MS-MS system comprising a CP-3900 gas chromatograph

(Walnut Creek, CA, USA) with a 1077 injector, a CP-8410 autosampler and an ion-trap mass spectrometer (Varian Saturn 2100). Separations were carried out using a ZB-5 (5%-phenyl-95%-dimethylpolysiloxane) fused-silica capillary column (30 m  $\times$  0.25 mm i.d., 0.25  $\mu\text{m}$  film thickness) from Phenomenex (Torrance, CA, USA). Oven temperature was programmed from 50 to 250  $^\circ\text{C}$  at 3  $^\circ\text{C min}^{-1}$ . The temperatures of the manifold, GC-MS interface and the ion trap were 70, 240 and 200  $^\circ\text{C}$ , respectively. Helium (99.999%) was used as carrier gas at a constant flow of 1.0  $\text{mL min}^{-1}$  and an injection volume of 1  $\mu\text{L}$  was employed (split ratio of 1:20). The MS scan parameters included electron impact ionization voltage of 70 eV, a mass range of 41-380  $m/z$  and a scan interval of 0.5 s.

A  $\text{C}_9\text{-C}_{21}$  *n*-alkanes mixture diluted in *n*-hexane was prepared for determination of the temperature programmed retention indices. Samples diluted in *n*-hexane were analyzed. Internal standards (*n*-alkanes) were then added to each sample to aid in the standardization of retention times and the samples were analyzed again. Retention indices (RI) for all compounds were determined according to the Van den Dool and Kratz.<sup>27</sup>

### Identification of essential oil constituents

The identification of the components was based on comparison of their mass spectra with those of NIST 2.0 and Saturn Libraries and those described by Adams,<sup>18</sup> as well as by comparison of their retention indices with literature data.<sup>18</sup>

### DPPH assay on TLC

Hydrogen atom- or electron-donation ability of the corresponding oils was measured from the bleaching of the purple-colored methanol solution of 2,2-diphenylpicrylhydrazyl (DPPH). Five microliters of a 1:10 dilution of the oils in hexane were applied to the TLC plates (aluminum sheets covered with silica gel 60 F<sub>254</sub>, Merck) and hexane-ethyl acetate (9:1) mixture was used as eluent. The plate was sprayed with a 0.2% DPPH reagent in methanol and left at room temperature for 30 min. Yellow spots formed from bleaching of the purple colour of DPPH reagent were evaluated as positive antioxidant activity.<sup>19</sup>

### Spectrophotometric DPPH assay

This spectrophotometric assay uses stable DPPH radical as a reagent.<sup>20,21</sup> Fifty microliters of various concentrations of the oils in methanol were added to 5 mL of a 0.004%

methanol solution of DPPH. After a 30 min incubation period at room temperature, the absorbance was read against a blank at 517 nm. Inhibition of DPPH free radical in percent (I%) was calculated in following way:

$$I \% = (A_{\text{blank}} - A_{\text{sample}}) \times 100$$

where  $A_{\text{blank}}$  is the absorbance of the control reaction (containing all reagents except the test compound), and  $A_{\text{sample}}$  is the absorbance of the test sample. Oil concentration providing 50% inhibition ( $IC_{50}$ ) was calculated from the graph plotting inhibition percentage against oil concentration. Tests were carried out in triplicate. Commercial standard antioxidant butylated hydroxytoluene (BHT) was also tested against DPPH and used as a reference.

#### Isolation of antioxidant fraction

For the isolation of the antioxidant compounds, 100 mg of the crude oil were submitted to preparative TLC (silicagel 60 GF<sub>254</sub>, Merck; hexane-EtOAc, 90:10). The antioxidant fraction was detected by application of the DPPH solution (0.2% in ethanol) on the lateral of the plate. This fraction was withdrawn and washed with EtOAc to give the antioxidant fraction (6 mg).

#### Isolation of 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4a-ol (**1**)

100 mg of crude essential oil of *C. urucurana* was further submitted to preparative TLC (SiO<sub>2</sub>; hexane-EtOAc, 85:15) afforded **1** (12 mg). Detection was achieved by UV light (254 nm) and by spraying with solutions of 10% H<sub>2</sub>SO<sub>4</sub> and 2% vanillin in ethanol/H<sub>2</sub>SO<sub>4</sub>, followed by heating. The structure of (**1**) was deduced chiefly from GC-MS analyses and <sup>1</sup>H/<sup>13</sup>C NMR experiments. The relative configuration was additionally confirmed by comparison of the spectral data and retention indices of isomers with those from literature.<sup>25</sup> (**1**):  $[\alpha]_D^{25}$ : + 4.9 (in CHCl<sub>3</sub>,  $c = 0.0075$ ); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  5.52 (brs, 1H), 4.73 (s, 1H), 4.68 (s, 1H), 2.47-2.51 (m, 2H), 2.22-2.26 (m, 2H), 2.01-2.04 (m, 2H), 1.80-1.85 (m, 2H), 1.71-1.74 (m, 1H), 1.67 (m, 1H), 1.59 (m, 1H), 1.22 (s, 3H), 0.98 (d,  $J$  2.7 Hz, 3H), 0.95 (d,  $J$  2.7 Hz, 3H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  21.2, 21.4, 24.0, 24.7, 30.0, 37.0, 37.4, 40.2, 45.0, 55.0, 80.6, 106.4, 121.3, 149.7, 153.9; EI-MS:  $m/z = 220$  [M]<sup>+</sup>, 202, 187, 159, 134, 119, 91.

#### Antimicrobial activity

The antibacterial activity of the crude oil was assayed using the broth micro dilution method. A collection of ten

microorganisms were used, including four Gram-positive bacteria: *Staphylococcus aureus* (ATCC 6538p), *Staphylococcus epidermidis* (ATCC 12228), *Pseudomonas aeruginosa* (ATCC27853) and *Bacillus subtilis* (ATCC 6633), three Gram-negative bacteria: *Klebsiella pneumoniae* (ATCC 10031), *Escherichia coli* (ATCC 11103) and *Salmonella setubal* (ATCC 19796), and three yeasts: *Saccharomyces cerevisiae* (ATCC 2601), *Cryptococcus neoformans* (ATCC 28952) and *Candida albicans* (ATCC 10231). Standard strains of microorganisms were obtained from American Type Culture Collection (ATCC), and standard antibiotics chloramphenicol and nistatine were used in order to control the sensitivity of the microbial test.<sup>22</sup> The minimal inhibitory concentration (MIC) was determined on 96 well culture plates by a micro dilution method using a microorganism suspension at a density of 10<sup>5</sup> CFU mL<sup>-1</sup> with Casein Soy Broth incubated for 24 h at 37 °C for bacteria, and Sabouraud Broth incubated for 72 h at 25 °C for yeasts. The cultures that did not present growth were used to inoculate plates of solid medium (Muller Hinton Agar and Sabouraud Agar) in order to determine the minimal letal concentration (MLC). Proper blanks were assayed simultaneously and samples were tested in triplicate. Technical data have been described previously.<sup>23,24</sup>

## Results and Discussion

In every extractions were used 100 g of *Croton urucurana* stem bark and the crude oil yield was 0.05%. The GC-MS analysis led to the identification and quantification of a total of 83 components (Table 1), accounting for 94.6% of the total components present in the crude essential oil of *C. urucurana*.

As shown in Table 1, borneol (14.7%), bornyl acetate (5.2%), *o*-cymene (3.2 %), terpineol (2.8%) and 1,8-cineole (1.8%) were the main monoterpenes identified in the *C. urucurana* stem bark essential oil. The analysis also showed that the essential oil is constituted by a high proportion of sesquiterpenes, dominated by sesquiceneole (10.5%), 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4a-ol (14.7%),  $\gamma$ -gurjunene epoxide (5.4%),  $\alpha$ -bisabolol (1.8%), elemene (1.8%) and  $\alpha$ -eudesmol (1.5%).

The crude essential oil of *C. urucurana* stem bark (100 mg) was subjected to repeated preparative thin layer chromatography on silicagel to afford the derivative of cadinane-type sesquiterpene 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4a-ol (**1**, 12 mg), that is the main sesquiterpene component of the essential oil. Compound **1** was obtained as an colorless oil and its structure was established on the basis of the <sup>1</sup>H and <sup>13</sup>C

**Table 1.** Percentage composition of the stem bark essential oil from *Croton urucurana* Bailon and of the antioxidant fraction

	Compounds <sup>a,b</sup>	Crude essential oil	Antioxidant fraction	RI <sup>c</sup>	Mass spectral data <sup>d</sup>
01	tricyclene	0.1	-	926	136 [M] <sup>+</sup> , 93
02	$\alpha$ -thujene	0.1	-	931	136 [M] <sup>+</sup> , 91
03	$\alpha$ -fenchene	0.6	-	951	136 [M] <sup>+</sup> , 93
04	sabinene	0.1	-	976	136 [M] <sup>+</sup> , 93
05	myrcene	0.2	-	991	136 [M] <sup>+</sup> , 41
06	$\alpha$ -terpinene	0.1	-	1018	136 [M] <sup>+</sup> , 121
07	<i>o</i> -cymene	3.2	-	1022	134 [M] <sup>+</sup> , 119
08	limonene	0.2	-	1031	136 [M] <sup>+</sup> , 67
09	1,8-cineole	1.8	-	1033	154 [M] <sup>+</sup> , 43
10	$\gamma$ -terpinene	0.3	-	1062	136 [M] <sup>+</sup> , 93
11	terpinolene	0.1	-	1087	136 [M] <sup>+</sup> , 93
12	linalool	0.3	-	1099	154 [M] <sup>+</sup> , 43
13	<i>trans</i> -thujone	0.1	-	1115	152 [M] <sup>+</sup> , 67
14	menth-2-en-1-ol	0.1	-	1120	154 [M] <sup>+</sup> , 43
15	camphor	0.5	-	1142	152 [M] <sup>+</sup> , 95
16	camphene hydrate	0.4	-	1146	154 [M] <sup>+</sup> , 43
17	tagetone	0.1	-	1152	152 [M] <sup>+</sup> , 67
18	isoborneol	0.2	-	1155	154 [M] <sup>+</sup> , 95
19	<i>cis</i> -chrysanthenol	0.2	-	1161	152 [M] <sup>+</sup> , 81
20	borneol	14.7	0.4	1164	154 [M] <sup>+</sup> , 95
21	terpinen-4-ol	2.8	2.5	1175	154 [M] <sup>+</sup> , 71
22	<i>p</i> -cymen-8-ol	0.1	1.0	1183	150 [M] <sup>+</sup> , 43
23	$\alpha$ -terpineol	1.2	-	1189	154 [M] <sup>+</sup> , 59
24	myrtenol	0.1	-	1195	152 [M] <sup>+</sup> , 79
25	isobornyl formate	0.4	-	1227	182 [M] <sup>+</sup> , 95
26	thymol methyl ether	0.2	-	1230	164 [M] <sup>+</sup> , 149
27	<i>E</i> -ocimene	0.1	-	1234	150 [M] <sup>+</sup> , 135
28	bornyl acetate	5.9	-	1285	196 [M] <sup>+</sup> , 43
29	thymol	0.2	2.7	1291	150 [M] <sup>+</sup> , 135
30	carvacrol	0.1	0.8	1300	150 [M] <sup>+</sup> , 135
31	$\delta$ -elemene	0.6	-	1337	204 [M] <sup>+</sup> , 121
32	cyclosativene	0.2	-	1367	204 [M] <sup>+</sup> , 105
33	$\alpha$ -copaene	0.1	-	1375	204 [M] <sup>+</sup> , 105
34	$\beta$ -elemene	0.3	-	1392	204 [M] <sup>+</sup> , 67
35	cyperene	0.3	-	1399	204 [M] <sup>+</sup>
36	$\alpha$ -cedrene	0.2	-	1408	204 [M] <sup>+</sup> , 119
37	$\delta$ -gurjunene	0.3	-	1413	204 [M] <sup>+</sup> , 105
38	$\beta$ -cedrene	1.3	-	1420	204 [M] <sup>+</sup> , 161
39	<i>cis</i> -thujopsene	2.4	-	1424	204 [M] <sup>+</sup> , 119
40	$\gamma$ -elemene	0.5	-	1430	204 [M] <sup>+</sup> , 121
41	geranyl acetone	0.4	-	1452	194 [M] <sup>+</sup> , 43
42	9- <i>epi</i> -caryophyllene	0.2	-	1465	204 [M] <sup>+</sup> , 41
43	$\gamma$ -gurjunene	0.5	-	1470	204 [M] <sup>+</sup> , 105
44	$\gamma$ -himachalene	0.3	-	1474	204 [M] <sup>+</sup> , 105
45	g-murolene	0.4	-	1477	204 [M] <sup>+</sup> , 161
46	curcumene	0.1	-	1483	204 [M] <sup>+</sup> , 119
47	$\beta$ -selinene	0.4	-	1486	204 [M] <sup>+</sup> , 93
48	$\alpha$ -selinene	1.4	-	1493	204 [M] <sup>+</sup> , 93
49	$\beta$ -himalachene	0.3	-	1500	204 [M] <sup>+</sup> , 119
50	cuparene	0.4	-	1505	202 [M] <sup>+</sup> , 132
51	sesquicineole	10.5	-	1514	222 [M] <sup>+</sup> , 139
52	$\delta$ -cadinene	0.8	-	1524	204 [M] <sup>+</sup> , 119
53	<i>trans</i> -calamenene	1.8	0.1	1534	202 [M] <sup>+</sup> , 159
54	$\alpha$ -calacorene	0.1	-	1543	200 [M] <sup>+</sup> , 157
55	nerolidol	0.4	2.4	1564	222 [M] <sup>+</sup> , 41
56	caryophyllene alcohol	1.1	1.5	1573	222 [M] <sup>+</sup> , 111
57	spathulenol	0.3	-	1577	220 [M] <sup>+</sup> , 43
58	sesquisabinene hydrate	0.3	-	1579	222 [M] <sup>+</sup> , 41
59	caryophyllene oxide	0.4	-	1583	220 [M] <sup>+</sup> , 41
60	khusimone	0.6	-	1592	204 [M] <sup>+</sup> , 119
61	elemenone	1.8	-	1597	219 [M] <sup>+</sup> , 107
62	guaial	0.4	8.2	1603	222 [M] <sup>+</sup> , 161

Table 1. cont.

	Compounds <sup>a,b</sup>	Crude essential oil	Antioxidant fraction	RI <sup>c</sup>	Mass spectral data <sup>d</sup>
63	humulane-1,6-dien-3-ol	0.3	-	1606	222 [M] <sup>+</sup> , 109
64	himachalene oxide	0.2	-	1609	220 [M] <sup>+</sup> , 95
65	1,10-di- <i>epi</i> -cubenol	0.8	2.3	1612	222 [M] <sup>+</sup> , 161
66	10- <i>epi</i> - $\gamma$ -eudesmol	0.5	-	1618	222 [M] <sup>+</sup> , 161
67	$\gamma$ -eudesmol	0.7	-	1625	222 [M] <sup>+</sup> , 161
68	1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4-ol*	14.7	5.8	1629	220 [M] <sup>+</sup> , 119
69	$\beta$ -acorenone		0.4	1634	222 [M] <sup>+</sup> , 119
70	cubenol	0.5	1.6	1642	222 [M] <sup>+</sup> , 161
71	vulgarone-B	0.5	4.0	1647	218 [M] <sup>+</sup> , 41
72	$\alpha$ -eudesmol	1.5	9.3	1652	222 [M] <sup>+</sup> , 59
73	$\alpha$ -bisabolol oxide B	-	0.4	1655	238 [M] <sup>+</sup> , 43
74	7- <i>epi</i> - $\alpha$ -eudesmol	-	0.3	1659	222 [M] <sup>+</sup> , 161
75	$\beta$ -bisabolol	-	1.8	1670	222 [M] <sup>+</sup> , 41
76	$\gamma$ -gurjunene epoxide	5.4	0.3	1673	220 [M] <sup>+</sup> , 107
77	khusinol	1.4	2.4	1676	220 [M] <sup>+</sup> , 41
78	$\alpha$ -bisabolol	1.8	38.3	1684	222 [M] <sup>+</sup> , 43
79	<i>epi</i> - $\alpha$ -bisabolol	-	0.6	1685	222 [M] <sup>+</sup> , 43
80	acorenone	0.5	-	1689	220 [M] <sup>+</sup> , 41
81	ni	0.2	-	1707	220 [M] <sup>+</sup> , 119
82	8-oxo-neoisolonlongifolene	0.1	-	1808	218 [M] <sup>+</sup> , 175
83	thunbergol	0.1	-	1825	290 [M] <sup>+</sup> , 43
	TOTAL	94.6	87.1		

<sup>a</sup>Compounds listed in order of elution from a ZB-5 column; <sup>b</sup>Identification: RI, retention indices, GC-MS, gas chromatography-mass spectroscopy; <sup>c</sup>Identified by RI, GC-MS and NMR (nuclear magnetic resonance); <sup>d</sup>Programmed temperature retention indices determined on apolar ZB-5 column (50-250 °C; 3 °C min<sup>-1</sup>); <sup>e</sup>Molecular ion [M]<sup>+</sup> and major fragment obtained from GC-MS analyses.

NMR spectra. This is the first report on the occurrence of **1** in *C. urucurana* and in the Euphorbiaceae family, but this compound has previously been isolated from the essential oil of *Aglaia odorata* Lour (Meliaceae) flowers.<sup>25</sup>

Next, DPPH free radical scavenging activity of crude essential oil was investigated. The model system of scavenging DPPH free radicals is a simple method to evaluate antioxidant activity. It is accepted that the DPPH free radical scavenging by antioxidants is due to their hydrogen-donating ability.<sup>26</sup> The essential oil of *C. urucurana* stem bark exhibited the IC<sub>50</sub> of 3.21 mg mL<sup>-1</sup> in the DPPH free radical scavenging activity spectrophotometric assay. This IC<sub>50</sub> value reveals that the tested essential oil is a weaker oxidant agent than the commercial standard antioxidant BHT, that presented IC<sub>50</sub> 0.18 mg mL<sup>-1</sup> in the same test system. The isolated sesquiterpene 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4a-ol (**1**) showed no antioxidant activity.

Since the crude essential oil of *C. urucurana* stem bark exhibited antioxidant capacity in the DPPH test, it was submitted to prep-TLC, affording the fraction (6.0 mg) that is responsible for the antioxidant activity in the oil. After isolation, this fraction was analyzed by GC-MS for determination of the components. The GC-MS analysis

of the antioxidant fraction of the essential oil of *C. urucurana* led to the identification and quantification of a total of the 21 major components (Table 1) accounting for 87.1% of the total components present and the main compounds are  $\alpha$ -bisabolol (38.3%),  $\alpha$ -eudesmol (9.3%) and guaiol (8.2%). These three compounds were present in the crude essential oil of *C. urucurana* stem bark in very lower yields.

The antioxidant fraction of the crude essential oil exhibited the IC<sub>50</sub> of 1.05 mg mL<sup>-1</sup> in the DPPH free radical scavenging assay. The measured IC<sub>50</sub> indicates that this oil is also a weaker antioxidant agent than BHT.

The antimicrobial activity of the essential oils was examined by broth microdilution susceptibility assay against a panel of 10 microorganisms. The results, presented in Table 2, reveal that the crude essential oil of *C. urucurana* stem bark inhibited the growth of all microorganisms and that *S. epidermidis* and *E. coli* (MIC = 1.25 mg mL<sup>-1</sup>) were the most sensitive, while *B. subtilis* and *C. albicans* were the most resistant microorganisms (MIC = 10 mg mL<sup>-1</sup>).


In comparison to the others essential oils from *Croton* species,<sup>9</sup> the oil of the *C. urucurana* was more complex in composition. The main compounds found in the essential oil from *C. urucurana*, such as borneol, bornyl



**Table 2.** Antimicrobial activity of the essential oil of *C. urucurana* stem bark (minimal inhibitory concentration (MIC) and the minimal lethal concentration (MLC) in mg mL<sup>-1</sup>)

Microorganisms <sup>a</sup>	Essential oil		Standard <sup>b,c</sup>
	MIC <sup>b</sup>	MLC <sup>b</sup>	MIC <sup>b</sup>
<i>Staphylococcus aureus</i>	2.5	>20	3.12 × 10 <sup>-3</sup>
<i>Staphylococcus epidermidis</i>	1.25	>20	3.12 × 10 <sup>-3</sup>
<i>Bacillus subtilis</i>	10	>20	1.56 × 10 <sup>-3</sup>
<i>Pseudomonas aeruginosa</i>	2.5	>20	3.12 × 10 <sup>-3</sup>
<i>Escherichia coli</i>	1.25	>20	3.12 × 10 <sup>-3</sup>
<i>Salmonella setubal</i>	2.5	>20	3.12 × 10 <sup>-3</sup>
<i>Klebsiella pneumoniae</i>	5	>20	1.56 × 10 <sup>-3</sup>
<i>Saccharomyces cerevisiae</i>	5	5	10.3 × 10 <sup>-3</sup>
<i>Candida albicans</i>	10	10	10.3 × 10 <sup>-3</sup>
<i>Cryptococcus neoformans</i>	5	5	5.15 × 10 <sup>-3</sup>

<sup>a</sup>ATCC (American Type Culture Collection); <sup>b</sup>Mean of 3 replicates in mg mL<sup>-1</sup>; <sup>c</sup>Standard antimicrobial agents: chloramphenicol against bacteria and nistatine against yeasts.

**Figure 1.** Structural formula of the derivative of cadinane-type sesquiterpene 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4-ol, main compound identified in the essential oil of stem bark from *Croton urucurana*.

acetate, sesquicineole,  $\gamma$ -gurjunene epoxide and the derivative of cadinane-type sesquiterpene (**1**) did not present in these others species.

## Supplementary Information

Supplementary data are available free of charge at <http://jbcs.sbq.org.br>, as PDF file.

## Acknowledgments

The authors are grateful to Fundação de Apoio ao Desenvolvimento de Ensino, Ciência e Tecnologia de Mato Grosso do Sul (FUNDECT/MS) and UFMS for the financial support. The authors also thank to CNPq for the IC, AT and DCR scholarships, to CAPES for the MS scholarship and Ubirazilda Resende for botanical support.

## References

1. Webster, G. L.; *Taxon* **1993**, *42*, 793.
2. Chen, Z. P.; Cai, Y.; Phillipson, J D.; *Planta Med.* **1994**, *60*, 541.
3. Pieters, L.; De Bruyne, T.; Van Poel, B.; Vingerhoets Totte, J.; Vanden Berghe, D.; Vlietinck, A.; *Phytomedicine* **1995**, *2*, 17.
4. Esmeraldino, L.E.; Souza, A.M.; Sampaio, S.V.; *Phytomedicine* **2005**, *12*, 570.
5. Gabriel, S. E.; Davenport, S. E.; Steagall, R. J.; Viaml, V.; Carlson, T.; Rozhon, R. J.; *Am. J. Physiol. Gastrointest. Liver Physiol.* **1999**, *276*, 58.
6. Holodniy, M.; Koch, J.; Mistal, M.; Schmidt, J. M.; Khandwala, A.; Pennington, J. E.; Porter, S. B.; *Am. J. Gastroenterol.* **1999**, *94*, 3267.
7. Miller, M.J.S.; MacNaughton, W. K.; Zhang, X. -J.; Thompson, J. H.; Charbonnet, R. M.; Bobrowski, P.; Lao, J.; Trentacosti, A.M.; Sandoval, M.; *Am. J. Physiol. Gastrointest. Liver Physiol.* **2000**, *279*, 192.
8. Jones, K.; *J. Altern. Complement. Med.* **2003**, *9*, 877.
9. Moraes, S. M. de; Catunda Júnior, F. E. A.; da Silva, A. R. A.; Neto, J. S. M.; Rondina, D.; Cardoso, J. H. L.; *Quim. Nova* **2006**, *29*, 907.
10. Gonzalez Torres, D. M.; *Catálogo de plantas Medicinales (Alimenticias y Útiles) Usadas en Paraguay.* Asuncion, Paraguay, 1970.
11. Gurgel, L. A.; Sidrim, J. J. C.; Martins, D. T.; Filho, V. C.; Rao, V. S.; *J. Ethnopharmacol.* **2005**, *97*, 409.
12. Gurgel, L. A.; Silva, R. M.; Santos, F. A.; Martins, D. T.; Matos, P. O.; Rao, V. S.; *Phytother. Res.* **2001**, *15*, 319.
13. Peres, M. T. L. P.; Delle Monache, F.; Pizzolatti, M. G.; Santos, A. R. S.; Beirith, A.; Calixto, J. B.; Yunes, R. A.; *Phytother. Res.* **1998**, *12*, 209.
14. Orlandi-Mattos, P. E.; Geremias, R.; Cordova, C. A. S.; Creczynski-Pasa, T. B.; Rebello, J. M.; Wilhem Filho, D.; Martins, D. T. O.; Llesuy, S.; Pedrosa, R. C.; *Free Radical Biol. Med.* **2002**, *33*, 645.
15. Peres, M.T.L.P.; Delle Monache, F.; Cruz, A.B.; Pizzolatti, M.G.; Yunes, R.A.; *J. Ethnopharmacol.* **1997**, *56*, 223.
16. Peres, M.T.L.P.; Pizzolatti, M.G.; Yunes, R.A.; Delle Monache, F.; *Phytochemistry* **1998**, *49*, 171.

17. Milo, B.; Risco, E.; Vila, R., Iglesias, J.; Canigueral, S.; *J. Nat. Prod.* **2002**, *65*, 1143.
18. Adams, R. P.; *Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy*, Allured Publishing: Illinois, 1995.
19. Mimica-Dukic, N.; Bozin, B.; Sokovic, M.; Mihajlovic, B.; Matavulj, B.; *Planta Med.* **2003**, *69*, 413.
20. Burits, M.; Bucar, F.; *Phytother. Res.* **2000**, *14*, 323.
21. Cuendet, M.; Hostettmann, K.; Potterat, O.; *Helv. Chim. Acta* **1997**, *80*, 1144.
22. FDA: *Code of Federal Regulation* **1991**, *21*, 300.
23. Hammer, K.A.; Carson, C.F.; Riley, T.V.; *J. Appl. Microbiol.* **1999**, *86*, 985.
24. National Committee for Clinical Laboratory Standards: *Reference Method of Broth Dilution Antifungal Susceptibility Testing of Yeasts*. Standard M77-T, NCCLS: Villanova, 1995.
25. Weyerstahl, P.; Marschall, H.; Son, P. T.; Giang, P. M.; *Flav. Fragr. J.* **1999**, *14*, 219.
26. Chen, C. W.; Ho, C. T.; *J. Food Lip.* **1995**, *2*, 35.
27. Van den Dool, H.; Kratz, P. D.; *J. Chromatogr.* **1963**, *11*, 463.
28. Salatino, A.; Salatino, M. L. F.; Negri, G.; *J. Braz. Chem. Soc.*, **2007**, *18*, 11.
29. Núñez-Sélles, A. J.; *J. Braz. Chem. Soc.* **2005**, *16*, 699.

Received: September 29, 2006

Web Release Date: July 20, 2007

## Chemical Composition and Evaluation of Antibacterial and Antioxidant Activities of the Essential oil of *Croton urucurana* Baillon (Euphorbiaceae) Stem Bark

Euclésio Simionatto,<sup>\*,a</sup> Vanderléa F. L. Bonani,<sup>a</sup> Ademir Farias Morel,<sup>c</sup> Nilva Ré Poppi,<sup>b</sup>  
Jorge Luiz Raposo Júnior,<sup>b</sup> Caroline Z. Stuker,<sup>c</sup> Gisele M. Peruzzo,<sup>a</sup> Marize T. L. P. Peres<sup>a</sup>  
and Sônia C. Hess<sup>a</sup>

<sup>a</sup>Departamento de Hidráulica e Transportes, Universidade Federal de Mato Grosso do Sul,  
79070-900 Campo Grande-MS, Brazil

<sup>b</sup>Departamento de Química, Universidade Federal de Mato Grosso de Sul,  
79070-900 Campo Grande-MS, Brazil

<sup>c</sup>Departamento de Química, Universidade Federal de Santa Maria,  
97105-900 Santa Maria-RS, Brazil




Figure S1. GC-MS chromatogram of the crude essential oil from stem bark of *Croton urucurana*.

\*e-mail: eusimionatto@yahoo.com.br






Figure S2. GC-MS chromatogram of the antioxidant fraction isolated from stem bark essential oil of *Croton urucurana*.




Figure S3.  $^{13}\text{C}$  NMR spectra (in  $\text{CDCl}_3$ , 75 MHz) of the sesquiterpene (1) isolated from stem bark essential oil of *Croton urucurana*.




Figure S4.  $^1\text{H}$  NMR spectra (in  $\text{CDCl}_3$ , 300 MHz) of the sesquiterpene (1) isolated from stem bark essential oil of *Croton urucurana*.




Figure S5. Mass Spectra of borneol.




Figure S6. Mass Spectra of bornyl acetate.




Figure S7. Mass Spectra of sesquicneote.






Figure S8. Mass Spectra of 1-isopropyl-7-methyl-4-methylene-1,3,4,5,6,8-hexahydro-2H-naphthalen-4-ol (1).

Figure S9. Mass Spectra of  $\alpha$ -bisabolol.Figure S10. Mass Spectra of  $\gamma$ -gurjunene epoxide.