

## Chemiluminescent 2,6-diphenylimidazo[1,2-a]pyrazin-3(7H)-ones: a new entry to *Cypridina* luciferin analogues†

Yuki Ishii, Chihiro Hayashi, Yoshihisa Suzuki and Takashi Hirano\*


An investigation of the chemiluminescent properties of 2,6-diphenylimidazo[1,2-a]pyrazin-3(7H)-one derivatives (**1**), having substituted phenyl groups, is described. Among the derivatives **1**, the 6-[4-(dimethylamino)phenyl] derivatives (**1a,d–f**) gave a high quantum yield ( $\Phi_{CL} \geq 0.0025$ ) in diglyme/acetate buffer, which is a model reaction condition for the *Cypridina* bioluminescence. Their efficient chemiluminescence is mainly caused by the electronic effect of the substituent at C6. In particular, the electron-donating 4-(dimethylamino)phenyl group at C6 of **1a,d–f** plays an essential role in increasing the chemi-excitation efficiency ( $\Phi_S$ ) by the charge transfer-induced luminescence (CTIL) mechanism. The results provide useful information for designing new *Cypridina* luciferin analogues showing efficient chemiluminescence.

Received 28th June 2013,  
Accepted 13th August 2013  
DOI: 10.1039/c3pp50197c  
www.rsc.org/ppps

### Introduction

*Cypridina* luciferin (**Ln**) is the substrate for the bioluminescence of the luminous ostracod *Cypridina* (Scheme 1), and produces light by a reaction with molecular oxygen ( $^3O_2$ ).<sup>1,2</sup> One of the characteristics of the bioluminescence is its high efficiency at light production ( $\Phi_{BL} \approx 0.3$ ).<sup>3</sup> The imidazo[1,2-a]pyrazin-3(7H)-one (imidazopyrazinone) ring is the core for the luminescence reactivity of **Ln**. Imidazopyrazinone derivatives, including **Ln**, exhibit chemiluminescence reactivity with  $^3O_2$  in aprotic solvents.<sup>4,5</sup>

To establish the mechanism of the bioluminescence, the chemiluminescence of imidazopyrazinone derivatives has been investigated as a model of the *Cypridina* bioluminescence reaction.<sup>5–9</sup> For this purpose, 6-aryl-2-methylimidazopyrazinones such as a *Cypridina* luciferin analogue (CLA) and its dimethylamino derivative (**dm-CLA**), whose  $\pi$ -electronic properties were modulated by a substituent at C6, have been employed (Scheme 1).<sup>5c,d,6</sup> The reason to choose 6-aryl imidazopyrazinone derivatives is that the 3-indolyl group at C6 on **Ln** is a  $\pi$ -electronically conjugated appendage. DMSO containing a base (DMSO/base)<sup>4</sup> and diglyme containing an acetate buffer (diglyme/acetate buffer)<sup>5</sup> have been used as the standard reaction conditions for imidazopyrazinone chemiluminescence. The diglyme/acetate buffer condition is predicted to be



**Scheme 1** *Cypridina* luciferin (**Ln**), its analogues CLA, **dm-CLA** and **1a–f**, and *Cypridina* oxyluciferin analogues **dm-OCLA** and **2a–f**.

Department of Engineering Science, Graduate School of Informatics and Engineering, The University of Electro-Communications, Chofu, Tokyo 182-8585, Japan.  
E-mail: hirano@pc.ucc.ac.jp

† In memory of the late Professor Nicholas J. Turro.

a favorable condition for the chemiluminescence reaction of **Ln** because it allows elementary processes similar to those of the bioluminescence reaction.<sup>2a,5</sup> Among the 6-aryl-2-methylimidazopyrazinones investigated, **dm-CLA** having an electron-donating 4-(dimethylamino)phenyl group gives the highest chemiluminescence quantum yield ( $\Phi_{\text{CL}} = 0.015$ ) in diglyme/acetate buffer.<sup>6c,d</sup> A mechanistic analysis of these substituent effects led to the conclusion that the 3-indolyl group of **Ln** plays an essential role as an electron-donating substituent that accelerates the oxygenation process and increases the chemiexcitation efficiency.

Despite the success described above, the chemiluminescence of an imidazopyrazinone still gives an efficiency lower than the  $\Phi_{\text{BL}}$  value of the *Cypridina* bioluminescence. In an attempt to solve this problem, new imidazopyrazinone derivatives **1**, with phenyl groups added at C2 and C6 to electronically modulate the chemiluminescence, were designed as new *Cypridina* luciferin analogues (Scheme 1). It has been found that benzamidopyrazine **2a**, which is the product of the chemiluminescence reaction of **1a**, functions as a fluorophore.<sup>10</sup> This encouraged us to study the chemiluminescence of 2,6-diphenylimidazopyrazinones. We investigated the chemiluminescent properties of a series of substituted 2,6-diphenyl derivatives **1** in DMSO/base and in diglyme/acetate buffer, and found that the derivatives having an electron-donating 4-(dimethylamino)phenyl group at C6 show efficient chemiluminescence in diglyme/acetate buffer. To understand the origin of the efficient chemiluminescence of the 6-[4-(dimethylamino)phenyl] derivatives, the factors that determine the  $\Phi_{\text{CL}}$  values, including the reaction efficiency, the chemiexcitation efficiency and the fluorescence quantum yield, were evaluated mechanistically. We report herein the chemiluminescent properties of **1**, which can provide a guideline for designing new *Cypridina* luciferin analogues with efficient chemiluminescence.

## Experimental

### General

Melting points were obtained using a Yamato MP-21 apparatus. IR spectra were measured using a Horiba FT-720 spectrometer. High-resolution electro-spray ionization (ESI) mass spectra were recorded using a JEOL JMS-T100LC mass spectrometer. <sup>1</sup>H NMR spectra were recorded using a JEOL ECA-500 instrument (500 MHz). UV/visible absorption spectra were measured using a Varian Cary 50 spectrophotometer (scan speed, 600 nm min<sup>-1</sup>; data interval, 1 nm). Fluorescence spectra and fluorescence quantum yields were obtained using a Hamamatsu Photonics Quantaaurus-QY absolute PL quantum yield measurement system. Chemiluminescence spectra and chemiluminescence quantum yields were measured using an ATTO AB-1850 luminescence detecting spectrophotometer. The intensity of the total light (400–700 nm) emitted from a chemiluminescence reaction was monitored using a luminometer with a Hamamatsu R5929 photomultiplier tube powered

by a Hamamatsu C4900 power supply. The signal from the luminometer was collected on a PC and the data were analyzed using the graphics program Igor Pro, Version 4.0 (Wave Metrics, Inc.). Spectroscopic measurements were made in a quartz cuvette (1 cm path length) at 25 ± 1 °C. Spectral-grade solvents were used for the measurement of UV/visible absorption, fluorescence and chemiluminescence. HPLC analyses were carried out using a JASCO 980 HPLC system with a Kanto Chemical Mightysil RP-18 GP-II column (250 mm × 4.6 mm, 5 μm). Density functional theory (DFT) calculations were performed using the Gaussian 09 program.<sup>11</sup> DFT includes Beck's three-parameter function combined with Lee, Yang and Parr's correlation function (B3LYP) along with the 6-31G(d) basis set.<sup>12</sup> Transition state structures were located using the Berny algorithm until the Hessian matrix had only one imaginary eigenvalue. Molecular graphics were generated using GaussView, Version 5.<sup>13</sup>

### Preparation of imidazopyrazinones **1** and amidopyrazines **2**

Imidazopyrazinone **1c** and amidopyrazine **2a** were prepared by previously reported procedures.<sup>10,14</sup> Other derivatives (**1a,b,d-f**) were synthesized from the corresponding 5-arylaminopyrazines and arylglyoxals as follows. Amidopyrazines **2b-f** were also synthesized by acylation of the corresponding 5-arylaminopyrazines.

**Synthesis of 1a.** To a solution of 5-[4-(dimethylamino)phenyl]aminopyrazine (40 mg, 0.20 mmol) and phenylglyoxal monohydrate (92 mg, 0.60 mmol) in 1,4-dioxane (2.0 mL) was added conc. HCl (0.2 mL) under Ar. The reaction mixture was heated at 100 °C for 1 h. After cooling, the mixture was concentrated under reduced pressure. The red residue was purified by silica gel column chromatography four times [CHCl<sub>3</sub>–methanol (20 : 1 and 5 : 1)], yielding **1a** (38 mg, 57%) as a brown powder: mp 256–257 °C.  $\delta_{\text{H}}$  (CD<sub>3</sub>OD) 3.00 (6 H, s), 6.83 (2 H, d, *J* 9.2 Hz), 7.37 (1 H, br t), 7.44 (2 H, br t), 7.52 (2H, d, *J* 8.6 Hz), 7.72 (1 H, br s), 8.00 (1 H, br s) and 8.41 (2 H, d, *J* 5.2 Hz).  $\nu$  (KBr)/cm<sup>-1</sup> 3056, 2927, 1608 and 1516. *m/z* (ESI) Found: 331.1597 ([M + H]<sup>+</sup>). C<sub>20</sub>H<sub>19</sub>N<sub>4</sub>O requires 331.1559.

**1b:** orange powder, mp 89–90 °C.  $\delta_{\text{H}}$  (CD<sub>3</sub>OD) 3.85 (3 H, s), 7.08 (2 H, d, *J* 8.6 Hz), 7.32 (2 H, t, *J* 7.2 Hz), 7.38 (1 H, t, *J* 7.5 Hz), 7.66 (2 H, d, *J* 8.6 Hz), 7.78 (1 H, s), 8.03 (1 H, br s) and 8.41 (2 H, d, *J* 7.5 Hz).  $\nu$  (KBr)/cm<sup>-1</sup> 3411, 3060, 1610 and 1508. *m/z* (ESI) Found: 318.1263 ([M + H]<sup>+</sup>). C<sub>19</sub>H<sub>16</sub>N<sub>3</sub>O<sub>2</sub> requires 318.1243.

**1d:** dark red powder, mp >300 °C.  $\delta_{\text{H}}$  (CD<sub>3</sub>OD) 3.28 (6 H, s), 3.87 (3 H, s), 7.09 (2 H, d, *J* 9.2 Hz), 7.53 (2 H, d, *J* 8.6 Hz), 7.98 (2 H, d, *J* 8.6 Hz), 8.22 (2 H, d, *J* 8.6 Hz), 8.28 (1 H, s) and 8.49 (1 H, s).  $\nu$  (KBr)/cm<sup>-1</sup> 3401, 1608 and 1508. *m/z* (ESI) Found: 361.1698 ([M + H]<sup>+</sup>). C<sub>21</sub>H<sub>21</sub>N<sub>4</sub>O<sub>2</sub> requires 361.1665.

**1e:** dark red powder, mp >300 °C.  $\delta_{\text{H}}$  (CD<sub>3</sub>OD) 3.32 (6 H, s), 3.82 (3 H, s), 3.94 (6 H, s), 7.64 (2 H, s), 7.66 (2 H, d, *J* 9.2 Hz), 8.04 (2 H, d, *J* 8.6 Hz), 8.33 (1 H, s) and 8.54 (1 H, s).  $\nu$  (KBr)/cm<sup>-1</sup> 3401, 1646, 1616 and 1508. *m/z* (ESI) Found: 421.1907 ([M + H]<sup>+</sup>). C<sub>23</sub>H<sub>25</sub>N<sub>4</sub>O<sub>4</sub> requires 421.1876.

**1f:** dark red powder, mp >300 °C.  $\delta_{\text{H}}$  (CD<sub>3</sub>OD) 3.26 (6 H, s), 3.92 (3 H, s), 3.95 (3 H, s), 3.97 (3 H, s), 7.02 (1 H, d, *J* 9.2 Hz), 7.52 (2 H, d, *J* 8.1 Hz), 7.98 (1 H, d, *J* 9.2 Hz), 8.12 (2 H, d, *J* 8.6 Hz) 8.69 (1 H, s) and 8.98 (1 H, s).  $\nu$  (KBr)/cm<sup>-1</sup> 2937, 1585 and 1499. *m/z* (ESI) Found: 421.1875 ([M + H]<sup>+</sup>). C<sub>23</sub>H<sub>25</sub>N<sub>4</sub>O<sub>4</sub> requires 421.1876.

**Synthesis of 2c.** A solution of 5-phenylaminopyrazine (69 mg, 0.41 mmol) and benzoyl chloride (70  $\mu$ L, 0.60 mmol) in pyridine (1 mL) was stirred at room temperature under Ar for 2.5 h. The reaction was quenched by the addition of brine (20 mL) and the product was extracted with chloroform (20 mL  $\times$  3). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by recrystallization from a mixture of CHCl<sub>3</sub> and ethyl acetate (10 : 1), yielding **2a** (50 mg, 45%) as colorless cubes, mp 210–211 °C.  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 7.46 (1 H, t, *J* 7.5 Hz), 7.52 (2 H, t, *J* 8.0 Hz), 7.55 (2 H, t, *J* 8.0 Hz), 7.62 (1 H, t, *J* 7.5 Hz), 7.97 (2 H, d, *J* 7.5 Hz), 8.01 (2 H, d, *J* 7.5 Hz), 8.53 (1 H, br s), 8.72 (1 H, br s) and 9.77 (1 H, d, *J* 0.9 Hz).  $\nu$  (KBr)/cm<sup>-1</sup> 3366, 3053, 1668, 1540 and 1501. *m/z* (ESI) Found: 276.1149 ([M + H]<sup>+</sup>). C<sub>17</sub>H<sub>14</sub>N<sub>3</sub>O requires 276.1137.

**2b:** yellow cubes, mp 189–190 °C.  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 3.88 (3 H, s), 7.02 (2 H, t, *J* 8.6 Hz), 7.53 (2 H, t, *J* 8.0 Hz), 7.61 (1 H, t, *J* 6.9 Hz), 7.95–7.97 (4 H, m), 8.49 (1 H, br s), 8.66 (1 H, d, *J* 1.8 Hz) and 9.71 (1 H, d, *J* 1.8 Hz).  $\nu$  (KBr)/cm<sup>-1</sup> 3369, 1668, 1610 and 1513. *m/z* (ESI) Found: 306.1266 ([M + H]<sup>+</sup>). C<sub>18</sub>H<sub>16</sub>N<sub>3</sub>O<sub>2</sub> requires 306.1243.

**Synthesis of 2d.** A solution of 5-[4-(dimethylamino)phenyl]-aminopyrazine (41 mg, 0.19 mmol), 4-methoxybenzoic acid (29 mg, 0.19 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (109 mg, 0.57 mmol) and 4-dimethylaminopyridine (23 mg, 0.19 mmol) in dichloromethane (2 mL) was heated under reflux under Ar for 3.5 h. The reaction was quenched by the addition of water (60 mL) and the product was extracted with chloroform (70 mL  $\times$  3). The organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by silica gel TLC [CHCl<sub>3</sub>–ethyl acetate (5 : 1)], yielding **2d** (18 mg, 28%) as a yellow powder: mp 219–220 °C.  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 3.04 (6 H, s), 3.89 (3 H, s), 6.81 (2 H, d, *J* 9.2 Hz), 7.00 (2 H, d, *J* 8.6 Hz), 7.92 (4 H, m), 8.35 (1 H, br s), 8.62 (1 H, d, *J* 1.7 Hz) and 9.65 (1 H, d, *J* 1.8 Hz).  $\nu$  (KBr)/cm<sup>-1</sup> 3392, 1670, 1608 and 1540. *m/z* (ESI) Found: 371.1516 ([M + Na]<sup>+</sup>). C<sub>20</sub>H<sub>20</sub>N<sub>4</sub>NaO<sub>2</sub> requires 371.1484.

**2e:** yellow cubes, mp 177–178 °C.  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 3.04 (6 H, s), 3.92 (3 H, s), 3.95 (6 H, s), 6.81 (2 H, d, *J* 8.6 Hz), 7.16 (2 H, s), 7.92 (2 H, d, *J* 8.6 Hz), 8.36 (1 H, br s), 8.64 (1 H, d, *J* 1.2 Hz) and 9.65 (1 H, br s).  $\nu$  (KBr)/cm<sup>-1</sup> 1646, 1610 and 1533. *m/z* (ESI) Found: 431.1732 ([M + Na]<sup>+</sup>). C<sub>22</sub>H<sub>24</sub>N<sub>4</sub>NaO<sub>4</sub> requires 431.1695.

**2f:** yellow cubes, mp 212–213 °C.  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 3.04 (6 H, s), 3.92 (3 H, s), 3.95 (3 H, s), 4.13 (3 H, s), 6.81 (2 H, d, *J* 9.2 Hz), 6.85 (1 H, d, *J* 9.2 Hz), 7.91 (2 H, d, *J* 9.2 Hz), 8.03 (1 H, d, *J* 9.2 Hz), 8.65 (1 H, d, *J* 1.7 Hz), 9.68 (1 H, d, *J* 1.2 Hz) and 10.52 (1 H, br s).  $\nu$  (KBr)/cm<sup>-1</sup> 3332, 2974, 2939 and 1657. *m/z* (ESI) Found: 431.1684 ([M + Na]<sup>+</sup>). C<sub>22</sub>H<sub>24</sub>N<sub>4</sub>NaO<sub>4</sub> requires 431.1695.

## Analysis of the chemiluminescence reactions of imidazopyrazinones

A small portion (20  $\mu$ L) of a stock solution of imidazopyrazinone **1** ( $1.0 \times 10^{-3}$  M) in methanol was added into a quartz cuvette placed in a luminometer and was mixed with aerated DMSO (2.0 mL) containing 0.10 M NaOH aqueous solution (1.0% v/v) or aerated diglyme (2.0 mL) containing 0.10 M of acetate buffer (acetic acid–sodium acetate buffer, pH 5.6, 0.66% v/v) at  $25 \pm 1$  °C. The chemiluminescence reactions of **1** were traced by monitoring the intensity (*I*) of the emitted light. The pseudo-first-order rate constant ( $k_{\text{CL}}$ ) was obtained by analysis using the equation:  $\ln I_t = -k_{\text{CL}}t + \ln I_0$ , where  $I_0$  and  $I_t$  are the intensities at  $t = 0$  and  $t$ , respectively. The  $\Phi_{\text{CL}}$  values were determined as values relative to the  $\Phi_{\text{CL}}$  (0.013) of luminol in DMSO containing *t*-BuOK/*t*-BuOH under air.<sup>15</sup>


The yields of amidopyrazines **2** obtained after chemiluminescence reactions of **1** in diglyme/acetate buffer were determined by HPLC. A spent solution (10 mL) obtained by a chemiluminescence reaction for 24 h was mixed with 2-ethylnaphthalene (1.0  $\mu$ L) as an internal standard, and was analyzed three times by reverse phase HPLC with a H<sub>2</sub>O–CH<sub>3</sub>CN (1 : 4) eluent (1.0 mL min<sup>-1</sup>). Yields of **2** were determined by relative peak integrations (monitoring the absorbance at 370 nm) and individual response factors against 2-ethylnaphthalene.

## Results and discussion

### Chemiluminescence of 1a–c in DMSO/NaOH (aq.) and in diglyme/acetate buffer

To investigate the chemiluminescent activity of a 2,6-diphenylimidazopyrazinone derivative, the substituent effect of the 6-phenyl group in **1a–c** was studied first. Chemiluminescence reactions of **1a–c** in aerated DMSO containing 0.10 M NaOH aqueous solutions were examined at 25 °C. The reactions of **1a–c** showed only a weak luminescence, and the  $\Phi_{\text{CL}}$  values were below  $10^{-4}$ . A chemiluminescence reaction of an imidazopyrazinone in basic DMSO gives light emission from the excited singlet (*S*<sub>1</sub>) state of the amide anion of an amidopyrazine product.<sup>5–7</sup> Fluorescence measurements of the amide anions of **2a–c** in DMSO containing 0.10 M NaOH aqueous solutions showed quantum yields ( $\Phi_{\text{F}}$ ) below 0.005. Therefore, the low luminescence of **1a–c** was due to the fact that the amide anions of **2a–c** were not fluorescent.

Next, chemiluminescence reactions of **1a–c** in aerated diglyme containing acetate buffer (pH 5.6, 0.66% v/v) were examined at 25 °C. While **1b,c** showed only a weak chemiluminescence ( $\Phi_{\text{CL}} < 10^{-4}$ ), **1a** gave an estimated  $\Phi_{\text{CL}}$  value of 0.0025. The chemiluminescence emission spectrum of **1a** showed the maximum ( $\lambda_{\text{CL}}$ ) at 521 nm. This coincides with the fluorescence spectrum of the neutral form of amidopyrazine **2a**, indicating that light emission occurred from the *S*<sub>1</sub> state of **2a**. The difference in chemiluminescence behavior between **1a–c** in diglyme/acetate buffer can be explained by the substituent effect of the 6-phenyl group with the reaction



Scheme 2 Reaction mechanism of the chemiluminescence of **1**.

mechanism reported previously (Scheme 2).<sup>6</sup> In DMSO/NaOH (aq.), the chemiluminescence reactions of **1a–c** give the  $S_1$  states of the amide anions of **2a–c** via anionic dioxetanones (**D<sup>-</sup>**). On the other hand, the chemiluminescence reaction of **1a** in diglyme/acetate buffer gives the  $S_1$  states of the neutral form of **2a** via neutral dioxetanone (**DH**), while the chemiluminescence reactions of **1b,c** still give the excited amide anions of **2b,c** via **D<sup>-</sup>**. The difference between **1a–c** is because of competition between the decomposition of **D<sup>-</sup>** to give the excited amide anion and the protonation of **D<sup>-</sup>** to give **DH**. Because the 4-(dimethylamino)phenyl group is a good electron-donating substituent, the basicity of the anionic N atom of **D<sup>-</sup>** is increased, which leads to the preferential protonation of **D<sup>-</sup>**. In the case of **1b,c**, the electron-donating properties of the 4-methoxyphenyl and phenyl groups are not sufficient for the preferential protonation of **D<sup>-</sup>**. Therefore, **1b,c** gave negligible chemiluminescence in diglyme/acetate buffer in a manner similar to that in DMSO/NaOH (aq.). Only **1a**, with the electron-donating 4-(dimethylamino)phenyl group, is chemiluminescent in diglyme/acetate buffer.

### Chemiluminescence of **1a,d–f** in diglyme/acetate buffer

Based on the chemiluminescent structure of **1a**, with the 4-(dimethylamino)phenyl group at C6, we investigated the effect on chemiluminescence of substituents on the phenyl group at C2 with **1d–f**, all of which showed chemiluminescence in diglyme/acetate buffer. The observed properties are summarized in Table 1, together with those of **1a** and **dm-CLA**, for comparison.

Chemiluminescence emission spectra for **1a,d–f** are shown in Fig. 1. The  $\lambda_{CL}$  values for **1d–f** are also coincident with the fluorescence emission maxima of the neutral forms of **2d–f**, as

Table 1 Chemiluminescent properties of **1a,d–f** and **dm-CLA** in aerated diglyme/acetate buffer<sup>a</sup> at 25 °C

Substrate	$\lambda_{CL}^b/\text{nm}$	$k_{CL}^c/10^{-4} \text{ s}^{-1}$	$\Phi_{CL}^d/10^{-2}$	$\Phi_R^e$	$\Phi_F^f$	$\Phi_S^g$
<b>1a</b>	521	1.3	0.25	0.89	0.16	0.018
<b>1d</b>	524	3.1	1.2	0.45	0.34	0.078
<b>1e</b>	523	1.2	0.62	0.32	0.29	0.067
<b>1f</b>	518	4.5	1.4	0.37	0.44	0.086
<b>dm-CLA</b> <sup>h</sup>	491	23	1.5	0.62	0.33	0.073

<sup>a</sup> Diglyme containing acetate buffer (0.10 M, pH 5.6, 0.66% v/v). <sup>b</sup> Emission maxima of the chemiluminescence spectra. <sup>c</sup> Pseudo-first-order rate constants. <sup>d</sup> Chemiluminescence quantum yields. <sup>e</sup> Chemical efficiencies to give the corresponding amidopyrazine products by the chemiluminescence reactions. <sup>f</sup> Fluorescence quantum yields of the corresponding amidopyrazines. <sup>g</sup> Chemiexcitation efficiencies. <sup>h</sup> Ref. 6d.





Fig. 1 Chemiluminescence emission spectra of **1a,d–f** ( $1.0 \times 10^{-5}$  M) in aerated diglyme/acetate buffer at 25 °C.

observed for **1a**, indicating that light emitters are the  $S_1$  states of **2d–f**. The  $\lambda_{CL}$  values for **1a,d–f** are similar to one another, indicating that the  $S_1$  states of **2a,d–f** have a similar  $\pi$ -electronic properties which are not affected by the structure of the acyl groups. These  $\pi$ -electronic properties of the  $S_1$  states of **2a,d–f** are consistent with our previous report.<sup>10</sup>

Decay curves of chemiluminescence intensities ( $I$ ) over time for **1a,d–f** gave pseudo-first-order rate constants ( $k_{CL}$ ) (Table 1). The chemiluminescence reaction rate of an imidazopyrazinone is regulated by the oxygenation process of an imidazopyrazinone anion as the rate-determining step (Scheme 2).<sup>5a,6b</sup> In particular, thermal electron transfer from an imidazopyrazinone anion to  $^3O_2$  determines the value of  $k_{CL}$ .<sup>6b</sup> Because the  $k_{CL}$  values of **1a,d–f** are smaller than that of **dm-CLA**, the  $\pi$ -electronic conjugation of the phenyl group at C2 of the anions of **1a,d–f** reduces the reaction rate. The difference in the  $k_{CL}$  value between **1a,d–f** is explained by the introduction of the methoxy group. The Hammett  $\sigma_p$  and  $\sigma_m$  constants of the methoxy group are  $-0.27$  and  $0.12$ , respectively.<sup>16</sup> Thus, methoxy groups at the *ortho* and *para* positions of a phenyl group are electron donating and a methoxy group at the *meta* position is electron withdrawing. The order of the strength of electron donation, 2,3,4-trimethoxyphenyl (**1f**) > 4-methoxyphenyl (**1d**) > 3,4,5-trimethoxyphenyl (**1e**)  $\approx$  phenyl (**1a**), corresponds to the order of the  $k_{CL}$  values, indicating that the




**Scheme 3** Three factors to make chemiluminescence quantum yield ( $\Phi_{CL}$ ).

electron-donating properties of the phenyl group at C2 of an imidazopyrazinone anion effectively modulate the oxygenation reactivity.

The  $\Phi_{CL}$  values for **1d–f** are in the range of 0.006–0.014 and are similar to that (0.015) for **dm-CLA** and higher than that (0.0025) for **1a** (Table 1). The value of  $\Phi_{CL}$  tends to increase with increasing electron donation from the phenyl group at C2 of **1a,d–f**. To confirm the origin of the high  $\Phi_{CL}$  values for **1a,d–f**, the elements of the  $\Phi_{CL}$  value were evaluated. A  $\Phi_{CL}$  value is a product of three efficiencies,  $\Phi_{CL} = \Phi_R \times \Phi_S \times \Phi_F$ , where  $\Phi_R$  is the chemical efficiency to produce **2** by an appropriate chemiluminescence reaction pathway,  $\Phi_S$  is the chemiexcitation efficiency for generating the  $S_1$  state of **2** from a neutral dioxetanone **DH** and  $\Phi_F$  is the fluorescence quantum yield of **2** (Scheme 3). The  $\Phi_R$  values were determined by product analyses of the chemiluminescence reactions of **1a,d–f** in diglyme/acetate buffer, which indicated that **2a,d–f** were obtained in 32–89% yields. The  $\Phi_F$  values of **2a,d–f** measured in diglyme/acetate buffer exceeded 0.16. Consequently, the  $\Phi_S$  values were estimated to be greater than 0.018. It is noteworthy that the chemiluminescence of **1d,f** in diglyme/acetate buffer gives higher  $\Phi_S$  values than that of **dm-CLA**.

#### Analysis of the chemiexcitation mechanism in the chemiluminescence of **1a,d–f**

To understand the cause of efficient chemiexcitation in the chemiluminescence of **1a,d–f**, we analyzed the chemiexcitation processes mechanistically. The thermal decomposition of a 1,2-dioxetanone for efficient chemiexcitation can be explained by the charge transfer-induced luminescence (CTIL) mechanism.<sup>6d,17</sup> In the CTIL mechanism, a high  $\Phi_S$  value requires that both the transition state (TS) of decomposition of **DH** for chemiexcitation and the  $S_1$  state of **2** have a strong intramolecular charge transfer (ICT) character. Electronic properties of **DH** and **TS** for the chemiluminescence of **1a,d–f** and **dm-CLA** were predicted by DFT calculations with the restricted (R) and unrestricted (U) B3LYP/6-31G(d) methods, respectively. Fig. 2 shows selected optimized geometries of **DH** and **TS**, and calculation data are summarized in Table 2. The value of  $\Delta E$  is the relative heat of formation of **TS** compared to that of the corresponding **DH**. The  $r_{CC}$  and  $r_{OO}$  values are the C–C and O–O bond lengths, respectively, and the  $q_{DO}$ ,  $q_{NH}$ ,  $q_{PY}$ ,  $q_{AR}$  and  $q_{AR'}$  values are the total Mulliken charge densities of the atoms constituting the dioxetanone ( $C_2O_3$ ), NH, pyrazine ( $C_4H_2N_2$ ) and two aryl parts, respectively (Scheme 4).




**Fig. 2** Optimized geometries of **DH[1a,e,f]** and **TS[1a,e,f]** calculated by the R- and UB3LYP/6-31G(d) methods, respectively.

The  $\Delta E$  values for **TS[1a,d–f]** are similar to that of **TS[dm-CLA]**. Because the value of  $\Delta E$  corresponds to the activation energy of the decomposition of **DH**, the reaction rates of the chemiexcitation processes from **DH[1a,d,e]** and **DH[dm-CLA]** will be similar to one another. The dipole moments, molecular geometries and charge distributions of **DH[1a,d,e]** and **TS[1a,d,e]** are also similar to those of **DH[dm-CLA]** and **TS[dm-CLA]**, respectively. On the other hand, **TS[1f]** has a characteristically long O–O bond (2.25 Å) and the negative and positive charges in the structure of **TS[1f]** are localized at the dioxetanone and 5-[4-(dimethylamino)phenyl]pyrazinylamino moieties, respectively. In fact, the  $q_{DO}$  value (−0.344) of **TS[1f]** is much less than that (−0.156) of **TS[1a]** and the summation of the  $q_{NH}$ ,  $q_{PY}$  and  $q_{AR}$  values (+0.335) for **TS[1f]** is much greater than that (+0.249) for **TS[1a]**. **TS[1f]** has a larger dipole moment (14.7 D) than the other species (8.4–10.0 D). These characteristic properties of **TS[1f]** originate from the strong electron-donating properties of the 2,3,4-trimethoxyphenyl group and the steric hindrance of its *ortho* substituent, yet the properties of **TS[1f]** do not cause a drastic change in the  $\Phi_S$  value. The differences in the  $q_{AR}$  values between **TS[1a,d–f]** and **DH[1a,d–f]** are small (around −0.05). Thus, the phenyl ( $Ar'$ ) groups at C2 of **1a,d–f** have only a small electronic effect on the properties of the corresponding **TS**. That is, **TS[1a,d–f]**


**Table 2** Relative energies ( $\Delta E$ ), dipole moments ( $\mu$ ), C–C and O–O bond distances ( $r_{CC}$  and  $r_{OO}$ ) and Mulliken charge densities ( $q$ ) for dioxetanones (DH) and transition states (TS) of the dioxetanone decompositions for chemiluminescence of **1a,d–f** and **dm-CLA** calculated by the R- and UB3LYP/6-31G(d) methods

Dioxetanone or transition state <sup>a</sup>	$\Delta E^b / \text{kcal mol}^{-1}$	$\mu / \text{D} (\Delta\mu)^c$	$r_{CC} / \text{\AA} (\Delta r_{CC})^c$	$r_{OO} / \text{\AA} (\Delta r_{OO})^c$	$q_{\text{Do}}^d (\Delta q)^c$	$q_{\text{NH}}^d (\Delta q)^c$	$q_{\text{Py}}^d (\Delta q)^c$	$q_{\text{Ar}}^d (\Delta q)^c$	$q_{\text{Ar}'}^d (\Delta q)^c$
DH[1a]	0.0	4.1	1.546	1.492	0.014	−0.321	0.119	0.099	0.088
TS[1a]	17.5	10.0 (+5.9)	1.565 (+0.019)	1.492 (+0.471)	−0.156 (−0.170)	−0.291 (+0.030)	0.196 (+0.077)	0.209 (+0.110)	0.042 (−0.046)
DH[1d]	0.0	3.4	1.545	1.492	0.009	−0.323	0.117	0.097	0.100
TS[1d]	17.6	8.4 (+5.0)	1.566 (+0.021)	1.492 (+0.470)	−0.156 (−0.165)	−0.293 (+0.030)	0.190 (+0.073)	0.197 (+0.100)	0.062 (−0.038)
DH[1e]	0.0	3.2	1.546	1.492	0.022	−0.322	0.120	0.097	0.082
TS[1e]	17.4	8.7 (+5.5)	1.565 (+0.019)	1.492 (+0.469)	−0.155 (−0.177)	−0.289 (+0.033)	0.199 (+0.079)	0.209 (+0.112)	0.036 (−0.046)
DH[1f]	0.0	3.8	1.551	1.493	0.041	−0.297	0.102	0.091	0.064
TS[1f]	18.7	14.7 (+11.0)	1.557 (+0.006)	2.250 (+0.757)	−0.344 (−0.385)	−0.194 (+0.103)	0.257 (+0.155)	0.272 (+0.182)	0.008 (−0.056)
DH[dm-CLA]	0.0	4.3	1.538	1.492	0.044	−0.313	0.115	0.099	0.056 <sup>e</sup>
TS[dm-CLA]	18.3	9.2 (+4.9)	1.558 (+0.020)	1.983 (+0.491)	−0.098 (−0.142)	−0.293 (+0.020)	0.179 (+0.064)	0.192 (+0.093)	0.019 <sup>e</sup> (−0.036) <sup>e</sup>

<sup>a</sup>DH and TS denote dioxetanone and the transition state of the dioxetanone decomposition, respectively. DH and TS were calculated by the R- and UB3LYP/6-31G(d) methods, respectively. <sup>b</sup>Energy of a transition state relative to that of the corresponding dioxetanone. <sup>c</sup>Differences between the values for TS and the corresponding DH. <sup>d</sup>Skeletal parts for the Mulliken charge density are shown in Scheme 4. <sup>e</sup>Mulliken charge densities of the methyl group.



**Scheme 4** Definitions of the C–C and O–O bond lengths ( $r_{CC}$  and  $r_{OO}$ ) and the total Mulliken charge densities ( $q$ ) of the skeletal parts of DH[1] and TS[1].



**Scheme 5** The CTIL mechanism for the chemiexcitation process from DH[1] to  $^12^*$  via TS[1] in the chemiluminescence of **1a,d–f**.

maintain a favorable ICT character, like TS[dm-CLA], for efficient chemiexcitation, because they have the common 4-(dimethylamino)phenyl group at the pyrazinylamino moiety. It has already been shown that **2a** has a large dipole moment ( $\mu$ , 16 D) in the  $S_1$  state.<sup>10</sup> Similarly, the  $S_1$  states of **2d–f** will have large  $\mu$  values. Thus, the  $S_1$  states of **2a,d–f**, like the  $S_1$  state of **dm-OCLA**, have a strong ICT character. Therefore, the decomposition of DH[1a,d–f] proceeds to give the  $S_1$  states of **2a,d–f** via TS[1a,d–f] efficiently by following the CTIL

mechanism in a manner similar to the decomposition of DH[dm-CLA] (Scheme 5).

## Conclusion

We have demonstrated that **1a** having the electron-donating 4-(dimethylamino)phenyl group at C6 shows moderate chemiluminescent activity in diglyme/acetate buffer, while **1a** gives

negligible chemiluminescence in DMSO/NaOH (aq.). The chemiluminescence reaction of **1a** proceeds to give the  $S_1$  state of the neutral form of **2a**. Because **1b,c** having the 4-methoxyphenyl and phenyl groups at C6 are non-chemiluminescent substrates in DMSO/NaOH (aq.) and in diglyme/acetate buffer, the introduction of the dimethylamino group to the phenyl group at C6 is essential for the chemiluminescent activity of 2,6-diphenyl derivatives **1** via the generation of the  $S_1$  state of a neutral amidopyrazine from a neutral dioxetanone **DH**. The derivatives **1d-f** having methoxy substituent(s) on the phenyl group at C2 together with the 4-(dimethylamino)phenyl group at C6 also show chemiluminescence in diglyme/acetate buffer, and their  $\Phi_{CL}$  values are similar to the high  $\Phi_{CL}$  value of **dm-CLA**. In the elements to determine the  $\Phi_{CL}$  values of **1a,d-f**, the  $\Phi_S$  values were greater than 0.018. The chemiexcitation processes for **1a,d-f** are explained by the efficient decompositions of **DH** to yield the  $S_1$  states of **2a,d-f** with a strong ICT character via ICT TS through the CTIL mechanism, as in the case of **dm-CLA**. While the  $\Phi_{CL}$  values of **1a,d-f** are still smaller than the  $\Phi_{BL}$  value (ca. 0.3),<sup>3</sup> **1a,d-f** are in the group of derivatives with a high  $\Phi_{CL}$  value greater than 0.001 in diglyme/acetate buffer like **dm-CLA**.<sup>6c,d</sup> Interestingly, the methoxy substituted phenyl groups at C2 of **1d-f** serves to maintain or slightly increase the  $\Phi_S$  value, therefore contributing to a high  $\Phi_{CL}$  value. This finding provides a guideline for designing a new Cypridina luciferin analogue with a  $\Phi_{CL}$  value close to the  $\Phi_{BL}$  value. To the 6-[4-(dimethylamino)phenyl]-2-phenylimidazopyrazinone structure, for instance, an alkoxy-arm group can be introduced to the phenyl group at C2, which will have a supramolecular photochemical function of intramolecularly regulating the reactivity of the imidazopyrazinone core.<sup>18</sup> Based on this idea, we are studying the reaction mechanism of an efficient Cypridina bioluminescence, in which Cypridina luciferase has supramolecular functions that regulate the luminescence reaction and increase the  $\Phi_{BL}$  value.

## Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research C (no. 22550031) from the Japan Society for the Promotion of Science (JSPS). We acknowledge technical assistance in computing the quantum chemical calculations from the Information Technology Center of UEC. We also thank Professors Mamoru Ohashi (UEC), Haruki Niwa (UEC) and Shojiro Maki (UEC) for their generous assistance and valuable discussion.

## References

- O. Shimomura, F. H. Johnson and T. Masugi, *Cypridina* bioluminescence – Light-emitting oxyluciferin-luciferase complex, *Nature*, 1969, **164**, 1299–1300.
- Reviews: (a) T. Goto, Chemistry of bioluminescence, *Pure Appl. Chem.*, 1968, **17**, 421–441; (b) O. Shimomura and F. H. Johnson, Introduction to the *Cypridina* system, *Methods Enzymol.*, 1978, **57**, 331–364; (c) *Bioluminescence: Chemical Principles and Methods*, ed. O. Shimomura, rev. edn, World Scientific, Singapore, 2012.
- (a) F. H. Johnson, O. Shimomura, Y. Saiga, L. C. Gershman, G. T. Reynolds and J. R. Waters, Quantum efficiency of Cypridina luminescence, with a note on that of Aequorea, *J. Cell. Comp. Physiol.*, 1962, **60**, 85–103; (b) O. Shimomura and F. H. Johnson, Mechanisms in quantum yield of *Cypridina* bioluminescence, *Photochem. Photobiol.*, 1970, **12**, 291–295.
- F. H. Johnson, H.-D. Stachel, E. C. Taylor and O. Shimomura, Chemiluminescence and fluorescence of Cypridina luciferin and of some new indole compounds in dimethylsulfoxide, *Bioluminescence in Progress*, ed. F. H. Johnson and Y. Haneda, Princeton University Press, Princeton, 1966, pp. 67–82.
- (a) T. Goto, S. Inoue and S. Sugiura, *Cypridina* bioluminescence 4. Synthesis and chemiluminescence of 3,7-dihydroimidazo[1,2-*a*]pyrazin-3-one and its 2-methyl derivative, *Tetrahedron Lett.*, 1968, 3873–3876; (b) T. Goto, S. Inoue, S. Sugiura, K. Nishikawa, M. Isobe and Y. Abe, *Cypridina* bioluminescence V. Structure of emitting species in the luminescence of Cypridina luciferin and its related compounds, *Tetrahedron Lett.*, 1968, 4035–4038; (c) T. Goto and H. Fukatsu, *Cypridina* bioluminescence 7. Chemiluminescence in micelle solutions – A model system for *Cypridina* bioluminescence, *Tetrahedron Lett.*, 1969, 4299–4302; (d) Y. Toya, T. Kayano, K. Sato and T. Goto, Synthesis and chemiluminescence properties of 6-(4-methoxyphenyl)-2-methylimidazo[1,2-*a*]pyrazin-3(7*H*)-one and 2-methyl-6-(2-naphthyl)imidazo[1,2-*a*]pyrazin-3(7*H*)-one, *Bull. Chem. Soc. Jpn.*, 1992, **65**, 2475–2479.
- (a) T. Hirano, Y. Gomi, T. Takahashi, K. Kitahara, F. Q. Chen, I. Mizoguchi, S. Kyushin and M. Ohashi, Chemiluminescence of coelenterazine analogs – Structure of emitting species, *Tetrahedron Lett.*, 1992, **39**, 5771–5774; (b) H. Kondo, T. Igarashi, S. Maki, H. Niwa, H. Ikeda and T. Hirano, Substituent effects on the kinetics for the chemiluminescence reaction of 6-arylimidazo[1,2-*a*]pyrazin-3(7*H*)-ones (Cypridina luciferin analogues): support for the single electron transfer (SET)-oxygenation mechanism with triplet molecular oxygen, *Tetrahedron Lett.*, 2005, **46**, 7701–7704; (c) Y. Takahashi, H. Kondo, S. Maki, H. Niwa, H. Ikeda and T. Hirano, Chemiluminescence of 6-aryl-2-methylimidazo[1,2-*a*]pyrazin-3(7*H*)-ones in DMSO/TMG and in diglyme/acetate buffer: support for the chemiexcitation process to generate the singlet-excited state of neutral oxyluciferin in a high quantum yield in the *Cypridina* (*Vargula*) bioluminescence mechanism, *Tetrahedron Lett.*, 2006, **47**, 6057–6061; (d) T. Hirano, Y. Takahashi, H. Kondo, S. Maki, S. Kojima, H. Ikeda and H. Niwa, The reaction mechanism for the high quantum yield of the *Cypridina* (*Vargula*) bioluminescence supported by chemiluminescence of 6-aryl-2-methylimidazo[1,2-*a*]pyrazin-3(7*H*)-ones (Cypridina luciferin analogues), *Photochem. Photobiol. Sci.*, 2008, **7**, 197–207.

- 7 F. McCapra and Y. C. Chang, Chemiluminescence of a Cypridina luciferin analogue, *J. Chem. Soc., Chem. Commun.*, 1967, 1011–1012.
- 8 O. Shimomura and F. H. Johnson, Mechanism of luminescent oxidation of Cypridina luciferin, *Biochem. Biophys. Res. Commun.*, 1971, **44**, 340–346.
- 9 K. Usami and M. Isobe, Low-temperature photooxygenation of coelenterate luciferin analog synthesis and proof of 1,2-dioxetanone as luminescence intermediate, *Tetrahedron*, 1996, **52**, 12061–12090.
- 10 T. Hirano, T. Nakagawa, A. Kodaka, S. Maki, H. Niwa and M. Yamaji, 5-[4-(Dimethylamino)phenyl]-2-benzamidopyrazines: fluorescent dyes based on Cypridina oxyluciferin, *Res. Chem. Intermed.*, 2013, **39**, 233–245.
- 11 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, Gaussian, Inc., Wallingford CT, 2009.
- 12 (a) A. D. Becke, Density-functional thermochemistry 3. The role of exact exchange, *J. Chem. Phys.*, 1993, **98**, 5648–5652; (b) C. T. Lee, W. T. Yang and R. G. Parr, Development of the Colle-Salvetti correlation-energy formula into a functional of the electron-density, *Phys. Rev. B: Condens. Matter*, 1988, **37**, 785–789; (c) P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, Ab-initio calculation of vibrational absorption and circular-dichroism spectra using density-functional force-fields, *J. Phys. Chem.*, 1994, **98**, 11623–11627.
- 13 R. Dennington, T. Keith and J. Millam, *Version 5*, Semi-chem Inc., Shawnee Mission KS, 2009.
- 14 S. Fujio, D. Hashizume, Y. Takamuki, M. Yasui, F. Iwasaki, S. Maki, H. Niwa, H. Ikeda and T. Hirano, Regioselective phenyl-substitution effects on the solvatochromism of 2-phenylimidazo[1,2-*a*]pyrazin-3(7*H*)-one derivatives: expansion of the color variation range of a visible indicator for the proton donor ability of solvents, *Tetrahedron Lett.*, 2004, **45**, 8531–8534.
- 15 J. Lee and H. H. Seliger, Quantum yields of luminol chemiluminescence reaction in aqueous and aprotic solvents, *Photochem. Photobiol.*, 1972, **15**, 227–237.
- 16 C. Hansch, A. Leo and R. W. Taft, A survey of Hammett substituent constant and resonance and field parameters, *Chem. Rev.*, 1991, **91**, 165–195.
- 17 H. Isobe, M. Okamura, S. Kuramitsu and K. Yamaguchi, Mechanistic insights in charge-transfer-induced luminescence of 1,2-dioxetanones with a substituent of low oxidation potential, *J. Am. Chem. Soc.*, 2005, **127**, 8667–8679.
- 18 *Modern Molecular Photochemistry of Organic Molecules*, ed. N. J. Turro, V. Ramamurthy and J. C. Scaiano, University Science Books, Sausalito, CA, 2010, pp. 925–1000.