

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Chemoselective Transfer Hydrogenation of Aldehydes with HCOONH<sub>4</sub> Catalyzed by RuCl(CNNPh)(PP) Pincer Complexes**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1693291> since 2019-02-19T10:35:20Z

*Published version:*

DOI:10.1002/cctc.201600892

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

*Revised*

*ChemCatChem*

## **Chemoselective Transfer Hydrogenation of Aldehydes with HCOONH<sub>4</sub> Catalyzed by RuCl(CNN<sup>Ph</sup>)(PP) Pincer Complexes**

Salvatore Baldino,<sup>\*[a]</sup> Sarah Facchetti,<sup>[b]</sup> Hans Günter Nedden,<sup>[b]</sup> Antonio Zanotti-Gerosa,<sup>[b]</sup> and Walter Baratta<sup>\*[a]</sup>

**Abstract:** Aldehydes are chemoselectively reduced to primary alcohols using HCOONH<sub>4</sub> as hydrogen donor, via transfer hydrogenation catalyzed by benzo[*h*]quinoline pincer complexes RuCl(CNN<sup>Ph</sup>)(PP) at S/C = 2000-20000. This practical reaction performed with aldehydes of commercial grade purity in a water / toluene biphasic system affords alcohols without formation of condensation or amination side products.


**Keywords:** aldehydes • ligands • formate • hydrogen transfer • ruthenium

The search of well-designed and productive catalysts for the selective hydrogenation<sup>[1]</sup> and transfer hydrogenation (TH)<sup>[2]</sup> of aldehydes to primary alcohols is a fundamental reaction of broad application in the industry.<sup>[3]</sup> As matter of fact, this catalytic route promoted by transition metals, results in a lower environmental impact and an easier work-up with respect to the classical approach entailing NaBH<sub>4</sub>, LiAlH<sub>4</sub>, boranes and Al alkoxides as reducing agents.<sup>[4]</sup> Among the transition metals used in catalytic homogeneous reductions, ruthenium, which is cheaper with respect to rhodium and iridium, has played a crucial role leading to a number of efficient catalysts for the aldehyde reduction using H<sub>2</sub> under pressure<sup>[5]</sup> or 2-propanol.<sup>[5e, 6]</sup> While the TH with 2-propanol is an equilibrium reaction that, on scale, requires the removal of acetone to drive the reaction to completion, the use of formic acid derivatives as hydrogen donors have the advantage of generating CO<sub>2</sub>, which is released from the reaction solution, driving the reaction and minimizing the reversibility problems.<sup>[7]</sup> Alkali formates HCOOM (M = Na, K) were employed in TH of aldehydes with RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub>,<sup>[8]</sup> [RuCl<sub>2</sub>(mtppps)<sub>2</sub>]<sub>2</sub> (mtppps = sodium 3-diphenylphosphinobenzenesulfonate),<sup>[9]</sup> CpRuCl(PPh<sub>3</sub>)(PN) (PN = diphenyl-2-pyridylphosphine),<sup>[10]</sup> RuCl<sub>2</sub>(PTA)<sub>4</sub> (PTA = 1,3,5-triaza-7-phosphaadamantane)<sup>[11]</sup> and RuCl<sub>2</sub>(PO)<sub>2</sub>

(PO=(2-methoxyethyl)diphenylphosphine)<sup>[12]</sup> catalysts at  $S/C \leq 1000$ , whereas the HCOOH-NEt<sub>3</sub> system has been used with [RuCl<sub>2</sub>(benzene)]<sub>2</sub>, but at a low  $S/C = 50$ .<sup>[13]</sup>

Ammonium formate is a cheap and readily accessible reducing agent widely employed in organic transformations. In addition to the Leuckart-Wallach reductive amination of carbonyl compounds,<sup>[14]</sup> the metal catalyzed version reaction has been reported by Kitamura<sup>[15]</sup> and Talwar<sup>[16]</sup> with Cp\* Rh(III) and Ir(III) complexes, respectively, whereas Kadyrov described the asymmetric version with RuCl<sub>2</sub>[(*R*)-tol-BINAP].<sup>[17]</sup> HCOONH<sub>4</sub> has also been widely used in Pd/C-catalyzed conversion of carbonyl compounds into alkanes,<sup>[18]</sup> nitro derivatives to amines<sup>[19]</sup> and in the hydrodehalogenation of aromatic chlorides.<sup>[20]</sup>


For the employment of HCOONH<sub>4</sub> in the TH of carbonyl compounds, recently Grainger<sup>[21]</sup> described the use of Wills' tethered Ru complexes in the reduction of ketones, while Nie<sup>[22]</sup> observed the concomitant TH and reductive amination of acetophenone with RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub>. Curiously, the TH of aldehydes with HCOONH<sub>4</sub> has only been described by Iyer in the reduction of 4-methoxybenzaldehyde and *trans*-cinnamaldehyde catalyzed by Ni[P(OPh)<sub>3</sub>]<sub>4</sub> at 110 °C, but at a very low  $S/C = 30$ .<sup>[23]</sup> Since the catalytically active metal-hydride species<sup>[24]</sup> are generated in basic conditions, where aldehydes, more than ketones, may undergo several side reactions (Claisen-Tishchenko,<sup>[25]</sup> Cannizzaro<sup>[26]</sup> and aldol condensation reactions<sup>[27]</sup>), the selective aldehyde reduction remains a delicate point. Recently, we described the preparation of the benzo[*h*]quinoline pincer complexes RuCl(CNN<sup>Ph</sup>)(PP) **1-3** which are active catalysts in the hydrogenation and TH with 2-propanol of ketones<sup>[28]</sup> and aldehydes<sup>[29]</sup> (Figure 1).



**Figure 1.** Benzo[*h*]quinoline pincer ruthenium complexes **1-3**.

In the TH of aldehydes in 2-propanol the reaction was carried out at low substrate concentration (0.1 M) and under weak basic conditions to inhibit the condensation reaction with the formed acetone, a side reaction which strongly depends on the type of the aldehyde.<sup>[29]</sup>

We report here the straightforward and chemoselective TH of aldehydes of commercial-grade purity to alcohols using ammonium formate as hydrogen donor in a water / toluene biphasic system, catalyzed by the pincer complexes **1-3** at S/C 2000-20000. Employment of HCOONH<sub>4</sub> with **1-3** allowed the clean reduction of aliphatic, conjugated and functionalized aromatic and heteroaromatic aldehydes, without formation of condensation and amination side products. Reaction of benzaldehyde **a** of commercial grade purity (assay 99%) in toluene (0.5 M) with 2 eq. of HCOONH<sub>4</sub> in water (1 M) and in the presence of complex **1** (S/C = 5000) affords selectively benzyl alcohol (76%) in 22 h at 90 °C (Table 1, entry 1), without formation of products of the Leuckart-Wallach reductive amination (Scheme 1).



**Scheme 1.** TH of aldehydes with HCOONH<sub>4</sub> catalyzed by **1-3**.

Complex **2** shows higher activity than **1** with 98 and 97% of alcohol in 9 and 15 h with S/C = 2000 and 5000, respectively (entries 2, 3). Complete conversion was also achieved at higher substrate concentration (1 M) with 2 and 4 eq. of formate (entries 4, 5). In addition, chemoselective TH of **a** to alcohol (96%) was attained with HCOONH<sub>4</sub> using a 2 M of substrate and at high S/C = 20000 in 48 h (entry 6). It is worth noting that performing the reaction at high aldehyde concentration is of

particular advantage for industrial applications and that the TH with 2-propanol requires lower aldehyde concentration (0.1 M) to avoid coupling reactions.<sup>[29]</sup>

Table 1. TH of benzaldehyde **a** to benzyl alcohol catalyzed by **1-3** using HCOONH<sub>4</sub>, HCOONa, and HCOOH / NEt<sub>3</sub> as hydrogen donors (DH<sub>2</sub>) in water / toluene at 90 °C.

Entry	Cat.	S/C	[S] <sup>[a]</sup>	[DH <sub>2</sub> ] <sup>[b]</sup>	DH <sub>2</sub> (eq.)	Time (h)	Conv. (%) <sup>[c]</sup>
1	<b>1</b>	5000	0.5	1 <sup>[d]</sup>	2	16	60
						22	76
2	<b>2</b>	2000	0.5	1 <sup>[d]</sup>	2	9	98
3	<b>2</b>	5000	0.5	1 <sup>[d]</sup>	2	15	97
4	<b>2</b>	5000	1	1 <sup>[d]</sup>	2	15	85
						24	96
5	<b>2</b>	5000	1	1 <sup>[d]</sup>	4	15	97
6	<b>2</b>	20000	2	2 <sup>[d]</sup>	4	15	58
						22	89
						48	96
7	<b>3</b>	5000	0.5	1 <sup>[d]</sup>	2	16	96
8	<b>3</b>	10000	0.5	1 <sup>[d]</sup>	2	20	86
						40	96
9	none	--	2	2 <sup>[d]</sup>	4	24	11
10	<b>2</b>	2000	0.5	2 <sup>[e]</sup>	4 <sup>[e]</sup>	24	50
11	<b>2</b>	5000	4.5	6.5 <sup>[f]</sup>	1.5 <sup>[f]</sup>	24	48
12	<b>2</b>	2000	0.5	2 <sup>[g]</sup>	4 <sup>[g]</sup>	14	2

[a] Substrate concentration in toluene. [b] DH<sub>2</sub> concentration in water. [c] The conversion and the purity were determined by GC and NMR analyses. [d] HCOONH<sub>4</sub>. [e] HCOOH / NEt<sub>3</sub> = 1 / 1. [f] Benzaldehyde was reacted in neat HCOOH / NEt<sub>3</sub> = 5 / 2. [g] HCOONa.

Complex **3** gave nearly quantitative reduction of **a** in 16 and 40 h with S/C = 5000 and 10000 (entries 7, 8). In absence of ruthenium catalyst, the reaction of **a** with 4 eq. of HCOONH<sub>4</sub> afforded 11% of alcohol after 24 h (entry 9), indicating that under these conditions **a** is reduced by HCOONH<sub>4</sub>, although at much lower rate with respect to the catalytic pathway. The use of HCOOH / NEt<sub>3</sub> = 5 / 2 and 1 / 1 mixtures (1.5 and 4 eq. of formic acid, respectively) in the presence of **2** at S/C = 5000 and 2000, gave 48-50% of alcohol after 24 h (entries 10, 11), whereas with HCOONa the conversion was 2% in 14 h (entry 12). These results indicate that the inexpensive HCOONH<sub>4</sub> can be employed as practical hydrogen donor for the selective reduction of **a** catalyzed by the robust pincer complexes **1-3**. Preliminary experiments with other media different than toluene / water

system, such as methanol / water mixtures or pure methanol led to poor conversion and poor selectivity, due to the formation of aminative condensation / reduction side-products.

To broaden the scope of the aldehyde TH with HCOONH<sub>4</sub>, aromatic, aliphatic, conjugated and heteroaromatic aldehydes were studied with complex **2**. Reduction of 4-bromobenzaldehyde **b** (2 M in toluene) with 4 eq. of HCOONH<sub>4</sub> afforded 97 and 98% of the corresponding alcohol in 10 and 24 h using **2** at S/C of 2000 and 10000, respectively (entries 1, 2, Table 2), whereas 75% alcohol was obtained at S/C = 20000 (entry 3).

Table 2. TH of aldehydes (2 M in toluene) with HCOONH<sub>4</sub> (4 eq., 2 M in water) catalyzed by complex **2** in toluene / water 90 °C.

Entry	Aldehyde	S/C	Time (h)	Conv. (%) <sup>[a]</sup>
1	<b>b</b>	2000	10	97
2	<b>b</b>	10000	24	98
3	<b>b</b>	20000	48	75
4	<b>c</b>	5000	15	96
5	<b>d</b>	2000	6	99, 87 <sup>[b]</sup>
6	<b>e</b>	2000	3.5	99, 70 <sup>[b]</sup>
7	<b>f</b>	2000	10	98 <sup>[c]</sup>
8	<b>f</b>	2000	4	99 <sup>[d]</sup>
9	<b>g</b>	10000	20	98
10	<b>h</b>	10000	24	97 <sup>[e]</sup>
11	<b>i</b>	2000	10	97 <sup>[f]</sup>
12	<b>i</b>	5000	48	97 <sup>[g]</sup>
13	<b>j</b>	5000	20	98 <sup>[e]</sup> , 88 <sup>[b]</sup>
14	<b>k</b>	5000	8	95
15	<b>l</b>	5000	18	99 <sup>[e]</sup>

[a] The conversion and the purity were determined by GC and <sup>1</sup>H-NMR analyses. [b] Isolated yield.

[c] Only the double-reduction product was detected. [d] With 1.5 eq. of HCOONH<sub>4</sub> a mixture of 4-(hydroxymethyl)benzaldehyde / 1,4-phenylenedimethanol in a 9 / 1 ratio was observed.

[e] [HCOONH<sub>4</sub>] = 4 M in water. [f] 91% of *trans*-cinnamol and 6% of the saturated alcohol 3-phenylpropan-1-ol. [g] 85% of *trans*-cinnamol and 12% of 3-phenylpropan-1-ol.

The electron-rich 4-(dimethylamino)benzaldehyde **c** was efficiently reduced to alcohol (96%) with  $S/2 = 5000$  in 15 h (entry 4). Conversely, the TH of the electron poor 4-nitrobenzaldehyde **d** and 4-cyanobenzaldehyde **e** afforded quantitative formation of the corresponding alcohols, isolated in 87 and 70% yield at  $S/2 = 2000$ , without reduction of the  $\text{NO}_2$  and CN functionalities or deactivation of the catalyst, i.e. by coordination at the metal center (entries 5, 6). Double TH was observed for 4-formylbenzaldehyde **f** leading to quantitative formation of 1,4-phenylenedimethanol in 10 h (entry 7). Interestingly, with a lower amount of  $\text{HCOONH}_4$  (1.5 eq.) the TH of **f** afforded the mono reduction product 4-(hydroxymethyl)benzaldehyde / 1,4-phenylenedimethanol in a 9 / 1 molar ratio, respectively (entry 8). Also the heteroaromatic 2-formylfuran **g** and 2-formylthiophene **h** were selectively reduced to alcohols 98 and 97% at  $S/2 = 10000$  in 20 and 24 h, respectively (entries 9, 10). Unsaturated *trans*-cinnamaldehyde **i** gave almost complete conversion (97%) to *trans*-cinnamol (91%) and 3-phenylpropan-1-ol (6%) at  $S/2 = 2000$  in 10 h, whereas at lower loading ( $S/2 = 5000$ ), formation of 85% of *trans*-cinnamol and 12% of the saturated alcohol was observed in 48 h, indicating that higher selectivity is achieved at higher catalyst loading and in shorter reaction time (entries 11, 12). On the other hand, *trans*- $\alpha$ -methylcinnamaldehyde **j** was chemoselectively transformed into *trans*- $\alpha$ -methylcinnamol, and isolated in 88% yield at  $S/2 = 5000$  in 20 h, with no reduction of the C=C double bond (entry 13). In addition, the aliphatic aldehydes hexanal **k** and *rac*-citronellal **l** were reduced to 1-hexanol and *rac*-citronellol 95 and 99%, with **2** at  $S/C = 5000$  (entries 14, 15). By contrast, with **2** ( $S/C = 2000$ ) the TH of vanillin and pyrrole-2-carboxaldehyde, displaying relatively acidic hydrogens, failed, leading to the unreacted starting material. It is worth noting that *cis*- $\text{RuCl}_2(\text{ampy})(\text{dppf})$  ( $S/C = 2000$ ), which is a complex related to **3** and efficiently catalyzes the TH of aldehydes with 2-propanol,<sup>[29]</sup> has been proven to be significantly less active in the TH of benzaldehyde (18 % conv., 15 h) with  $\text{HCOONH}_4$  (4 eq.). All spectral data were in the agreement with the literature, as all the obtained compounds are known.

As regards the mechanism, it is likely that the pincer complex  $\text{RuCl}(\text{CNN}^{\text{Ph}})(\text{PP})$  in the presence of  $\text{HCOONH}_4$  leads to the formate complex  $\text{Ru}(\text{O}_2\text{CH})(\text{CNN}^{\text{Ph}})(\text{PP})$ , with formation of the hydride  $\text{RuH}(\text{CNN}^{\text{Ph}})(\text{PP})$ <sup>[30]</sup> by elimination of  $\text{CO}_2$ . The subsequent reaction with the RCHO substrate gives the alkoxide  $\text{Ru}(\text{OCH}_2\text{R})(\text{CNN}^{\text{Ph}})(\text{PP})$  which is protonated by  $\text{HCOONH}_4$ , affording the alcohol product, ammonia and the formate complex, closing the cycle, as inferred from TCD-gas analysis, showing that  $\text{CO}_2$  and  $\text{NH}_3$  are evolved during the reaction. Since  $\text{HCOONH}_4$  displays better performances with respect to  $\text{HCOONa}$  and the  $\text{HCOOH} / \text{NEt}_3$  system, it is reasonable that the elimination of  $\text{NH}_3$  during the catalysis has positive effects, shifting the reaction toward the alcohol product, preventing a significant increase of the  $\text{OH}^-$  concentration, thus disfavoring the base catalyzed aldehyde side reactions. As a matter of fact, control experiments carried out during



the catalytic reduction of benzaldehyde and *rac*-citronellal showed that the pH values of the aqueous phase were in the range 7.5-8.5.

In conclusion, simple and functionalized aldehydes have been chemoselectively reduced to primary alcohols using HCOONH<sub>4</sub> as hydrogen donor with the benzo[*h*]quinoline pincer complexes RuCl(CNN<sup>Ph</sup>)(PP) at S/C up to 20000. This straightforward reaction carried out with aldehydes of commercial grade purity at high substrate concentration (2 M) in a water / toluene biphasic system gives alcohols without formation of condensation or amination side products. These reaction conditions are experimentally simple and provide significant options for industrial applications with respect to the use of 2-propanol as reducing agent. Further studies on ruthenium catalyzed transfer hydrogenation reactions are underway.

### Acknowledgments

We thank Dr. P. Martinuzzi (Università di Udine) for NMR assistance.

[a] Dr. S. Baldino, Prof. W. Baratta

Dipartimento DI4A

Università di Udine, Via Cottonificio 108, 33100 Udine (Italy)

Fax: +39-0432-558803

[b] Dr. S. Facchetti, Dr. A. Zanotti-Gerosa, Dr. H. G. Nedden

Johnson Matthey Fine Chemicals Division

28 Cambridge Science Park, Milton Road

Cambridge, CB4 0FP, United Kingdom

E-mails: [salvatore.baldino@uniud.it](mailto:salvatore.baldino@uniud.it), [walter.baratta@uniud.it](mailto:walter.baratta@uniud.it)

### References and notes

- [1] a) *The Handbook of Homogeneous Hydrogenation, Vols. 1-3* (Eds. J. G. de Vries, C. J. Elsevier) Wiley-VCH, Weinheim, **2007**; b) *Transition Metals for Organic Synthesis*, 2nd ed. (Eds. M. Beller, C. Bolm) Wiley-VCH, Weinheim, **2004**, p. 29.
- [2] a) D. Wang, D. Astruc, *Chem. Rev.* **2015**, *115*, 6621; b) W. Baratta, P. Rigo, *Eur. J. Inorg. Chem.* **2008**, 4041; c) J. S. M. Samec, J. E. Bäckvall, P. G. Andersson, P. Brandt, *Chem. Soc. Rev.*, 2006, **35**, 237.
- [3] J. Magano, J. R. Dunetz, *Org. Process Res. Dev.* **2012**, *16*, 1156.

- [4] a) J. Seyden-Penne, *Reductions by the Alumino- and Borohydrides in Organic Synthesis*, VCH, New York, **1991**; b) J. S. Cha, *Org. Process Res. Dev.* **2006**, *10*, 1032.
- [5] a) X. Tan, G. Wang, Z. Zhu, C. Ren, J. Zhou, H. Lv, X. Zhang, L. W. Chung, L. Zhang, X. Zhang, *Org. Lett.* **2016**, *18*, 1518; b) L. Bonomo, L. Kermorvan, P. Dupau, *ChemCatChem.* **2015**, *7*, 907; c) T. Miyada, E. H. Kwan, M. Yamashita, *Organometallics* **2014**, *33*, 6760; d) P. Dupau, L. Bonomo, L. Kermorvan, *Angew. Chem. Int. Ed.* **2013**, *125*, 11557; e) E. Putignano, G. Bossi, P. Rigo, W. Baratta, *Organometallics* **2012**, *31*, 1133; f) K. E. Jolley, A. Zanotti-Gerosa, F. Hancock, A. Dyke, D. M. Grainger, J. A. Medlock, H. G. Nedden, J. J. M. Le Paih, S. J. Roseblade, A. Seger, V. Sivakumar, I. Prokes, D. J. Morris, M. Wills, *Adv. Synth. Catal.* **2012**, *354*, 2545.
- [6] a) N. Kharat, A. Bakhoda, B. T. Jahromi, *Inorg. Chem. Commun.* **2011**, *14*, 1161; b) B. Deb, P. P. Sarmah, D. K. Dutta, *Eur. J. Inorg. Chem.* **2010**, 1710; c) M. Zhao, Z. Yu, S. Yan, Y. Li, *Tetrahedron Lett.* **2009**, *50*, 4624; d) B. Deb, B. J. Borah, B. J. Sarmah, B. Das, D. K. Dutta, *Inorg. Chem. Commun.* **2009**, *12*, 868; e) W. Baratta, K. Siega, P. Rigo, *Adv. Synth. Catal.* **2007**, *349*, 1633.
- [7] X. Zhou, X. Wub, B. Yang, J. Xiao, *J. Mol. Catal A: Chem.* **2012**, *357*, 133.
- [8] R. Bar, L. K. Bar, Y. Sasson, J. Blum, *J. Mol. Catal.* **1985**, *33*, 161.
- [9] I. Szatmári, G. Papp, F. Joó, Á. Kathó, *Catal. Today* **2015**, *247*, 14.
- [10] P. Kumar, A. K. Singh, S. Sharma, D. S. Pandey, *J. Organomet. Chem.* **2009**, *694*, 3643.
- [11] D. J. Darensbourg, F. Joo, M. Kannisto, A. Katho, J. H. Reibenspies, *Organometallics* **1992**, *11*, 1990.
- [12] S. Sabata, J. Vcelak, J. Hetflejs, *Collect. Czech. Chem. Commun.* **1995**, *60*, 127.
- [13] H. Cheng, R. Liu, J. Hao, Q. Wang, Y. Yu, S. Cai, F. Zhao, *Appl. Organomet. Chem.* **2010**, *24*, 763.
- [14] a) R. Leuckart, *Ber. Dtsch. Chem. Ges.* **1885**, *18*, 2341; b) O. Wallach, *Ber. Dtsch. Chem. Ges.* **1891**, *24*, 3992; c) M. L. Moore, *Org. React.* **1949**, *5*, 301; d) H. W. Gibson, *Chem. Rev.* **1969**, *69*, 673.
- [15] M. Kitamura, D. Lee, S. Hayashi, S. Tanaka, M. Yoshimura, *J. Org. Chem.* **2002**, *67*, 8685.
- [16] D. Talwar, N. P. Salguero, C. M. Robertson, J. Xiao, *Chem. Eur. J.* **2014**, *20*, 245.
- [17] R. Kadyrov, T. H. Riermeier, *Angew. Chemie Int. Ed.* **2003**, *42*, 5472.
- [18] S. Ram, L. D. Spicer, *Tetrahedron Lett.* **1988**, *29*, 3741.


- [19] a) S. Ram, R. E. Ehrenkauffer, *Tetrahedron Lett.* **1988**, 29, 5733; b) A. Saha, B. Ranu, *J. Org. Chem.* **2008**, 73, 6867.
- [20] R. Nakao, H. Rhee, Y. Uozumi, *Org. Lett.* **2005**, 7, 163.
- [21] D. M. Grainger, A. Zanotti-Gerosa, K. P. Cole, D. Mitchell, S. A. May, P. M. Pollock, J. R. Calvin, *ChemCatChem.* **2013**, 5, 1205.
- [22] H. Nie, H. Zhou, X. Li, Y. Li, J. Wang, *Chin. J. Org. Chem.* **2013**, 33, 2412.
- [23] S. Iyer, A. K. Sattar, *Synth. Commun.* **1998**, 28, 1721.
- [24] a) S. E. Clapham, A. Hadzovic, R. H. Morris, *Coord. Chem. Rev.* **2004**, 248, 2201; b) P. Espinet, A. C. Albéniz, *Fundamentals of Molecular Catalysis, Current Methods in Inorganic Chemistry, Vol. 3* (Eds. H. Kurosawa, A. Yamamoto) Elsevier, Amsterdam, **2003**, Chap. 6, p. 328.
- [25] a) A. M. P. Koskinen, A. O. Kataja, *Org. React.* **2015**, Chap. 2, p. 105; b) T. Werner, J. Koch, *Eur. J. Org. Chem.* **2010**, 6904; c) K. Ekoue-Kovi, C. Wolf, *Chem. Eur. J.* **2008**, 14, 6302.
- [26] T. A. Geissman, *Org. React.* **1944**, 2, 94.
- [27] M. B. Smith, J. March, *Advanced Organic Chemistry 5th ed.*; New York, Wiley Interscience, **2001**, p. 1218.
- [28] a) S. Facchetti, V. Jurcik, S. Baldino, S. Giboulot, H. G. Nedden, A. Zanotti-Gerosa, A. Blackaby, R. Bryan, A. Boogaard, D. B. McLaren, E. Moya, S. Reynolds, K. S. Sandham, P. Martinuzzi, W. Baratta, *Organometallics* **2016**, 35, 277; b) W. Baratta, M. Ballico, S. Baldino, G. Chelucci, E. Herdtweck, K. Siega, S. Magnolia, P. Rigo, *Chem. Eur. J.* **2008**, 14, 9148; c) G. Chelucci, S. Baldino, W. Baratta, *Coord. Chem. Rev.* **2015**, 300, 29.
- [29] S. Baldino, S. Facchetti, A. Zanotti-Gerosa, H. G. Nedden, W. Baratta, *ChemCatChem.* **2016**, 8, 2279.
- [30] a) W. Baratta, S. Baldino, M. J. Calhorda, P. J. Costa, G. Esposito, E. Herdtweck, S. Magnolia, C. Mealli, A. Messaoudi, S. A. Mason, L. F. Veiros, *Chem. Eur. J.* **2014**, 20, 13603. b) W. Baratta, M. Ballico, A. Del Zotto, E. Herdtweck, S. Magnolia, R. Peloso, K. Siega, M. Toniutti, E. Zangrando, P. Rigo, *Organometallics* **2009**, 28, 4421.

*Graphical abstract*

**Chemoselective Transfer Hydrogenation of Aldehydes with HCOONH<sub>4</sub> Catalyzed by RuCl(CNN<sup>Ph</sup>)(PP) Pincer Complexes**

Salvatore Baldino,<sup>[a]</sup> Sarah Facchetti,<sup>[b]</sup> Hans G. Nedden,<sup>[b]</sup> Antonio Zanotti-Gerosa,<sup>[b]</sup> and Walter Baratta<sup>[a]</sup>

Ammonium formate efficiently reduces commercial-grade aldehydes to alcohols, via transfer hydrogenation, catalyzed by benzo[*h*]quinoline pincer ruthenium complexes without formation of condensation or amination side products.



**S/C up to 20000**