



Chikungunya: Resurgent Arbovirus and its Impact in Brazil

Max Willian Lisboa Gomes^{1,2}, Claudio Cesar Cirne-Santos^{1,2}, Caroline de Souza Barros^{1,2} and Izabel Christina Palmer Paixão^{1,2*}

¹Graduate Program in Science and Biotechnology, Fluminense Federal University, Brazil

²Department of Cellular and Molecular Biology, Molecular Virology and Biotechnology Laboratory, Brazil

***Corresponding author:** Izabel Christina Palmer Paixão, Department of Cellular and Molecular Biology, Institute of Biology, Fluminense Federal University, Brazil.

Received Date: February 21, 2020

Published Date: March 04, 2020

Introduction

The Chikungunya virus (CHIKV) is an alphavirus transmitted by the bite of blood-sucking mosquitoes and causes a significant public health problem in tropical and subtropical regions [1]. The disease carried out by the virus is generally characterized by acute fever in the first days, the presence of skin rashes and polyarthralgia accompanied by headaches, joint swelling and inflammatory conditions such as conjunctivitis. Hematophagous arthropods vectors such as: *Aedes aegypti* and *Aedes albopictus* are the most common transmitters of Chikungunya fever to humans [2].

Since its discovery in 1952, four CHIKV genotypes have been identified and characterized. The East-Central-South African (ECSA) and West Africa genotypes began their endemic process in Sub-Saharan Africa. The Asian genotype is circulating from urban transmission in Southeast Asia. The viral lineage of the Indian Ocean (IOL) caused explosive epidemics in the period 2005 and 2011 in the Indian Ocean and Asia [3]. This strain harbors mutations that increase its infectivity and transmission capacity in *Ae. Albopictus*, making transmission to humans also reach the temperate regions. As of October 31, 2014, there were reports of more than 1,200,000 suspected cases of CHIKV in the Americas, and cases of indigenous infections were confirmed in 50 territories in the region [4].

In 2014, we were able to observe the emergence of cases of CHIKV infections in the Brazilian territory. On September 13, 2014, the first indigenous case was confirmed in the State of Amapá. Soon after the first confirmation, other cases also arose in the State of Bahia, until the Ministry of Health released on October 18, information on the confirmation of 682 indigenous cases in the country. Imported cases of the disease were a major related problem the increase in infected patients in all Brazilian states.

The whole reality of the Brazilian territory was a factor in favoring the introduction and expansion of the virus. The vectors can be located in more than 4,000 municipalities (*Aedes aegypti*) and in the case of *Ae. Albopictus*, was observed in 3,285 municipalities. A wide flow of people, a high vector dispersion and the population's susceptibility to infection are also important variables for the construction of the disease's infectivity [5].

A scenario marked by the coexistence of arboviruses has been outlined in Brazil since the introduction of the disease. The impacts were observed in several sectors, from health services to the costs of services targeting vectorized diseases, in order to minimize their effects on society [6].

Figure 1: Number of confirmed cases of CHIKV in the epidemiological weeks for 2018 and 2019. Source: Sinan Online (updated database 2018/2019 - Adapted from Health Surveillance Secretariat - Ministry of Health of Brazil). There was a significant increase in the number of cases and in the notifications of patients infected with CHIKV in 2018 to 2019 (Figure 1).

These figures reflect the confirmed cases of the disease, which may still be due to the notifications. The growing number of cases can be associated with the entire basic sanitation infrastructure, as well as poor health services in low-income populations [7].

There are still no antiviral treatments and vaccines against CHIKV. Many research groups are working on tests with drugs that are candidates for an effective antiviral. The treatment currently offered to the population consists of supporting the symptoms caused by the disease [8]. Currently, many substances are being tested against CHIKV. Synthetic, semi-synthetic and natural

products are tested in in vitro and in vivo models [9-13].

During the outbreak from 2014 to 2016, there was an immense difficulty in characterizing the disease in clinical services, precisely because of the problem of the availability of specific tests for

diagnosis within the public health system. After the characterization of the tests as RT-PCR and other immunological tests, the accuracy of notifications of confirmed cases of CHIKV was further increased [4,14].




Figure 1: Number of confirmed cases of CHIKV in the epidemiological weeks for 2018 and 2019. Source: Sinan Online (updated database 2018/2019 - Adapted from Health Surveillance Secretariat - Ministry of Health of Brazil).

Conclusion

Chikungunya fever is currently a public health problem that affects many people in Brazil. The development of new technologies involving diagnostic tests favored the characterization of the disease state throughout the national territory. There is a growing need for the discovery of effective therapeutic alternatives against CHIKV. The impact of this disease in Brazil consists of a Health, Social and Economic issue.

Acknowledgement

None.

Conflicts of Interest

No conflicts of interest.

References

1. Powers AM, Logue CH (2007) Changing patterns of chikungunya virus: re-emergence of a zoonotic arbovirus. *Journal of General Virology* 88(9): 2363-2377.
2. Maniero VC, Santos MO, Ribeiro RL, De Oliveira PA, Da Silva TB, et al. (2016) Dengue, chikungunya and zika virus in Brazil: epidemiological situation, clinical aspects and preventive measures. *Multidisciplinary research almanac* 1.
3. Donalisio MR, Freitas ARR (2015) Chikungunya in Brazil: an emerging challenge. *Brazilian Journal of Epidemiology* 18: 283-285.
4. Beltran-Silva S, Chacon-Hernandez S, Moreno-Palacios E, Pereyra-Molina J (2018) Clinical and differential diagnosis: Dengue, chikungunya and Zika. *Revista Médica del Hospital General de Mexico* 81: 146-153.
5. Azevedo Rdo So, Oliveira CS, Vasconcelos PF (2015) Risk of chikungunya for Brazil. *Public Health Journal* 49: 58.
6. Grandadam M, Caro V, Plumet S, Thiberge JM, Souarès Y, et al. (2011) Chikungunya virus, southeastern France. *Emerging infectious diseases* 17(5): 910-913.
7. Nunes J (2020) The everyday political economy of health: community health workers and the response to the 2015 Zika outbreak in Brazil. *Review of International Political Economy* 27: 146-166.
8. De Lamballerie X, Ninove L, Charrel RN (2009) Antiviral treatment of chikungunya virus infection. *Infectious Disorders-Drug Targets (Formerly Current Drug Targets-Infectious Disorders)* 9(2): 101-104.
9. Atitaya H, Sarawut K, Nitwara W, Sittiruk R, Sutee Y, et al. (2020) Evaluation of the antiviral activity of orlistat (tetrahydrolipstatin) against dengue virus, Japanese encephalitis virus, Zika virus and chikungunya virus. *Scientific Reports Nature Publisher Group* 10.
10. Esteves PO, De Oliveira MC, De Souza Barros C, Cirne-Santos CC, Laneuvlille VT, et al. (2019) Antiviral Effect of Caulerpin Against Chikungunya. *Natural Product Communications* 14: 10.
11. Cirne-Santos CC, De Souza Barros C, Nogueira CCR, Amorim LdSC, De Mendonça Campos R, et al. (2018) Antiviral effect of the seaweed *Osmundaria obtusiloba* against the Zika virus. *Journal of Medicinal Plants Research* 12: 387-395.
12. Haese N, Powers J, Streblov DN, (2020) Small Molecule Inhibitors Targeting Chikungunya Virus. *Curr Top Microbiol Immunol*.
13. Lani R, Hassandarvish P, Chiam CW, Moghaddam E, Chu JJH, et al. (2015) Antiviral activity of silymarin against chikungunya virus. *Scientific reports* 5: 11421.
14. Cirne-Santos CC, Barros C dS, Azevedo RC, Nogueira C, Yamamoto K, et al. (2019) Inhibition by marine algae of chikungunya virus isolated from patients in a recent disease outbreak in Rio de Janeiro. *Frontiers in microbiology* 10: 2426.