



Chiral Scale and Conformal Invariance in 2D Quantum Field Theory

Citation

Hofman, Diego M., and Andrew E. Strominger. 2011. Chiral scale and conformal invariance in 2D quantum field theory. *Physical Review Letters* 107(16): 161601.

Published Version

doi:10.1103/PhysRevLett.107.161601

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:8156535>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Chiral Scale and Conformal Invariance in 2D Quantum Field Theory

Diego M. Hofman* and Andrew Strominger*

Center for the Fundamental Laws of Nature, Harvard University,
Cambridge, MA 02138, USA

It is well known that a local, unitary Poincare-invariant 2D QFT with a global scaling symmetry and a discrete non-negative spectrum of scaling dimensions necessarily has both a left and a right local conformal symmetry. In this paper we consider a chiral situation beginning with only a left global scaling symmetry and do not assume Lorentz invariance. We find that a left conformal symmetry is still implied, while right translations are enhanced either to a right conformal symmetry or a left $U(1)$ Kac-Moody symmetry.

I. INTRODUCTION

A two-dimensional (2D) Poincare and scale invariant quantum field theory (QFT) has at least four global symmetries under which the light cone coordinates transform as

$$\begin{aligned} x^- &\rightarrow x^- + a, & x^+ &\rightarrow x^+ + b, \\ x^- &\rightarrow \lambda^- x^-, & x^+ &\rightarrow \lambda^+ x^+. \end{aligned} \quad (1)$$

If in addition one posits that the theory is unitary and that the spectrum of the dilation operator for $\lambda^+ = \lambda^-$ is discrete and non-negative, then it was shown in [1] that the four global symmetries are enhanced to left and right infinite-dimensional conformal symmetries. Explicit counterexamples indicate that the enhancement need not occur if the dilation spectrum is not discrete.

In recent years, interest in 2D QFTs with other types of global scaling symmetries has arisen in a variety of contexts ranging from condensed matter to string theory. In this paper we consider the special case with three global symmetries:

$$x^- \rightarrow x^- + a, \quad x^+ \rightarrow x^+ + b, \quad x^- \rightarrow \lambda x^-, \quad (2)$$

comprising two translational symmetries and a chiral “left” dilational symmetry. Our assumptions, detailed in the next section, include locality, unitarity and a discrete non-negative dilational spectrum but not Lorentz invariance. In an argument parallel to the one in [1], we find that these three global symmetries are sufficient to conclude that there are (at least) two infinite-dimensional sets of local symmetries. One of these is a left local conformal symmetry which enhances the left dilational symmetry. The other enhances the right translational symmetry and can be either a right conformal symmetry or a left current algebra.

It is surprising that such a powerful conclusion can be reached from such minimal assumptions. However the element of surprise is potentially reduced by the fact that at this point there is no definite example of

a QFT which non-trivially satisfies all of our stated assumptions¹. Hence our powerful conclusions may apply to an empty set. On the other hand, possible interesting examples are suggested by the recent appearance of warped AdS_3 geometries in a variety of string theoretic investigations including the holographic duals of the so-called dipole deformations of 2D gauge theories [3–8] and the near-horizon geometries of extreme Kerr black holes [9–11], see also [12–16]. Warped AdS_3 has an $SL(2, R) \times U(1)$ isometry group which contains (2). As these spaces are continuous deformations of AdS_3 spaces with CFT_2 duals, we expect that their holographic duals exist and are deformed CFT_2 s with the symmetries (2). However at present it is not clear to what extent these duals obey all the assumptions stated below. In this paper we concentrate on the pure field theory analysis and leave these interesting issues to future investigations. It is also of interest to try to adapt our analysis to other types of 2D scaling symmetries such as, for example, Lifshitz scaling $x \rightarrow \lambda x, t \rightarrow \lambda^z t$.

II. FROM DILATIONS TO VIRASORO

We wish to consider local, unitary, translationally-invariant quantum field theories in 2D flat Minkowski space with a (linearly-realized) chiral global scale invariance. The assumed symmetries act on light-cone coordinates $x^\pm = t \pm x$ as

$$x^- \rightarrow x^- + a, \quad x^+ \rightarrow x^+ + b, \quad x^- \rightarrow \lambda x^-, \quad (3)$$

for constant $\lambda > 0, a, b$. We do not assume Lorentz invariance, an action or a conserved symmetric stress tensor. The operators generating left-moving (i.e. x^-) translations and dilations will be denoted H and D , while right-moving translations are denoted \bar{P} , where the bar in general denotes right-moving charges. By assumption these operators annihilate the vacuum. Their commutation relations are:

$$i[D, H] = H, \quad i[D, \bar{P}] = 0, \quad i[H, \bar{P}] = 0. \quad (4)$$

* Email: dhofman@physics.harvard.edu, andy@physics.harvard.edu

¹ One possible example might be given by the continuum limit of the large N chiral Potts model, as discussed in [2].

We moreover assume, following [1], that the eigenvalue spectrum λ_i of D is discrete and non-negative² and there exists a complete basis of local operators Φ_i such that

$$\begin{aligned} i[H, \Phi_i] &= \partial_- \Phi_i, & i[\bar{P}, \Phi_i] &= \partial_+ \Phi_i, \\ i[D, \Phi_i] &= x^- \partial_- \Phi_i + \lambda_i \Phi_i \end{aligned} \quad (5)$$

and $\int_C d\Phi_i = 0$ for any closed or complete space-like contour C . Note that ‘‘local operators’’ as here defined do not involve explicit functions of x^\pm . We will refer to λ_i as the weight of the operator Φ_i . Translational plus dilational invariance implies the vacuum two-point functions of the $\{\Phi_i\}$ obey

$$\langle \Phi_i(x^-, x^+) \Phi_j(x'^-, x'^+) \rangle = \frac{f_{ij}(x^+ - x'^+)}{(x^- - x'^-)^{\lambda_i + \lambda_j}} \quad (6)$$

for some a priori unknown functions f_{ij} .

Noether’s theorem implies that each of the operators H, D, \bar{P} is associated to a conserved Noether current with components denoted h_\pm, d_\pm and p_\pm whose dual contour integral, e.g.

$$H = \int dx^+ h_+ - \int dx^- h_- \quad (7)$$

then gives the global charges. A proof in the present context is reviewed in appendix A. All of these currents have an ambiguity under shifts of the form $\pm \partial_\pm O$, where O is a more general type of operator potentially involving explicit functions of x^\pm

$$O(x^+, x^-) = \sum_i f_i(x^+, x^-) \Phi_i(x^+, x^-). \quad (8)$$

We also show in appendix A that the shifts can be chosen so that the currents satisfy canonical commutation relations, viz:

$$\begin{aligned} i[H, h_\pm] &= \partial_- h_\pm, & i[H, p_\pm] &= \partial_- p_\pm, \\ i[H, d_\pm] &= \partial_- d_\pm - h_\pm, \end{aligned} \quad (9)$$

$$\begin{aligned} i[\bar{P}, h_\pm] &= \partial_+ h_\pm, & i[\bar{P}, p_\pm] &= \partial_+ p_\pm, \\ i[\bar{P}, d_\pm] &= \partial_+ d_\pm. \end{aligned} \quad (10)$$

This implies that h_\pm and p_\pm are local operators, but the term proportional to h_\pm in $i[H, d_\pm]$ implies that d_\pm must have explicit dependence on the x^- coordinate. Appendix A demonstrates - relying crucially on the discreteness of the spectrum λ_i - that the currents can be chosen to be eigenoperators of D . The weights of the global charges (4) then imply

$$\begin{aligned} i[D, h_-] &= x^- \partial_- h_- + 2h_-, \\ i[D, h_+] &= x^- \partial_- h_+ + h_+, \end{aligned} \quad (11)$$

$$\begin{aligned} i[D, p_-] &= x^- \partial_- p_- + p_-, \\ i[D, p_+] &= x^- \partial_- p_+, \end{aligned} \quad (12)$$

$$i[D, d_-] = x^- \partial_- d_- + d_-, \quad i[D, d_+] = x^- \partial_- d_+. \quad (13)$$

We see that d_+ and p_+ are weight 0, d_- , h_+ and p_- are weight 1 and h_- is weight 2.³

As we mentioned, d_\pm must have explicit coordinate dependence. Let us find it and write the current in terms of local operators. Defining s_\pm by

$$d_\pm = x^- h_\pm + s_\pm \quad (14)$$

and using (9) we find that

$$i[H, s_\pm] = \partial_- s_\pm, \quad i[\bar{P}, s_\pm] = \partial_+ s_\pm. \quad (15)$$

We conclude (s_+, s_-) are local operators with weights $(0, 1)$. Conservation of the dilation current yields

$$\begin{aligned} \partial_+ d_- + \partial_- d_+ &= \\ x^- (\partial_+ h_- + \partial_- h_+) + h_+ + \partial_- s_+ + \partial_+ s_- &= 0. \end{aligned} \quad (16)$$

h_\pm conservation then gives

$$h_+ = -\partial_- s_+ - \partial_+ s_- \quad (17)$$

We have not at this point fixed all the shift freedom in the currents. In particular, we may shift away s_- :

$$h_\pm \rightarrow h_\pm \mp \partial_\pm s_-, \quad d_\pm \rightarrow d_\pm \mp \partial_\pm (x^- s_-), \quad (18)$$

which remains consistent with the commutators (9-13) as well as current conservation. Equations (14) and (17) now take the simpler form

$$d_+ = x^- h_+ + s_+, \quad d_- = x^- h_- \quad (19)$$

and

$$h_+ = -\partial_- s_+ \quad (20)$$

Now, we can use the general form of the two-point functions given by (6). Bearing in mind the fact that s_+ is a local operator of weight 0, we must have

$$\langle s_+ s_+ \rangle = f_{s_+}(x^+) \quad (21)$$

which implies

$$\partial_- s_+ = h_+ = 0. \quad (22)$$

Conservation of h_\pm then reduces to $\partial_+ h_- = 0$ or, equivalently,

$$h_- = h_-(x^-). \quad (23)$$

This fact immediately leads to the existence of an infinite set of conserved charges. Define:

$$T_\xi = - \int dx^- \xi(x^-) h_-, \quad \bar{J}_\chi = \int dx^+ \chi(x^+) s_+ \quad (24)$$

² For sigma-models this means the target space must be compact.

³ In an ordinary 2D CFT, d_+ and h_+ and p_- all vanish, $p_+ = T_{++}$, $d_- = x^- T_{--}$ and $h_- = T_{--}$.

where $\xi(x^-)$ and $\chi(x^+)$ are smooth functions. In particular, $H = T_1$ and $D = \bar{J}_1 + T_{x^-}$. Notice that, while h_- can't vanish if we are to have a non trivial H operator, s_+ could be identically zero. $s_+ \neq 0$ leads to the existence of even more local symmetries unrelated to the originally posited global symmetries. Whether or not it is zero, we show in appendix B that

$$i[\bar{J}_\chi, T_\xi] = 0. \quad (25)$$

This means we are free to calculate the algebra spanned by the conserved charges T_ξ without worrying about the action of \bar{J}_χ . We henceforth concentrate on the minimal case $s_+ = 0$.

Let us now work out the algebra spanned by T_ξ . The action of $H = T_1$ and $D = T_{x^-}$ on h_- imply

$$i[T_1, T_\xi] = T_{-\xi'}, \quad i[T_{x^-}, T_\xi] = T_{\xi - \xi' x^-}, \quad \xi' \equiv \partial_- \xi. \quad (26)$$

This in turn implies that the action of T_ξ on h_- is

$$i[T_\xi, h_-] = \xi \partial_- h_- + 2\xi' h_- + \partial_-^2 O_\xi. \quad (27)$$

The scaling symmetry plus locality implies O_ξ must be of the form $O_\xi = \xi O_1 + \xi' O_0$ with O_1 a local operator of weight one. As it cannot depend on x^+ , O_0 must be a weight zero constant. Integrating both sides with respect to $dx^- \zeta(x^-)$ gives

$$i[T_\xi, T_\zeta] = T_{\xi' \zeta - \zeta' \xi} + \int dx^- \zeta \partial_-^2 O_\xi. \quad (28)$$

Antisymmetry under the exchange of $\xi \leftrightarrow \zeta$ then requires that $O_1 = 0$ and :

$$\partial_-^2 O_\xi = O_0 \partial_-^3 \xi. \quad (29)$$

Defining $c = 24\pi O_0$ we end up with the following commutations relations for the charges T_ξ :

$$i[T_\xi, T_\zeta] = T_{\xi' \zeta - \zeta' \xi} + \frac{c}{48\pi} \int dx^- (\xi'' \zeta' - \zeta'' \xi'). \quad (30)$$

We recognize this as the algebra of the left-moving conformal generators on the Minkowski plane with central charge c .

III. ENHANCEMENT OF RIGHT MOVING TRANSLATIONS TO A LOCAL SYMMETRY

Notice that up until now we have not made much use of the fact that our theory is translationally invariant in the x^+ dimension as well. In particular the above results also apply when we do not possess this symmetry. What happens now that we add p_\pm to the game?

In this case, the key observation is that p_+ is a zero weight local operator, as implied by (9-13). This means

$$\langle p_+ p_+ \rangle = f_{p_+}(x^+). \quad (31)$$

Acting with ∂_- on both insertions and using the fact that a hermitian operator with a vanishing two-point function

is trivial we learn that $\partial_- p_+ = 0$. Current conservation then implies $\partial_+ p_- = 0$. It follows that

$$p_+ = p_+(x^+), \quad p_- = p_-(x^-). \quad (32)$$

We cannot have both $p_+ = 0 = p_-$ as the charge \bar{P} is generically nonzero, although from what we have seen so far one of them could vanish. We now discuss all possibilities.

A. $p_- = 0 \Rightarrow$ right-moving Virasoro algebra

In this case we have infinitely many charges given by

$$\bar{T}_\xi = \int dx^+ \xi(x^+) p_+. \quad (33)$$

Since $\bar{T}_1 = \bar{P}$ we have

$$i[\bar{T}_1, \bar{T}_\xi] = -\bar{T}_{\xi'}. \quad (34)$$

This, in turn, constraints the action of \bar{T}_ξ on p_+ to be

$$i[\bar{T}_\xi, p_+] = \xi \partial_+ p_+ + 2\xi' p_+ + \partial_+ \bar{O}_\xi. \quad (35)$$

If we compare this expression with (27) we see that we are very close to the previous situation for T_ξ . Multiplying by $\zeta(x^+)$ and integrating both sides of this equation we get

$$i[\bar{T}_\xi, \bar{T}_\zeta] = \bar{T}_{\xi' \zeta - \zeta' \xi} + \int dx^+ \zeta \partial_+ \bar{O}_\xi. \quad (36)$$

Antisymmetry with respect to exchange of ξ and ζ then implies that \bar{O}_ξ is an even number of derivatives of ξ . The term with no derivatives can be eliminated by a constant shift of p_+ . Terms with four or more derivatives would violate the Jacobi identity. We conclude (shifting p_+ by a constant if needed)

$$i[\bar{T}_\xi, \bar{T}_\zeta] = \bar{T}_{\xi' \zeta - \zeta' \xi} + \frac{\bar{c}}{48\pi} \int dx^+ (\xi'' \zeta' - \zeta'' \xi'). \quad (37)$$

We recognize this as the algebra of the right-moving conformal generators on the Minkowski plane with central charge \bar{c} .⁴

Of course the vacuum will not in general be invariant under the global $SL(2, R)_R$ subgroup. Acting with D and H on \bar{T}_ξ we can check that $i[\bar{T}_\xi, h_-] = \partial_-^2 \Phi_\xi$. But Φ_ξ must be a weight zero operator, so we are left with $i[\bar{T}_\xi, h_-] = 0$. The upshot is that $[\bar{T}_\xi, T_\zeta] = 0$, as expected.

B. $p_+ = 0 \Rightarrow$ left-moving current algebra

In this case we have infinitely many left-moving charges

$$J_\chi = - \int dx^- \chi(x^-) p_-(x^-). \quad (38)$$

⁴ Interestingly, it is this right moving Virasoro that gives the entropy in Kerr-CFT[9].

Because the zero mode J_1 acts as ∂_+ , we must have $i[J_1, p_-] = 0$. This implies $[J_1, J_\chi] = 0$ and hence

$$i[J_\chi, p_-] = \partial_- M_\chi \quad (39)$$

where the operator M_χ is, by locality, a linear function of χ . Now we are in position to repeat the arguments used around (27). Multiplying by $\psi(x^-)$, integrating over x^- and invoking antisymmetry and the Jacobi identity we find

$$i[J_\chi, J_\psi] = \frac{k}{4\pi} \int dx^- (\psi' \chi - \chi' \psi) \quad (40)$$

where the constant k parameterizes the central element. This is a $U(1)$ Kac-Moody current algebra.

We also need the $[T_\xi, J_\chi]$ commutator. The fact that $[J_1, T_\xi] = 0$ implies

$$[T_\xi, p_-] = \xi \partial_- p_- + \xi' p_- + \partial_- N_\xi \quad (41)$$

with the operator N_ξ linear in ξ . The Jacobi identity with a third operator T_ζ then implies $N_\xi = f \partial_- \xi$ for some constant f [17, 18]. If f is nonzero, the current p_- is not a dimension one chiral current. However we may then shift h_- by a k -dependent multiple of $\partial_- p_-$ so that p_- is a good dimension one current. This shift affects the central charge of T_ξ . Performing this transformation leaves us with the standard commutator

$$i[T_\xi, J_\chi] = J_{-\xi \chi'}. \quad (42)$$

It may seem rather strange to have a left-moving Kac-Moody current algebra whose zero mode generates right translations. However reminiscent structures have appeared before. In the KK circle reduction of AdS_3 to AdS_2 , one begins with two Virasoros and ends with a single left-moving Virasoro and current algebra associated to the KK $U(1)$ [19]. The left current algebra zero mode J_0 descends from right Virasoro zero mode in AdS_3 . Related structures have appeared in Kerr/CFT, where left Virasoro and right current algebra zero modes are sometimes identified [20], as well as in the study of asymptotic symmetries of warped AdS_3 [21].

C. Non-minimal $p_- \neq 0, p_+ \neq 0$

In this case left and right currents decouple. The commutators $i[\bar{P}, \bar{T}_\xi] = \bar{T}_{-\xi'}$ implies $i[\bar{T}_\xi, p_-] = \partial_- \Phi_\xi$, for some local Φ_ξ . Furthermore, because \bar{T}_ξ does not transform under D and p_- is a weight 1 operator, Φ_ξ must be weight 0. Therefore

$$i[\bar{T}_\xi, J_\chi] = 0. \quad (43)$$

implying that the conserved charges can be analyzed separately as above.

In summary, left translational and dilational symmetries together with right translations imply the existence of (at least) two sets of infinite dimensional algebras. On the left we always find a local conformal symmetry, while the right translational current is enhanced either to a local right conformal symmetry or a left $U(1)$ current algebra.

ACKNOWLEDGMENTS

We are delighted to acknowledge fruitful discussions with C. Bachas, A. Castro, M. Guica, T. Hartman, S. Hartnoll, J. Maldacena, A. Maloney, J. McGreevy, J. Polchinski, S. Sachdev, N. Seiberg and W. Song. This work was supported in part by DOE grant DE-FG0291ER40654 and the Fundamental Laws Initiative at Harvard.

Appendix A: Noether's theorem

In this section we will prove Noether's theorem for H and P and show the Noether currents can be put in a canonical "diagonal" form. We assume the existence of a unitary hamiltonian \mathcal{H} whose commutator with any operator obeys

$$i[\mathcal{H}, O] = \frac{dO}{dt} - \frac{\partial O}{\partial t}, \quad (A1)$$

where the last derivative acts on any explicit coordinate dependence in O . Conserved charges are defined as any operator \mathcal{Q} such that $\frac{d\mathcal{Q}}{dt} = 0$. Locality implies

$$\mathcal{Q} = \int_{-\infty}^{\infty} dx q_t(x, t). \quad (A2)$$

Charge conservation is then

$$\frac{d\mathcal{Q}}{dt} = \int dx \frac{q_t}{dt} = 0, \quad (A3)$$

implying $\frac{q_t}{dt} = -\frac{q_x}{dx}$, for some q_x . (q_x, q_t) is the sought after conserved Noether current associated to \mathcal{Q} .

In this paper we further assume the existence of a conserved momentum charge \mathcal{P} commuting with \mathcal{H} and obeying

$$i[\mathcal{P}, O] = \frac{dO}{dx} - \frac{\partial O}{\partial x} \quad (A4)$$

for any operator O . From these we construct left and right translation charges $2H = \mathcal{H} - \mathcal{P}$ and $2\bar{P} = \mathcal{H} + \mathcal{P}$. If a conserved charge \mathcal{Q} commutes with both H and \bar{P} the associated Noether current must obey

$$i[H, q_\pm] = \partial_- q_\pm \pm \partial_\pm F, \quad (A5)$$

$$i[\bar{P}, q_\pm] = \partial_+ q_\pm \pm \partial_\pm G, \quad (A6)$$

where ∂_\pm are total derivatives with respect to $x^\pm = t \pm x$ and F and G can be expanded

$$F = \sum_i f_i(x^+, x^-) \Phi_i, \quad G = \sum_i g_i(x^+, x^-) \Phi_i. \quad (A7)$$

The Jacobi identity relates the coefficients f_i and g_i

$$[[H, \bar{P}], q_\pm] = [H, [\bar{P}, q_\pm]] - [\bar{P}, [H, q_\pm]] = 0. \quad (A8)$$

This translates into an integrability condition implying the existence of a set of functions r_i such that $g_i = \partial_+ r_i$ and $f_i = \partial_- r_i$. Now let us use the shift freedom $q_\pm \rightarrow$

$q_{\pm} \mp \partial_{\pm} R$, with $R = \sum_i r_i(x^+, x^-) \Phi_i$. We then obtain the canonical form of the commutators

$$i[H, q_{\pm}] = \partial_- q_{\pm}, \quad i[\bar{P}, q_{\pm}] = \partial_+ q_{\pm}. \quad (\text{A9})$$

We see that any current associated to a symmetry generating charge that commutes with the hamiltonian and the momentum operator can be chosen so as to have no explicit dependence on the coordinates. In particular, this applies to the h_{\pm} and p_{\pm} current themselves. It is worth pointing out that, for any local (coordinate independent) operator Φ we can still shift $q_{\pm} \rightarrow q_{\pm} \mp \partial_{\pm} \Phi$ and preserve the above commutation relations.

Now we show, following [1], that the currents can also be made dilation eigenoperators. Current conservation and the commutation relations (4) imply:

$$\begin{aligned} i[D, h_{\pm}] &= x^- \partial_- h_{\pm} + \lambda(h_{\pm}) h_{\pm} \pm \partial_{\pm} O_h, \\ i[D, p_{\pm}] &= x^- \partial_- p_{\pm} + \lambda(p_{\pm}) p_{\pm} \pm \partial_{\pm} O_p, \end{aligned} \quad (\text{A10})$$

where $\lambda(p_+, p_-, h_+, h_-) = (0, 1, 1, 2)$. The Jacobi identity can be used to show that O_h and O_p are local operators with no explicit coordinate dependence. This means they are expandable in the (by assumption) discrete basis (5)

$$O_q = \sum_i a_i \Phi_i \quad (\text{A11})$$

for $q = h, p$ where Φ_i has weight λ_i . Let us now shift

$$q_{\pm} \rightarrow q_{\pm} \mp \partial_{\pm} \sum_i b_i \Phi_i, \quad b_i = \frac{a_i}{w(q_{\pm}) - \lambda_i}, \quad (\text{A12})$$

for $\lambda_i \neq w(q_{\pm})$. This shift eliminates all the Φ_i in O_q with weights $\lambda_i \neq w(q_{\pm})$. Operators with weight equal to $w(q_{\pm})$ cannot appear in O_q by the assumption (5) that the spectrum of D is discrete and diagonalizable. We are therefore left with the canonical form of the commutators:

$$\begin{aligned} i[D, h_{\pm}] &= x^- \partial_- h_{\pm} + \lambda(h_{\pm}) h_{\pm}, \\ i[D, p_{\pm}] &= x^- \partial_- p_{\pm} + \lambda(p_{\pm}) p_{\pm}. \end{aligned} \quad (\text{A13})$$

It was shown above that there exists a conserved current d_{\pm} associated to D . A slight variant of the preceding arguments shows that we can exploit the shift freedom to set

$$i[H, d_{\pm}] = \partial_- d_{\pm} - h_{\pm}, \quad i[H, d_{\pm}] = \partial_+ d_{\pm}. \quad (\text{A14})$$

The non trivial commutator $i[D, H] = H$ requires the extra term proportional to h_{\pm} , which implies that the dilation current cannot be independent of the coordinates.

Finally, the action of the dilation charge on its own associated current can be obtained as above and shifted to the canonical form

$$i[D, d_{\pm}] = x^- \partial_- d_{\pm} + \lambda(d_{\pm}) d_{\pm}, \quad (\text{A15})$$

where $\lambda(d_+, d_-) = (0, 1)$. We note that demanding that the current/charge commutators take this canonical form does not fix all the ambiguity in the former. Some of the remaining shift freedom is exploited in this work around equation (18) to shift away s_- .

Appendix B: Decoupling of s_+

In this appendix we show that in the non-minimal case $s_+ \neq 0$, $[\bar{J}_{\chi}, T_{\xi}] = 0$. Given that s_+ is independent of x^+ and transforms as a zero weight operator under D , we have

$$\begin{aligned} i[H, s_+] &= 0 \rightarrow i[H, \bar{J}_{\chi}] = 0 \\ i[D, s_+] &= 0 \rightarrow i[D, \bar{J}_{\chi}] = 0. \end{aligned} \quad (\text{B1})$$

The Jacobi identity implies that the commutator of an operator annihilated by H (such as \bar{J}_{χ}) and a local field (such as h_-) must be a local field itself. Therefore, using (B1), we have

$$i[\bar{J}_{\chi}, h_-] = \partial_-^2 \Phi_{\chi}, \quad (\text{B2})$$

where Φ_{χ} is a local operator of weight zero. This implies, $\partial_- \Phi_{\chi} = 0$ as an operator equation. Immediately we get

$$i[\bar{J}_{\chi}, T_{\xi}] = 0. \quad (\text{B3})$$

-
- [1] J. Polchinski, "Scale and Conformal Invariance in Quantum Field Theory," Nucl. Phys. B **303**, 226 (1988).
[2] J. L. Cardy, "Critical exponents of the chiral Potts model from conformal field theory," Nucl. Phys. **B389**, 577-586 (1993). [hep-th/9210002].
[3] O. Lunin and J. M. Maldacena, "Deforming field theories with $U(1) \times U(1)$ global symmetry and their gravity duals," JHEP **0505**, 033 (2005) [arXiv:hep-th/0502086].
[4] M. Guica, K. Skenderis, M. Taylor and B. C. van Rees, "Holography for Schrodinger backgrounds," JHEP **1102**, 056 (2011) [arXiv:1008.1991 [hep-th]].
[5] A. Bergman and O. J. Ganor, "Dipoles, twists and non-commutative gauge theory," JHEP **0010**, 018 (2000) [arXiv:hep-th/0008030].
[6] A. Bergman, K. Dasgupta, O. J. Ganor, J. L. Karczmarek and G. Rajesh, "Nonlocal field theories and their gravity duals," Phys. Rev. D **65**, 066005 (2002) [arXiv:hep-th/0103090].
[7] J. Maldacena, D. Martelli and Y. Tachikawa, "Comments on string theory backgrounds with non-relativistic conformal symmetry," JHEP **0810**, 072 (2008) [arXiv:0807.1100 [hep-th]].
[8] M. Guica and S. El-Showk, in progress; W. Song and A. Strominger in progress.
[9] M. Guica, T. Hartman, W. Song and A. Strominger, "The Kerr/CFT Correspondence," Phys. Rev. D **80**,

- 124008 (2009) [arXiv:0809.4266 [hep-th]].
- [10] M. Guica and A. Strominger, “Microscopic Realization of the Kerr/CFT Correspondence,” *JHEP* **1102**, 010 (2011) [arXiv:1009.5039 [hep-th]].
- [11] J. M. Bardeen, G. T. Horowitz, “The Extreme Kerr throat geometry: A Vacuum analog of $AdS(2) \times S^{*2}$,” *Phys. Rev.* **D60**, 104030 (1999). [hep-th/9905099].
- [12] S. Detournay, D. Israel, J. M. Lapan and M. Romo, “String Theory on Warped AdS_3 and Virasoro Resonances,” *JHEP* **1101**, 030 (2011) [arXiv:1007.2781 [hep-th]].
- [13] M. J. Duff, H. Lu, C. N. Pope, “ $AdS(5) \times S^{*5}$ untwisted,” *Nucl. Phys.* **B532**, 181-209 (1998). [hep-th/9803061].
- [14] D. Anninos, “Hopfing and Puffing Warped Anti-de Sitter Space,” *JHEP* **0909**, 075 (2009). [arXiv:0809.2433 [hep-th]].
- [15] R. Dijkgraaf, “Chiral deformations of conformal field theories,” *Nucl. Phys.* **B493**, 588-612 (1997). [hep-th/9609022].
- [16] R. N. Caldeira Costa and M. Taylor, “Holography for chiral scale-invariant models,” *JHEP* **1102**, 082 (2011) [arXiv:1010.4800 [hep-th]].
- [17] A. M. Polyakov, A. B. Zamolodchikov, “Fractal Structure of Two-Dimensional Supergravity,” *Mod. Phys. Lett.* **A3**, 1213-1219 (1988).
- [18] G. A. T. F. da Costa, L. O’Raifeartaigh, “Nontrivial noncanonical W algebras from Kac-Moody reductions,” *Phys. Lett.* **B333**, 353-363 (1994). [hep-th/9405115].
- [19] A. Strominger, “ $AdS(2)$ quantum gravity and string theory,” *JHEP* **9901**, 007 (1999) [arXiv:hep-th/9809027].
- [20] I. Bredberg, T. Hartman, W. Song, A. Strominger, “Black Hole Superradiance From Kerr/CFT,” *JHEP* **1004**, 019 (2010). [arXiv:0907.3477 [hep-th]].
- [21] G. Compere, S. Detournay, “Semi-classical central charge in topologically massive gravity,” *Class. Quant. Grav.* **26**, 012001 (2009). [arXiv:0808.1911 [hep-th]].