

2009-05-09

CHOP mediates endoplasmic reticulum stress-induced apoptosis in Gimap5-deficient T cells

Steven C. Pino
University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: <https://escholarship.umassmed.edu/oapubs>



Part of the [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)

Repository Citation

Pino SC, O'Sullivan-Murphy B, Lidstone EA, Yang C, Lipson KL, Jurczyk A, Diiorio PJ, Brehm MA, Mordes JP, Greiner DL, Rossini AA, Bortell R. (2009). CHOP mediates endoplasmic reticulum stress-induced apoptosis in Gimap5-deficient T cells. Open Access Publications by UMass Chan Authors. <https://doi.org/10.1371/journal.pone.0005468>. Retrieved from <https://escholarship.umassmed.edu/oapubs/2081>

This material is brought to you by eScholarship@UMassChan. It has been accepted for inclusion in Open Access Publications by UMass Chan Authors by an authorized administrator of eScholarship@UMassChan. For more information, please contact Lisa.Palmer@umassmed.edu.

CHOP Mediates Endoplasmic Reticulum Stress-Induced Apoptosis in Gimap5-Deficient T Cells

Steven C. Pino¹, Bryan O'Sullivan-Murphy¹, Erich A. Lidstone¹, Chaoxing Yang¹, Kathryn L. Lipson¹, Agata Jurczyk¹, Philip dilorio¹, Michael A. Brehm¹, John P. Mordes¹, Dale L. Greiner^{1,2}, Aldo A. Rossini^{1,2}, Rita Bortell^{1*}

¹ Department of Medicine, University of Massachusetts Medical School, Worcester, Massachusetts, United States of America, ² Molecular Medicine, University of Massachusetts Medical School, Worcester, Massachusetts, United States of America

Abstract

Gimap5 (GTPase of the immunity-associated protein 5) has been linked to the regulation of T cell survival, and polymorphisms in the human *GIMAP5* gene associate with autoimmune disorders. The BioBreeding diabetes-prone (BBDP) rat has a mutation in the *Gimap5* gene that leads to spontaneous apoptosis of peripheral T cells by an unknown mechanism. Because Gimap5 localizes to the endoplasmic reticulum (ER), we hypothesized that absence of functional Gimap5 protein initiates T cell death through disruptions in ER homeostasis. We observed increases in ER stress-associated chaperones in T cells but not thymocytes or B cells from *Gimap5*^{-/-} BBDP rats. We then discovered that ER stress-induced apoptotic signaling through C/EBP-homologous protein (CHOP) occurs in *Gimap5*^{-/-} T cells. Knockdown of CHOP by siRNA protected *Gimap5*^{-/-} T cells from ER stress-induced apoptosis, thereby identifying a role for this cellular pathway in the T cell lymphopenia of the BBDP rat. These findings indicate a direct relationship between Gimap5 and the maintenance of ER homeostasis in the survival of T cells.

Citation: Pino SC, O'Sullivan-Murphy B, Lidstone EA, Yang C, Lipson KL, et al. (2009) CHOP Mediates Endoplasmic Reticulum Stress-Induced Apoptosis in Gimap5-Deficient T Cells. PLoS ONE 4(5): e5468. doi:10.1371/journal.pone.0005468

Editor: Dominik Hartl, LMU University of Munich, Germany

Received: February 25, 2009; **Accepted:** April 5, 2009; **Published:** May 8, 2009

Copyright: © 2009 Pino et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Research supported in part by research grants DK49106 (D.L.G.), DK25306 (A.A.R.), and DK066447 (J.P.M.) from the National Institutes of Health, an Autoimmunity Prevention Center grant supported by the NIAID (A.A.R.), Diabetes Endocrinology Research Center (DERC) Grant DK32520, a grant from the American Diabetes Association (J.P.M.), and a grant funded jointly by The Iacocca Foundation and the JDRF (D.L.G. and R.B.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: rita.bortell@umassmed.edu

Introduction

The expression products of the GTPase of the immunity-associated gene family (gimaps; formerly known as IANs or immune-associated nucleotide-binding proteins) have been implicated in the regulation of T cell survival through modulation of T cell receptor (TCR) signaling [1,2]. A member of this family, *GIMAP5*, has an important role in human immune modulation as a polyadenylation polymorphism in the human *GIMAP5* gene increases systemic lupus erythematosus risk [3] and is also associated with elevated islet autoantibodies in type 1 diabetics [4]. Gimap protein may also play a significant role in T cell homeostasis because peripheral T cell survival is impaired in both *gimap5* null mice [5] and in BioBreeding diabetes-prone (BBDP) rats, which have a frameshift mutation in the *Gimap5* gene [6–8]. In the BBDP rat, Gimap5 is a truncated protein in which the carboxy-terminal 215 amino acids are replaced by 19 other amino acids. Absence of functional wild type Gimap5 protein results in mitochondrial dysfunction and apoptosis *in vitro*, and severe T cell lymphopenia *in vivo* [9–11]. Knockdown of human Gimap5 in Jurkat T cells *in vitro* recapitulates the natural apoptosis that occurs in T cells from the BBDP rat [11]. However, the mechanism by which *Gimap5* promotes T cell survival is not understood.

Gimap5 has been reported to localize mainly to the endoplasmic reticulum (ER) [12,13] but is also detected in mitochondria,

the Golgi apparatus, and centrosomes [13,14]. The ER is a multifunctional organelle which plays a vital role in the regulation of numerous cellular processes, including synthesis, folding, and assembly of secretory and transmembrane proteins [15,16], calcium (Ca²⁺) signaling regulation, vesicle trafficking, drug metabolism, and lipid biogenesis [17–19]. As the main organelle involved in protein processing within a cell, the ER has evolved numerous signaling pathways that monitor its protein folding capacity and ensure that these pathways do not become dysfunctional [16]. Pathological disturbances to normal cellular functioning, including disruption of protein folding or alterations in Ca²⁺ homeostasis [20], interfere with the functioning of the ER and cause stress to the organelle.

Normally, stress within the ER triggers an adaptive cellular mechanism known as the unfolded protein response or ER stress response that attempts to return the cell to homeostasis [21,22]. Activation of the ER stress response is accomplished by specific ER transmembrane proteins that function as ER stress transducers [23]. Glucose-regulated protein 78 (GRP78) is the central regulator of the ER stress response. It normally binds to and inhibits signaling from ER stress transducers. However, during ER stress GRP78 is released from those transducers, thereby activating the ER stress response [24]. If ER stress persists and cellular homeostasis cannot be restored, the ER stress response can initiate cell death stimuli that lead to ER stress-induced apoptosis [18].

Initiation of ER stress-induced apoptosis through ER stress response signaling involves transcriptional activation of the bZIP transcription factor *chop* (C/EBP-homologous protein) [25]. CHOP acts to repress the promoter of the *bcl-2* gene, thus downregulating antiapoptotic Bcl-2 protein and rendering cells sensitive to the proapoptotic effects of BH3-only proteins [26]. Because *Gimap5* localizes primarily to the ER [12,13], we hypothesized that *Gimap5* protein in T cells is necessary to maintain ER homeostasis. In this report, we show that ER stress response signaling occurs in *Gimap5*^{-/-} T cells as evidenced by an increase in ER chaperone expression. We further show that initiation of ER stress-induced apoptotic signaling in *Gimap5*^{-/-} T cells is associated with an increase of CHOP protein. Knockdown of CHOP expression protects *Gimap5*^{-/-} T cells from ER stress-induced cell death, thus revealing an essential role for *Gimap5* in the maintenance of ER homeostasis and T cell survival.

Results

Increased expression of ER stress response proteins in *Gimap5*^{-/-} BBDR rat lymphocytes

Gimap5 is normally expressed in the thymus and in both T and B cells of lymphoid tissues [11]. However, depletion of functional *Gimap5*, whether by RNAi, genetic knockout or mutation, leads to spontaneous apoptosis only in T cells [5,11]. To examine the potential role of ER stress in T cell death of *Gimap5*^{-/-} rats, we compared the expression of ER chaperone proteins in lymph node cells and thymocyte lysates from *Gimap5*^{-/-} BBDR and *Gimap5*^{+/+} BBDR rats. Western blot analyses revealed that GRP94, GRP78, and ER protein 72 (ERp72) levels were increased in lymphocytes from *Gimap5*^{-/-} BBDR rats when compared to that in *Gimap5*^{+/+} BBDR rats (Fig. 1). However, ER chaperone expression was similar in thymocytes from *Gimap5*^{-/-} BBDR and *Gimap5*^{+/+} BBDR rats. These results indicate that the ER stress response leading to increased ER chaperone expression is specifically upregulated in lymph node cells and not thymocytes in *Gimap5*^{-/-} BBDR rats.

To determine if ER stress-induced apoptosis was also activated in *Gimap5*^{-/-} lymph node cells, we examined expression of the ER stress apoptotic factor, CHOP. CHOP expression is minimal and difficult to detect under normal homeostatic conditions [27], and upregulation of CHOP has been reported to signal the activation of ER stress-mediated apoptotic signals [25]. These signals ultimately culminate in cell death through proteolytic cleavage of caspase-3 [28]. We found that CHOP protein is abundantly expressed in lymph node cells from *Gimap5*^{-/-} BBDR rats, but is not detectable in BBDR thymocytes or in thymocytes or lymphocytes from *Gimap5*^{+/+} BBDR rats (Fig. 1). Furthermore, the presence of activated (cleaved) caspase-3, the executor of apoptosis downstream of ER stress-mediated apoptotic signaling, was only detected in *Gimap5*^{-/-} BBDR lymphocytes. As expected, *Gimap5* protein was not detected in lymph node cells from *Gimap5*^{-/-} BBDR rats, but was expressed at comparable levels in both lymph node cells and thymocytes from *Gimap5*^{+/+} BBDR rats (Fig. 1). These data provide evidence of an ER stress response and ER stress-induced apoptotic signaling that is specific to lymph node cells from *Gimap5*^{-/-} BBDR rats.

Increased GRP78 and CHOP in both CD4⁺ and CD8⁺ *Gimap5*^{-/-} BBDR T cells

Because BBDR rats have a peripheral T cell lymphopenia that, in transgenic rescue experiments [29], has been proven to be due to its mutation in the *Gimap5* gene [9,10,30], we hypothesized that the increased levels of ER stress proteins found in BBDR rat lymph




Figure 1. The expression of ER chaperones increases in lymphocytes from the *Gimap5*^{-/-} BBDR rat. Western blot analyses of ER chaperone proteins GRP94, GRP78, and ERp72; the ER stress apoptotic factor CHOP; and cleaved caspase-3 (CASP-3) in lymphocytes and thymocytes from BBDR and BBDR rats. *Gimap5* protein expression was confirmed in lysates from BBDR rats and actin was used as a loading control. Data shown are representative of three independent experiments.

doi:10.1371/journal.pone.0005468.g001

node cells preparations (Fig. 1) would be associated specifically with the T cell compartment. As expected, we found a decreased percentage of TCR⁺ cells in the lymph nodes of BBDR rats ($22.8 \pm 0.4\%$) as compared to *Gimap5*^{+/+} BBDR rats ($68.5 \pm 1.4\%$, $n = 4$; $P < 0.001$, Fig. 2A). We also observed a concomitant enrichment of B cells (TCR⁻CD45RA⁺) in BBDR ($69.1 \pm 0.7\%$) vs BBDR rats ($27.5 \pm 0.9\%$, $n = 4$; $P < 0.001$, Fig. 2B). In cervical and mesenteric lymph nodes, the absolute number of T cells in the BBDR rat ($23.9 \pm 2.2 \times 10^6$) was significantly less than in the *Gimap5*^{+/+} BBDR rats ($123.3 \pm 15.0 \times 10^6$, $n = 4$; $P < 0.01$), whereas B cell numbers were significantly increased ($75.6 \pm 5.2 \times 10^6$ vs $50.0 \pm 6.2 \times 10^6$, $n = 4$; $P < 0.05$). To determine the T cell-specific levels of ER stress response and ER stress-induced apoptotic signaling, we performed intracellular flow cytometry. As measured by the mean fluorescence intensity (MFI), both GRP78 and CHOP levels were higher in TCR⁺ lymphocytes from *Gimap5*^{-/-} BBDR rats than from *Gimap5*^{+/+} BBDR rats (Fig. 2A). In contrast, B lymphocytes (TCR⁻CD45RA⁺ cells) exhibited no difference in GRP78 and CHOP expression between the two rat strains (Fig. 2B).




Figure 2. *Gimap5*^{-/-} BBBDP rat T cell populations exhibit enhanced ER stress response signaling. Representative analyses showing the gating of TCR⁺ (A), TCR⁻CD45RA⁺ (B), and CD4⁺ or CD8⁺ (C) lymphocytes from BBBDP and BBDR rats. Above the representative gate is shown the average percentage for that cell type from 4 rats of each strain (\pm SEM). Representative histograms show the percentage of cells expressing intracellular GRP78 and CHOP protein at the indicated levels in BBBDP (black line) and BBDR (shaded region) rat T cells (A), B cells (B), and CD4⁺ or CD8⁺ T cells (D,E). The corresponding isotype control is depicted as a dotted line in each histogram. Bar graphs display the MFI in arbitrary units (AU) of intracellular GRP78 and CHOP protein in each cell type (\pm SEM). Data shown are representative of three independent experiments (** P <.01, *** P <.001).

doi:10.1371/journal.pone.0005468.g002

We next examined ER stress response activation within peripheral CD4⁺ and CD8⁺ T cells. Consistent with previous observations [30], we found that the CD8⁺ T cell subset in BBBDP rats ($1.2 \pm 0.07\%$ vs $16.9 \pm 1.4\%$ in BBDR, $n = 3$) was reduced to a much greater degree than was the CD4⁺ subset ($37.0 \pm 1.5\%$ vs $57.0 \pm 0.6\%$ in BBDR, $n = 4$) (Fig. 2C). Absolute CD4⁺ T cell number in lymph node cells were $40.4 \pm 4.0 \times 10^6$ cells in the BBBDP rat vs $102.5 \pm 11.7 \times 10^6$ in the BBDR ($n = 4$; $P < 0.01$). Absolute CD8⁺ T cell number in lymph node cells were $1.4 \pm 0.1 \times 10^6$ in the BBBDP rat vs $30.6 \pm 4.5 \times 10^6$ in the BBDR, ($n = 4$; $P < 0.01$).

To determine whether ER stress levels are higher in the CD8⁺ subset, we again analyzed GRP78 and CHOP by intracellular flow cytometry. We observed that GRP78 and CHOP expression was similar in both CD4⁺ and CD8⁺ T cells from the *Gimap5*^{-/-} BBBDP rat, but both subsets of BBBDP T cells had GRP78 and CHOP levels that were higher than their counterparts from the *Gimap5*^{+/+} BBDR rat (Fig. 2D,E). These data suggest that the ER stress response and ER stress-induced apoptotic signaling are activated in both CD4⁺ and CD8⁺ T cells in the BBBDP rat.

Increased ER stress response signaling occurs when *Gimap5*^{-/-} T cells first exit the thymus

Thymocyte development in the *Gimap5*^{-/-} BBBDP rat is generally normal [31], with typical differentiation of double positive CD4⁺CD8⁺ cells to single positive thymocyte precursors [30]. Consistent with previous reports [30], our flow cytometry analyses revealed similar percentages of CD4⁺CD8⁺, CD4⁺CD8⁻, and CD4⁻CD8⁺ thymocytes in both BBBDP and BBDR rats (BBBDP: $89.8 \pm 1.0\%$, $5.2 \pm 0.6\%$, $4.8 \pm 1.0\%$; BBDR: $88.3 \pm 0.6\%$, $6.4 \pm 0.4\%$, $4.9 \pm 0.5\%$, respectively, $n = 3$; Fig. 3A). Western blot analyses (Fig. 1) revealed no substantial difference in ER stress proteins in thymocytes of BBBDP vs BBDR rats. To determine if enhanced ER stress signaling might be initiated in only a small subpopulation of thymocytes, we employed flow cytometry. Comparison of thymocytes isolated from BBBDP and BBDR rats revealed no significant difference in GRP78 and CHOP levels within the CD4⁺CD8⁺, CD4⁺CD8⁻, and CD4⁻CD8⁺ thymocyte subsets (Fig. 3B,C). These data suggest that enhanced ER stress response signaling is *not* initiated in *Gimap5*^{-/-} thymocytes prior to thymic export.




Figure 3. ER stress response signaling is similar in thymocytes from *Gimap5*^{-/-} BBDP and *Gimap5*^{+/+} BBDR rats. (A) Representative flow cytometric dot plots depicting CD8 expression (horizontal axis) and CD4 expression (vertical axis) on thymocytes. Numbers represent the percentage of cells (\pm SEM of triplicate samples) in each gate shown. Representative histograms depicting the expression of intracellular GRP78 (B) and CHOP (C) protein in thymocyte populations from BBDP (black line) and BBDR (shaded region) rats. Isotype control for GRP78 or CHOP staining (dotted line) is shown in each histogram. Bar graphs display the MFI in arbitrary units (AU) of intracellular GRP78 or CHOP protein expression (\pm SEM). Data shown are representative of three independent experiments. doi:10.1371/journal.pone.0005468.g003

To determine the precise stage in CD4⁺ and CD8⁺ T cell development at which enhanced ER stress signaling is initiated, we next compared GRP78 and CHOP levels in mature T cells and recent thymic emigrants (RTEs, identified by the expression of CD90/Thy-1). Flow cytometry demonstrated that the preponderance of CD4⁺ and CD8⁺ T cells from *Gimap5*^{-/-} BBDP rats still express CD90 on the cell surface, indicating that few T cells survive to the mature CD90⁻ stage (Fig. 4A). In contrast, the majority of CD4⁺ and CD8⁺ T cells from the *Gimap5*^{+/+} BBDR rats mature and down-regulate CD90 (Fig. 4B). CD4⁺ and CD8⁺ RTEs from BBDP rats initiate significantly enhanced ER stress signals as compared to BBDR rats, as evidenced by increased GRP78 and CHOP expression (Fig. 4C,D). These increased levels of GRP78 and CHOP continue throughout T cell development into the small population of mature CD4⁺ and CD8⁺ CD90⁻ T cells that survive in the *Gimap5*^{-/-} BBDP rat (Fig. 4C,D). Similar findings were also observed in congenic WF.*ART2a.Gimap5*^{-/-} rats that harbor the same recessive mutation in the *Gimap5* gene as do BBDP rats, but do not develop spontaneous diabetes (Figure S1). All T cell populations, except for TCR⁺CD8⁺CD90⁺ cells from WF.*ART2a.Gimap5*^{-/-} rats, contained increased expression of GRP78 and CHOP, as compared to control WF.*ART2a.Gimap5*^{+/+} congenic rats. We attribute the difference in ER stress response signaling in the TCR⁺CD8⁺CD90⁺ population to genetic background differences which occur between the BB and WF rat strains. These results indicate that ER stress response signaling is

not a secondary consequence of diabetes development. Overall, our data reveal a central role for *Gimap5* protein in ER homeostasis during post-thymic development when T cells first enter the periphery.

Enhanced GRP78 and CHOP expression is not dependent on T cell activation

BBDP rat T cells have been reported to exhibit an unusual state of partial activation following thymic export [2,31]. Our data reveal a significant increase in the percentages of CD4⁺ ($n = 3$; $P < 0.001$) and CD8⁺ ($n = 3$; $P = 0.01$) RTEs and mature CD4⁺ ($n = 3$; $P < 0.001$) and CD8⁺ ($n = 3$; $P < 0.01$) T cells from the *Gimap5*^{-/-} BBDP rat that express CD25, a marker of activation (see Fig. 5A,B for representative data). Because we and others have shown that activation of T cells causes ER stress response signaling [32,33], we wanted to determine if there was a link between cellular activation status and elevated GRP78 and CHOP expression in BBDP T cells. In the BBDP rat, CD4⁺ and CD8⁺ RTEs and mature T cells (CD90⁻) that express CD25 have significantly increased GRP78 expression as compared to cells with the same phenotype in the BBDR rat (Fig. 5C). However, the CD25⁻ T cells that survive in the BBDP rat also have significantly increased GRP78 levels as compared with CD25⁻ T cells from the BBDR rat (Fig. 5D). Expression of CHOP within CD25⁺ and CD25⁻ T cell populations was similar to that of GRP78 with respect to activation with one exception. Only




Figure 4. Enhanced ER stress response signaling is initiated in RTEs of the *Gimap5*^{-/-} BBDR rat and continues during maturation to the CD90⁻ stage. (A,B) Representative flow cytometric dot plots depicting CD90 expression (horizontal axis) and forward scattering (vertical axis) of lymphocytes within gated CD4⁺ and CD8⁺ T cell populations in BBDR or BBDR rats. Numbers represent the corresponding average of CD90⁻ and CD90⁺ T cells (\pm SEM of triplicate samples). The expression of intracellular GRP78 (C) or CHOP (D) protein in representative histograms (BBDR, black line; BBDR, shaded region; isotype control staining, dotted line). The MFI of GRP78 or CHOP protein expression is indicated in bar graphs with error bars representing the SEM. Data shown are representative of three independent experiments (* P <.05; ** P <.01; *** P <.001). doi:10.1371/journal.pone.0005468.g004

CD4⁺CD90⁻CD25⁺ and CD8⁺CD90⁻CD25⁺ T cells from BBDR rats expressed significantly higher CHOP levels than did cells with the same phenotype from BBDR rats (Fig. 5E,F). These data indicate that, although ER stress levels do increase with activation status as previously reported, absence of functional *Gimap5* protein is by itself sufficient to initiate ER stress response and ER stress-induced apoptotic signaling without concomitant T cell activation.

Knockdown of CHOP protein expression in *Gimap5*^{-/-} rat T cells inhibits apoptosis and cell death

CHOP protein plays an important role in ER stress-induced programmed cell death and was found to be increased only in *Gimap5*^{-/-} BBDR rat T cells (Figures 2,4). To investigate a role for CHOP in apoptotic signaling of these T cells, we purified and transfected BBDR T cells with siRNAs targeted to CHOP (siCHOP) or non-specific control siRNA (siControl). Western blot analyses revealed a significant reduction in CHOP protein expression in T cells transfected with siCHOP as compared to siControl transfected T cells (Fig. 6A). The effects of reducing CHOP protein expression on induction of ER stress-induced apoptosis were examined by flow cytometry. Annexin V and 7AAD staining revealed three distinct populations of T cells: viable (annexinV⁻7AAD⁻, lower left quadrant), early apoptotic (annexinV⁺7AAD⁻, upper left quadrant), and late apoptotic and necrotic (annexinV⁺7AAD⁺, upper right quadrant) (Fig. 6B). As compared to T cells transfected with siControl, CHOP depletion significantly decreased the percentage of T cells in late apoptosis, with a

concomitant increase in the percentage of non-apoptotic T cells (Fig. 6B). In contrast, the percentage of cells that were in the early stages of apoptosis were similar in T cells transfected with siCHOP or siControl (Fig. 6B).

To ensure that the results of CHOP depletion were specific to T cells, the effects of siCHOP transfection were also examined in TCR⁻ lymphocytes. In contrast to T cells, there were no statistical differences in the percentages of viable, early apoptotic, and late apoptotic TCR⁻ lymphocytes between siCHOP and siControl transfected cells (Fig. 6C). In their aggregate, these data suggest that CHOP protein actively participates in apoptotic signaling in *Gimap5*^{-/-} T cells, and reduction of this ER stress apoptotic factor inhibits T cell death.

Discussion

In this study we analyzed the ER stress response in T cells from *Gimap5*^{-/-} BBDR rats. We demonstrate that absence of functional full length *Gimap5* protein in T cells leads to ER stress response signaling as evidenced by increased GRP78 protein expression. In addition, our data show that BBDR T cells initiate ER stress-induced apoptotic signaling through upregulation of CHOP protein. Finally, decreasing the expression of this ER stress apoptotic factor, protected BBDR T cells from ER stress-induced death. These observations suggest that *Gimap5* protein plays an essential role in the maintenance of ER homeostasis and is indispensable for T cell survival.

Lymphopenia in the BBDR rat was originally linked to a recessive mutation in a diabetes susceptibility locus designated




Figure 5. Enhanced ER stress response signaling in *Gimap5*^{-/-} BBDP rat T cells is not dependent upon activation status. Representative flow dot plots depicting CD25 expression (horizontal axis) and forward scatter (vertical axis) of lymphocytes within gated CD4⁺ and CD8⁺ RTEs and mature T cells from BBDP (A) and BBDR (B) rats. Numbers represent the percentage (\pm SEM of triplicate samples) of corresponding CD25⁻ and CD25⁺ T cells. Bar graphs displaying the MFI of intracellular GRP78 and CHOP expression in CD25⁺ (C,E) and CD25⁻ (D,F) T cells. Error bars represent the SEM; data shown are representative of three independent experiments (* P <.05; ** P <.01; *** P <.001). doi:10.1371/journal.pone.0005468.g005

lyp/Idm1, which encodes for *Gimap5* protein [9,10]. Although spontaneous diabetes development in the BBDP rat depends on the presence of *lyp/Idm1*, diabetes susceptibility and lymphopenia are traits that can be inherited independently [34]. To ensure that ER stress response signaling in T cells from the BBDP rat is due to absence of functional *Gimap5* protein and not a byproduct of diabetes development, we only used nondiabetic BBDP rats for studies. Furthermore, flow cytometry

analyses on *Gimap5*^{-/-} T cells from congenic *WF.ART2a.Gimap5*^{-/-} rats that do not develop spontaneous diabetes but carry the *Gimap5* null mutation have increased expression of ER stress response proteins, including GRP78 and CHOP (Figure S1E,F). Overall, this indicates that ER stress response signaling in T cells from the BBDP rat is a result of the absence of functional *Gimap5* protein product and not a secondary consequence of spontaneous diabetes development.




Figure 6. Knockdown of CHOP protein expression enhances the viability of T cells from *Gimap5*^{-/-} rats. (A) Western blot analyses of the ER stress apoptotic factor CHOP in purified T cells transfected with either CHOP siRNA (siCHOP) or non-specific control siRNA (siControl). Shown are results of two separate knockdown experiments. Actin was used as a loading control. (B,C) Representative flow dot plots depicting cells positive for annexin V (vertical axis) and 7AAD staining (horizontal axis) on gated TCR⁺ or TCR⁻ lymphocytes. Bar graphs display the percentage of TCR⁺ or TCR⁻ lymphocytes that were positive for both annexin V and 7AAD staining (left), positive for annexin V staining alone (middle), and negative for both staining (right). Error bars represent the SEM of quadruplicate samples; data shown are representative of two independent experiments (***) $P < .001$.

doi:10.1371/journal.pone.0005468.g006

Within the immune system, the establishment of immunological tolerance involves mechanisms to delete self-reactive T cells in order to avoid autoimmune diseases such as type 1 diabetes [35]. The interaction of TCRs on thymocytes with thymic stromal cells are important for proper T cell development and eventually allow for CD4⁺CD8⁺ double positive thymocytes to differentiate into mature CD4⁺CD8⁻ or CD4⁻CD8⁺ single positive thymocytes [36,37]. Our data reveal that the percentage of double positive and single positive thymocytes does not differ between BBDR and BBBD rats, despite the absence of *Gimap5* in the BBBD rat. Furthermore, ER stress response signaling is similar in the thymocyte populations from both BBDR and BBBD rats indicating the main ER regulatory functions for *Gimap5* protein may occur during T cell post-thymic development. In general, our data indicate that ER stress response and ER stress-mediated signaling may only occur minimally during thymocyte development as levels of ER chaperones remained constant as double positive thymocytes matured to the single positive stage.

Following thymic development, T cells still expressing CD90, known as RTEs, enter the circulation. In the BBBD rat, a limited number of RTEs survive, down-regulate CD90 expression and become mature T cells, as others have reported and we have confirmed (Fig. 4A) [38]. Because RTEs undergo apoptosis rapidly in the *Gimap5*^{-/-} BBBD rat, we initially hypothesized that ER stress protein levels would be higher in RTEs than in CD90⁻

mature T cells from the same animal. Our data suggest that ER chaperone expression is similar in all CD90⁺ and CD90⁻ *Gimap5*^{-/-} T cell subpopulations in the BBBD rat. Our data further demonstrate that the percentage of the peripheral CD90⁺ and CD90⁻ T cells that express the activation marker CD25 is higher in the BBBD rat than in the BBDR rat (Fig. 5A,B). Although activation of T cells is known to enhance ER stress [32,33], CD25⁻ T cell populations from the *Gimap5*^{-/-} BBBD rat expressed ER stress levels significantly higher than their counterparts from the *Gimap5*^{+/+} BBDR rat. These data indicate a global absence of the protein is sufficient to initiate ER stress response signaling throughout peripheral T cell development and that activation signals are not required for ER stress induction.

ER stress response signaling has been shown to be required for numerous physiological functions in several cell types [32,33,39,40]. In addition, pathological signals from the ER have been attributed to cell death and apoptosis and linked to many diseases, including diabetes [18,41]. In this report, we show that absence of functional *Gimap5* protein leads to these pathological signals from the ER and subsequent ER stress-induced apoptosis in T cells from the BBBD rat. We suggest that triggering of the ER stress-induced apoptotic pathway in the T cells of BBBD rats is the underlying mechanism for their lymphopenia. This inference is supported by our data demonstrating that decreasing CHOP protein expression reduces the number of apoptotic T cells.

Interestingly, although B cells normally express *Gimap5*, we found that ER stress response signaling in B cells is similar between *Gimap5*^{+/+} BBDR and *Gimap5*^{-/-} BBDR rats (Fig. 2B). These data suggest that modulation of ER stress response signaling by *Gimap5* is required for the maintenance of T cell survival.

As documented above, this report demonstrates that ER stress-induced apoptosis occurs in *Gimap5*^{-/-} T cells. However, we and others have previously shown a role for *Gimap5* protein in the maintenance of mitochondrial integrity for T cell survival [11,42]. These two *Gimap5*-dependent processes are both linked through CHOP, a protein that is both highly expressed in cells undergoing ER stress-induced cell death and also capable of sensitizing mitochondria to numerous apoptotic factors [43]. This sensitization occurs via downregulation of antiapoptotic Bcl-2 protein, activation of JNK and its associated proapoptotic downstream kinases [18], and activation of caspase-12 [44]. We hypothesize that CHOP protein through its role in ER stress-induced apoptotic signaling may lead to the disruption of mitochondrial integrity that is observed in *Gimap5*^{-/-} T cells [11,42]. We propose that loss of mitochondrial integrity in *Gimap5*^{-/-} T cells may be secondary to ER dysfunction and subsequent ER stress-induced apoptotic signaling.

The intracellular distribution of *Gimap5* protein remains controversial, but recent research indicates localization to the ER [12]. Our data support an intimate link between wild type *Gimap5* and the ER due to ER stress response signaling in *Gimap5*^{-/-} T cells. Accumulation of intracellular proteins is known to cause ER stress and can lead to cell death through CHOP signaling [45]. This association has been demonstrated in pancreatic beta cells of the Akita mouse in which accumulation of mutant insulin causes ER stress and eventual apoptosis through CHOP protein induction [46]. We observed an accumulation of $\alpha\beta$ TCR protein within *Gimap5*^{-/-} BBDR rat lymphocytes that correlates with increased GRP78 levels (unpublished observations).

In the BBDR rat there is the frameshift mutation in *Gimap5* which results in a truncated protein product [12]. We can not rule out the possibility that the presence of this truncated protein product or increased intracellular $\alpha\beta$ TCR protein may accumulate in the ER and lead to the pathological ER stress that is observed in *Gimap5*^{-/-} T cells. In either case, disruption of CHOP was effective in delaying apoptosis and preventing cell death, thus indicating the importance of CHOP in ER stress-associated T cell death signaling. Interestingly, overexpression of either wild type rat *Gimap5* or the truncated BBDR rat *Gimap5* in a rat T cell line is associated with enhanced apoptosis [42]. In this report, the apoptosis-inducing effect of mutant *Gimap5* was much greater than that of the overexpressed wild type protein. In the context of our data, this may indicate that the expression of *Gimap5* in T cells is both required for survival and tightly regulated.

In conclusion, our findings suggest a role for ER stress in mediating the spontaneous apoptosis found in T cells from *Gimap5*^{-/-} rats. It will be of interest to determine if the same mechanisms are responsible for the T cell lymphopenia and hepatic dysfunction observed in *gimap5*^{-/-} mice [5]. We provide the first evidence of a link between *Gimap5* protein and the maintenance of ER homeostasis and cellular integrity. Our results indicate that proper functioning of the ER in T cells is critical for their survival. Investigations into the transient lymphopenia which occurs following virus infections in humans [47] and in the blood of patients with type 1 diabetes [48] may reveal a role for ER regulatory pathways in mediating human immunity. Further studies of wild type and mutant *Gimap5* protein are needed to precisely define the mechanisms by which they affect T cell

survival and function. Such studies may uncover novel pathways capable of modulating the survival and function of T cells and other tissues.

Materials and Methods

Animals

Gimap5^{-/-} BBDR and *Gimap5*^{+/+} BBDR rats were obtained from Biomedical Research Models (Worcester, MA). BBDR rats are T cell lymphopenic in both the CD4 and to an even greater extent in the CD8 compartments; BBDR rats have a normal peripheral lymphocyte phenotype [30]. Eight week-old male or female rats were used, and all rats were nondiabetic at the time of study. Congenic *WF.ART2a.Gimap5*^{-/-} rats were also developed by us by crossing WF (Charles River Breeding Laboratories, Wilmington, MA) and BBDR rats. The intercross was followed by 8 backcrosses to WF before intercrossing to establish homozygosity for both the *Iddm14* [49] and *lyp* (*Gimap5*) loci on chromosome 4 and the *ART2a* locus [50] on chromosome 1. The *WF.ART2a.Gimap5*^{-/-} rat bears no BB markers except in the regions of chromosome 4 bounded by *DARat101-DARat58* and chromosome 1 bounded by *D1Rat29* and *D1Rat287*, and it does not develop spontaneous diabetes (unpublished observations). Control *WF.ART2a* congenic rats were developed by us and differ from ordinary non-lymphopenic, non-diabetic WF animals in that they express the “a” rather than the “b” allotype of the ART2 T cell alloantigen on chromosome 1 [50,51]. *WF.ART2a* are nondiabetic and homozygous for wild type *Gimap5*. All animals were housed in a viral antibody-free facility and maintained in accordance with the guidelines of the University of Massachusetts Medical School Institutional Animal Care and Use Committee and the *Guide for the Care and Use of Laboratory Animals* (Institute of Laboratory Animal Resources, National Research Council, National Academy of Sciences, 1996).

Thymi and lymph node cell preparation

Thymi and cervical and mesenteric lymph nodes were removed from BBDR and BBDR rats, processed aseptically, and extruded through mesh sieves. Single cell suspensions for flow cytometry were prepared as described below.

Western blotting

Thymi, lymphocytes, or purified T cells were lysed [52] and protein concentrations determined by bicinchoninic acid (BCA) protein assay (Sigma-Aldrich, St. Louis, MO). Protein (30 μ g) was mixed with 4 \times SDS-PAGE loading buffer and Western blot analyses were performed as described [52]. Actin was used as a loading control.

Antibodies

Anti-rat CD8a-PE (clone Ox-8), FITC or PerCp-conjugated anti-rat TCR $\alpha\beta$ (clone R73), anti-rat CD90-FITC (clone HIS51), Biotin-conjugated anti-rat CD25 (clone OX-39), and Streptavidin-conjugated APC-Cy7 mAbs were obtained from BD Biosciences (San Diego, CA). Purified GRP78 mAbs (clone 40) and isotype control Abs were also obtained from BD Biosciences. A Zenon Mouse IgG2a or IgG1 Labeling Kit, Alexa Fluor 647 (Invitrogen, Carlsbad, CA), was used to label purified GRP78 or CHOP mAbs per the manufacturer’s directions. Anti-rat CD4-Pacific Blue mAb and its corresponding isotype control Ab were obtained from Serotec (Raleigh, NC). Anti-actin (clone C4) antibody was obtained from Chemicon International (Temecula, CA). Anti-ERp72 and anti-GRP94 were obtained from Calbiochem (San Diego, CA) and Stressgen (San Diego, CA), respectively. Cleaved

capase-3 antibody was obtained from Cell Signaling Technology (Danvers, MA). Polyclonal antiserum to Gimap5 was generated by us as described [11]. Anti-CHOP antibody and anti-mouse IgG horseradish peroxidase (HRP) conjugate were obtained from Santa Cruz Biotechnology (Santa Cruz, CA).

Flow cytometry

Single cell suspensions from cervical and mesenteric lymph node cells were washed and suspended in PBS containing 1% fetal clone serum (HyClone, Logan, UT) and 0.1% sodium azide (Sigma-Aldrich). Samples were washed, incubated for 20 min with fluorescent mAbs to various cell-surface markers as described in Results. To detect intracellular GRP78 or CHOP protein, cells were permeabilized using Cytotfix/Cytoperm (BD Biosciences) according to the manufacturer's directions. Cells were washed and incubated with Alexa Fluor 647-conjugated GRP78 or CHOP mAb for 20 min. Labeled cells were washed, fixed with 1% paraformaldehyde (Polysciences, Warrington, PA) in PBS and analyzed using an LSR II instrument (BD Biosciences, San Jose, CA) and FlowJo Software (PC version 7.2.2; Tree Star, Ashland, OR). Lymphoid cells were gated according to their light-scattering properties.

T cell transfection

T cells from BBDP rats were purified as described [52]. T cells were plated at 6×10^6 cells per well in 3 mL of RPMI medium (Sigma-Aldrich) containing 10% FBS (Hyclone), 1% Pen/Strep/Glut (Gibco, Carlsbad, CA), and 0.1% β -mercaptoethanol (Gibco) at 37°C in the presence of 10 ng/mL PMA (Calbiochem) and 100 ng/mL ionomycin (Calbiochem). After 12 h, cells were resuspended in 100 μ L of Nucleofector solution using human T cell Nucleofector kit (VPA-1002, Amaxa Biosystems, Gaithersburg, MD) with 100 nM rat CHOP siRNA (catalogue no. L-088282-01, Dharmacon, Lafayette, CO) or 100 nM control siRNA (catalogue no. D-001210-02, Dharmacon) and nucleofected using program U-014 in the Amaxa Nucleofector apparatus. Following nucleofection, T cells were plated in 3 mL complete media and harvested at 30 h for protein preparation and flow cytometry analysis.

Apoptosis Detection

To measure cellular apoptotic signaling, transfected T cells were harvested, washed twice with PBS, and reacted with PE-conjugated anti-annexin V and 7-amino-actinomycin D (7AAD)

according to the manufacturer's directions (BD Pharmingen). At least 20,000 cells per sample were analyzed using a FACSCalibur (BD Biosciences) and FlowJo software.

Statistics

Statistical analyses were performed with GraphPad Prism software (Graphpad Software, San Diego, CA). Parametric data are presented as the mean \pm the standard error of measurement (SEM) and were analyzed using two-tailed unpaired *t*-tests. Values of $p < 0.05$ were considered statistically significant.

Supporting Information

Figure S1 T cell populations from congenic WF.*ART2a.Gimap5*^{-/-} rats that do not develop spontaneous diabetes exhibit enhanced ER stress response signaling. Representative analyses indicating the gating of TCR⁺ (A) and CD4⁺ or CD8⁺ (B) lymphocytes within congenic WF.*ART2a.Gimap5*^{-/-} (Type 31) and control congenic WF.*ART2a.Gimap*^{+/+} (Type 20) rats. Representative flow dot plots depicting CD90 expression (horizontal axis) and forward scattering (vertical axis) of lymphocytes within gated CD4⁺ and CD8⁺ T cell populations from Type 31 (C) or Type 20 (D) rats. Numbers represent the percentage of cells (\pm SEM of triplicate samples) in each gate shown. Intracellular GRP78 (E) and CHOP (F) expression in RTEs (CD90⁺) and mature T cells (CD90⁻) from Type 31 (black line) and Type 20 rats (shaded region). Depicted in each histogram is the isotype control (dotted line). The MFI of GRP78 or CHOP protein expression is displayed in bar graphs with error bars representing the SEM of duplicate samples. Data shown are representative of two independent experiments (* $P < .05$; ** $P < .01$; *** $P < .001$). Found at: doi:10.1371/journal.pone.0005468.s001 (0.94 MB TIF)

Acknowledgments

We thank Linda Leehy, Linda Paquin, Mary Lively, and Elaine Norowski for their technical assistance.

Author Contributions

Conceived and designed the experiments: SCP BOM EAL CY AJ Pd DLG RB. Performed the experiments: SCP BOM EAL CY AJ. Analyzed the data: SCP KLL Pd MAB JPM DLG AAR RB. Contributed reagents/materials/analysis tools: JPM. Wrote the paper: SCP KLL MAB JPM DLG AAR RB.

References

- Nitta T, Nasreen M, Seike T, Goji A, Ohgashi I, et al. (2006) IAN family critically regulates survival and development of T lymphocytes. *PLoS Biol* 4: 593–605.
- Nitta T, Takahama Y (2007) The lymphocyte guard-IANs: regulation of lymphocyte survival by IAN/GIMAP family proteins. *Trends Immunol* 28: 58–65.
- Hellquist A, Zucchelli M, Kivinen K, Saarialho-Kere U, Koskenmies S, et al. (2007) The human GIMAP5 gene has a common polyadenylation polymorphism increasing risk to systemic lupus erythematosus. *J Med Genet* 44: 314–321.
- Shin JH, Janer M, McNeney B, Blay S, Deutsch K, et al. (2007) IA-2 autoantibodies in incident type I diabetes patients are associated with a polyadenylation signal polymorphism in GIMAP5. *Genes Immun* 8: 503–512.
- Schulteis RD, Chu H, Dai X, Chen Y, Edwards B, et al. (2008) Impaired survival of peripheral T cells, disrupted NK/NKT cell development, and liver failure in mice lacking gimap5. *Blood* 112: 4905–4915.
- Krucken J, Schroetel RM, Muller IU, Saidani N, Marinovski P, et al. (2004) Comparative analysis of the human gimap gene cluster encoding a novel GTPase family. *Gene* 341: 291–304.
- Ramanathan S, Norwich K, Poussier P (1998) Antigen activation rescues recent thymic emigrants from programmed cell death in the BB rat. *J Immunol* 160: 5757–5764.
- Schnell S, Demolliere C, van den BP, Jacobs H (2006) Gimap4 accelerates T-cell death. *Blood* 108: 591–599.
- Hornum L, Romer J, Markholst H (2002) The diabetes-prone BB rat carries a frameshift mutation in Ian4, a positional candidate of Iddm1. *Diabetes* 51: 1972–1979.
- MacMurray AJ, Moralejo DH, Kwitek AE, Rutledge EA, Van Yserloo B, et al. (2002) Lymphopenia in the BB rat model of type 1 diabetes is due to a mutation in a novel immune-associated nucleotide (Ian)-related gene. *Genome Res* 12: 1029–1039.
- Pandarpurkar M, Wilson-Fritch L, Corvera S, Markholst H, Hornum L, et al. (2003) Ian4 is required for mitochondrial integrity and T cell survival. *Proc Natl Acad Sci U S A* 100: 10382–10387.
- Dalberg U, Markholst H, Hornum L (2007) Both Gimap5 and the diabetogenic BBDP allele of Gimap5 induce apoptosis in T cells. *Int Immunol* 19: 447–453.
- Sandal T, Aumo L, Hedin L, Gjertsen BT, Doskeland SO (2003) Irod/Ian5: an inhibitor of gamma-radiation- and okadaic acid-induced apoptosis. *Mol Biol Cell* 14: 3292–3304.
- Daheron L, Zenz T, Siracusa LD, Brenner C, Calabretta B (2001) Molecular cloning of Ian4: a BCR/ABL-induced gene that encodes an outer membrane mitochondrial protein with GTP-binding activity. *Nucleic Acids Res* 29: 1308–1316.

15. Kaneko M, Nomura Y (2003) ER signaling in unfolded protein response. *Life Sci* 74: 199–205.
16. Kaufman RJ (2002) Orchestrating the unfolded protein response in health and disease. *J Clin Invest* 110: 1389–1398.
17. Borgese N, Francolini M, Snapp E (2006) Endoplasmic reticulum architecture: structures in flux. *Curr Opin Cell Biol* 18: 358–364.
18. Boyce M, Yuan J (2006) Cellular response to endoplasmic reticulum stress: a matter of life or death. *Cell Death Differ* 13: 363–373.
19. Ron D (2002) Translational control in the endoplasmic reticulum stress response. *J Clin Invest* 110: 1383–1388.
20. Rao RV, Peel A, Logvinova A, Del Rio G, Hermel E, et al. (2002) Coupling endoplasmic reticulum stress to the cell death program: role of the ER chaperone GRP78. *FEBS Lett* 514: 122–128.
21. Harding HP, Calfon M, Urano F, Novoa I, Ron D (2002) Transcriptional and translational control in the mammalian unfolded protein response. *Annu Rev Cell Dev Biol* 18: 575–599.
22. Yoshida H, Oku M, Suzuki M, Mori K (2006) pXBP1(U) encoded in XBP1 pre-mRNA negatively regulates unfolded protein response activator pXBP1(S) in mammalian ER stress response. *J Cell Biol* 172: 565–575.
23. Zhang K, Kaufman RJ (2004) Signaling the unfolded protein response from the endoplasmic reticulum. *J Biol Chem* 279: 25935–25938.
24. Lee AS (2005) The ER chaperone and signaling regulator GRP78/BiP as a monitor of endoplasmic reticulum stress. *Methods* 35: 373–381.
25. Sato N, Urano F, Yoon LJ, Kim SH, Li M, et al. (2000) Upregulation of BiP and CHOP by the unfolded-protein response is independent of presenilin expression. *Nat Cell Biol* 2: 863–870.
26. Puthalakath H, O'Reilly LA, Gunn P, Lee L, Kelly PN, et al. (2007) ER stress triggers apoptosis by activating BH3-only protein Bim. *Cell* 129: 1337–1349.
27. Wang XZ, Lawson B, Brewer JW, Zinszner H, Sanjay A, et al. (1996) Signals from the stressed endoplasmic reticulum induce C/EBP-homologous protein (CHOP/GADD153). *Mol Cell Biol* 16: 4273–4280.
28. Medigeshi GR, Lancaster AM, Hirsch AJ, Briese T, Lipkin WI, et al. (2007) West Nile virus infection activates the unfolded protein response, leading to CHOP induction and apoptosis. *J Virol* 81: 10849–10860.
29. Michalkiewicz M, Michalkiewicz T, Eittinger RA, Rutledge EA, Fuller JM, et al. (2004) Transgenic rescue demonstrates involvement of the *Ian5* gene in T cell development in the rat. *Physiol Genomics* 19: 228–232.
30. Jung CG, Kamiyama T, Agui T (1999) Elevated apoptosis of peripheral T lymphocytes in diabetic BB rats. *Immunology* 98: 590–594.
31. Lang JA, Kominski D, Bellgrau D, Scheinman RI (2004) Partial activation precedes apoptotic death in T cells harboring an *IAn* gene mutation. *Eur J Immunol* 34: 2396–2406.
32. Pino SC, O'Sullivan-Murphy B, Lidstone EA, Thornley TB, Jurczyk A, et al. (2008) Protein kinase C signaling during T cell activation induces the endoplasmic reticulum stress response. *Cell Stress Chaperones* 13: 421–434.
33. Scheu S, Stetson DB, Reinhardt RL, Leber JH, Mohrs M, et al. (2006) Activation of the integrated stress response during T helper cell differentiation. *Nat Immunol* 7: 644–651.
34. Mordes JP, Bortell R, Blankenhorn EP, Rossini AA, Greiner DL (2004) Rat models of type 1 diabetes: genetics, environment, and autoimmunity. *ILAR J* 45: 278–291.
35. Gallegos AM, Bevan MJ (2006) Central tolerance: good but imperfect. *Immunol Rev* 209: 290–296.
36. Kisielow P, von Boehmer H (1995) Development and selection of T cells: facts and puzzles. *Adv Immunol* 58: 87–209.
37. Willerford DM, Swat W, Alt FW (1996) Developmental regulation of V(D)J recombination and lymphocyte differentiation. *Curr Opin Genet Dev* 6: 603–609.
38. Iwakoshi NN, Goldschneider I, Tausche F, Mordes JP, Rossini AA, et al. (1998) High frequency apoptosis of recent thymic emigrants in the liver of lymphopenic diabetes-prone BioBreeding rats. *J Immunol* 160: 5838–5850.
39. Gass JN, Gifford NM, Brewer JW (2002) Activation of an unfolded protein response during differentiation of antibody-secreting B cells. *J Biol Chem* 277: 49047–49054.
40. Nakanishi K, Sudo T, Morishima N (2005) Endoplasmic reticulum stress signaling transmitted by ATF6 mediates apoptosis during muscle development. *J Cell Biol* 169: 555–560.
41. Oyadomari S, Araki E, Mori M (2002) Endoplasmic reticulum stress-mediated apoptosis in pancreatic beta-cells. *Apoptosis* 7: 335–345.
42. Keita M, Leblanc C, Andrews D, Ramanathan S (2007) GIMAP5 regulates mitochondrial integrity from a distinct subcellular compartment. *Biochem Biophys Res Commun* 361: 481–486.
43. Kim R, Emi M, Tanabe K, Murakami S (2006) Role of the unfolded protein response in cell death. *Apoptosis* 11: 5–13.
44. Faitova J, Krekac D, Hrstka R, Vojtesek B (2006) Endoplasmic reticulum stress and apoptosis. *Cell Mol Biol Lett* 11: 488–505.
45. Kim I, Xu W, Reed JC (2008) Cell death and endoplasmic reticulum stress: disease relevance and therapeutic opportunities. *Nat Rev Drug Discov* 7: 1013–1030.
46. Araki E, Oyadomari S, Mori M (2003) Impact of endoplasmic reticulum stress pathway on pancreatic beta-cells and diabetes mellitus. *Exp Biol Med* 228: 1213–1217.
47. Rossini AA (2004) Autoimmune diabetes and the circle of tolerance. *Diabetes* 53: 267–275.
48. Kaaba SA, Al-Harbi SA (1995) Abnormal lymphocyte subsets in Kuwaiti patients with type-1 insulin-dependent diabetes mellitus and their first-degree relatives. *Immunol Lett* 47: 209–213.
49. Mordes JP, Cort L, Norowski E, Leif J, Fuller JM, et al. (2009) Analysis of the rat *Iddm14* diabetes susceptibility locus in multiple rat strains: identification of a susceptibility haplotype in the *Tcrb-V* locus. *Mamm Genome*, In press.
50. Mordes JP, Leif J, Novak S, DeScipio C, Greiner DL, et al. (2002) The *iddm4* locus segregates with diabetes susceptibility in congenic WF.*iddm4* rats. *Diabetes* 51: 3254–3262.
51. Blankenhorn EP, Rodemich L, Martin-Fernandez C, Leif J, Greiner DL, et al. (2005) The rat diabetes susceptibility locus *Iddm4* and at least one additional gene are required for autoimmune diabetes induced by viral infection. *Diabetes* 54: 1233–1237.
52. Rigby MR, Bortell R, Greiner DL, Czech MP, Klarlund JK, et al. (1996) The rat T-cell surface protein RT6 is associated with src family tyrosine kinases and generates an activation signal. *Diabetes* 45: 1419–1426.