

## Chromate Reduction by Cell-Free Extract of *Bacillus firmus* KUCr1

GOPI BALLAV SAU, SWAGATA CHATTERJEE and SAMIR KUMAR MUKHERJEE\*

Department of Microbiology, University of Kalyani, Kalyani 741235, India

Received 9 February 2010, revised 10 June 2010, accepted 20 June 2010

### Abstract

Microbial enzymatic reduction of a toxic form of chromium [Cr(VI)] has been considered as an effective method for bioremediation of this metal. This study reports on the *in vitro* reduction of Cr(VI) using cell-free extracts from a Cr(VI) reducing *Bacillus firmus* KUCr1 strain. Chromium reductase was found to be constitutive and its activity was observed both in soluble cell fractions ( $S_{12}$  and  $S_{150}$ ) and membrane cell fraction ( $P_{150}$ ). The reductase activity of  $S_{12}$  fraction was found to be optimal at 40  $\mu\text{M}$  Cr(VI) with enzyme concentration equivalent to 0.493 mg protein/ml. Enzyme activity was dependent on NADH or NADPH as electron donor; optimal temperature and pH for better enzyme activity were 70°C and 5.6, respectively. The  $K_m$  value of the reductase was 58.33  $\mu\text{M}$  chromate having a  $V_{max}$  of 11.42  $\mu\text{M}/\text{min}/\text{mg}$  protein. The metabolic inhibitor like sodium azide inhibited reductase activity of membrane fraction of the cell-free extract. Metal ions like  $\text{Cu}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$  and  $\text{As}^{3+}$  stimulated the enzyme but others, such as  $\text{Ag}^+$ ,  $\text{Hg}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$ , inhibited Cr(VI) reductase activity.

**Key words:** *Bacillus firmus*, cell-free extract, chromium reductase

### Introduction

Application of Cr-resistant bacteria to detoxify Cr(VI) has been considered as an effective bioremediation method (Ohtake and Silver, 1994; Shakoori *et al.*, 1999; Ganguli and Tripathi, 2002; Cheung and Gu, 2007; Sau *et al.*, 2008). Bioremediation of Cr(VI) can occur by enzymatic reduction of Cr(VI) to Cr(III) via Cr(V) and Cr(IV) intermediates (Camargo *et al.*, 2003; Xu *et al.*, 2004; Xu *et al.*, 2005; Pal *et al.*, 2005; Cheung *et al.*, 2006) or through immobilization (Losi *et al.*, 1994). In the course of aerobic reduction, the cytosolic fractions of most chromium resistant bacteria exhibit Cr(VI) reductase activity (Campos *et al.*, 1995; Cervantes *et al.*, 2001), while under anaerobic conditions,  $\text{CrO}_4^{2-}$  is used as a terminal electron acceptor and is reduced in the cell membrane (McLean and Beveridge, 2001).

Requirement of electron supply is obvious during the reaction process, considering the process a redox reaction. The enzyme responsible for Cr(VI) reduction has not yet been well characterized, though there are reports on reduction kinetics of Cr(VI) reduction by cell-free extracts (CFE) with varied results (Ishibashi *et al.*, 1990; Suzuki *et al.*, 1992; McLean and Beveridge, 2001; Park *et al.*, 2000; Camargo *et al.*,

2003; Pal *et al.*, 2005). Studies using partially purified Cr(VI) reductase from *Pseudomonas ambigua* G-1 (Suzuki *et al.*, 1992) and *P. putida* MK1 (Park *et al.*, 2000) have indicated Cr(VI) reduction using NADH or NADPH as a source of electrons. In *Bacillus* sp. ES29, Cr(VI) reductase was found in the soluble fraction of CFE, which utilizes NADH as electron donor (Camargo *et al.*, 2003).

Cr(VI) reducing *Bacillus firmus* KUCr1 has been reported and its potential to reduce Cr(VI) using whole cells under culture condition was documented earlier (Sau *et al.*, 2008). This study reports on the reduction of Cr(VI) by CFE of this strain, effects of different electron donors, metal ions, inhibitors, pH and temperature on its Cr(VI) reducing activity, and cellular localization of Cr(VI) reductase.

### Experimental

#### Materials and Methods

**Microorganism and growth condition.** A Cr(VI) resistant *Bacillus firmus* KUCr1 (NCBI GenBank 16S rDNA sequence Accession No EU784699) was earlier reported (Sau *et al.*, 2008) and used in this study.

\* Corresponding author: S.K. Mukherjee, Department of Microbiology, University of Kalyani, Kalyani 741235, India; phone: (+91) 3325827315; fax: (+91) 3325828282; e-mail: [dr.samirmukherjee@klyuniv.ac.in](mailto:dr.samirmukherjee@klyuniv.ac.in)

Cells were grown aerobically in PYG medium (peptone 10 g/l; yeast extract 5 g/l; glucose 3 g/l; pH 7.2) supplemented with or without (control) 0.1 mM Cr(VI) as  $K_2CrO_4$  under continuous shaking on a rotary shaker at 35°C for 24 h.

**Preparation of cell-free extract (CFE) and cellular fractions.** Cell-free extract (CFE) and cellular fractions were prepared following McLean and Beveridge (2001) with little modification. Cells grown in PYG medium were harvested at the exponential phase, washed three times in 0.2 M phosphate buffer (pH 7.2) by centrifugation ( $4000 \times g$  at 4°C), resuspended in the same buffer to have a 10 ml suspension and kept in an ice bath. Cells were disrupted with an ultrasonicator (Hielscher Ultrasound Technology, UP50H). Power was applied five times in 1 min pulses with 100% amplitude. After sonication the suspension was centrifuged at  $12\,000 \times g$  for 10 min at 4°C to obtain a soluble fraction ( $S_{12}$ ). Five ml of  $S_{12}$  fraction was centrifuged at  $150\,000 \times g$  for 90 min at 4°C to obtain  $S_{150}$  fraction. The pellet fraction was washed twice with 0.2 M phosphate buffer (pH 7.2) to remove soluble chromium reductase, if any and was resuspended in same buffer (5 ml) to obtain membrane fraction ( $P_{150}$ ). Equivalent volume of  $S_{12}$ ,  $S_{150}$  and  $P_{150}$  fraction were assayed for studying the cellular localization of Cr(VI) reductase enzyme.

**Cr(VI) reductase assay.** Cr(VI) reductase activity of CFE was assayed following the procedure of Park *et al.* (2000). The reaction mixture (1 ml) for the enzyme assay contained 0.2 mM  $K_2CrO_4$ , 0.2 mM NADH and 400  $\mu$ l of CFE in 0.2 M phosphate buffer (pH 7.2). The reaction was stopped by adding 0.5 ml of 20% trichloroacetic acid (Horitsu *et al.*, 1987). Reduction of Cr(VI) was measured by estimating the decrease in Cr(VI) in the reaction mixture after 30 min of incubation at 35°C and quantified colorimetrically using 1,5-diphenylcarbazide as the complexing reagent (Urone, 1955). To eliminate the abiotic reduction of Cr(VI), if any, a control set was prepared through out the course of the study without putting any biotic component including bacterial cell. Whenever required, abiotic reduction of Cr(VI) has been subtracted from the total reduction to get the exact influence on chromium reduction by the biotic factors. Amount of protein in the CFE was estimated by the folin-phenol method (Lowry *et al.*, 1951) using bovine serum albumin as the standard. One unit of Cr(VI) reductase activity was defined as the amount of enzyme which decreased 1.0  $\mu$ M Cr(VI) per min at 35°C. The effect of pH and temperature on Cr(VI) reductase were measured at different pH values (4 to 10.6) of the reaction mixture at 35°C and at different reaction temperature (20 to 100°C) at pH 7.2 respectively.

## Results and Discussions

The reductase activities of the soluble fraction ( $S_{12}$ ) from both the induced and uninduced cells of *B. firmus* KUCr1 were almost similar with regard to time course (Fig. 1). This study revealed that the chromate reductase in this strain is constitutive, which supports earlier reports on the enzymatic reduction of Cr(VI) under aerobic conditions (Bopp and Ehrlich, 1988; Campos *et al.*, 1995; Wang and Xiao, 1995; McLean and Beveridge, 2001; Pal *et al.*, 2005), though an inducible reductase in the soluble fraction of CFE of *Ochrobactrum* sp. was reported (Thacker and Datta, 2005). As the reductase activity was found to be constitutive, thus for further experiments CFE ( $S_{12}$ ) was prepared from cells grown in Cr(VI)-free medium for characterization of chromate reductase activity.

The effect of initial concentration of Cr(VI) on reductase activity of  $S_{12}$  fraction was determined at a concentration range of 0 to 80  $\mu$ M of Cr(VI). Specific activity increased with increasing initial concentration of chromate up to 40  $\mu$ M, after that it slowed down but reduction continued (Fig. 2a) with enzyme equivalent to 0.493 mg protein/ml. The saturation kinetics of Cr(VI) reduction of  $S_{12}$  fraction fit with the linearized Lineweaver-Burk plot, and the apparent Michaelis-Menten constant ( $K_m$ ) was found at 58.33  $\mu$ M chromate and  $V_{max}$  was 11.42  $\mu$ M per min/mg protein (Fig. 2b). The  $K_m$  and  $V_{max}$  values differed from the enzyme activity of the CFE of *Bacillus* sp. ES 29 (Camargo *et al.*, 2003), *B. subtilis* (Garbisu *et al.*, 1998), *B. Sphaericus* AND 303 (Pal *et al.*, 2005), *P. putida* (Ishibashi *et al.*, 1990; McLean and Beveridge




Fig. 1. Cr(VI) reduction by the extracts ( $S_{12}$ ) from both induced [0.1 mM Cr(VI) in the medium] and non-induced cells of *B. firmus* KUCr1. The reaction mixture contained 0.2 mM Cr(VI) and 0.2 mM NADH as electron donor in 0.2 M phosphate buffer (pH 7.2) and the reaction temperature was 35°C.




Fig. 2. Kinetics of Cr(VI) reduction of cell-free extract ( $S_{12}$ ) of *B. firmus* KUCr1 at different Cr(VI) concentrations. Reaction time was 30 min at 35°C (a). Linearized Lineweaver-Burk plot for Cr(VI) reduction of cell-free extract ( $S_{12}$ ) (b).

2001; Park *et al.*, 2000) and in *P. ambigua* G-1 (Suzuki *et al.*, 1992). Lower  $K_m$  value of Cr(VI) reductase suggests higher affinity for the substrate, at least with what was found with the cell-free extract in this strain.

The effect of pH on reductase activity was assessed at a pH range of 4.0 to 10.6 using 0.2 M of citrate buffer (pH 4.0 to 5.6), 0.2 M of phosphate buffer (pH 6.0 to 7.6), 0.2 M of tris-HCl buffer (pH 8.4 to 8.8) and 0.2 M of glycine-NaOH buffer (pH 9.2 to 10.6) separately. The reductase activity achieved a maximum at pH 5.6 (Fig. 3). Similarly, the optimum temperature for highest Cr(VI) reduction was found at 70°C (Fig. 4). These results varied from other earlier reports with *B. sphaericus* AND 303, where they were 30°C and 6.0 respectively (Pal *et al.*, 2005) and with *P. putida* MK1 (Park *et al.*, 2000) or *P. ambigua* G-1 (Suzuki *et al.*, 1992), the optimal temperature and pH being 80°C, 50°C, and pH 5.0, 8.6 respectively.

The effect of electron donors, inhibitors and metal ions on chromate reduction by CFE ( $S_{12}$ ) of *B. firmus* KUCr1 was determined. 0.2 mM each of NADH, NADPH, glutathione, D-glucose, and D-fructose were




Fig. 3. Effect of pH on chromate reductase activity ( $S_{12}$ ) of *B. firmus* KUCr1 at 35°C for 30 min of incubation.




Fig. 4. Effect of temperature on chromate reductase activity ( $S_{12}$ ) of *B. firmus* KUCr1. The assay was conducted in 0.2 M phosphate buffer (pH 7.2) for 30 min.

used as electron donors. Among the electron donors used, only NADH showed a significant effect on chromate reductase activity (72% activity over control). NADPH gave 32% less activity than NADH (Table I).

Table I  
Effect of electron donor on Cr(VI) reductase activity in cell-free extracts ( $S_{12}$ ) of *B. firmus* KUCr1

Electron donor (0.2 mM)	Specific activity <sup>a</sup> (U/mg protein)	Relative specific activity <sup>b</sup> (%)
None (control)	2.40 (±0.01)	27.74
NADH	8.65 (±0.06)	100.00
NADPH	5.87 (±0.08)	67.86
Glutathione	2.53 (±0.02)	29.24
D-glucose	2.89 (±0.01)	33.41
D-fructose	2.70 (±0.07)	31.21

<sup>a</sup> Data are the mean of three replications plus standard error. The reaction mixture contained 0.2 mM Cr(VI) in 0.2 M phosphate buffer (pH 7.2) and was incubated for 30 min at 35°C.

<sup>b</sup>  $\{(\text{Specific activity}) / (\text{specific activity})_{\text{NADH}}\} \times 100$




Fig. 5. Effect of azide ( $\text{NaN}_3$ , 0.2 mM) and some selected metals (0.2 mM) on Cr(VI) reductase activity in the cell-free extracts ( $\text{S}_{12}$ ) of *B. firmus* KUCr1. Data are the mean of three replications with error bars.

In the presence of glutathione, D-glucose, and D-fructose, the non-enzymatic reductants of Cr(VI), the activity of the reductase was almost equal to that of the control.

The cell-free enzyme of *B. firmus* KUCr1 required NADH or NADPH as an electron donor for better enzymatic Cr(VI) reduction. The reductase became sharply more active in the presence of NADH than NADPH, suggesting the requirement of a cofactor for catalytic activity. NADH dependent Cr(VI) reduction was also advocated by several researchers earlier in *Bacillus* (Garbisu *et al.*, 1998; Camargo *et al.*, 2003) and in *Pseudomonas* (Suzuki *et al.*, 1992; Park *et al.*, 2000). Our study also supports the earlier observations on the role of nonenzymatic reductants like glutathione, D-glucose and D-fructose on Cr(VI) reduction by CFE (Branca *et al.*, 1990; Shi and Dalal, 1990; Suzuki *et al.*, 1992).

Cr(VI) reduction by CFE was significantly inhibited in the presence of 0.2 mM sodium azide (Fig. 5). The metal cations,  $\text{Ag}^+$ ,  $\text{Zn}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Mn}^{2+}$ , and  $\text{Hg}^{2+}$  inhibited reductase activity by more than 50% over the control in the reaction mixture (Fig. 5). However,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{As}^{3+}$  and  $\text{Cu}^{2+}$  stimulated the activity of CFE. The order of stimulation by these metal cations in reductase activity was found to be  $\text{As}^{3+} > \text{Ni}^{2+} > \text{Co}^{2+} > \text{Cu}^{2+}$ .

In this study, Cr(VI) reductase activity was found to be inhibited by a respiratory inhibitor, sodium azide (0.2 mM). Though chromate reductase in CFE was reported to be unaffected by azide in *Escherichia coli* ATCC 33456 (Shen and Wang, 1993), *Bacillus* sp. ES29 (Camargo *et al.*, 2003) and in *B. megaterium* TKW3 (Cheung *et al.*, 2006) by 1.0 mM  $\text{NaN}_3$ . However, inhibition of Cr(VI) reduction due to  $\text{NaN}_3$  in live cells of a *Bacillus subtilis* strain was reported (Garbisu *et al.*, 1998). Inhibition of a cytoplasmic membrane associated Cr(VI) reductase by azide was

also reported in *Shewanella putrefaciens* MR-1 (Myers *et al.*, 2000). The interference of azide in Cr(VI) reduction in this study and views from earlier reports suggest the possible coexistence of Cr(VI) reductase in the cytosol and membrane as well. However, further investigations are required to elucidate this particular feature.

The cations  $\text{Hg}^{2+}$  and  $\text{Ag}^+$  (0.2 mM) inhibited reductase activity. The noncompetitive inhibitory effect of  $\text{Hg}^{2+}$  and  $\text{Ag}^+$  on Cr(VI) reduction in *P. putida* was reported earlier (Ishibashi *et al.*, 1990). The inhibitory effect of  $\text{Hg}^{2+}$  is expected, because of its affinity for ligands containing thiol (-SH) group of a variety of enzymatic proteins. Enzymatic reduction of Cr(VI) by CFE of *B. firmus* KUCr1 was stimulated by  $\text{Cu}^{2+}$ ,  $\text{As}^{3+}$ ,  $\text{Ni}^{2+}$  and  $\text{Co}^{2+}$  at 0.2 mM concentration separately, though Pal *et al.* (2005) reported the inhibition of Cr(VI) reductase activity of *B. sphaericus* AND 303 by  $\text{Ni}^{2+}$  and  $\text{Co}^{2+}$  at 100  $\mu\text{M}$  concentration. The reductase activity was found to be unaffected by  $\text{As}^{3+}$  in *P. putida* MK1 (Park *et al.*, 2000). Stimulation of enzyme activity by  $\text{Cu}^{2+}$  might be due to its nature as a prosthetic group of many reductase enzymes and also indirectly involved in the protection of chromate reductase from  $\text{O}_2$ , for oxygen-sensitive enzyme (Ettinger 1984; Camargo *et al.*, 2003). Abe *et al.* (2001) reported that  $\text{Cu}^{2+}$  acting as electron-transport protection or acting as a single electron redox center and as a shuttle for electron between protein subunits.  $\text{Zn}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$  and  $\text{Mn}^{2+}$  inhibited the reductase activity at 0.2 mM. Inhibitory effects by  $\text{Cd}^{2+}$  and  $\text{Zn}^{2+}$  support the earlier views (Park *et al.*, 2000; Pal *et al.*, 2005; Desai *et al.*, 2008). On the contrary, Camargo *et al.* (2003) showed slight stimulatory effect of  $\text{Mn}^{2+}$  on Cr(VI) reductase in *Bacillus* sp. ES 29. These variations seem to be due to the different functional nature of the Cr(VI) reductase in *B. firmus* KUCr1, which warrants further investigation.

Cr(VI) reductase activities by different cellular fractions are presented in Table II. In the presence of electron donor (0.2 mM NADH)  $\text{S}_{150}$  fraction showed higher activity than fractions  $\text{S}_{12}$  and  $\text{P}_{150}$ .

Table II  
Localization of Cr(VI) reductase in cell fractions of *B. firmus* KUCr1 and its catalytic activity

Cellular fraction <sup>a</sup>	% Cr(VI) reduction <sup>b</sup>	Specific activity <sup>c</sup> (U/mg protein)
$\text{S}_{12}$	14.21	9.12 ( $\pm 0.03$ )
$\text{S}_{150}$	15.66	14.68 ( $\pm 0.04$ )
$\text{P}_{150}$	6.98	7.34 ( $\pm 0.04$ )

<sup>a</sup> S stands for soluble cellular fractions and P stands for membrane fractions (see Materials and Methods).

<sup>b</sup> Reaction mixture contained 0.2 mM Cr(VI) in phosphate buffer (pH 7.2) and was incubated for 30 min at 35°C.

<sup>c</sup> Data are the mean of three repetitions plus standard error.

The percentage of Cr(VI) reduction by fraction P<sub>150</sub> is significantly less compared to other fractions, but the specific activity signifies its catalytic function, suggesting the occurrence of some membrane-bound protein responsible for Cr(VI) reduction. In *Bacillus* QC1-2 (Campos *et al.*, 1995), *B. sphaericus* AND 303 (Pal *et al.*, 2005) and *P. putida* (Ishibashi *et al.*, 1990; Park *et al.*, 2000) the chromium reductase activities were reported to be associated with the cytosolic and soluble fractions. In *Enterobacter cloacae* (Wang *et al.*, 1990), *S. putrefaciens* MR-1 (Myers *et al.*, 2000), *B. megaterium* TKW3 (Cheung *et al.*, 2006) and *P. fluorescences* (Bopp and Ehrlich, 1988) Cr(VI) reductase appears to be membrane associated. Lovley and Phillips (1994) reported that both the soluble and membrane fractions reduced chromate in *Desulfovibrio vulgaris* but soluble protein fraction reduced Cr(VI) faster than the membrane fraction did. In *Bacillus cereus* S-6 (Iftikhar *et al.*, 2007) Cr(VI) reduction occur both in cytosolic and membrane fraction of this strain but percentage of reduction in cytosolic fraction is higher than membrane fraction. It seems that KUCr1 harbors Cr(VI) reductase constitutively both in membrane and cytosol. Thus further investigations to visualize the exact pathway of Cr(VI) reduction in this particular strain are required.

Enzyme inhibition by respiratory inhibitors like NaN<sub>3</sub> and Cr(VI) reductase activity by the P<sub>150</sub> fraction suggest the co-existence of a membrane-bound, respiratory-chain-linked reductase activity in this strain; study with purified proteins of both the cytosolic and membrane fractions would provide more knowledge about the exact pathway of Cr(VI) in this particular strain. Moreover, the feature of higher temperature optima of Cr(VI) reductase of the cytosolic soluble fraction in KUCr1 suggests the possibility to clone the gene into a thermophilic bacteria, so that it could offer a decisive advantage in chromium bioremediation under conditionally elevated temperature or in a bioreactor system using immobilized enzyme.

#### Acknowledgement

This work was supported by the grant received from the University of Kalyani, India.

#### Literature

- Abe F., T. Miura, T. Nagahama, A. Inoue, R. Usami and K. Horikoshi.** 2001. Isolation of a highly copper-tolerant yeast, *Cryptococcus* sp., from the Japan trench and the induction of superoxide dismutase activity by Cu<sup>2+</sup>. *Biotechnol. Lett.* 23: 2027–2034.
- Bopp L.H. and H.L. Ehrlich.** 1988. Chromate resistance and reduction in *Pseudomonas fluorescens* strain LB300. *Arch. Microbiol.* 150: 426–431.
- Branca M., A. Dessi, H. Koziowski, G. Micera and J. Swiatek.** 1990. Reduction of chromate ions by glutathione tripeptide in the presence of sugar ligands. *J. Inorg. Biochem.* 39: 217–226.
- Camargo F.A.O., B.C. Okeke, F.M. Bento and W.T. Frankenberger.** 2003. *In vitro* reduction of hexavalent chromium by a cell-free extract of *Bacillus* sp. ES29 stimulated by Cu. *Appl. Microbiol. Biotechnol.* 62: 569–573.
- Campos J., M. Martinez-Pacheco and C. Cervantes.** 1995. Hexavalent-chromium reduction by a chromate-resistant *Bacillus* sp. strain. *Antonie van Leeuwenhoek* 68: 203–208.
- Cervantes C., J.C. Garcia, S. Devars, F.G. Corona, H.L. Tavera, J.C. Guzman and R.M. Sanchez.** 2001. Interactions of chromium with microorganisms and plants. *FEMS Microbiol. Rev.* 25: 335–347.
- Cheung K.H. and J.D. Gu.** 2007. Mechanism of hexavalent chromium detoxification by microorganisms and bioremediation application potential: A review. *Int. Biodet. Biodegrad.* 59: 8–15.
- Cheung K.H., H.Y. Lai and J.D. Gu.** 2006. Membrane-associated hexavalent chromium reductase of *Bacillus megaterium* TKW3 with induced expression. *J. Microbiol. Biotechnol.* 16: 855–862.
- Desai C., J. Kunal and M. Datta.** 2008. Hexavalent chromate reductase activity in cytosolic fractions of *Pseudomonas* sp. G1DM21 isolated from Cr(VI) contaminated industrial landfill. *Process Biochem.* 43: 713–721.
- Ettinger M.J.** 1984. Copper metabolism and diseases of copper metabolism. pp. 175–230. In: R. Lontie (ed). *Copper Proteins and Copper Enzymes*. CRC, Boca Raton.
- Ganguli A. and A.K. Tripathi.** 2002. Bioremediation of toxic chromium from electroplating effluent by chromate-reducing *Pseudomonas aeruginosa* A2Chr in two bioreactors. *Appl. Microbiol. Biotechnol.* 58: 416–420.
- Garbisu C., I. Alkorta, M.J. Llama and J.L. Serra.** 1998. Aerobic chromate reduction by *Bacillus subtilis*. *Biodegrad.* 9: 133–141.
- Horitsu H., S. Futo, Y. Miyazawa, S. Ogai and K. Kawai.** 1987. Enzymatic reduction of hexavalent chromium by hexavalent chromium tolerant *Pseudomonas ambigua* G-1. *Agric. Biol. Chem.* 51: 2417–2420.
- Iftikhar S., M. Faisal and S. Hasnain.** 2007. Cytosolic reduction of toxic Cr(VI) by indigenous microorganism. *Res. J. Env. Sci.* 1: 77–81.
- Ishibashi Y., C. Cervantes and S. Silver.** 1990. Chromium reduction in *Pseudomonas putida*. *Appl. Environ. Microbiol.* 56: 2268–2270.
- Losi M.E., C. Amrhein and W.T. Frankenberger.** 1994. Environmental biochemistry of chromium. *Rev. Environ. Contam. Toxicol.* 36: 91–121.
- Lovley D.R. and E.J.P. Phillips.** 1994. Reduction of chromate by *Desulfovibrio vulgaris* and its *c<sub>3</sub>* cytochrome. *Appl. Environ. Microbiol.* 60: 726–728.
- Lowry O.H., N.J. Resebrough and A.L. Farr.** 1951. Protein measurement with the folin-phenol reagent. *J. Biol. Chem.* 193: 265–275.
- McLean J. and T.J. Beveridge.** 2001. Chromate reduction by a pseudomonad isolated from a site contaminated with chromated copper arsenate. *Appl. Environ. Microbiol.* 67: 1076–1084.
- Myers C.R., B.P. Carstens, W.E. Antholine and J.M. Myers.** 2000. Chromium (VI) reductase activity is associated with the cytoplasmic membrane of anaerobically grown *Shewanella putrefaciens* MR-1. *J. Appl. Microbiol.* 88: 98–106.
- Ohtake H. and S. Silver.** 1994. Bacterial detoxification of toxic chromate. pp. 403–415. In: G.R. Choudhuri (ed). *Biological Degradation and Bioremediation of Toxic Chemicals*. Discorides Press, Portland.
- Pal A., S. Dutta and A.K. Paul.** 2005. Reduction of hexavalent chromium by cell-free extract of *Bacillus sphaericus* AND 303 isolated from serpentine soil. *Curr. Microbiol.* 66: 327–330.
- Park C.H., B. Keyhan, B. Wielinga, S. Fendorf and A. Matin.** 2000. Purification to homogeneity and characterization of a novel

- Pseudomonas putida* chromate reductase. *Appl. Environ. Microbiol.* 66: 1788–1795.
- Sau G.B., S. Chatterjee, S. Sinha and S.K. Mukherjee.** 2008. Isolation and characterization of a Cr(VI) reducing *Bacillus firmus* strain from industrial effluents. *Polish J. Microbiol.* 57: 327–332.
- Shakoori A.R., S. Tahseen and R.U. Haq.** 1999. Chromium-tolerant bacteria isolated from industrial effluents and their use in detoxification of hexavalent chromium. *Folia Microbiol.* 44: 50–54.
- Shen H. and Y.T. Wang.** 1993. Characterization of enzymatic reduction of hexavalent chromium by *Escherichia coli* ATCC 33456. *Appl. Environ. Microbiol.* 59: 3771–3777.
- Shi X. and N.S. Dalal.** 1990. On the hydroxyl radical formation in the reaction between hydrogen peroxide and biologically generated chromium(V) species. *Arch. Biochem. Biophys.* 277: 342–350.
- Suzuki T., N. Miyata, H. Horitsu, K. Kawai, K. Tsakamizawa, Y. Tai and M. Okazaki.** 1992. NAD(P)H dependent chromium (VI) reductase of *Pseudomonas ambigua* G-1: A Cr(V) intermediate is formed during the reduction of Cr(VI) to Cr(III). *J. Bacteriol.* 174: 5340–5345.
- Thacker U. and M. Datta.** 2005. Reduction of toxic chromium and partial localization of chromium reductase activity in bacterial isolate DM1. *World J. Microbiol. Biotechnol.* 21: 891–899.
- Urone F.E.** 1955. Stability of colorimetric reagent for chromium. S-diphenylcarbazides in various solvents. *Anal. Chem.* 27: 1354–1355.
- Wang P.C., T. Mori, K. Toda and H. Ohtake.** 1990. Membrane associated chromate reductase activity from *Enterobacter cloacae*. *J. Biotechnol.* 172: 1670–1672.
- Wang Y.T. and C. Xiao.** 1995. Factors affecting hexavalent chromium reduction in pure cultures of bacteria. *Water Res.* 29: 2467–2474.
- Xu X.R., H.B. Li and J.D. Gu.** 2004. Reduction of hexavalent chromium by ascorbic acid in aqueous solutions. *Chemosphere* 57: 609–613.
- Xu X.R., H.B. Li, J.D. Gu and X.Y. Li.** 2005. Kinetics of the reduction of chromium (VI) by vitamin C. *Environ. Toxicol. Chem.* 24: 1310–1314.