

Chromosomal Damage Risk Assessment to Benzene Exposure among Gasoline Station Workers in Bangkok Metropolitan, Thailand

Tanasorn Tunsaringkarn^{1*} Panthira Ketkaew² Jamsai Suwansaksri³ Wattasit Siriwigong¹

Anusorn Rungsiyothin¹ Kalaya Zapuang¹ Mark Gregory Robson^{4,5,6}

1. College of Public Health Sciences, Chulalongkorn University, Institute Building 2-3, Soi Chulalongkorn 62 Phyathai Rd, Bangkok 10330, Thailand
2. Department of chemistry, Faculty of Science, King Mongkut's University of Technology Thonburi, Bangkok 10140, Thailand
3. Department of Clinical Chemistry, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand
4. School of Environmental and Biological Sciences, Rutgers University, 59 Dudley Rd, Rm 204A Foran Hall New Brunswick, New Jersey 08854, USA
5. UMDNJ-School of Public Health, Piscataway, 65 Bergen St, Suite 701 Newark, New Jersey 08854, USA
6. Thai Fogarty ITREOH Center, Chulalongkorn University, Bangkok 10330, Thailand

* E-mail of the corresponding author: tkalayan@chula.ac.th

Abstract

This study was a cross-sectional survey study to assess relative risk (RR) of chromosomal damage through benzene exposure in 45 gasoline stations workers compared to 30 controls in central area of Bangkok. Sister chromatid exchange (SCE) is as genotoxic biomarker, performed in white blood cells, and blood benzene level (BBL) is a biological marker of benzene exposure was performed by gas chromatography-flame ionization detector (GC-FID) using modified headspace solid-phase micro-extraction (HS-SPME) technique. The results showed that the average blood benzene level of these workers was significantly higher than in the controls ($p < 0.001$) as well as the frequency of sister chromatid exchange. The sister chromatid exchange was strongly and positively associated with blood benzene level of gasoline workers ($p < 0.001$) with the chromosomal damage relative risk at 2.50 ($p < 0.001$).

Keywords: gasoline worker, benzene, sister chromatid exchange, chromosomal damage

1. Introduction

Air pollution has become to be a serious health problem in Bangkok, Thailand, especially among occupational workers such as gasoline workers. Exposure to gasoline vapors is classified by the International Agency for Research on Cancer as possible cancer risk in humans, mainly on the basis of the established carcinogenicity of some chemical components such as benzene (IARC 1989). The mechanism of benzene toxicity, particularly its leukemogenic effects, is far from being fully understood. Contamination of the environment with volatile organic compounds (VOCs) has become an important issue, since many of these compounds are toxic and may pose health risks of various concerns. To assess the biological risks caused by gasoline vapor, by biological monitoring using biological marker of oxidative chromosomal damage and repairing capacity (Lambert *et al.* 1982; Carrano *et al.* 1983), as frequency of sister chromatid exchange (SCE) in gasoline workers, may provide useful information about the genotoxic risk associated with exposure to this carcinogenic agent that is benzene. In major cities in Asia, the levels of ambient air benzene are relatively high compared with those in Europe or in the United States (Leong & Laortanakul 2003) and there is high prevalence of cancer and leukemia related to petrol station density (Yimrungruang *et al.* 2008; Chang *et al.* 2009; Weng *et al.* 2009). The purpose of this study was to evaluate the relative risk

(RR) of chromosomal damage cause by benzene exposure in gasoline station workers in central area of Bangkok, Pathumwan district, comparing to controls.

2. Research Methods

2.1 Population study

A cross sectional surveyed at 11 gasoline stations in Pathumwan district, Bangkok, Thailand, and collected blood samples analysis from 45 gasoline workers, which 33 non-smokers and 12 smokers, compared to 30 non-smoker controls from April to June 2009. All subjects have given informed consent before the study. The Ethical Review Committee for Research Involving Human Research Subjects, Health Science Group, Chulalongkorn University, approved the study. All subjects were healthy and had worked more than six months.

2.2 Sample collection

The venous blood samples were drawn from subjects during the 6-8 h shift work, 2 x 2 mL, using plastic heparinized vacuum blood tube, stored at 4 °C, one SCE frequency analysis within 6 h and the other one stored at -20 °C before benzene analysis within 1 month.

2.3 Sample analyses

The blood examination for SCE frequency was modified in manner suggested by Tucker and Preston (1996). Aliquots of 0.5 mL of heparin zed blood samples, both worker and control groups, added to 5.0 mL of the culture medium containing Roswell Park Memorial Institute medium (RPMI 1640, Hyclone, Utah, USA), supplemented with 15% fetal bovine serum, 2.5% phytohemagglutinine (PHA, Sigma, Germany) and 1% penicillin-streptomycin. Afterwards 100 µL of 1.3 mg/L 5'-bromodeoxyuridine (BrdU, Sigma Chemical Co.) was added to the medium and additionally incubated in the dark room temperature for 96 hours. Immediately, added colchicine (0.2 µg/L, Sigma Chemical Co.), collected cultured cells and treated with 0.075 mol/L potassium chloride (KCl) at 37 °C for 10 min to be fixed with methanol-acetic acid (3:1). Standard harvest procedure was performed by a drop of harvested cell pellets spread on clean glass slide and stained by Hoechst No.22358 plus Giemsa technique (Koto *et al.* 1975). Finally, the slides were examined by a light microscope (Nikon E200) in regard to SCE frequency/metaphase cell. The total of 15 well-spread metaphases was evaluated in worker and control groups. Counting of SCE frequency was done by using oil immersion.

Blood benzene determination was performed by GC-FID using modified HS-SPME technique (Tunsaringkarn *et al.* 2004). Briefly, to 0.5 mL of blood sample in glass cap bottle was added 0.2 g of sodium chloride, shaking vortex for 15 s, then controlled and absorbed by SPME in water bath at 50 °C with vibrator for 20 min. Injected in GC (Varian CP 3800) at 220 °C with flame ionization detector at 220 °C (used column CP-SIL5 CB, split less). Oven temperature started at 50 °C for 10 min, then increased 5 °C/min until 90 °C and finally increased by 30 °C/min until 250 °C for 17 min. The quantity of blood benzene was analyzed under relative intensity of chromatographic signal for 40 min. The Limit of Detection (LOD) of benzene was 10.00 µg/L (ppb) and the average coefficient of determination (r^2) was 0.999657.

Statistical analyses were carried out with the SPSS 17.0 statistical software (SPSS Inc., Chicago, IL, USA). Descriptive statistics were used for BBL and frequency of SCE in gasoline worker and control groups which were presented as mean and standard error of the estimate (Mean \pm SE). The comparison between control and worker parameters were analyzed by independent-t test with a value $p < 0.05$ limitation. The association between parameters and SCE was the estimated relationship of them by multiple linear regression, which SCE as dependent variable, BBL, age, sex and cigarette smoking as independent variables. The relative risk (RR) of chromosomal damage was calculated by comparing the proportion of

workers with frequency of SCE higher than the mean plus standard error value of control to proportion of workers with frequency of SCE lower than the mean plus standard error value of the control group. Chi-square analysis was used to determine differences between groups in frequency distribution with 95% confidence interval (CI).

3. Analysis results

A total of 75 subjects, 30 controls and 45 gasoline station workers, were included in this study. The average age of worker and control groups was 31.2 ± 1.4 and 31.2 ± 1.8 years. The gasoline workers were 60% male and 26.7% smoker while the controls were 60% male and all of them were non-smokers. The frequency of SCE in workers and controls was 13.62 ± 0.24 and 6.97 ± 0.20 SCE/cell (Table 1 and Figure 1). Both, smoking and non-smoking workers, SCE frequencies were significantly higher than those of controls ($p < 0.001$). SCE frequency in smokers had higher level than in non-smokers but there was no significant difference. The average BBL of workers and controls was 302.84 ± 20.54 and 27.28 ± 15.31 $\mu\text{g/L}$, but workers had significantly higher levels than controls ($p < 0.001$) as shown in Table 1. But SCE and BBL of male and female were not difference in each group. In addition, the benzene exposure risk ratio (BBL of workers/controls) was 11.83 while the chromosomal damage risk ratio (SCE of workers/controls) was 1.94.

The association between BBL and SCE frequency was analyzed by multiple linear regression with SCE as dependent variable and BBL, age, sex and cigarette smoking as independent variables (Table 2). The frequency of SCE was positively and significantly associated with BBL ($p = 0.001$). Age, sex and cigarette smoking, were not significantly associated with SCE ($p > 0.05$). The frequencies of SCE were dichotomized into high- and low-frequency groups based on their means plus standard error of estimate (SE) values of control group at $7.17 (6.97 + 0.2)$. The chromosomal damage relative risk of gasoline station workers compared to controls was 2.50 (χ^2 -test, 95% CI = 1.17-5.34, $p < 0.001$).

4. Discussion

Volatile organic compound in gasoline is a common source of benzene which is classified as carcinogen. It can cause serious health effects including genotoxicity. Biomarkers of response indicate biological or biochemical changes in target tissue or surrogate from chemical action. The biomarkers commonly used are chromosome aberration (CA), micronucleus (MN) and SCE. This study used SCE as genotoxic biomarker and BBL as exposure biomarker. The results of this study showed significantly higher genotoxic or chromosomal damage in gasoline workers than in controls which supported the previous studies (Vijayalaxmi & Evans 1982; Celik & Akba 2005; Calderón-Ezquerro *et al.* 2007). The SCE frequency was not effective for low smoker workers with less than 10 cigarettes a day (average number of cigarette smoking 5.2 cigarette/day), but there was a trend of higher chromosomal damage in smokers than in non-smokers. The comparison between smokers and non-smokers should be related to the number of cigarettes smoked a day with a significant difference in moderate and heavy smoker (Vijayalaxmi & Evans 1982; Celik & Akba 2005; Calderón-Ezquerro *et al.* 2007). The BBL of all workers was significantly higher than in controls ($p < 0.001$), but BBL of smokers trending upwards compared to non-smokers, but not significantly different, most non-smokers being women with higher heart rate than men (Ryan *et al.* 1994; Stein *et al.* 1997). As the results of women were higher in BBL than in men with same rate of benzene clearance, it may influence its internal exposure and cancer development. Gender affected more BBL than benzene exposure from low cigarette smoking. In addition, BBL of all gasoline workers were higher than the biological monitoring of occupationally exposed persons, an exposure equivalent for carcinogenic working material (EKA-value) of $54 \mu\text{g/L}$ (Angerer *et al.* 1991). It should be considered that BBL in workers was 5.6 folds of limited level, corresponding to high level of benzene exposure (11.83 folds of exposure risk ratio) and this marker indicating that cells have been exposed to mutagen or carcinogen (Keretse *et al.* 2008). The frequency of SCE provided information of cumulative effects of carcinogens which was associated with increased risk of cancer (1.9 folds of chromosomal damage risk ratio). The results of this study indicated that the frequency of SCE was strongly associated to BBL ($p < 0.001$). With

every increase of one year in age, the average SCE decreased by 0.25 whilst every increasing of BBL level 1 µg/L, the average SCE increased by 0.44. In fact, the average age of control group was not higher than the average in the worker group. Age, cigarette smoking and sex were not significantly associated to chromosomal damage. Some studies showed that no correlation between SCE frequency, duration of exposure, smoking habit, and age (Hoet *et al.* 2009; Ulker *et al.* 2008). The gasoline workers had an increased relative risk of chromosomal damage at 2.50 compared to controls (χ^2 -test, 95% CI = 1.17-5.34, $p < 0.001$), which benzene exposure and chromosomal damage risk ratios of the workers were 11.93 and 1.94, respectively.

However, biomonitoring in peripheral lymphocytes served as an early indicator of chromosomal damage. Biomonitoring of benzene exposure by using BBL among the gasoline workers should show the association of cancer risk development in gasoline workers prompting to plan a screening program and primary prevention for them to minimize the risk of cancer. Future studies of genotoxicity in smokers and non-smokers comparison should cover a large population and take into consideration confounders such as age, gender, smoking habits, alcohol consumption and family history of cancer.

Acknowledgments

This study was supported by Thai Fogarty ITREOH Center (D43 TW007849 NIH FIC), College of Public Health Sciences, and Chulalongkorn University Centenary Academic Development Project, Bangkok, Thailand. And, many thanks go to the gasoline station owners and their workers for the agreeable collaboration. Last, but not least, thanks to Dr. Karl J. Neeser for his advice.

References

- Angerer, J., Seherer, G., Sehallar, K.H., & Müller, J. (1991), "The determination of benzene in human blood as an indicator of environmental exposure to volatile aromatic compounds", *Fresenius Journal of Analytical Chemistry* **339**, 740-742.
- Calderón-Ezquerro, C., Sánchez-Reyes, A., Sansores, R.H., Villalobos-Pietrini, R., Amador-Muñoz, O., Guerrero-Guerra, C., Calderón-Segura, M.E., Uribe-Hernández, R., Gómez-Arroyo, S. (2007), "Cell proliferation kinetics and genotoxicity in lymphocytes of smokers living in Mexico City", *Human Experimental Toxicology* **26**(9), 715-722.
- Carrano, A.V., & Natarajan A.T. (1988), "Considerations for population monitoring using cytogenetic techniques", *Mutation Research* **204**, 379-406.
- Celik, K.A., & Akba S.E. (2005), "Evaluation of sister chromatid exchange and chromosomal aberration frequencies in peripheral blood lymphocytes of gasoline station attendants", *Ecotoxicology and Environmental Safety* **60**(1), 106-112.
- Chang, C.C., Tsai, S.S., Chiu, H.F., Wu, T.N., & Yang C.Y. (2009), "Traffic air pollution and lung cancer in females in Taiwan: petrol station density as an indicator of disease development", *Journal of Toxicology and Environmental Health, Part A* **72**(10), 651-657.
- Hoet, P. De Smedt, E., Ferrari, M., Imbriani, M., Maestri, L., Negri, S., De Wilde, P., Lison, D., & Haufroid, V. (2009), "Evaluation of urinary biomarkers of exposure to benzene: correlation with blood benzene and influence of confounding factors", *International Archives of Occupational and Environmental Health* **82**, 985-995.
- IARC (International Agency For Research On Cancer). (1989), "Occupational exposures in petroleum refining; crude oil and major petroleum fuels", Monographs on the evaluation of carcinogenic risks to humans. Vol. 45, Lyon, France.
- Keretetse, G.S., P. J. Laubscher, P.J., Du Plessis, J.L., Pretorius, P.J., Van Der Westhuizen, F.H., Van Deventer, E., Van Dyk, E., Eloff, F.C., Van Aarde, M.N., & Du Plessis, L.H. (2008), "DNA damage and repair detected by the comet assay in lymphocytes of African petrol attendants: A pilot study". *Annals of Occupational Hygiene* **52**(7), 653-662.

Koto, K., Akematsu, T., Shimadzu, H., & Sugiyama, T. (1975), "Simple differential Giemsa staining of sister chromatid after exposure to light and the mechanism of staining", *Chromosoma (Berl.) Springer-Verlag* **53**, 223-230.

Lambert, B.A., Lindblad K., Holmberg K., & Francesconi, S. (1982), "The use of sister chromatid exchanges to monitor human populations for exposure to toxicologically harmful agents", in: Wolff, S., (Ed.), *Sister Chromatid Exchange*, New York, Wiley-Interscience, pp. 149-182.

Leong, S.T., & Laortanakul, P. (2003), "Indicators of benzene emissions and exposure in Bangkok street", *Environmental Research* **92**(3), 173-181.

Pitarque, M., Carbonell, E., Lapeña, N., Marsá, M., Valbuena, A., Creus, A., & Marcos, R. (1997), "SCE analysis in peripheral blood lymphocytes of a group of filling station attendants" *Mutation Research* **390**(1-2), 153-159.

Ryan, S.M., Goldberger, A.L., Pincus, S.M., Mietus, J., & Lipsitz, L.A. (1994), "Gender- and age-related differences in heart rate dynamics: are women more complex than men?", *Journal of the American College of Cardiology* **24**(7), 1700-1707.

Soogarun, S., Suwansakri, J., & Wiwanitkit, V. (2006), "High sister chromatid exchange among a sample of traffic policemen in Bangkok, Thailand", *Southeast Asian Journal of Tropical Medicine and Public Health* **37**(3), 578-580.

Stein, P.K., Kleiger, R.E., & Rottman, J.N. (1997), "Differing effects of age on heart rate variability in men and women", *American Journal of Cardiology* **80**(3), 302-305.

Tucker, J.D., & Preston R.J. (1996), "Chromosome aberrations, micronuclei, aneuploidy, sister chromatid exchanges, and cancer risk assessment", *Mutation Research* **363**, 147-159.

Tunsaringkarn, T., Choochat, N., & Theppitaksak, B. (2004), "Headspace-Solid Phase Micro-extraction for determination of benzene, toluene, ethylbenzene, xylene and mTBE in blood", *Thai Journal of Health Research* **18**(1), 50-59.

Ulker, O.C., Ustundag, A., Duydu, Y., Yucesoy, B., & Karakaya, A. (2008), "Cytogenetic monitoring of coal workers and patients with coal workers' pneumoconiosis in Turkey", *Environmental and Molecular Mutagenesis* **49**, 232-237.

Vijayalaxmi., & Evans, H.J. (1982), "In vivo and in vitro effects of cigarette smoke on chromosomal damage and sister-chromatid exchange in human peripheral blood lymphocytes", *Mutation Research* **92**(1-2), 321-322.

Weng, H.H., Tsai, S.S., Chiu, H.F., Wu, T.N., & Yang, C.Y. (2009), "Childhood leukemia and traffic air pollution in Taiwan: petrol station density as an indicator", *Journal of Toxicology and Environmental Health, Part A* **72**(2), 83-87.

Yimrungruang, D., Cheevaporn, V., Boonphakdee, T., Watchalayann, P., & Helander, H.F. (2008), "Characterization and health risk assessment of volatile organic compounds in gas service station workers", *EnvironmentAsia* **2**, 21-29.

Zhou, X., Li, L., Cui, M., Yu, R.F., Li, L., & Yan, Z.A. (1986), "Cytogenetic monitoring of petrochemical workers", *Mutation Research* **175**, 237-242.

Tanasorn Tunsaringkarn received her M.Sc. (Biochemistry) degree from Chulalongkorn University (CU) in 1978. From 1978-2007, she worked as researcher for Institute of Health Research, CU and from 2007-present as a faculty member, Assistant Dean and an executive board member at College of Public Health Sciences (CPHS), CU. She is a member of the Asia Pacific Academic Consortium of Public Health (APACPH). Her research interests include environmental health and toxicology.

Panthira Ketkaew is the Head of Department of chemistry, Faculty of Science, King Mongkut's University of Technology Thonburi. She received her Bachelor of Science in Chemistry from Chiangmai

University in 1975 and Doctoral degree in Chemistry Oklahoma State University in 1982. Her research includes analytical environmental analyses and passive sampler.

Jamsai Suwansaksri received her M.Sc. (Toxicology) degree from Mahidol University in 1971, B.Sc. (Occupational Health and safety) from SukhothaiThammathirat University in 2003 and B.Sc. (Medical Technology) from Chiangmai University. She worked as lecturer/ assistant professor at Department of Clinical Chemistry, Faculty of Medical Technology, Khonkaen University from 1969-1994. Now, she is an associate professor at Department of Clinical Chemistry, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok, Thailand.

Wattasit Siriwong is a faculty member, Assistant Dean, and an executive board member of the College of Public Health Sciences (CPHS), Chulalongkorn University. He received a Bachelor of Science in Biotechnology from Mahidol University, Master of Science degree in Environmental Science from Thammasat University and his Doctoral degree in Environmental Management from Chulalongkorn University. Dr. Wattasit is an active researcher in the field of environmental health. He is also a member of the editorial board of International Journal of Occupational and Environmental Health and Journal of Health Research.

Anusorn Rungsiyothin received his B.Ed. degree in biology from Srinakharinwirot University in 1974. He is a research-staff in laboratory of College of Public Health Sciences (CPHS), Chulalongkorn University.

Kalaya Zapuang received her B.Ed. degree from Chandrakasem Rajabhat University in 1974. She is a research-staff at College of Public Health Sciences (CPHS), Chulalongkorn University.


Mark Gregory Robson Mark Gregory Robson is the Dean of Agricultural and Urban Programs and Professor of Entomology at Rutgers University-School of Environmental and Biological Sciences and Professor of Environmental and Occupational Health the University of Medicine and Dentistry of New Jersey (UMDNJ) School of Public Health (SPH). He graduated with a B.S. with High Honors (1977) M.S. (1979) and Ph.D. (1988) from Rutgers University and M.P.H. (1995) from UMDNJ SPH. Dr. Robson received an honorary doctoral degree in Public Health from Chulalongkorn University in 2010.

*Statistical difference (independent-t test) between control and gasoline workers					
Parameter	Control		Gasoline Worker		p - value*
	N	(Mean ± SE)	N	(Mean ± SE)	
Age (years)	30	31.2 ± 1.8	45	31.2 ± 1.4	< 0.001
Frequency of SCE (SCE/cell)	30	6.97 ± 0.20	45	13.62 ± 0.24	< 0.001
Sex					
Female	12	7.28 ± 0.13	18	13.41 ± 0.33	< 0.001
Male	18	6.77 ± 0.31	27	13.76 ± 0.34	< 0.001
Cigarette Smoking					
Non Smoker	30	6.97 ± 0.20	33	13.47 ± 0.26	< 0.001
Smoker		-	12	14.07 ± 0.59	
Blood Benzene (µg/L)	30	27.28 ± 15.31		302.84 ± 20.54	0.001
Sex					
Female	12	36.65 ± 36.65	18	322.26 ± 0.26	0.001
Male	18	21.03 ± 12.10	27	291.05 ± 20.91	0.001
Cigarette Smoking					
Non Smoker	30	27.28 ± 15.31	33	201.97 ± 24.78	0.001
Smoker	-	-	12	321.33 ± 25.96	
Benzene Exposure Risk Ratio (BBL of workers/controls) = 11.83					
Chromosomal Damage Risk Ratio (SCE of workers/controls) = 1.94					

Table 1. BBL and frequency of SCE in controls and gasoline workers

Dependent Variable: SCE					
*Statistical relation between SCE and parameters					
Parameter	Unstandardized		Standardized Coefficients	95% CI Lower to Upper	p - value*
	B	SE			
Blood Benzene Level	0.008	0.002	0.439	0.004 to 0.012	0.001
Age	-0.064	0.033	-0.249	-0.131 to -0.004	0.063
Sex	-0.101	0.700	-0.016	-0.1506 to 1.305	0.886
Cigarette Smoking	1.460	0.879	0.202	-0.222 to 3.142	0.087

Table 2. Association between SCE frequency and parameters



(a) Control group

(b) Gasoline worker group

The symbol (↔) indicated sister chromatid separation.

Figure 1. Sister chromatid exchanges of control and gasoline worker groups

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. **Prospective authors of IISTE journals can find the submission instruction on the following page:**

<http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a fast manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

