

Chronic Peripheral Inflammation is Associated With Cognitive Impairment in Schizophrenia: Results From the Multicentric FACE-SZ Dataset

Ewa Bulzacka^{1,2}, Laurent Boyer^{1,3}, Franck Schürhoff^{1,4}, Ophélie Godin^{1,5}, Fabrice Berna^{1,6}, Lore Brunel^{1,2}, Méja Andrianarisoa^{1,2}, Bruno Auizerate^{1,7,8}, Delphine Capdevielle^{1,9}, Isabelle Chéreau-Boudet^{1,4}, Gabrielle Chesnoy-Servanin^{1,10}, Jean-Marie Danion^{1,6}, Caroline Dubertret^{1,11}, Julien Dubreucq^{1,12}, Catherine Faget^{1,13}, Franck Gabayet^{1,12}, Tifenn Le Gloahec^{1,2}, Pierre-Michel Llorca^{1,4}, Jasmina Mallet^{1,11}, David Misdrahi^{1,7,14}, Romain Rey^{1,10}, Raphaëlle Richieri^{1,13}, Christine Passerieux^{1,15}, Paul Roux^{1,15}, Hanan Yazbek^{1,9}, Marion Leboyer^{1,2}, and Guillaume Fond^{*,1,2}, the FACE-SZ (FondaMental Academic Centers of Expertise for Schizophrenia) Group[†]

¹Fondation FondaMental, RTRS santé mentale, Créteil, France; ²INSERM U955, Translational Psychiatry Team, Créteil, France, Paris Est University, DHU Pe-PSY, Pôle de Psychiatrie des Hôpitaux Universitaires H Mondor, Créteil, France; ³Pôle psychiatrie universitaire, CHU Sainte-Marguerite, F-13274 Marseille cedex 09, France; ⁴CMP B, CHU, EA 7280 Faculté de Médecine, Université d'Auvergne, BP 69 63003 Clermont-Ferrand Cedex 1, France; ⁵Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, F-75013, Paris, France, INSERM, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, F-75013, Paris, France; ⁶Hôpitaux Universitaires de Strasbourg, Université de Strasbourg, INSERM U1114, Fédération de Médecine Translationnelle de Strasbourg, Strasbourg, France; ⁷Centre Hospitalier Charles Perrens, Université de Bordeaux, F-33076 Bordeaux, France; ⁸Inserm, Neurocentre Magendie, Physiopathologie de la Plasticité Neuronale, U862, F-33000 Bordeaux, France; ⁹Service Universitaire de Psychiatrie Adulte, Hôpital la Colombière, CHRU Montpellier, Université Montpellier 1, Inserm 1061, Montpellier, France; ¹⁰Université Claude Bernard Lyon 1/Centre Hospitalier Le Vinatier Pole Est BP 300 39—95 bd Pinel—69678 BRON Cedex, France; ¹¹AP-HP, Department of Psychiatry, Louis Mourier Hospital, Colombes, Inserm U894, Université Paris Diderot, Sorbonne Paris Cité, Faculté de médecine, France; ¹²Centre Référent de Réhabilitation Psychosociale, CH Alpes Isère, Grenoble, France; ¹³Assistance Publique des Hôpitaux de Marseille (AP-HM), pôle universitaire de psychiatrie, Marseille, France; ¹⁴CNRS UMR 5287 - Institut de Neurosciences cognitives et intégratives d'Aquitaine (INICIA), Bordeaux, France; ¹⁵Service de psychiatrie d'adulte, Centre Hospitalier de Versailles, UFR des Sciences de la Santé Simone Veil, Université Versailles Saint-Quentin en Yvelines, Versailles, France

*To whom correspondence should be addressed; Centre Expert Schizophrénie, Pôle de Psychiatrie et d'Addictologie des Hôpitaux Universitaires Henri Mondor, 40 rue de Mesly, 94 000 Créteil, France; tel: 33-1-49-81-32-90, fax: 33-1-49-81-30-99, e-mail: guillaume.fond@gmail.com

[†]FACE-SCZ Group members and their affiliations are provided in the Acknowledgments section.

Objectives: Inflammation, measured by abnormal blood C-reactive protein (CRP) level, has been described in schizophrenia (SZ), being inconsistently related to impaired cognitive functions. The aim of the present study is to investigate cognitive impairment associated with abnormal CRP levels in a large multi-centric sample of community-dwelling SZ patients, using a comprehensive neuropsychological battery. **Method:** Three hundred sixty-nine community-dwelling stable SZ subjects (76.2% men, mean age 32.7 y) were included and tested with a comprehensive battery of neuropsychological tests. Abnormal CRP level was defined as $>3\text{mg/L}$. **Results:** Multiple factor analysis revealed that abnormal CRP levels, found in 104 patients (28.2%), were associated with impaired General Intellectual Ability and Abstract Reasoning (aOR = 0.56, 95% CI 0.35–0.90, $P = .014$), independently of age, sex, education level, psychotic symptomatology, treatments, and addiction

comorbidities. Abnormal CRP levels were also associated with the decline of all components of working memory (respectively effect size [ES] = 0.25, $P = .033$; ES = 0.27, $P = .04$; ES = 0.33, $P = .006$; and ES = 0.38, $P = .004$) and a wide range of other impaired cognitive functions, including memory (ES = 0.26, $P = .026$), learning abilities (ES = 0.28, $P = .035$), semantic memory (ES = 0.26, $P = .026$), mental flexibility (ES = 0.26, $P = .044$), visual attention (ES = 0.23, $P = .004$) and speed of processing (ES = 0.23, $P = .043$). **Conclusion:** Our results suggest that abnormal CRP level is associated with cognitive impairment in SZ. Evaluating the effectiveness of neuroprotective anti-inflammatory strategies is needed in order to prevent cognitive impairment in SZ.

Key words: C-reactive protein (CRP)/schizophrenia/cognition/cognitive impairment/inflammation

Introduction

The contribution of chronic inflammation to schizophrenia (SZ) has received considerable attention in the last decade.^{1,2} In clinical practice, chronic inflammation is primarily measured by an elevated blood level of C-reactive protein (CRP).³ A recent meta-analysis revealed an increased rate of abnormal CRP levels $\geq 5\text{mg/L}$ in 28% of SZ subjects compared to healthy controls,⁴ with abnormal CRP levels being recently found to be a risk factor for late-onset SZ.⁵ Chronic inflammation was also suggested to be one of the biological substrates of several major psychiatric disorders including depression^{6,7} and bipolar disorders.⁸

Cognitive symptoms are thought to have a higher impact on the outcome of SZ than any other symptoms, and are considered as a core feature of this disorder.⁹ Approximately 25% of people with SZ have a poor long-term outcome due to impaired cognition.¹⁰ However the association between abnormal CRP levels and impaired cognitive function in SZ is unclear. In non-SZ populations, increased CRP levels has been associated with a wide range of brain dysfunctions including age-related cognitive decline,^{11,12} all-cause dementia,¹³ and stroke.¹⁴

In SZ, only 2 studies exploring the impact of inflammation on cognition have been carried out to date. In the first study, abnormal CRP levels were associated with cognitive impairment.¹⁵ This study was well performed although the cognitive assessment comprised a 30 minutes-long battery, which provided only a brief screening of different cognitive domains associated with abnormal CRP levels. In a second study using the MATRICS battery, no association between CRP levels and cognitive decline was found.¹⁶ However, the sample size ($N = 88$) was small and the absence of association may be due to a lack of statistical power. Thus, given the potential impact of cognitive impairment prevention and treatment in SZ, the association between abnormal CRP levels and cognitive function deserves further explorations in this population.²

The objective of the present study was to investigate if cognitive impairment was associated with elevated CRP levels in a large multi-centric sample of community dwelling stabilized subjects with SZ, using an exhaustive neuropsychological battery.

Materials and Methods

Study Population

The FACE-SZ (FondaMental Academic Centers of Expertise for Schizophrenia) cohort is based on a French national network of 10 Schizophrenia Expert Centers (Bordeaux, Clermont-Ferrand, Colombes, Créteil, Grenoble, Lyon, Marseille, Montpellier, Strasbourg, Versailles), set up by a French scientific cooperation foundation, FondaMental Foundation (www.fondamental.org) and created by the French Ministry of Research in order to build a platform that links systematic clinical assessment to research.¹⁷

[fondamental.org](http://www.fondamental.org)) and created by the French Ministry of Research in order to build a platform that links systematic clinical assessment to research.¹⁷

Inclusion Criteria

Consecutive clinically stable patients (defined by no hospitalization and no treatment changes during the 4wk before evaluation) with a DSM-IV-TR diagnosis of SZ or schizoaffective disorder were consecutively included in the study. Diagnosis was confirmed by 2 trained psychiatrists of the Schizophrenia Expert Centres network. All subjects were referred by their general practitioner or psychiatrist who subsequently received a detailed evaluation report with suggestions for personalized interventions.

Noninclusion Criteria

Patients with a history of neurological disorders (including stroke, epilepsy, and head injury) or all non-psychiatric concurrent illnesses affecting the central nervous system and inflammation (especially auto-immune illnesses, such as lupus and rheumatoid arthritis) were excluded from the present study.

Measurements

Clinical and Sociodemographic Measures. All patients were interviewed by members of the specialized multidisciplinary team of the Expert Centre. All subjects were interviewed by 2 independent psychiatrists with the Structured Clinical Interview for Mental Disorders (SCID 1.0) to confirm the diagnosis. Information about education, onset and course of the illness, family history, somatic diseases and comorbidities were recorded. Schizophrenic symptomatology was assessed using the Positive And Negative Syndrome Scale (PANSS).¹⁸ Current depressive symptoms were evaluated using the Calgary Depression Rating Scale for Schizophrenia (CDRS).¹⁹ Manic symptoms were assessed with Young Mania Rating Scale (YMRS).²⁰ Ongoing psychotropic treatment, current cannabis consumption and tobacco smoking were also recorded.

Neuropsychological Measures. The National Adult Reading Test²¹ provides an estimate of premorbid intellectual ability. French version of the National Adult Reading Test was used in our analysis.²²

Wechsler Adult Intelligence Scale—3rd Edition²³ provides a measure of general intellectual function in older adolescents and adults. Seven subtest short form²⁴ was used to estimate the Full Scale IQ (FSIQ), Verbal IQ (VIQ) and Performance IQ (PIQ), and allowed exploration of the following cognitive areas: Picture Completion (visual exploration and detail perception), Digit-Symbol Coding (visual-motor coordination, motor and mental speed), Similarities (abstract verbal reasoning), Arithmetic

(mathematical problem solving), Matrix Reasoning (non-verbal abstract problem solving, inductive spatial reasoning), Digit span (attention, working memory, mental control), Information (general information acquired from culture, semantic memory). An additional subtest, Letter-Number Sequencing was administered, which along with 2 other primary subtests, Digit Span and Arithmetic, allowed the calculation of a Working Memory Index (WMI; auditory working memory and mental control).

Trail Making Test²⁵ reflects the control of attention, visual exploration, speed and mental flexibility. The subject is asked to connect, by making pencil lines, encircled numbers randomly arranged on a page in proper order (Part A) and encircled numbers and letters in alternating order (Part B). A French version of the normative data was used.²⁶

California Verbal Learning Test²⁷ is designed to measure verbal learning and memory using a multiple-trial list-learning task. The examiner reads the word list and records the patient's oral responses verbatim in the order in which they are given. Learning efficiency, strategies, interference management and learning bias are measured. A French version of the task was used in this study.²⁸

Doors test²⁹ is a visual recognition memory test in which participants view photographs of 12 doors for 3 seconds each. Immediately thereafter, participants are presented with 12 arrays of 4 doors each, and are asked to identify the door from the previous list. In the second part new photographs of doors are displayed, but the recognition stimuli are rather similar to the key list.

The Continuous Performance Test—Identical Pairs (CPT-IP) is a computerized measure of sustained, focused attention or vigilance. This version is a part of MCCB-Matrices Consensus Cognitive Battery and involves monitoring a series of multiple digits and responding with a button press each time that 2 stimuli in a row are identical.³⁰

Biological Measures. High sensitivity CRP (hs-CRP) was measured with an assay using nephelometry (Dade Behring). Abnormal CRP level was defined as >3 mg/L according to the ("The Emerging Risk Factors Collaboration"; 2010). Patients with hs-CRP levels >30 mg/L, which corresponds to an acute inflammation, were not included in the analyses ("The Emerging Risk Factors Collaboration"; 2010).

Ethical Concerns

The study was carried out in accordance with ethical principles for medical research involving humans (WMA, Declaration of Helsinki). The assessment protocol was approved by the relevant ethical review board (CPP-Ile de France IX, January 18, 2010). All data were collected anonymously. As this study include data coming from regular care assessments, a non-opposition form was signed by all participants.

Statistical Analysis

Sociodemographics, clinical and neuropsychological characteristics and treatments are presented using measures of means and dispersion (SD) for continuous data and frequency distribution for categorical variables. The data were examined for normal distribution with the Shapiro-Wilk test and for homogeneity of variance with the Levene test. Univariate associations between demographic, neuropsychological and clinical characteristics of patients with abnormal CRP levels were performed using the chi-square test for categorical variables. Continuous variables were analyzed with Student *t*-tests for normally distributed data and Mann-Whitney tests in case of non-normal distributions. Effect size—defined as the mean difference divided by the SD of the whole sample—were computed for each cognitive tests.³¹ According to Samsa et al,³² an effect size of at least 0.2 is considered as clinically relevant. An effect size of 0.2 is considered a small effect, 0.5 a moderate effect, and 0.8 a large effect.³³

A multivariate logistic regression was performed to estimate the adjusted Odds Ratio (aOR) and its corresponding 95% CI for an association between cognition and CRP. In order to eliminate multicollinearity associated with cognitive tests, we performed a principal component analysis (PCA) with varimax rotation to reduce the large number of cognitive tests to a smaller number of uncorrelated cognitive component scores. The number of components was chosen based on the Kaiser stopping criterion (ie, all components with eigenvalues greater than 1) and the scree test. The use of component scores as the independent variables in multivariate models is considered as relevant in the case of multicollinearity.³⁴ The cognitive scores were thus used for multiple regression analysis, associated to a set of confounding factors selected from the univariate analysis, with selection based on a threshold *P*-value < .20 (ie, age, years of education, illness duration). Sex and PANSS score were also included because of their sociodemographic and clinical interest. CRP levels were also analyzed in a quantitative manner using a multiple linear regression to check the robustness of our findings.

All of the tests were 2-sided. Statistical significance was defined as *P* < .05. Statistical analysis was performed using the SPSS version 18.0 software package (SPSS Inc). This study was a confirmatory analysis. The hypothesis was that abnormal CRP levels were associated with cognitive impairment, based on the previous preliminary data.³⁵ No correction for multiple testing has been therefore carried out, consistently with recommendations.³⁶

Results

A sample of 369 community-dwelling stable SZ subjects enrolled in FACE-SZ cohort was included in this study. Table 1 shows demographical, clinical and neuropsychological characteristics of the sample, as well as associations with abnormal CRP levels. The majority

Table 1. Association Between Peripheral Inflammation (Defined by Abnormal C-Reactive Protein [CRP] Level [>3 mg/L]) and Neuropsychological Impairment ($N = 369$)

	Whole Sample ($N = 369$)	Abnormal CRP			Statistic	P			
		Abnormal CRP		Statistic					
		No ($N = 265$)	Yes ($N = 104$)						
Sociodemographic characteristics									
Sex (male)	281	76.2%	203	76.6%	78	75.0%	Chi-square	.745	
Mean age (y) (SD)	32.72	10.03	32.29	10.00	33.80	10.08	Student <i>t</i>	.195	
Mean years of education (SD)	12.37	2.86	12.49	2.85	12.08	2.90	U/Mann-Whitney	.130	
Clinical variables									
Age at onset, mean (SD)	21.68	6.44	21.52	6.12	22.07	7.21	Student <i>t</i>	.476	
Age at first antipsychotic treatment, mean (SD)	22.89	6.54	22.74	6.71	23.23	6.12	Student <i>t</i>	.531	
Illness duration (y), mean (SD)	11.01	8.61	10.78	8.60	11.64	8.64	U/Mann-Whitney	.194	
Psychotic symptomatology (PANSS total score), mean (SD)	70.84	18.05	70.97	18.60	70.50	16.66	Student <i>t</i>	.822	
Positive symptoms (PANSS positive score), mean (SD)	14.69	5.33	14.73	5.39	14.60	5.21	U/Mann-Whitney	.939	
Negative symptoms (PANSS negative score), mean (SD)	20.96	7.09	21.11	7.31	20.60	6.54	Student <i>t</i>	.539	
General psychopathology (PANSS general score), mean (SD)	35.14	9.82	35.13	10.10	35.17	9.14	Student <i>t</i>	.972	
Depressive symptoms (CDSS score), mean (SD)	3.97	4.29	3.97	4.21	3.97	4.48	U/Mann-Whitney	.807	
Manic symptoms (YMRS score), mean (SD)	2.73	4.31	2.87	4.49	2.38	3.81	U/Mann-Whitney	.253	
Current alcohol disorder, mean (SD)	16	4.3%	13	4.9%	3	2.9%	Chi-square	.391	
Current substance use disorder, mean (SD)	4	1.4%	3	1.4%	1	1.3%	Chi-square	.924	
Current daily tobacco smoking	192	53.0%	133	51.4%	59	53.0%	Chi-square	.308	
Current cannabis consumption disorder	23	19.2%	19	21.1%	4	13.3%	Chi-square	.349	
Treatment									
Second generation antipsychotics	266	88.7%	191	89.3%	75	87.2%	Chi-square	.614	
First generation antipsychotics	80	26.7%	53	24.8%	27	31.4%	Chi-square	.240	
Cholinergic drugs	50	16.7%	35	16.4%	15	17.4%	Chi-square	.819	
Abnormal CRP									
		Whole sample ($N = 369$)		No ($N = 265$)		Yes ($N = 104$)			
	Mean range ^a							Statistic	
Current and premorbid intellectual ability								Effect size ^b	
fNART based premorbid IQ, mean (SD)	[80–119]	104.61	8.45	105.01	8.37	103.50	8.63	Student <i>t</i>	0.18
Full Scale IQ, mean (SD)	[80–119]	86.17	15.38	87.72	15.64	82.45	14.15	Student <i>t</i>	0.34
Verbal IQ, mean (SD)	[80–119]	90.27	14.64	91.68	14.53	86.89	14.43	Student <i>t</i>	0.006
Performance IQ, mean (SD)	[80–119]	83.26	17.45	84.76	18.12	79.76	15.27	Student <i>t</i>	0.016
Working memory									
Digit Span (standard score), mean (SD)	[7–13]	8.16	2.79	8.37	2.81	7.67	2.69	Student <i>t</i>	0.033
Letter-Number Sequencing (standard score), mean (SD)	[7–13]	8.03	2.74	8.23	2.72	7.49	2.73	Student <i>t</i>	0.040
Arithmetic (standard score), mean (SD)	[7–13]	7.29	3.08	7.58	3.16	6.57	2.78	Student <i>t</i>	0.006
Working Memory Index (WMI), mean (SD)	[80–119]	87.08	14.88	88.59	14.94	82.91	13.97	Student <i>t</i>	0.004

Table 1. Continued

	Whole sample (N = 369)		Abnormal CRP		Statistic	Effect size ^b
	Mean range ^a	No (N = 265)	Yes (N = 104)			
Learning abilities, episodic and semantic memory						
CVLT short delay free recall, mean (SD)	19–15.5]	3.51	3.34	8.58	U Mann-Whitney	0.26
CVLT short delay cued recall, mean (SD)	[10.5–15.5]	3.27	3.29	9.71	U Mann-Whitney	0.09
CVLT long delay free recall, mean (SD)	[10–15.5]	3.43	3.32	9.44	U Mann-Whitney	0.13
CVLT long delay cued recall, mean (SD)	[11–15.5]	3.34	3.30	9.76	U Mann-Whitney	0.11
CVLT recognition, mean (SD)	[14–16]	3.13	3.25	14.38	U Mann-Whitney	0.098
Doors Test (A&B), mean (SD)	[16–22]	4.58	4.31	14.39	Student t	0.035
Information (standard score), mean (SD)	[7–13]	8.98	9.23	8.38	Student t	0.026
Executive functions and problem solving						
Trail Making Test B (time), mean (SD)	[40–81]	104.37	100.16	115.67	U Mann-Whitney	0.26
Trail Making Test B (errors), mean (SD)	[0]	0.69	1.32	0.68	U Mann-Whitney	0.01
Matrix Reasoning (standard score), mean (SD)	[7–13]	7.97	8.28	7.26	Student t	0.29
Similarities (standard score), mean (SD)	[7–13]	8.56	8.73	8.17	Student t	0.18
Visual attention and speed of processing						
Trail Making Test A (time), mean (SD)	[19–44]	42.55	41.15	46.03	U Mann-Whitney	0.25
Trail Making Test A (errors), mean (SD)	[0]	0.21	0.17	0.29	U Mann-Whitney	0.004
Digit Symbol Coding (standard score), mean (SD)	[7–13]	5.90	6.07	5.52	Student t	0.19
Picture Completion (standard score), mean (SD)	[7–13]	7.56	7.75	7.12	Student t	0.18
CPT-IP d-prime, mean (SD)	[2.25–4]	2.31	2.34	2.20	Student t	0.199

Note: Statistically significant in bold ($P < .05$). PANSS, Positive and Negative Syndrome Scale; YMRS, Young Mania Rating Scale; CPT-IP, Continuous Performance Test—Identical Pairs; CDSS, Calgary Depression rating Scale for Schizophrenia; FNART, French National Adult Reading Test; CVLT, California Verbal Learning Test.

^aThe provided scores are issued from the normative data used for clinical interpretation of the neuropsychological tests. The scores provided correspond the mean age and education of the clinical sample and are therefore reported here for informative purpose only. The normal range is defined following usual conventions: Z scores ranging from -1.33 to $+1.33$; Percentiles ranging from 10 to 90, Standard scores ranging from 7 to 13 and IQ scores ranging from 80 to 119.

^bEffect size—defined as the mean difference divided by the SD of the whole sample—were computed for each cognitive tests.³² According to Samsa et al.,³³ an effect size of at least 0.2 is considered as clinically relevant. An effect size of 0.2 is considered a small effect, 0.5 a moderate effect and 0.8 a large effect.³⁴

of the subjects in the sample (76.2%) were men and the mean age of the patients was 32.7 ± 10 years. Overall, 104 patients (28.2%) had abnormal CRP levels. The mean age at SZ onset was 21.7 ± 6.4 years, the mean duration of illness was 11 ± 8.6 years and the mean PANSS total score was 70.8 ± 18 . Current tobacco smoking, current cannabis consumption, current alcohol disorder and current substance abuse disorders were reported in table 1. None of these variables was significantly associated with abnormal CRP levels (all $P > .05$).

The impaired cognitive functions associated with abnormal CRP levels are presented in table 1. The following cognitive functions were found to be significantly impaired in SZ subjects with abnormal CRP levels compared to those without abnormal CRP levels: current intellectual ability (FSIQ, effect size [ES] = 0.34, $P = .004$; VIQ, ES = 0.33, $P = .006$; PIQ, ES = 0.29, $P = .016$), working memory (all measures, ES = 0.25–0.38, $P < .05$), memory and learning abilities (short delay free verbal recall, ES = 0.26, $P = .026$; visual recognition, ES = 0.28, $P = .035$), semantic memory (Information subtest, ES = 0.26, $P = .026$), mental flexibility (TMT B, time, ES = 0.26, $P = .044$), abstract thinking (Matrix reasoning subtest, ES = 0.29, $P = .016$), visual attention (TMT A, errors, ES = 0.23, $P = .004$) and speed of processing (TMT A, time, ES = 0.25, $P = .043$). These results were independent of sociodemographic variables, illness duration and severity (psychotic symptomatology) in multivariate analysis.

Four components emerged in the PCA. All 4 components had eigenvalues >1 and in total they accounted for 67.03 % of the variance (table 2). The content of each component was considered as relevant and meaningful by 2 experts in psychology (E.B. and G.F.) and

named: “General intellectual ability/ Abstract thinking,” “Working memory/Attention,” “Processing speed,” and “Memory and learning.” In the multivariate analyses reported in table 3, the relationship between factor 1 (“General intellectual ability/ Abstract thinking”) score and CRP level remained significant after adjusting for sociodemographic and clinical variables. The component scores are presented in patients with normal and abnormal CRP levels in figure 1. These findings were confirmed using CRP in a quantitative manner. The component 1 remained significant (standardized $\beta = -.20$; $P = .04$).

Discussion

The major results of this study may be summarized as follows: peripheral sub-inflammation, assessed here by abnormal CRP levels, is associated with a wide array of small to moderate cognitive impairments in a large sample of community-dwelling SZ patients, independently of SZ characteristics, sociodemographic variables, treatments, tobacco smoking and cannabis consumption. Multivariate analysis revealed that General Intellectual Ability/Abstract Reasoning was the most specifically impaired cognitive factor associated with abnormal CRP levels, being independent of age, sex, education level, psychotic symptomatology, treatments, and addiction comorbidities. Elevated CRP levels were also associated with an impairment of all components of working memory and a wide range of other impaired cognitive functions including memory and learning abilities, as well as semantic memory, mental flexibility, visual attention and speed of processing. Firstly, the data presented here indicate peripheral inflammation to be associated with a global impairment of current intellectual ability

Table 2. Principal Component Analysis After Varimax Rotation

	Component 1, General Intellectual Ability/ Abstract Thinking	Component 2, Working Memory/ Attention	Component 3, Processing Speed	Component 4, Memory and Learning
Information (standard score)	0.876			
Full Scale IQ	0.766			
Similarities (standard score)	0.638			
fNART IQ	0.632			
Picture Completion (standard score)	0.552			
Matrix Reasoning (standard score)	0.480			
Digit Span (standard score)		0.772		
Letter-Number Sequencing (standard score)		0.737		
CPT-IP mean d-prime		0.681		
Arithmetic (standard score)		0.620		
Trail Making Test A (time)			-0.808	
Trail Making Test B (time)			-0.761	
Digit Symbol Coding (standard score)			0.560	
CVLT recognition				0.812
CVLT short delay free recall				0.797
Doors Test (A&B)				0.454

Downloaded from https://academic.oup.com/schizophreniabulletin/article/42/5/1290/24139110 by guest on 20 August 2022

Table 3. Factors Associated With High CRP: Multivariate Analysis

Factors	Adjusted Odds Ratio	95% CI	P-value
Sex (male)	1.213	0.473–3.110	.756
Component 1: “General intellectual ability / Abstract thinking”	0.560	0.347–0.904	.014
Component 2: “Working memory/Attention”	0.841	0.565–1.251	.271
Component 3: “Processing speed”	0.721	0.471–1.104	.227
Component 4: “Memory and learning”	1.120	0.763–1.644	.516
PANSS (total score)	0.986	0.963–1.011	.212
Age (y)	1.024	0.961–1.091	.621
Illness duration (y)	1.002	0.936–1.073	.984
Years of education	0.962	0.813–1.140	.776

Note: Statistically significant in bold.




Fig. 1. Cognitive impairment associated with chronic peripheral inflammation in schizophrenia (measured by C-Reactive protein [CRP] blood levels ≥ 3 mg/L).

in SZ subjects. This suggests a major role for peripheral inflammation in the debilitating course of SZ, even in young patients (mean age was 32 y, with a mean illness duration of 11 y). Such data overlaps with data in older populations, where elevated CRP levels were associated with age-related cognitive decline (“inflammaging”),^{11,12} as well as with all-cause dementia.¹³

The cognitive impairment following the onset of SZ has been well documented (see recent meta-analyses).^{37–40} This life span neurodegenerative perspective is not new as SZ was initially termed “dementia praecox” in 1891 by the German psychiatrist Arnold Pick (1851–1924), due to the cognitive decline during the first 2 decades of the illness. From a neuroanatomical point of view, abnormal CRP is known to be associated with dysexecutive syndrome and a cerebral microstructure disintegration in frontal pathways,^{41,42} a crucial cortical area in SZ pathophysiology.^{43–45} CRP is one of the triggering molecules of peripheral inflammation that may also activate brain inflammation. A growing body of literature has further underscored the potential role of microglial activation in SZ, including an infection-triggered immune reaction.^{46,47} Altogether, our results combined with literature suggest

that abnormal CRP may be considered as a peripheral biomarker of cognitive impairment in SZ.

As noted, the association between inflammation and cognition is not limited to SZ, however the data presented here indicate some specificity. Inconsistent results have been found in depression and bipolar disorder, as to inflammation and cognitive decline. Lower cognitive functioning in individuals with bipolar disorder has been associated with elevated levels of CRP.⁸ The authors found an inverse relationship between CRP level and performance on RBANS battery, which was significant on immediate memory, attention and language factors, as well as psychomotor speed. In major depression, except psychomotor speed and some executive components, most of the neuropsychological variables (verbal and visual memory, working memory, attention, language) were not associated with higher CRP levels.^{48,49} In summary, CRP is associated with lowered psychomotor speed in a non-specific (transnosographic) manner. But our results further suggest that CRP-associated impaired general intellectual ability, abstract reasoning, working memory, semantic memory and mental flexibility is a more diagnosis-specific aspect of peripheral inflammation in SZ. It

may be hypothesized that these discrepancies may be due to several factors: genetic brain vulnerabilities, origin of chronic inflammation (infections, perivisceral fat, antidepressant consumption⁵⁰), different immunological disturbances across illnesses. Further studies should determine if cognition is differently impacted according to inflammation etiological factors, and if these factors may explain the differences between SZ and bipolar disorders.

Second, peripheral inflammation was found in our study to be associated with a global impairment of all assessed working memory components. This association was already described in other non-psychiatric populations (including cancer, coronary heart disease and lupus patients, as well as middle-age healthy subjects),⁵¹⁻⁵⁴ but not in SZ. This suggests a deleterious effect of high CRP levels on working memory across illnesses. The role of working memory is crucial to temporarily maintain and store information. This system supports thought mechanisms by providing an interface between perception, long-term memory and action.⁵⁵ All working memory measures were significantly lower in patients with higher levels of CRP. Targeting CRP may thus be decisive for improving working memory in SZ subjects.

Thirdly, elevated CRP was associated with impaired verbal and visual episodic learning abilities, as well as impaired semantic memory. This is consistent with previous findings in non-SZ populations, where healthy-aging subjects with higher CRP had lower memory performance and smaller left temporal lobe volumes⁵⁶ as well as impaired frontal⁴² and hippocampal functioning.⁵⁷ The last 2 areas are essential for memory formation and its biological underpinnings that are thought to arise from long-term potentiation. CRP-associated impaired memory was also described in type II diabetes patients.⁵⁸ However, no association was observed between abnormal CRP and long-term verbal recall impairment in SZ. Memory impairment related to inflammatory processes is less clear in depression. In 1 study, no association has been found between memory functions and CRP in moderately depressed elderly.⁵⁹ Instead, encoding and recall were inversely associated with interleukin (IL)-6 across the groups. While the results in bipolar patients⁸ are comparable to those in this study, this was inconsistent with previous findings in healthy aging population.^{60,61} Several explanations may be suggested to explain these discrepancies, including variations in the delay of recall across studies. Also, the wide range of cognitive impairment in SZ may have erased the CRP-long-term-verbal memory association, which was found in healthy-aging subjects. It may be hypothesized that SZ subjects with normal CRP levels may have decreased verbal memory due to other causes.

Fourthly, peripheral inflammation was found here to be associated with lower performance in inductive reasoning and mental flexibility. Along with working memory and abstract thinking, these functions form a broad

executive control and problem solving system that is necessary to adapt to novel contexts, as well as to initiate and regulate autonomous behaviors. Executive functions correlate with the ability to solve interpersonal problems,⁶² vocational functioning, daily management of personal and professional activities,^{63,64} and quality of life.⁶⁶ Our results are consistent with those found in middle-aged healthy subjects,⁵³ patients with diabetes,⁵⁸ and healthy aging.^{42,66-68} In patients suffering from depression, with elevated CRP levels, there seems to be a tendency for poorer executive function as indicated on the Wisconsin card sorting test measures: number of completed categories,⁴⁸ and design fluency.⁴⁹

Fifthly, peripheral inflammation was associated with impaired components of visual attention and processing speed. Similarly, increased levels of hsCRP were associated with lower psychomotor speed (Trail making A, Finger Tapping test) in depressed outpatients.^{48,49} This association is also observed in nondepressed subjects with higher CRP levels related to arterial disease.⁵² A potential mediator, as discussed might be the cytokine-induced hypermetabolism in the basal ganglia leading to a psychomotor slowing.

Our major results were inconsistent with the Joseph et al study,¹⁶ and consistent with Dickerson et al.¹⁵ This is probably due to the higher mean age of SZ subjects (49 y, mean illness duration 23 y) in Joseph et al study vs 40 years (mean illness duration 19 y) and 32 years (mean illness duration 11 y), respectively in Dickerson et al study and ours. Overall, these studies indicate that CRP mostly impacts cognitive functioning in the first 2 decades following SZ onset. Future longitudinal studies are required to investigate this hypothesis.

The absence of association between abnormal CRP levels and psychotic symptoms in our sample was consistent with Dickerson et al¹⁵ results. However a recent meta-analysis suggested an association between abnormal CRP and positive symptoms, which may be due to the inclusion of untreated patients in this meta-analysis.^{1,69,70} In our sample, all patients were treated by antipsychotics, which possibly contributes to the non-replication of this association in the data presented here.

Contrary to Dickerson et al and Joseph et al studies, we found no sex differences. Our sex ratio (76% male) was comparable to Dickerson et al¹⁵ (70%), although not with Joseph et al¹⁶ (39%). The mean age at illness onset was also comparable (respectively 21 y here and in the Dickerson et al study, vs 26 y for Joseph et al) possibly due to the later onset in SZ female patients. Future meta-analyses should explore such sex differences to determine if CRP levels are significantly altered in SZ females compared to males.

The neurobiological underpinnings of CRP-associated cognitive impairment in SZ are not fully understood to date. Inflammation in the central nervous system is closely related to neurodegeneration. Blood CRP levels may be

seen as an indirect peripheral common marker of central neuroinflammation. CRP levels are raised by the increase of pro-inflammatory cytokines, including IL-1 β , TNF- α and IL-6. These pro-inflammatory cytokines have individual effects on neurogenesis. IL-1 β induces focal and sustained hippocampal inflammation, resulting in severe depletion of developing neuroblasts.⁷¹ IL-1 β was also found to suppress cell proliferation in the dentate gyrus.⁷² IL-1 β generally contributes to memory formation at the physiological level. However, when IL-1 β is expressed excessively, it inhibits hippocampus-mediated memory formation in a manner of an inverted U shape.⁷³ Activated microglia-derived TNF- α was found to enhance the death of hippocampal progenitor cells.⁷⁴ IL-6 may be the most important cytokine involved in microglial activity and inflammatory responses. Overexpressed IL-6 inhibits hippocampal neurogenesis.⁷⁵ All these mechanisms may be involved in CRP-associated cognitive impairment.

Elevation in CRP levels may have direct therapeutic relevance to the prevention of cognitive impairment, within a personalised approach to the management of SZ. A recent meta-analysis highlighted the clinical utility of anti-inflammatory add-on therapies in SZ, perhaps especially aspirin.⁷⁶ These interventions are described as potentially more effective early in the course of SZ, reinforcing the need to develop an early prevention-focused perspective, perhaps especially in patients with elevated CRP levels.⁷⁷ CRP is also associated with both increased atheroma genesis⁷⁸ and increased risk of stroke.¹⁴ As the prevalence of metabolic syndrome is high in young SZ subjects,⁷⁹ it may be reasonably suggested that atheroma may contribute to the mechanisms of cognitive impairment in SZ.⁸⁰ Aspirin may therefore be doubly helpful, by its anti-inflammatory and anti-atheroma properties.⁸¹ Future studies should assess the pro-cognitive effects of anti-inflammatory agents that have been proposed as beneficial in aged populations.^{82,83} Beyond anti-inflammatory agents, an “anti-inflammatory diet” was also suggested to improve cognitive function in healthy participants and should be evaluated in SZ patients with both inflammatory disturbances and impaired cognitive functions.^{82–84} The impact of lifestyle hygiene factors (such as psychological stress management or physical activity) on the reduction of inflammatory response should also be taken into consideration.^{85,86} In summary, abnormal CRP levels in SZ may be a biomarker for cognitive dysfunction and therefore an indication for anti-inflammatory strategies as well as early cognitive remediation. Future studies are needed to investigate the effectiveness of anti-inflammatory add-on therapies for cognitive improvement in SZ patients with peripheral inflammation.

The purpose of the present study was not to determine the etiological factors of chronic peripheral inflammation in SZ. However, unraveling etiological factors may help guiding anti-inflammatory and preventive strategies for the future. For example, abnormal CRP levels have been independently associated with both antidepressant

consumption and abdominal obesity in SZ, but not with tobacco or cannabis consumption.⁸⁷ The lack of association between tobacco smoking or cannabis consumption in the present study is consistent with previous results.⁸⁸ Future studies should determine if different etiological factors of peripheral inflammation may be associated with different cognitive outcomes.

Strengths

The present study has clear strengths, particularly the use of homogenous and exhaustive standardized diagnostic protocols and neuropsychological assessment in a large national multicentric study. Important confounding factors were taken into account in our analyses, especially sociodemographic characteristics, antipsychotic treatments, tobacco smoking and cannabis consumption. Our sample was relatively young (mean age 32 y, mean illness duration 11 y).

Limits

Due to the cross-sectional design of this study, no longitudinal data of abnormal CRP on cognitive trajectory outcome were obtained. First, an important methodological problem is the definition of our groups (normal CRP level and high CRP). Indeed, there are no generally accepted criteria. However, we have chosen the most consensual definition in the recent scientific literature, which is recognized as the cut-off point for high cardiovascular risk.^{89–91} Second, CRP was the sole marker of inflammation in this study. Although CRP is strongly associated with IL-6 activity, we did not directly assess any pro-inflammatory cytokines. Future studies with extensive assessment of inflammatory markers may be required. Third, we have only collected CRP at one time point, and repeated testing has been recommended to confirm elevated plasma levels⁸⁹ especially as concentrations can be affected by acute infection. However, patients with acute infections were removed and were not included in this study. Finally, although the experimental protocol takes into account a large set of potentially confounding variables, additional factors might have been interesting to consider. In particular, cognitive impairment, a core symptom of SZ, is associated with metabolic syndrome⁹² and more specifically with perivisceral fat.⁷⁹ Alcohol consumption or substance abuse have not been described as a risk factor for peripheral inflammation in SZ to date, however this data may have been explored in the present study as a confounding factor. The purpose of the present study was not to determine the potential etiological factors associated with abnormal CRP levels, which has been done elsewhere⁸⁷ and would require larger samples.

Conclusion

Altogether, our results yielded important findings. We confirmed that peripheral inflammation was associated

with a general loss of intellectual abilities in SZ, which strongly impacts on functional outcomes, and is independent of psychotic symptomatology, age, sex, treatments and addiction co-morbidities. Therapeutic trials are needed to determine if anti-inflammatory add-on strategies are effective in preventing cognitive impairment in SZ, especially at the beginning of the illness.

Funding

This work was funded by Assistance Publique des Hôpitaux de Paris (AP-HP), Fondation FondaMental (RTRS Santé Mentale), by the Investissements d'Avenir program managed by the ANR under reference ANR-11-IDEX-0004-02 and ANR-10-COHO-10-01, and by Institut National de la Santé et de la Recherche Médicale (INSERM).

Acknowledgments

FACE-SCZ Group: Méja Andrianarisoa^{a,b}, Bruno Aouizerate^{b,c}, Fabrice Berna^{b,d}, Olivier Blanc^{b,e}, Lore Brunel^{a,b}, Ewa Bulzacka^{a,b}, Delphine Capdevielle^{b,f}, Isabelle Chereau-Boudet^{b,e}, Gabrielle Chesnoy-Servanin^{b,g}, Jean-Marie Danion^{b,d}, Thierry D'Amato^{b,g}, Deloge Arnaud^{b,h}, Claire Delorme^{b,i}, Hélène Denizot^{b,e}, Jean-Michel Dorey^{b,g}, Caroline Dubertret^{b,j}, Julien Dubreucq^{b,i}, Catherine Faget^{b,k}, Cécile Fluttaz^{b,i}, Guillaume Fond^{a,b}, Sandrine Fonteneau^{b,l}, Franck Gabayet^{b,i}, Elisabeth Giraud-Baro^{b,i}, Marie-Christine Hardy-Baylé^{b,l}, Delphine Lacelle^{b,e}, Christophe Lançon^{b,k}, Hakim Laouamri^b, Marion Leboyer^{a,b}, Tifenn Le Gloahec^{a,b}, Yann Le Strat^{b,j}, Pierre-Michel Llorca^{b,e}, Jasmina Mallet^{j,k}, Emeline Metairie^{b,k}, David Misdrahi^{b,h}, Isabelle Offerlin-Meyer^{b,d}, Christine Passerieux^{b,l}, Pauline Peri^{b,k}, Sylvie Pires^{b,e}, Céline Portalier^{b,j}, Romain Rey^{b,g}, Céline Roman^{b,i}, Mathilde Sebilliau^{b,l}, Aurélie Schandrin^{b,f}, Franck Schurhoff^{a,b}, Arnaud Tessier^{b,h}, Anne-Marie Tronche^{b,e}, Mathieu Urbach^{b,l}, Florence Vaillant^{b,k}, Aurélie Vehier^{b,g}, Pierre Vidailhet^{b,d}, Estelle Vilà^{b,h}, Hanan Yazbek^{b,f}, and Anna Zinetti-Bertschy^{b,d}. *Affiliations:* ^aAP-HP, DHU Pe-PSY, Pôle de Psychiatrie et d'addictologie des Hôpitaux Universitaires H Mondor, INSERM U955, Eq 15 Psychiatrie Translationnelle, Université Paris Est-Créteil, Créteil, France; ^bFondation Fondamental; ^cCentre Hospitalier Charles Perrens, F-33076 Bordeaux, France; Université de Bordeaux, Inserm, Neurocentre Magendie, Physiopathologie de la Plasticité Neuronale, U862, F-33000 Bordeaux, France; ^dHôpitaux Universitaires de Strasbourg, Université de Strasbourg, INSERM U1114, Fédération de Médecine Translationnelle de Strasbourg, Strasbourg, France; ^eCMP B, CHU, EA 7280 Faculté de Médecine, Université d'Auvergne, BP 69 63003 Clermont-Ferrand

Cedex 1, France; ^fService Universitaire de Psychiatrie Adulte, Hôpital la Colombière, CHRU Montpellier, Université Montpellier 1, Inserm 1061, Montpellier, France; ^gUniversité Claude Bernard Lyon 1/Centre Hospitalier Le Vinatier Pole Est BP 300 39 - 95 bd Pinel - 69678 BRON Cedex, France; ^hCentre Hospitalier Charles Perrens, F-33076 Bordeaux, France; Université de Bordeaux, CNRS UMR 5287-INCIA; ⁱCentre Référent de Réhabilitation Psychosociale, CH Alpes Isère, Grenoble, France; ^jAP-HP, Department of Psychiatry, Louis Mourier Hospital, Colombes, Inserm U894 Université Paris Diderot, Sorbonne Paris Cité, Faculté de médecine, France; ^kAssistance Publique des Hôpitaux de Marseille (AP-HM), pôle universitaire de psychiatrie, Marseille, France; ^lService de psychiatrie d'adulte, Centre Hospitalier de Versailles, Le Chesnay, EA 4047 HANDIRESP, UFR des Sciences de la Santé Simone Veil, Université Versailles Saint-Quentin en Yvelines, Versailles, France. We express all our thanks to the nurses, and to the patients who were included in the present study. We thank Hakim Laouamri, and his team (Stéphane Beaufort, Seif Ben Salem, Karmène Souyris, Victor Barteau, and Mohamed Laaidi) for the development of the FACE-SZ computer interface, data management, quality control, and regulatory aspects. The authors have declared that there are no conflicts of interest in relation to the subject of this study.

References

1. Fernandes BS, Steiner J, Bernstein H-G, et al. C-reactive protein is increased in schizophrenia but is not altered by antipsychotics: meta-analysis and implications. *Mol Psychiatry*. 2015;21:554–564. doi:10.1038/mp.2015.87.
2. Fond G, d'Albis M-A, Jamain S, et al. The promise of biological markers for treatment response in first-episode psychosis: a systematic review. *Schizophr Bull*. 2015;41:559–573.
3. Lopresti AL, Maker GL, Hood SD, Drummond PD. A review of peripheral biomarkers in major depression: the potential of inflammatory and oxidative stress biomarkers. *Prog Neuropsychopharmacol Biol Psychiatry*. 2014;48:102–111.
4. Miller BJ, Culppepper N, Rapaport MH. C-reactive protein levels in schizophrenia. *Clin Schizophr Relat Psychoses*. 2014;7:223–230.
5. Dickerson F. Elevated C reactive protein in adults predicts the later development of late-onset or very-late-onset schizophrenia. *Evid Based Ment Health*. 2015;18:e2–e2.
6. Strawbridge R, Arnone D, Danese A, Papadopoulos A, Vives AH, Cleare AJ. Inflammation and clinical response to treatment in depression: a meta-analysis. *Eur Neuropsychopharmacol*. 2015;25:1532–1543.
7. Valkanova V, Ebmeier KP, Allan CL. CRP, IL-6 and depression: a systematic review and meta-analysis of longitudinal studies. *J Affect Disord*. 2013;150:736–744.
8. Dickerson F, Stallings C, Origoni A, Vaughan C, Khushalani S, Yolken R. Elevated C-reactive protein and cognitive deficits in individuals with bipolar disorder. *J Affect Disord*. 2013;150:456–459.

9. Elvevag B, Goldberg TE. Cognitive impairment in schizophrenia is the core of the disorder. *Crit Rev Neurobiol*. 2000;14:21. doi:10.1615/CritRevNeurobiol.v14.i1.10.
10. Zipursky RB, Reilly TJ, Murray RM. The myth of schizophrenia as a progressive brain disease. *Schizophr Bull*. 2013;39:1363–1372.
11. Laurin D, Curb JD, Masaki KH, White LR, Launer LJ. Midlife C-reactive protein and risk of cognitive decline: a 31-year follow-up. *Neurobiol Aging*. 2009;30:1724–1727.
12. Mancinella A, Mancinella M, Carpinteri G, et al. Is there a relationship between high C-reactive protein (CRP) levels and dementia? *Arch Gerontol Geriatr*. 2009;49:185–194.
13. Kravitz BA, Corrada MM, Kawas CH. Elevated C-reactive protein levels are associated with prevalent dementia in the oldest-old. *Alzheimers Dement*. 2009;5:318–323.
14. Kuo H-K, Yen C-J, Chang C-H, Kuo C-K, Chen J-H, Sorond F. Relation of C-reactive protein to stroke, cognitive disorders, and depression in the general population: systematic review and meta-analysis. *Lancet Neurol*. 2005;4:371–380.
15. Dickerson F, Stallings C, Origoni A, Boronow J, Yolken R. C-reactive protein is associated with the severity of cognitive impairment but not of psychiatric symptoms in individuals with schizophrenia. *Schizophr Res*. 2007;93:261–265.
16. Joseph J, Depp C, Martin AS, et al. Associations of high sensitivity C-reactive protein levels in schizophrenia and comparison groups. *Schizophr Res*. 2015;168:456–460.
17. Schürhoff F, Fond G, Berna F, et al. A National network of schizophrenia expert centres: an innovative tool to bridge the research-practice gap. *Eur Psychiatry*. 2015;30: 728–735.
18. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull*. 1987;13:261–276.
19. Addington D, Addington J, Maticka-Tyndale E, Joyce J. Reliability and validity of a depression rating scale for schizophrenics. *Schizophr Res*. 1992;6:201–208.
20. Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. *Br J Psychiatry*. 1978;133:429–435.
21. Nelson HE, O'Connell A. Dementia: the estimation of premorbid intelligence levels using the New Adult Reading Test. *Cortex J Devoted Study Nerv Syst Behav*. 1978;14:234–244.
22. Mackinnon A, Mulligan R. [The estimation of premorbid intelligence levels in French speakers]. *L'Encéphale*. 2005;31:31–43.
23. Wechsler D, Coalson DL, et Raiford SE. *WAIS-IV: Wechsler adult intelligence scale*. San Antonio, TX: Pearson; 2008.
24. Ryan JJ, Ward LC. Validity, reliability, and standard errors of measurement for two seven-subtest short forms of the Wechsler Adult Intelligence Scale—III. *Psychol Assess*. 1999;11:207.
25. Reitan RM. Validity of the Trail Making Test as an indicator of organic brain damage. *Percept Mot Skills*. 1958;8:271–276.
26. Godefroy O. Fonctions exécutives et pathologies neurologiques et psychiatriques: évaluation en pratique clinique. *Groupe de Boeck*. 2008.
27. Delis DC, Kramer JH, Kaplan E, Ober BA. *California Verbal Learning Test*. San Antonio, TX: The Psychological Corporation. Harcourt Assessment Company; 2000.
28. Poitrenaud J, Deweer B, Kalafat M, Van der Linden M. Adaptation en langue française du California Verbal Learning Test. *Paris Éditions Cent Psychol Appliquée*. 2007.
29. Baddeley AD, Emslie H, Nimmo-Smith I. Doors and people: a test of visual and verbal recall and recognition. *Thames Valley Test Company*. 1994.
30. Nuechterlein KH, Green MF, Kern RS, et al. The MATRICS Consensus Cognitive Battery, part I: test selection, reliability, and validity. *Am J Psychiatry*. 2008;165:203–213.
31. Thalheimer W, Cook S. How to calculate effect sizes from published research: a simplified methodology. *Work-Learn Res*. 2002;1–9.
32. Samsa G, Edelman D, Rothman ML, Williams GR, Lipscomb J, Matchar D. Determining clinically important differences in health status measures. *Pharmacoeconomics*. 1999;15:141–155.
33. Kazis LE, Anderson JJ, Meenan RF. Effect sizes for interpreting changes in health status. *Med Care*. 1989;27:S178–189.
34. Tabachnick BG, Fidell LS. *Using Multivariate Statistics*. Boston, MA: Allyn and Bacon Pearson Education Company; 2001.
35. Dickerson F, Stallings C, Origoni A, Boronow J, Yolken R. C-reactive protein is associated with the severity of cognitive impairment but not of psychiatric symptoms in individuals with schizophrenia. *Schizophr Res*. 2007;93:261–265.
36. Bender R, Lange S. Adjusting for multiple testing—when and how? *J Clin Epidemiol*. 2001;54:343–349.
37. Aas M, Dazzan P, Mondelli V, Melle I, Murray RM, Pariante CM. A systematic review of cognitive function in first-episode psychosis, including a discussion on childhood trauma, stress, and inflammation. *Front Psychiatry*. 2014;4:182.
38. Chung YS, Barch D, Strube M. A meta-analysis of mentalizing impairments in adults with schizophrenia and autism spectrum disorder. *Schizophr Bull*. 2014;40:602–616.
39. Fioravanti M, Bianchi V, Cinti ME. Cognitive deficits in schizophrenia: an updated metanalysis of the scientific evidence. *BMC Psychiatry*. 2012;12:64.
40. Schaefer J, Giangrande E, Weinberger DR, Dickinson D. The global cognitive impairment in schizophrenia: consistent over decades and around the world. *Schizophr Res*. 2013;150:42–50.
41. Najjar S, Pearlman DM. Neuroinflammation and white matter pathology in schizophrenia: systematic review. *Schizophr Res*. 2015;161:102–112.
42. Wersching H, Duning T, Lohmann H, et al. Serum C-reactive protein is linked to cerebral microstructural integrity and cognitive function. *Neurology*. 2010;74:1022–1029.
43. Fraguas D, Díaz-Caneja CM, Pina-Camacho L, Janssen J, Arango C. Progressive brain changes in children and adolescents with early-onset psychosis: a meta-analysis of longitudinal MRI studies. *Schizophr Res*. 2014.
44. Tse MT, Piantadosi PT, Floresco SB. Prefrontal cortical gamma-aminobutyric acid transmission and cognitive function: drawing links to schizophrenia from preclinical research. *Biol Psychiatry*. 2015;77:929–939.
45. Zhou Y, Ma X, Wang D, et al. The selective impairment of resting-state functional connectivity of the lateral sub-region of the frontal pole in schizophrenia. *PloS One*. 2015;10:e0119176.
46. Davis J, Moylan S, Harvey BH, Maes M, Berk M. Neuroprogression in schizophrenia: pathways underpinning clinical staging and therapeutic corollaries. *Aust N Z J Psychiatry*. 2014;48:512–529.
47. Najjar S, Pearlman DM. Neuroinflammation and white matter pathology in schizophrenia: systematic review. *Schizophr Res*. 2015;161:102–112.

48. Chang HH, Lee IH, Gean PW, et al. Treatment response and cognitive impairment in major depression: association with C-reactive protein. *Brain Behav Immun*. 2012;26:90–95.
49. Krogh J, Benros ME, Jørgensen MB, Vesterager L, Elfving B, Nordentoft M. The association between depressive symptoms, cognitive function, and inflammation in major depression. *Brain Behav Immun*. 2014;35:70–76.
50. Fond G, Godin O, Brunel L, et al. Peripheral sub-inflammation is associated with antidepressant consumption in schizophrenia. Results from the multi-center FACE-SZ data set. *J Affect Disord*. 2016;191:209–215.
51. Amidi A, Wu LM, Agerbaek M, et al. Cognitive impairment and potential biological and psychological correlates of neuropsychological performance in recently orchiectomized testicular cancer patients. *Psychooncology*. 2015;24:1174–1180.
52. Mangiafico RA, Sarnataro F, Mangiafico M, Fiore CE. Impaired cognitive performance in asymptomatic peripheral arterial disease: relation to C-reactive protein and D-dimer levels. *Age Ageing*. 2006;35:60–65.
53. Marsland AL, Gianaros PJ, Kuan DC-H, Sheu LK, Krajina K, Manuck SB. Brain morphology links systemic inflammation to cognitive function in midlife adults. *Brain Behav Immun*. 2015;48:195–204.
54. Shucard JL, Gaines JJ, Ambrus J, Shucard DW. C-reactive protein and cognitive deficits in systemic lupus erythematosus. *Cogn Behav Neurol Off J Soc Behav Cogn Neurol*. 2007;20:31–37.
55. Baddeley A. Working memory: looking back and looking forward. *Nat Rev Neurosci*. 2003;4:829–839.
56. Bettcher BM, Wilhelm R, Rigby T, et al. C-reactive protein is related to memory and medial temporal brain volume in older adults. *Brain Behav Immun*. 2012;26:103–108.
57. Semmler A, Okulla T, Sastre M, Dumitrescu-Ozimek L, Heneka MT. Systemic inflammation induces apoptosis with variable vulnerability of different brain regions. *J Chem Neuroanat*. 2005;30:144–157.
58. Akrivos J, Ravona-Springer R, Schmeidler J, et al. Glycemic control, inflammation, and cognitive function in older patients with type 2 diabetes. *Int J Geriatr Psychiatry*. 2015;30:1093–1100.
59. Elderkin-Thompson V, Irwin MR, Hellemann G, Kumar A. Interleukin-6 and memory functions of encoding and recall in healthy and depressed elderly adults. *Am J Geriatr Psychiatry*. 2012;20:753–763.
60. Noble JM, Manly JJ, Schupf N, Tang MX, Mayeux R, Luchsinger JA. Association of C-reactive protein with cognitive impairment. *Arch Neurol*. 2010;67:87–92.
61. Teunissen CE, Van Boxtel MPJ, Bosma H, et al. Inflammation markers in relation to cognition in a healthy aging population. *J Neuroimmunol*. 2003;134:142–150.
62. Green MF, Kern RS, Braff DL, Mintz J. Neurocognitive deficits and functional outcome in schizophrenia. *Schizophr Bull*. 2000;26:119–136.
63. Kessler RK, Giovannetti T, MacMullen LR. Everyday action in schizophrenia: performance patterns and underlying cognitive mechanisms. *Neuropsychology*. 2007;21:439.
64. Penadés R, Catalán R, Puig O, et al. Executive function needs to be targeted to improve social functioning with Cognitive Remediation Therapy (CRT) in schizophrenia. *Psychiatry Res*. 2010;177:41–45.
65. Tolman AW, Kurtz MM. Neurocognitive predictors of objective and subjective quality of life in individuals with schizophrenia: a meta-analytic investigation. *Schizophr Bull*. 2012;38:304–315.
66. Gunstad J, Bausserman L, Paul RH, et al. C-reactive protein, but not homocysteine, is related to cognitive dysfunction in older adults with cardiovascular disease. *J Clin Neurosci Off J Neurosurg Soc Australas*. 2006;13:540–546.
67. Heringa SM, van den Berg E, Reijmer YD, et al. Markers of low-grade inflammation and endothelial dysfunction are related to reduced information processing speed and executive functioning in an older population - the Hoorn Study. *Psychoneuroendocrinology*. 2014;40:108–118.
68. Schram MT, Euser SM, de Craen AJM, et al. Systemic markers of inflammation and cognitive decline in old age. *J Am Geriatr Soc*. 2007;55:708–716.
69. Faugere M, Micoulaud-Franchi JA, Alessandrini M, et al. Quality of life is associated with chronic inflammation in schizophrenia: a cross-sectional study. *Sci Rep*. 2015;5:10793. doi:10.1038/srep10793.
70. Micoulaud-Franchi J-A, Faugere M, Boyer L, et al. Elevated C-reactive protein is associated with sensory gating deficit in schizophrenia. *Schizophr Res*. 2015;165:94–96.
71. Wu MD, Hein AM, Moravan MJ, Shaftel SS, Olschowka JA, O'Banion MK. Adult murine hippocampal neurogenesis is inhibited by sustained IL-1 β and not rescued by voluntary running. *Brain Behav Immun*. 2012;26:292–300.
72. Koo JW, Duman RS. IL-1 β is an essential mediator of the antineurogenic and anhedonic effects of stress. *Proc Natl Acad Sci*. 2008;105:751–756.
73. Goshen I, Kreisel T, Ounallah-Saad H, et al. A dual role for interleukin-1 in hippocampal-dependent memory processes. *Psychoneuroendocrinology*. 2007;32:1106–1115.
74. Cacci E, Claasen J-H, Kokaia Z. Microglia-derived tumor necrosis factor- α exaggerates death of newborn hippocampal progenitor cells in vitro. *J Neurosci Res*. 2005;80:789–797.
75. Aravena O, Pesce B, Soto L, et al. Anti-TNF therapy in patients with rheumatoid arthritis decreases Th1 and Th17 cell populations and expands IFN- γ -producing NK cell and regulatory T cell subsets. *Immunobiology*. 2011;216:1256–1263.
76. Sommer IE, van Westrhenen R, Begemann MJH, de Witte LD, Leucht S, Kahn RS. Efficacy of anti-inflammatory agents to improve symptoms in patients with schizophrenia: an update. *Schizophr Bull*. 2013, sbt139.
77. Fond G, Hamdani N, Kapczynski F, et al. Effectiveness and tolerance of anti-inflammatory drugs' add-on therapy in major mental disorders: a systematic qualitative review. *Acta Psychiatr Scand*. 2014;129:163–179.
78. Schaub N, Reichlin T, Meune C, et al. Markers of plaque instability in the early diagnosis and risk stratification of acute myocardial infarction. *Clin Chem*. 2012;58:246–256.
79. Godin O, Leboyer M, Gaman A, et al. Metabolic syndrome, abdominal obesity and hyperuricemia in schizophrenia: results from the FACE-SZ cohort. *Schizophr Res*. 2015.
80. Vincenzi B, Stock S, Borba CPC, et al. A randomized placebo-controlled pilot study of pravastatin as an adjunctive therapy in schizophrenia patients: effect on inflammation, psychopathology, cognition and lipid metabolism. *Schizophr Res*. 2014;159:395–403.
81. Fond G, Girerd N, Clavel F, Tamouza R, Leboyer M. Recently discovered properties of aspirin may be doubly helpful in bipolar disorders. *Med Hypotheses*. 2014;82:640–641.
82. Kelley BJ, McClure LA, Unverzagt FW, et al. Regular aspirin use does not reduce risk of cognitive decline. *J Am Geriatr Soc*. 2015;63:390–392.

83. Wichmann MA, Cruickshanks KJ, Carlsson CM, et al. NSAID use and incident cognitive impairment in a population-based cohort. *Alzheimer Dis Assoc Disord*. 2015.
84. Nilsson A, Tovar J, Johansson M, Radeborg K, Björck I. A diet based on multiple functional concepts improves cognitive performance in healthy subjects. *Nutr Metab*. 2013;10:49.
85. Creswell JD, Irwin MR, Burklund LJ, et al. Mindfulness-Based Stress Reduction training reduces loneliness and pro-inflammatory gene expression in older adults: a small randomized controlled trial. *Brain Behav Immun*. 2012;26:1095–1101.
86. Wolf SA, Melnik A, Kempermann G. Physical exercise increases adult neurogenesis and telomerase activity, and improves behavioral deficits in a mouse model of schizophrenia. *Spec Issue Adapt Immun Cent Nerv Syst Funct*. 2011;25:971–980.
87. Fond G, Godin O, Llorca PM, Leboyer M. Abnormal C-reactive protein (CRP) levels in schizophrenia and schizoaffective disorders. Results from the FACE-SZ dataset. *Eur Psychiatry*. 2015;30:S112.
88. Barzilay R, Lobel T, Krivoy A, Shlosberg D, Weizman A, Katz N. Elevated C-reactive protein levels in schizophrenia inpatients is associated with aggressive behavior. *Eur Psychiatry*. 2016;31:8–12.
89. Pearson TA, Mensah GA, Alexander RW, et al. Markers of inflammation and cardiovascular disease application to clinical and public health practice: a statement for health-care professionals from the centers for disease control and prevention and the American Heart Association. *Circulation*. 2003;107:499–511.
90. Sicras-Mainar A, Rejas-Gutiérrez J, Navarro-Artieda R, Blanca-Tamayo M. C-reactive protein as a marker of cardiovascular disease in patients with a schizophrenia spectrum disorder treated in routine medical practice. *Eur Psychiatry*. 2013;28:161–167.
91. Wysokinski A, Margulska A, Strzelecki D, Kloszewska I. Levels of C-reactive protein (CRP) in patients with schizophrenia, unipolar depression and bipolar disorder. *Nord J Psychiatry*. 2015;69:346–353.
92. Boyer L, Richieri R, Dassa D, et al. Association of metabolic syndrome and inflammation with neurocognition in patients with schizophrenia. *Psychiatry Res*. 2013;210:381–386.