

Research article

Open Access

Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. by decreasing the DNA-binding activity of the quorum sensing response regulator LuxR

Gilles Brackman¹, Tom Defoirdt^{2,3}, Carol Miyamoto⁴, Peter Bossier³, Serge Van Calenbergh⁵, Hans Nelis¹ and Tom Coenye*¹

Address: ¹Laboratory of Pharmaceutical Microbiology, Ghent University, Harelbekestraat 72, B-9000 Ghent, Belgium, ²Laboratory of Microbial Ecology and Technology, Ghent University, Coupure Links 653, 9000 Ghent, Belgium, ³Laboratory of Aquaculture and Artemia Reference Center, Ghent University, Rozier 44, 9000 Ghent, Belgium, ⁴Department of Biochemistry, McGill University, McIntyre Medical Sciences Building, Room 818, 3655 Promenade Sir William Osler, Montreal, H3G 1Y6, Canada and ⁵Laboratory of Medicinal Chemistry, Ghent University, Harelbekestraat 72, B-9000 Ghent, Belgium

Email: Gilles Brackman - gilles.brackman@ugent.be; Tom Defoirdt - tom.defoirdt@ugent.be; Carol Miyamoto - carol.miyamoto@mcgill.ca; Peter Bossier - peter.bossier@ugent.be; Serge Van Calenbergh - serge.vancalenbergh@ugent.be; Hans Nelis - hans.nelis@ugent.be; Tom Coenye* - Tom.Coenye@UGent.be

* Corresponding author

Published: 16 September 2008

Received: 23 April 2008

BMC Microbiology 2008, 8:149 doi:10.1186/1471-2180-8-149

Accepted: 16 September 2008

This article is available from: <http://www.biomedcentral.com/1471-2180/8/149>

© 2008 Brackman et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: To date, only few compounds targeting the AI-2 based quorum sensing (QS) system are known. In the present study, we screened cinnamaldehyde and substituted cinnamaldehydes for their ability to interfere with AI-2 based QS. The mechanism of QS inhibition was elucidated by measuring the effect on bioluminescence in several *Vibrio harveyi* mutants. We also studied *in vitro* the ability of these compounds to interfere with biofilm formation, stress response and virulence of *Vibrio* spp. The compounds were also evaluated in an *in vivo* assay measuring the reduction of *Vibrio harveyi* virulence towards *Artemia* shrimp.

Results: Our results indicate that cinnamaldehyde and several substituted derivatives interfere with AI-2 based QS without inhibiting bacterial growth. The active compounds neither interfered with the bioluminescence system as such, nor with the production of AI-2. Study of the effect in various mutants suggested that the target protein is LuxR. Mobility shift assays revealed a decreased DNA-binding ability of LuxR. The compounds were further shown to (i) inhibit biofilm formation in several *Vibrio* spp., (ii) result in a reduced ability to survive starvation and antibiotic treatment, (iii) reduce pigment and protease production in *Vibrio anguillarum* and (iv) protect gnotobiotic *Artemia* shrimp against virulent *Vibrio harveyi* BB120.

Conclusion: Cinnamaldehyde and cinnamaldehyde derivatives interfere with AI-2 based QS in various *Vibrio* spp. by decreasing the DNA-binding ability of LuxR. The use of these compounds resulted in several marked phenotypic changes, including reduced virulence and increased susceptibility to stress. Since inhibitors of AI-2 based quorum sensing are rare, and considering the role of AI-2 in several processes these compounds may be useful leads towards antipathogenic drugs.

Background

Vibriosis, caused by *Vibrio* spp., is a major disease of marine fish and shellfish and is an important cause of economic loss in aquaculture [1,2]. Until recently prophylactic antibiotics were extensively used in aquaculture [3,4]. However, overuse of antibiotics resulted in increased rates of resistance so that novel approaches are required to manage vibriosis [5]. A possible novel target is the bacterial communication system. Bacteria monitor their population densities through the production and sensing of small signal molecules called autoinducers, a process entitled as quorum sensing (QS). To date three types of QS systems have been identified in *Vibrio* spp. [6]. Acylated homoserine lactones (AHL) are used as signal molecules in the LuxM/N QS system [7], while in the CqsA/S system, (S)-3-hydroxytridecan-4-one ("Cholera autoinducer 1", CAI-1) is used [8]. A third QS system appears to be shared by many Gram-positive and Gram-negative bacteria and is based on a mixture of interconvertible molecules collectively referred to as autoinducer-2 (AI-2) [9]. A key enzyme in the production of AI-2 is LuxS. LuxS catalyzes the cleavage of S-ribosylhomocysteine to homocysteine and 4,5-dihydroxy-2,3-pentanedione (DPD) [10]. DPD will subsequently undergo spontaneous derivatizations, forming a mixture of molecules, including (2R,4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran (R-THMF) and (2S,4S)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran-borate (S-THMF-borate) [11]. Although not all QS systems are present in all *Vibrio* spp., most of them contain the AI-2 based QS system [12]. In *Vibrio* spp. AI-2 binds to LuxP, a periplasmic AI-2 receptor that is associated with the LuxQ sensor kinase-phosphatase. At low population density only basal amounts of diffusible signal molecules are produced, and in this situation LuxQ will act as a kinase resulting in a phosphorylation of the response regulator LuxO through a cascade involving LuxU. Phosphorylation activates LuxO resulting in the production of small regulatory RNAs [13]. These small RNAs, together with the chaperone protein Hfq, will destabilize mRNA encoding the response regulator LuxR. However, when population density is sufficiently high, AI-2 will bind to LuxP and as a result LuxQ will act as a phosphatase, leading to a dephosphorylation of LuxO [14]. Since dephosphorylated LuxO is inactive, no small regulatory RNAs will be formed and the LuxR mRNA remains stable, resulting in the production of LuxR and ultimately an altered gene expression pattern. AI-2 based QS was found to play an important role in regulating the production of several virulence factors, biofilm formation and stress responses in several *Vibrio* spp. [15-17] and it was found to be associated with virulence as shown in several *in vivo* assays [18,19]. In contrast, in *Vibrio cholerae*, CAI-1 was found to be the principle signal molecule in virulence regulation [8]. Because of this involvement in virulence, inhibitors of AI-2 based QS have been proposed as

novel antipathogenic agents. While there is a growing interest in and evidence for the use of these antipathogenic substances to interfere with interspecies QS in the control of virulence and biofilm formation, only a few inhibitors of AI-2 based QS are known, including halogenated furanones and cinnamaldehyde [20-23]. Halogenated furanones have previously been shown to disrupt AHL and AI-2 based quorum sensing in *Vibrio* spp. by decreasing the DNA-binding activity of the response regulator LuxR [24-26]. Halogenated furanones can attenuate the virulence of several *Vibrio* spp. in gnotobiotic brine shrimp *Artemia franciscana* and their use results in a reversal of the negative effects of *Vibrio harveyi* BB120 towards rotifers [27,28]. Unfortunately, the toxicity of halogenated furanones towards both brine shrimp and rotifers limits their use. In contrast, cinnamaldehyde is a non-toxic synthetic flavouring substance that is widely used in food, beverages, chewing gum, and the perfume and food chemistry, and is generally recognised as safe [29]. Cinnamaldehyde concentrations in food range from 4 ppm to more than 300 ppm [30]. Although cinnamaldehyde is known to be a QS-inhibitor [21], its exact mechanism of action remains to be elucidated. The goal of the present study was to determine the mechanism of action of cinnamaldehyde and to evaluate its effect on virulence of *Vibrio* spp. *in vitro* and *in vivo*.


Results and discussion

Effect of cinnamaldehyde and cinnamaldehyde derivatives on microbial growth

When used in concentrations up to 150 μ M, cinnamaldehyde and most cinnamaldehyde derivatives (Fig. 1) had no inhibitory effect on the growth of strains in the present study (data not shown). The same was true for 4-NO₂-cinnamaldehyde, but only in concentrations up to 50 μ M. In all further experiments, 100 μ M was used (except for 4-NO₂-cinnamaldehyde, 25 μ M), unless otherwise mentioned.

Effect of cinnamaldehyde and 2-NO₂-cinnamaldehyde on bioluminescence

To rule out direct interference with the bioluminescence system of *Vibrio harveyi*, a constitutively bioluminescent strain was constructed. A plasmid containing *luxCDABE* genes under *lacZ* promotion was conjugated into *Escherichia coli* DH5 α (a strain defective in AI-2 production). The bioluminescence was not inhibited by cinnamaldehyde and cinnamaldehyde derivatives (data not shown) and these results indicate that the enzymes of *Vibrio harveyi* involved in bioluminescence are not inhibited by cinnamaldehyde or cinnamaldehyde derivatives.



| Compound | abbreviation | R1 | R2 | MW (g/mol) |
|---------------------------------|------------------------|--------------------|-------------------|------------|
| Cinnamaldehyde | CA | H | H | 132.16 |
| 4-methoxy-cinnamaldehyde | 4-MeO-CA | OMe- | H | 162.19 |
| 2-methoxy-cinnamaldehyde | 2-MeO-CA | H | OMe- | 162.19 |
| 4-nitro-cinnamaldehyde | 4-NO ₂ -CA | NO ₂ - | H | 177.16 |
| 2-nitro-cinnamaldehyde | 2-NO ₂ -CA | H | NO ₂ - | 177.16 |
| 4-dimethyl-amino-cinnamaldehyde | 4-Me ₂ N-CA | Me ₂ N- | H | 175.23 |

Figure 1
Cinnamaldehyde and cinnamaldehyde derivatives used in this study.

Effect of cinnamaldehyde and cinnamaldehyde derivatives on AI-2 based QS

Since bioluminescence is a QS regulated phenotype in *Vibrio harveyi*, we evaluated the effect of the different compounds on bioluminescence in this species. In a first screening we used *Vibrio harveyi* BB170. It was observed

that all of the compounds blocked the AI-2 QS system in a concentration-dependent way (Fig. 2). At 100 μ M, cinnamaldehyde and 2-NO₂-cinnamaldehyde were found to be the most active compounds, yielding an inhibition of $65 \pm 13\%$ and $62 \pm 7\%$, respectively. 2-MeO-cinnamaldehyde, 4-MeO-cinnamaldehyde and 4-Me₂N-cinnamaldehyde




Figure 2
Effect of cinnamaldehyde and cinnamaldehyde derivatives on AI-2 based QS. Bioluminescence in *Vibrio harveyi* BB170 as a function of the concentration of cinnamaldehyde and cinnamaldehyde derivatives. Bioluminescence measurements were performed 6 h after the addition of the compounds. Bioluminescence of the control (without addition of compound) was set at 100% and the responses for other samples were normalised accordingly. The error bars represent the standard deviation.

hyde were found to be less active at this concentration, with inhibitions of $14 \pm 5\%$, $34 \pm 9\%$ and $17 \pm 1\%$, respectively. The effect of 4-NO₂-cinnamaldehyde was only evaluated at lower concentrations because of its growth inhibitory effect. It was found to be the most active compound at concentrations of 25 and 50 μM , with inhibitions of $12 \pm 11\%$ and $33 \pm 7\%$, respectively. In general, the QS inhibition assay detected several active QS inhibitors and some striking structure-activity relationships. The inhibitory effect was highly dependent on the substitution pattern of the aromatic ring. Replacement of the dimethylamine (Me₂N) substituent with a methoxy (MeO) or a nitro (NO₂) group enhanced the activity. In both the methoxy and the nitro series the activity dropped (approximately $\pm 10\text{--}20\%$) upon moving the substituent from the para to the ortho position. In general, no cinnamaldehyde derivative was found to be more active than the unsubstituted cinnamaldehyde at concentrations of 100 μM and only one compound, 2-NO₂-cinnamaldehyde, was found to result in the same level of inhibition. At lower concentrations, 4-NO₂-cinnamaldehyde was significantly more active than the unsubstituted cinnamaldehyde, but the growth inhibitory effect of this compound prohibited its testing at higher concentrations.

Effect of cinnamaldehyde and cinnamaldehyde derivatives on the bioluminescence of *Vibrio harveyi* QS mutants

Bioluminescence in *Vibrio harveyi* BB170 is mainly controlled by AI-2, as this strain is not responsive to AHL stimulation [7]. Hence we limited the possible target of cinnamaldehyde and cinnamaldehyde derivatives to the AI-2 QS system. To determine the molecular target within the AI-2 QS pathway we measured the effect of cinnamaldehyde and cinnamaldehyde derivatives on the bioluminescence in different QS mutants. *Vibrio harveyi* MM30 has a mutation in the *luxS* gene, making it incapable of producing AI-2. However, this strain will react to exogenously added AI-2 with activation of the QS transduction system leading to bioluminescence. Inhibition of bioluminescence in this mutant would suggest the absence of an inhibitory effect on LuxS. Further we evaluated the effect of the test compounds on the production of AI-2 in *Escherichia coli* K12. The *Vibrio harveyi* JAF553 and JAF483 mutants contain a point mutation in the *luxU* and *luxO* genes, respectively, thereby preventing their phosphorelay capacity. *Vibrio harveyi* BNL258 has a Tn5 insertion in the *hfq* gene, resulting in a non-functional Hfq protein. *Vibrio harveyi* strains JAF553, JAF483 and BNL258 are all constitutively luminescent and inhibition of bioluminescence in one of these indicates that the cinnamaldehyde compounds act downstream of the mutated protein. Cinnamaldehyde and 2-NO₂-cinnamaldehyde were found to block bioluminescence in *Vibrio harveyi* MM30 (Fig. 3),




Figure 3

Effect of cinnamaldehyde and 2-NO₂-cinnamaldehyde on the bioluminescence of wild type *Vibrio harveyi* BB120 and the different *Vibrio harveyi* QS mutants. The percentage of bioluminescence of the *Vibrio harveyi* wild type BB120 and the mutants MM30, JAF553, JAF483 and BNL258 with 100 μM cinnamaldehyde (white bars) or 100 μM 2-NO₂-cinnamaldehyde (black bars) are presented. Measurements were performed 6 h after the addition of the compounds. Bioluminescence of the control (without addition of compound) was set at 100% and the response for the other samples were normalised accordingly. The error bars represent the standard deviation.

suggesting that these compounds do not exert their effect at the level of AI-2 production but rather at the level of the QS transduction system. Affirmatively, the supernatants of *Escherichia coli* K12 treated with cinnamaldehyde and cinnamaldehyde derivatives revealed no difference in AI-2 activity compared to the control (data not shown). Cinnamaldehyde and 2-NO₂-cinnamaldehyde were found to block bioluminescence to the same extent in all other mutants tested (Fig. 3). This suggests that the target of cinnamaldehyde and cinnamaldehyde derivatives is the downstream component of the AI-2 signalling transduction pathway, the transcriptional regulatory protein LuxR.

Effect of cinnamaldehyde on LuxR protein levels and on LuxR DNA-binding activity

Using purified LuxR protein, the presence of 0.19 mM and 0.75 mM cinnamaldehyde resulted in a maximal difference in LuxR DNA shift compared to the untreated control (Fig. 4a). These data indicate that *in vitro* binding of the transcriptional regulator LuxR to its promoter sequence is affected in the presence of cinnamaldehyde. Surprisingly, when adding higher concentrations of cinnamaldehyde (1.9 mM) no difference in shift could be observed anymore. This inconsistency was also observed for high

furanone concentrations and may be due to aspecific interactions with DNA and/or protein, although the exact reasons for this remain unknown (C. Miyamoto, unpublished data). Purified LuxR was also used to test whether cinnamaldehyde resulted in protein degradation. Three samples of LuxR containing varying amounts of cinnamaldehyde (0.19, 0.75 and 1.9 mM) and an untreated control were stained following electrophoresis on a 10% SDS-PAGE gel and were shown not to have been affected by cinnamaldehyde (Fig. 4b). To test whether the DNA-binding ability was also altered *in vivo*, lysates of *Vibrio harveyi* cells that were grown in the presence and absence of various cinnamaldehyde concentrations were also tested for their ability to cause a mobility shift of LuxR DNA (data not shown). Surprisingly, no effects were observed with concentrations < 1 mM. Using 1 mM cinnamaldehyde, there was about 4-fold less shift of LuxR DNA for the same amount of total protein in the lysate of *Vibrio harveyi* BB120 treated with cinnamaldehyde. There are several possible explanations for this apparent contradiction in terms of cinnamaldehyde concentrations required to cause a band shift. First of all, there may be considerable differences between the extra- and intracellular cinnamaldehyde concentrations, possibly explaining why we observed a shift with 0.19 mM cinnamaldehyde when purified LuxR protein was used but that higher concentrations were required when cell lysates were used. Secondly, there are no data on how much inhibition of binding of LuxR to its promoter is required in order to observe phenotypic changes (e.g. changes in bioluminescence). It may very well be that relatively minor changes in LuxR DNA binding (caused by relatively low cinnamaldehyde concentrations) are sufficient to cause reductions in bioluminescence but would go unnoticed in the gel shift assay. Combined, our data indicate that in the presence of cinnamaldehyde binding of the transcriptional regulator LuxR to its promoter sequence is affected, while leaving the protein intact. However, further research is needed to explain the differences between the *in vitro* and *in vivo* situation in terms of the cinnamaldehyde concentration required to observe this effect. Interestingly, the best-studied QS inhibitors, halogenated furanones, also interfere with binding of LuxR to its promoter sequence without degrading the protein [26].




Figure 4
LuxR DNA-binding as determined by mobility shifts and LuxR protein levels as determined by SDS-PAGE. A. Autoradiograph after 5% polyacrylamide gel electrophoresis of ³²P-labelled LuxR promoter DNA containing the LuxR binding sites, mixed with purified LuxR in the presence (0.19, 0.75 and 1.9 mM) and absence of cinnamaldehyde. B. SDS-PAGE of purified LuxR protein in the presence (0.19, 0.75 and 1.9 mM) or absence of cinnamaldehyde.

Effect of cinnamaldehyde and cinnamaldehyde derivatives on *Vibrio anguillarum* protease and pigment production

Cinnamaldehyde and 2-NO₂-cinnamaldehyde were found to decrease protease activity by 34 ± 2% and 49 ± 5%, respectively after 24 h (Fig. 5). 4-MeO-cinnamaldehyde was the only other cinnamaldehyde derivative to cause a significant decrease in protease activity (25 ± 6%) (Fig. 5). A time dependent inhibition of pigment production was found for cinnamaldehyde and 2-NO₂-cinnamaldehyde. After 48 h, inhibition in pigment




Figure 5
Effect of cinnamaldehyde and cinnamaldehyde derivatives on the protease activity of *Vibrio anguillarum* LMG 441 I. Cinnamaldehyde and cinnamaldehyde derivatives were tested at 100 μM, except 4-NO₂-cinnamaldehyde (25 μM). The effect of cinnamaldehyde or cinnamaldehyde derivatives on protease activity was compared to an untreated control. The error bars represent the standard deviation. *: Signal significantly different from the control (p < 0.05).

production was 25 ± 7% and 40 ± 2% for cinnamaldehyde and 2-NO₂-cinnamaldehyde (Fig. 6). In contrast, none of the other cinnamaldehyde derivatives were able to signif-

icantly reduce pigment production after 48 h (data not shown). Previously, it was shown that several virulence factors in *Vibrio anguillarum*, including pigment and pro-




Figure 6
Effect of cinnamaldehyde and 2-NO₂-cinnamaldehyde on the pigment production of *Vibrio anguillarum* LMG 441 I. Cinnamaldehyde and 2-NO₂-cinnamaldehyde were tested at 100 μM. *Vibrio anguillarum* LMG 441 I was allowed to produce pigment in the absence (solid symbol) or presence of cinnamaldehyde (open symbol, square) or 2-NO₂-cinnamaldehyde (open symbol, triangle). Three ml samples were taken at multiple time points. The effect of cinnamaldehyde or 2-NO₂-cinnamaldehyde on pigment production was estimated by measuring the absorbance at 405 nm. The error bars represent the standard deviation.

tease production, were regulated by QS. It was found that a mutation in *vanT* (the *luxR* homologue in *Vibrio anguillarum*) resulted in a significant decrease in total protease activity due to loss of expression of the metalloprotease EmpA [16]. Loss of protease activity could have several implications for the virulence of *Vibrio* spp. The protease Vvp of *Vibrio vulnificus*, which is homologous to EmpA, is thought to play an essential role in the colonisation of mucosal surfaces [31]. In addition, EmpA protease from *Vibrio anguillarum* is important for virulence during infection of the Atlantic salmon (*Salmo salar*) and contributes to hemorrhagic skin damage [32,33]. Several other phenotypes, including pigment production, were also found to be affected in a *Vibrio anguillarum vanT* mutant [16].

Effect of cinnamaldehyde and cinnamaldehyde derivatives on biofilm formation

Cinnamaldehyde was previously shown to inhibit *Escherichia coli* biofilms. Since cinnamaldehyde significantly reduced swimming motility in *Escherichia coli* it was hypothesized that reduced biofilm formation could be explained in part by an inability of the strain to reach the substratum [34]. However it should be noticed that no link with QS was described and cinnamaldehyde was used in high concentrations (> 2000 μ M). Cinnamaldehyde and some cinnamaldehyde derivatives decreased biofilm formation in *Vibrio anguillarum* LMG 4411 and *Vibrio vul-*

nificus LMG 16867 (Fig. 7). Cinnamaldehyde reduced total biomass (as measured by crystal violet staining, CV) with $26 \pm 7\%$ and $27 \pm 13\%$ in *Vibrio anguillarum* LMG 4411 and *Vibrio vulnificus* LMG 16867, respectively. 2-NO₂-cinnamaldehyde and 4-MeO-cinnamaldehyde resulted in a significant decrease in biomass of *Vibrio anguillarum* LMG 4411 (decrease of $34 \pm 16\%$ and $20 \pm 12\%$, respectively). No effect of cinnamaldehyde derivatives on *Vibrio vulnificus* LMG 16867 biomass was observed (Fig. 7). The cell-viability assay revealed no significant decrease in the number of metabolically active cells in *Vibrio anguillarum* LMG 4411 and *Vibrio vulnificus* LMG 16867 biofilm following treatment. In summary, cinnamaldehyde has an effect on total biofilm biomass but not on the number of viable cells. This suggests that cinnamaldehyde may have an effect on the production and/or accumulation of the exopolysaccharide (EPS) matrix (which is also stained with CV). To investigate this hypothesis, EPS was stained using Calcofluor white. Calcofluor white is a fluorescent dye which binds β 1-3 and β 1-4 carbohydrate linkages and which has been used to study EPS in a variety of organisms [35-37]. The staining was carried out on biofilms treated with cinnamaldehyde as this compound overall had most effect on biofilm biomass as assessed using CV. The use of cinnamaldehyde resulted in a lower fluorescent signal compared to an untreated control ($81 \pm 13\%$ and $69 \pm 27\%$ in *Vibrio*




Figure 7

Effect of cinnamaldehyde and cinnamaldehyde derivatives on *Vibrio* spp. biofilms. Biomass was quantified through CV staining. Cell-viability was quantified through CTB staining. CV signals are presented as a percentage compared to 100% control not receiving treatment (black bars = *Vibrio vulnificus*; white bars = *Vibrio anguillarum*). CTB signals are presented as a percentage compared to a 100% control not receiving treatment (vertical striped bars = *Vibrio vulnificus*; horizontal striped bars = *Vibrio anguillarum*). *: Signal significantly different compared to 100% control ($p < 0.05$).

anguillarum LMG 4411 and *Vibrio vulnificus* LMG 16867, respectively). These data support the hypothesis that the effect of cinnamaldehyde on biofilm formation can be explained by reduced EPS production and/or accumulation.

Protection of *Artemia* from *Vibrio harveyi*

For many pathogenic *Vibrio* spp., the production of protease, pigment and their capacity to form biofilms contribute to their virulence [31-33]. We investigated the ability of cinnamaldehyde and 2-NO₂-cinnamaldehyde, the two most active inhibitors, to protect *Artemia* shrimp against the virulent *Vibrio harveyi* BB120 strain. To this end, we followed the survival of *Artemia* after exposure to *Vibrio harveyi* BB120, with and without addition of compounds (Fig. 8). Cinnamaldehyde and 2-NO₂-cinnamaldehyde alone had no effects on *Artemia* shrimp (data not shown). As expected, high mortality rates were observed when exposing *Artemia* to *Vibrio harveyi* BB120. In contrast, cinnamaldehyde and 2-NO₂-cinnamaldehyde were able to completely protect *Artemia* against virulent *Vibrio harveyi* BB120 when used at concentrations of 100 μM and 150 μM (Fig. 8). At these concentrations, there was no effect on the growth of *Vibrio harveyi* BB120, ruling out that the protective effect of cinnamaldehyde and 2-NO₂-cinnamaldehyde was due to inhibition of the bacterial pathogen. These results suggest that cinnamaldehyde and

cinnamaldehyde derivatives may be useful as antipathogenic compounds.

Effect of cinnamaldehyde on the starvation response

The effect of cinnamaldehyde on the starvation response of *Vibrio vulnificus* LMG 16867 and *Vibrio anguillarum* LMG 4411 was investigated. In the control experiment no decrease in the number of culturable cells after 24 h of starvation was observed (Fig. 9). Upon treatment with cinnamaldehyde, however, cell numbers were significantly reduced (53 ± 3% and 57 ± 7% for *Vibrio vulnificus* LMG 16867 and *Vibrio anguillarum* LMG 4411, respectively) ($p < 0.05$). After 48 h, cell numbers were even further reduced in the cinnamaldehyde treated cultures (87 ± 3% and 63 ± 18% for *Vibrio vulnificus* LMG 16867 and *Vibrio anguillarum* LMG 4411, respectively), while there was only a 77 ± 5% and 4 ± 28% reduction in number of culturable cells in the control for *Vibrio vulnificus* LMG 16867 and *Vibrio anguillarum* LMG 4411, respectively. Bacteria are known for their ability to survive and respond to changes in their surroundings. One of these adaptations is the starvation response found in many marine bacteria. *Vibrio* spp. are known to survive for a long time without the addition of supplemental nutrition and this starvation response allows cells to survive adverse conditions. QS is thought to play a role in this response to stress conditions [38]. Our data indicate that inhibition of AI-2 based QS




Figure 8

Effect of cinnamaldehyde and 2-NO₂-cinnamaldehyde on the survival of *Artemia*. White bars represent the survival of *Artemia* without challenge with *Vibrio harveyi* BB120. Black bars represent the percentage survival of *Artemia* after challenge with *Vibrio harveyi* BB120 in untreated conditions. Striped bars represent the percentage of survival of *Artemia* after challenge with *Vibrio harveyi* BB120 when treated with cinnamaldehyde or 2-NO₂-cinnamaldehyde (horizontal: 100 μM stripes; vertical stripes: 150 μM, respectively). *: Survival significantly different from the treatment with pathogen alone ($p < 0.01$).




Figure 9
Effect of cinnamaldehyde on *Vibrio* spp. starvation response. The cells were allowed to starve in the presence (horizontal striped bars = *Vibrio vulnificus*; vertical striped bars = *Vibrio anguillarum*) and absence (black bars = *Vibrio vulnificus*; white bars = *Vibrio anguillarum*) of cinnamaldehyde. The number of CFU/ml was determined after 24 h and 48 h on TSA plates containing 2% NaCl. Data are presented as a percentage of the initial count. Error bars represent standard deviations.




Figure 10
Effect of cinnamaldehyde on antibiotic susceptibility of *Vibrio vulnificus* LMG 16867. Effects of chloramphenicol (squares) and doxycycline (triangles) on the growth of *Vibrio vulnificus* LMG 16867 in the presence (open symbols) and absence (solid symbols) of cinnamaldehyde (100 µM) are presented. The absorbance at 590 nm was measured after 24 h of growth. Error bars represent standard deviations.

suppresses the starvation response and makes cells more susceptible to starvation-associated stress conditions. This is in agreement with a previously published study [17] in which starvation survival in *Vibrio vulnificus* was reduced by mutation of LuxR in the QS system and by halogenated furanones.

Effect of cinnamaldehyde on antibiotic susceptibility

We have examined the association between QS and antibiotic susceptibility in two *Vibrio* spp. Two antibiotics with a different mode of action were chosen. Chloramphenicol, previously used as prophylactic in aquaculture, targets the 50S ribosomal subunit [39,40]. Doxycycline, an antibiotic targeting the 30S ribosomal subunit, is the recommended antibiotic therapy for *Vibrio vulnificus* infections [41]. *Vibrio vulnificus* LMG 16867 showed an increased antibiotic susceptibility when treated with cinnamaldehyde (Fig. 10). This difference was most pronounced when using chloramphenicol. In contrast, in *Vibrio anguillarum* LMG 4411, no differences were observed between cinnamaldehyde treatment and control (data not shown). Previously, it was found that QS inhibition could alter the susceptibility of a strain towards antimicrobial agents. *Vibrio cholerae* strains with various mutations in the AI-2 signal transduction system appeared to be more sensitive to treatment with hydrogen peroxide [42]. Similarly, a *Streptococcus anginosus* LuxS mutant was found to be more susceptible towards ampicillin and erythromycin than the wild type strain [43].

Conclusion

Cinnamaldehyde and several derivatives were shown to interfere with AI-2 based QS by decreasing the ability of LuxR to bind to its target promoter sequence. These compounds, used in sub-inhibitory concentrations, did not only affect *in vitro* the production of multiple virulence factors and biofilm formation, but also reduced *in vivo* the mortality of *Artemia* shrimp exposed to *Vibrio harveyi* BB120. In addition, cinnamaldehyde reduced the ability to cope with stress factors like starvation and exposure to antibiotics. Our results indicate that cinnamaldehyde and cinnamaldehyde derivatives are potentially useful antipathogenic lead compounds for treatment of vibriosis.

Methods

Cinnamaldehyde and cinnamaldehyde derivatives

Cinnamaldehyde (Sigma-Aldrich, Bornem, Belgium) and cinnamaldehyde derivatives [4-MeO-cinnamaldehyde (VWR International, West Chester, USA), 2-MeO-cinnamaldehyde (Wako Pure Chemical Industries, Osaka, Japan), 4-NO₂-cinnamaldehyde, 2-NO₂-cinnamaldehyde and 4-Me₂N-cinnamaldehyde (Acros Organics, Geel, Belgium)] (Fig. 1), were diluted in DMSO (0.5% v/v). The stock solutions were stored at -20°C. Control solutions

(CS) contained the same amount of DMSO, without cinnamaldehyde or cinnamaldehyde derivatives.

Bacterial strains, plasmid and growth conditions

The strains and plasmid used in this study are shown in Table 1. All strains were routinely cultured in Marine-Broth (MB) (BD, Sparks, MD, USA) in the presence of appropriate antibiotics, except for *Escherichia coli* DH5 α and K12, which were grown in Luria-Bertani broth (LB) (BD). The medium was supplemented with 100 μ g/ml ampicillin (Sigma-Aldrich) for *Escherichia coli* DH5 α containing the pBlueLux plasmid. *Vibrio anguillarum* LMG 4411, *Vibrio vulnificus* LMG 16867 and the various *Vibrio harveyi* strains were cultured overnight at 30°C on a rotary shaker. *Escherichia coli* DH5 α and K12 were cultured overnight at 30°C and 37°C, respectively, without agitation. Minimal inhibitory concentrations were determined for each compound by using a microdilution assay in flat bottomed 96-well microtiter plates (TPP, Trasadingen, Switzerland), using MB and LB medium for all vibrios and both *Escherichia coli* strains, respectively. The plates were incubated for 24 h and the absorption at 590 nm was measured using a Victor Wallac² multilabel counter (Perkin Elmer Life and Analytical Sciences, Boston, MA, USA).

Effect of cinnamaldehyde and cinnamaldehyde derivatives on bioluminescence

To determine whether any of the compounds had an effect on bioluminescence not related to inhibition of QS, *Escherichia coli* DH5 α was transformed with the pBlueLux plasmid, containing *luxCDABE* under control of a *lacZ* promoter and the effect on bioluminescence was measured. The pBlueLux plasmid was transformed in *Escherichia coli* DH5 α as follows. Overnight cultures were suspended in a 50 mM CaCl₂ solution at 0°C. The pBlueLux plasmid was added and the solution was incubated for 15 min. After this, the solution was transferred to 42°C for 90 sec and the cell suspension was plated on Trypton soy agar (TSA) (Oxoid, Basingstoke, Hampshire, UK) containing 100 μ g/ml ampicillin (Sigma-Aldrich) for selection of transformants. For the bioluminescence assay an overnight culture was diluted to OD_{590 nm} of approximately 0.1 and 100 μ l of cell suspension was added to each well of a black 96-well microtiter plate (Perkin Elmer). The effect on bioluminescence for the active compounds was compared to controls not receiving the active molecules.

Bioassay for LuxS inhibition

In order to determine whether any of the compounds tested had an effect on production of AI-2, AI-2 activity was measured in supernatants of *Escherichia coli* K12 cultures grown for 16 h with or without compounds. Overnight cultures of *Escherichia coli* K12 were centrifuged (5000 rpm, 5 min, room temperature) and filter sterilised (0.22 μ m, Whatman GmbH, Dassel, Germany). The

Table 1: Strains and plasmid used in this study.

| Strain/plasmid | Relevant features | Reference or source |
|---------------------------------|--|---------------------|
| <i>Vibrio harveyi</i> strains | | |
| BB120 | Wild type from which strains BB152, BB170, MM30, JAF553, JAF483 and BNL258 are derived | [9] |
| BB170 | <i>luxN::Tn5</i> | [7] |
| MM30 | <i>luxS::Tn5</i> | [10] |
| JAF553 | <i>luxU</i> H58A linked to Kan ^R | [48] |
| JAF483 | <i>luxO</i> D47A linked to Kan ^R | [14] |
| BNL258 | <i>hfq::Tn5lacZ</i> | [49] |
| <i>Vibrio anguillarum</i> | | |
| LMG 4411 | Isolated from young sea trout (<i>Salmo trutta</i>) | BCCM/LMG |
| <i>Vibrio vulnificus</i> | | |
| LMG 16867 | Isolated from tankwater from eelfarm | BCCM/LMG |
| <i>Escherichia coli</i> strains | | |
| DH5 α | AI-2 ⁻ strain | [23] |
| K12 | AI-2 ⁺ strain | [23] |
| ATCC 25404 | | |
| Plasmid | | |
| pBlueLux | pBlueLux polylinker and <i>luxCDABE</i> | S. Atkinson |

BCCM/LMG: Belgian Co-ordinated Collections of Micro-organisms/Laboratory of Microbiology collection (Ghent University, Belgium); ATCC: American Type Culture Collection

supernatants were used immediately or stored at -20°C. AI-2 levels were determined in a *Vibrio harveyi* BB170 assay as described previously [21]. In brief, an overnight culture of the reporter strain was diluted 1:5000 into fresh sterile MB medium and 90 μ l of this cell suspension was added to the wells of a black 96-well microtiter plate (Perkin Elmer). Ten μ l of the appropriate sterile supernatants was then added to the wells, the microtiter plates were incubated at 30°C and bioluminescence was measured hourly using a Wallac Victor² multilabel counter (Perkin Elmer). Bioluminescence was expressed as the fraction of bioluminescence measured in the positive control reaction. Confirmation of these results was obtained using *Vibrio harveyi* MM30, a Δ *LuxS* mutant, instead of *Vibrio harveyi* BB170. The effect on bioluminescence for the active compounds was compared to controls not receiving the active molecules.

Other *Vibrio harveyi* bioassays

Using *Vibrio harveyi* strains BB120, JAF553, JAF483 and BNL258, we determined whether the molecular target of our compounds was located in the AI-2 signalling transduction pathway. The bioluminescence assay as described above was used with minor modifications. In brief, the

positive control reaction received 10 μ l of *Escherichia coli* K12 supernatant, without addition of the test molecule. Negative control reactions received 10 μ l sterile MB-medium. Other wells received 10 μ l of *Escherichia coli* K12 supernatants (containing AI-2) and appropriate amounts of the test molecule.

Effect of cinnamaldehyde on *LuxR* protein levels and DNA-binding activity

Mobility shift assays and SDS-PAGE assays were performed as described previously [26] with minor modifications. *Vibrio harveyi* BB120 cells were grown in the presence and absence of cinnamaldehyde and all cell lysates were taken at different optical densities (OD_{600 nm} = 1.2, 1.6, 1.8 and 2.1). Previously purified *LuxR* [44] was used for mobility shift and SDS-PAGE assay. For SDS-PAGE the following protein standard (Bio-rad) was used: 250, 150, 100, 75, 50, 37, 25, 20, 15 and 10 kDa.

Quantification of protease activity

Vibrio anguillarum LMG 4411 was grown overnight in MB. Protease activity was quantified following inoculation of cultures into medium containing 2.0% Bacto agar (Oxoid), 2.0% NaCl (Novolab, Geraardsbergen, Belgium)

and 3.0% Skim Milk powder (Oxoid). Appropriate amounts of test compounds and CS were added to the mixtures, 0.5 ml of these mixtures was added to the wells of a 24-well microtiter plate (TPP, Trasadingen, Switzerland) and the plate was incubated at 30°C. Clearing was measured spectrophotometrically with a Wallac Victor² multilabel counter after 24 h.

Quantification of pigment production

Vibrio anguillarum LMG 4411 was grown overnight at 30°C in MB. The overnight culture was then diluted to OD_{590 nm} = 0.05 in Tryptone Soy Broth (TSB) (Oxoid) containing 5 mM L-tyrosine (Sigma-Aldrich) with or without test compound and incubated at 30°C with shaking. At various time points, 3 ml samples were taken from the cultures and supernatants were collected by centrifugation (5000 rpm, 4 min, room temperature), followed by filter sterilisation (0.22 µm). Pigment production was followed by measuring the absorbance at 405 nm.

Biofilm formation assay

Vibrio anguillarum LMG 4411 (doubling time T_d: 3.2 h) and *Vibrio vulnificus* LMG 16867 (T_d: 5.3 h) were grown overnight in MB, centrifuged, resuspended in double concentrated Marine Broth (2xMB) and diluted to an OD_{590 nm} = 0.1 in 2xMB. Fifty µl of the diluted bacterial suspension was transferred to the wells of a round-bottomed 96-well microtiter plate (TPP). Negative controls received 50 µl of CS. Positive controls received 50 µl of the test compound in appropriate concentrations. Bacteria were allowed to adhere and grow without agitation for 4 h at 30°C. After 4 h, plates were emptied and washed with sterile physiological saline (PS). After this washing step, negative control wells were filled with 50 µL 2xMB and 50 µl CS. Other wells were filled with 50 µl 2xMB and 50 µl compound solution and the plate was incubated for 24 h at 30°C. Biofilm biomass was quantified by crystal violet (CV) staining, as described previously [45]. In brief, plates were rinsed with sterile PS, biofilms were fixed by adding 100 µl 99% methanol for 15 min, after which the methanol was removed and plates were air-dried. Biofilms were then stained with 100 µl CV (Pro-lab Diagnostics, Richmond Hill, ON, Canada). After 20 min, CV was removed and wells were filled with 150 µl 33% acetic acid (Sigma-Aldrich). The absorbance was measured at 590 nm using a Wallac Victor² multilabel counter and results were expressed as the percentages compared to the signal of the control not receiving treatment. For quantification of the number of metabolically active (i.e. living) cells in the biofilm, a resazurin assay was used [45]. In brief, wells were rinsed after 24 h biofilm formation and 100 µl PS was added, followed by addition of 20 µl CellTiter-Blue (CTB) (Promega, Leiden, The Netherlands) solution. After 60 min, fluorescence (ex_{560 nm}/em_{590 nm}) was measured using a Wallac Victor² multilabel counter. For the quanti-

fication of EPS, a Calcofluor white staining (Sigma-Aldrich) was used. In brief, wells were rinsed after 24 h biofilm formation and 100 µl phosphate buffered saline (PBS) containing 0.5 µl 5 mM CFW was added to the wells. After 60 min, fluorescence (ex_{405 nm}/em_{500 nm}) was measured using a Wallac Victor² multilabel counter.

Artemia Challenge tests

All experiments were performed with high quality hatching cysts of *Artemia franciscana* (EG[®] Type, batch 6940, INVE Aquaculture, Baasrode, Belgium). 200 mg of cysts were hydrated in 18 ml of tap water during 1 h. Sterile cysts and nauplii were obtained via decapsulation as described previously [46]. Challenge tests were performed as described previously [18] with minor modifications. Briefly, after hatching, groups of 20 nauplii were transferred to new sterile 50 ml tubes that contained 20 ml of 0.22 µm filtered and autoclaved artificial seawater. *Vibrio harveyi* BB120 was washed in filtered and autoclaved artificial seawater after incubation and added to the *Artemia* culture water at a concentration of approximately 10⁵ CFU/ml. A suspension of autoclaved LVS3 bacteria in filtered and autoclaved artificial seawater was added as feed in a concentration of approximately 10⁷ CFU/ml culture water. After the addition of 100 µM or 150 µM of cinnamaldehyde or 2-NO₂-cinnamaldehyde (or an appropriate volume of solvent), the falcon tubes were put back on the rotor and kept at 28°C. *Artemia* cultures to which only autoclaved LVS3 bacteria were added were used as controls. The survival of *Artemia* was scored 48 h after the addition of the strains. All manipulations were done under a laminar flow hood in order to maintain sterility of the cysts and nauplii. Each treatment was done in triplicate.

Starvation assay

Vibrio anguillarum LMG 4411 and *Vibrio vulnificus* LMG 16867 strains were grown overnight in MB, the cells were collected by centrifugation (5000 rpm, 4 min), washed in PS and resuspended in artificial seawater (ASW) [47] containing 0.1% MB (with and without test compound). These suspensions were incubated at 30°C without shaking. At various time points, 1 ml samples were taken and the number of culturable cells was determined by plating serial dilutions on TSA (Oxoid) plates containing 2% NaCl. Results were expressed as the percentage survival compared to the untreated control.

Effect of cinnamaldehyde on antibiotic resistance

Fifty µL of double concentrated TSB (2xTSB) containing 4% NaCl with or without chloramphenicol (Sigma-Aldrich) or doxycycline (Sigma-Aldrich) (added in the range of 0.001 µg/ml – 25 µg/ml) were dispensed into flat-bottomed 96-well microtiter plates (TPP). An equal amount of cinnamaldehyde was added (final concentra-

tion of 100 μM). For the controls, equal amounts of CS were added to the wells. *Vibrio vulnificus* LMG 16867 or *Vibrio anguillarum* LMG 4411 was added in a final concentration of 10^5 CFU/ml. The plates were incubated overnight at 30°C and growth was evaluated after 24 h by absorbance measurements at 590 nm using a Wallac Victor²multilabel counter.

Statistics

Independent samples t-tests were performed using the SPSS software, version 15.0 (SPSS, Chicago, IL, USA).

Authors' contributions

GB carried out most of the experiments, analysed the data, provided figures and tables and drafted the manuscript. TD and PB carried out the in vivo virulence assay. CM carried out the mobility shift and SDS-PAGE assays. SVC coordinated and participated in the selection of the compounds. TC and HN coordinated the study and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors like to thank Dr. S. Atkinson for kindly providing the pBlueLux plasmid. This work was supported by the 'Instituut voor de aanmoediging van Innovatie door Wetenschap en Technologie in Vlaanderen' (GB) and FWO-Vlaanderen (TC).

References

- Austin B, Zhang XH: **Vibrio harveyi: a significant pathogen of marine vertebrates and invertebrates.** *Letts Appl Microbiol* 2006, **43**:119-124.
- APEC/FAO/NACA/SEMARNAP: **Report of a joint APEC/FAO/NACA/SEMARNAP ad-hoc expert consultation on trans-boundary aquatic animal pathogen transfer and development of harmonised standards on aquaculture health management.** Puerto Vallarta, Jalisco, Mexico. 24-28 July 2000
- Holmström K, Gräslund S, Wahlström A, Pongshompoo S, Bengtsson BE, Kautsky N: **Antibiotic use in shrimp farming and implications for environmental impacts and human health.** *Int J Food Sci Technol* 2003, **38**:255-266.
- Tendencia EA, Dela Pena M: **Antibiotic resistance of bacteria from shrimp ponds.** *Aquaculture* 2001, **195**:193-204.
- Karunasagar I, Pai R, Malahti GR, Karunasagar I: **Mass mortality of Penaeus monodon larvae due to antibiotic-resistant Vibrio harveyi infection.** *Aquaculture* 1994, **128**:203-209.
- Xavier KB, Bassler BL: **LuxS quorum sensing: more than just a numbers game.** *Curr Opin Microbiol* 2003, **6**:191-197.
- Bassler BL, Wright M, Showalter RE, Silverman MR: **Intercellular signalling in Vibrio harveyi: sequence and function of genes regulating expression of luminescence.** *Mol Microbiol* 1993, **9**:773-786.
- Higgins DA, Pomianek ME, Kraml CM, Taylor RK, Semmelhack MF, Bassler BL: **The major Vibrio cholerae autoinducer and its role in virulence factor production.** *Nature* 2007, **450**:883-88.
- Bassler BL, Greenberg EP, Stevens AM: **Cross-species induction of luminescence in the quorum-sensing bacterium Vibrio harveyi.** *J Bacteriol* 1997, **179**:4043-4045.
- Surette MG, Miller MB, Bassler BL: **Quorum sensing in Escherichia coli, Salmonella typhimurium, and Vibrio harveyi: a new family of genes responsible for autoinducer production.** *Proc Natl Acad Sci* 1999, **96**:1639-1644.
- Miller ST, Xavier KB, Campagna SR, Taga ME, Semmelhack MF, Bassler BL, Hughson FM: **Salmonella typhimurium recognizes a chemically distinct form of the bacterial quorum-sensing signal AI-2.** *Mol Cell* 2004, **15**:677-687.
- Milton DL: **Quorum sensing in Vibrio spp.: complexity for diversification.** *Int J Med Microbiol* 2006, **296**:61-71.
- Bassler BL, Wright M, Silverman MR: **Sequence and function of LuxO, a negative regulator of luminescence in Vibrio harveyi.** *Mol Microbiol* 1994, **12**(3):403-412.
- Freeman JA, Bassler BL: **A genetic analysis of the function of LuxO, a two component response regulator involved in quorum sensing in Vibrio harveyi.** *Mol Microbiol* 1999, **31**:665-667.
- Larsen MH, Blackburn N, Larsen JL, Olsen JE: **Influences of temperature, salinity and starvation on the motility and chemotactic response of Vibrio anguillarum.** *Microbiology* 2004, **150**:1283-1290.
- Croxatto A, Chalker VJ, Lauritz J, Jass J, Hardman A, Williams P, Camara M, Milton DL: **VanT, a homologue of Vibrio harveyi LuxR, regulates serine, metalloprotease, pigment, and biofilm production in Vibrio anguillarum.** *J Bacteriol* 2002, **184**:1617-1629.
- McDougald D, Rice SA, Kjelleberg S: **SmcR-dependent regulation of adaptive phenotypes in Vibrio vulnificus.** *J Bacteriol* 2001, **183**:758-762.
- Defoirdt T, Bossier P, Sorgeloos P, Verstraete W: **The impact of mutations in the quorum sensing system of Aeromonas hydrophila, Vibrio anguillarum and Vibrio harveyi on their virulence towards gnotobiotically cultured Artemia franciscana.** *Environ Microbiol* 2005, **7**:1239-1247.
- Lee JH, Rhee JE, Park U, Ju HM, Lee BC, Kim TS, Jeong HS, Choi SH: **Identification and functional analysis of Vibrio vulnificus SmcR, a novel global regulator.** *J Microbiol Biotechnol* 2007, **17**:325-334.
- Alfaro JF, Zhang T, Wynn DP, Karschner EL, Zhou ZS: **Synthesis of LuxS inhibitors targeting bacterial cell-cell communication.** *Org Lett* 2004, **6**:3043-3046.
- Niu C, Alfere S, Gilbert ES: **Subinhibitory concentrations of cinnamaldehyde interfere with quorum sensing.** *Letts Appl Microbiol* 2006, **43**:489-494.
- Novak JS, Fratamico PM: **Evaluation of ascorbic acid as a quorum sensing analogue to control growth, sporulation, and enterotoxin production in Clostridium perfringens.** *J Food Sci* 2004, **69**:72-78.
- Ren D, Bedzyk LA, Ye RW, Thomas SM, Wood TK: **Differential gene expression shows natural brominated furanones interfere with the autoinducer-2 bacterial signalling system of Escherichia coli.** *Biotech Bioeng* 2004, **88**:630-642.
- Ren D, Sims JJ, Wood TK: **Inhibition of biofilm formation and swarming of Escherichia coli by (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2-(5H)-furanone.** *Environ Microbiol* 2001, **3**:731-736.
- Hentzer M, Givskov M: **Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections.** *J Clin Invest* 2003, **112**:1300-1307.
- Defoirdt T, Miyamoto CM, Wood TK, Meighen EA, Sorgeloos P, Verstraete W, Bossier P: **The natural furanone (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2-(5H)-furanone disrupts quorum sensing-regulated gene expression in Vibrio harveyi by decreasing the DNA-binding activity of the transcriptional regulator protein luxR.** *Environ Microbiol* 2007, **9**:2486-2495.
- Defoirdt T, Crab R, Wood TK, Sorgeloos P, Verstraete W, Bossier P: **Quorum sensing-disrupting brominated furanones protect the gnotobiotic brine shrimp Artemia franciscana from pathogenic Vibrio harveyi, Vibrio campbellii, and Vibrio parahaemolyticus isolates.** *Appl Environ Microbiol* 2006, **72**:6419-6423.
- Tinh NT, Linh ND, Wood TK, Dierckens K, Sorgeloos P, Bossier P: **Interference with the quorum sensing systems in a Vibrio harveyi strain alters the growth rate of gnotobiotically cultured rotifer Brachionus plicatilis.** *J Appl Microbiol* 2007, **103**:194-203.
- Adams TB, Cohen SM, Doull J, Feron VJ, Goodman JJ, Marnett LJ, Munro IC, Portoghese PS, Smith RL, Waddell WJ, Wagner BM: **The FEMA GRAS assessment of cinnamyl derivatives used as flavour ingredients.** *Food Chem Toxicol* 2004, **42**:157-185.
- Friedman M, Kozukue N, Harden LA: **Cinnamaldehyde content in foods determined by gas chromatography-mass spectrometry.** *J Agric Food Chem* 2000, **48**:5702-5709.
- Valiente E, Lee CT, Hor LI, Fouz B, Amaro C: **Role of the metalloprotease Vvp and the virulence plasmid pR99 of Vibrio vulnificus serovar E in surface colonization and fish virulence.** *Environ Microbiol* 2007, **10**:328-338.

32. Denkin SM, Nelson DR: **Regulation of *Vibrio anguillarum* EmpA metalloprotease expression and its role in virulence.** *Appl Environ Microbiol* 2004, **70**:4193-4204.
33. Miyoshi S: ***Vibrio vulnificus* infection and metalloprotease.** *J Dermatol* 2006, **33**:589-595.
34. Niu C, Gilbert ES: **Colorimetric method for indentifying plant essential oil components that affect biofilm formation and structure.** *Appl Environ Microbiol* 2004, **70**:6951-6956.
35. Ledebouer NA, Jones BD: **Exopolysaccharide sugars contribute to biofilm formation by *Salmonella enterica* serovar Typhimurium on Hep-2 cells and chicken intestinal epithelium.** *J Bacteriol* 2005, **187**:3214-3226.
36. Wells DH, Long SR: **The *Sinorhizobium meliloti* stringent response affects multiple aspects of symbiosis.** *Mol Microbiol* 2005, **43**:1115-1127.
37. Zogaj X, Nimtz M, Rohde M, Bokranz W, Romling U: **The multicellular morphotypes of *Salmonella typhimurium* and *Escherichia coli* produce cellulose as the second component of the extracellular matrix.** *Mol Microbiol* 2001, **39**:1452-1463.
38. McDougald D, Srinivasan S, Rice SA, Kjelleberg S: **Signal-mediated cross-talk regulates stress adaptation in *Vibrio* species.** *Microbiology* 2003, **149**:1923-1933.
39. Defoirdt T, Boon N, Sorgeloos P, Verstraete W, Bossier P: **Alternatives to antibiotics to control bacterial infections: luminescent vibriosis in aquaculture as an example.** *Trends Biotechnol* 2007, **25**:472-479.
40. Sutcliffe JA: **Improving on nature: antibiotics that target the ribosome.** *Curr Opin Microbiol* 2005, **8**:534-542.
41. Bross MH, Soch K, Morales R, Mitchell RB: ***Vibrio vulnificus* infection: diagnosis and treatment.** *Am Fam Physician* 2007, **76**:539-544.
42. Joelsson A, Kan B, Zhu J: **Quorum sensing enhances the stress response in *Vibrio cholerae*.** *Appl Environ Microbiol* 2007, **73**:3742-3746.
43. Ahmed NA, Petersen FC, Scheie AA: **AI-2 quorum sensing affects antibiotic susceptibility in *Streptococcus anginosus*.** *J Antimicrob Chemother* 2007, **60**:49-53.
44. Swartzman E, Meighen EA: **Purification and characterization of a poly-(dA-dT) lux-specific DNA-binding protein from *Vibrio harveyi* and identification as LuxR.** *J Biol Chem* 1993, **268**:16706-16716.
45. Peeters E, Nelis HJ, Coenye T: **Comparison of multiple methods for quantification of microbial biofilms grown in microtiter plates.** *J Microbiol Methods* 2008, **72**:157-165.
46. Marques A, Francois JM, Dhont J, Bossier P, Sorgeloos P: **Influence of yeast quality on performance of gnotobiotically grown *Artemia*.** *J Exp Mar Biol Ecol* 2004, **310**:247-264.
47. Bang W, Drake MA, Jaykus LA: **Recovery and detection of *Vibrio vulnificus* during cold storage.** *Food Microbiol* 2007, **24**:664-670.
48. Freeman JA, Bassler BL: **Sequence and function of LuxU: a two component phosphorelay protein that regulates quorum sensing in *Vibrio harveyi*.** *J Bacteriol* 1999, **181**:899-906.
49. Lenz DH, Mok KC, Lilley BN, Kulkarni RV, Wingreen NS, Bassler BL: **The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in *Vibrio harveyi* and *Vibrio cholerae*.** *Cell* 2004, **118**:69-82.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

