



Circulating tumor DNA-based predictive biomarkers in breast cancer clinical trials: a narrative review

Oraianthi Fiste[^], Michael Liontos[^], Konstantinos Koutsoukos[^], Evangelos Terpos[^], Meletios A. Dimopoulos[^], Flora Zagouri[^]

Department of Clinical Therapeutics, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece

Contributions: (I) Conception and design: O Fiste, F Zagouri; (II) Administrative support: All authors; (III) Provision of study materials or patients: O Fiste; (IV) Collection and assembly of data: O Fiste; (V) Data analysis and interpretation: O Fiste, F Zagouri; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Oraianthi Fiste, MD, MSc. Department of Clinical Therapeutics, School of Medicine, National and Kapodistrian University of Athens, Alexandra Hospital 80 Vasilissis Sophias, 11528 Athens, Greece. Email: ofiste@med.uoa.gr.

Abstract: Breast carcinoma is the most frequent and the second leading cause of cancer mortality in women worldwide. Current treatment decisions are based on tumor profiling of the initial tissue biopsy. Cancer though evolves both spatially and temporarily in a significant percentage of patients during treatment. However, sequential biopsies from the primary tumor or its metastatic sites are not either convenient or feasible in the majority of cases. In the era of precision medicine, analysis of circulating blood-based biomarkers in the field of liquid biopsies provides an insight into the dynamic molecular profiling of the primary tumor and its metastases, in a relatively non-invasive way. The latter permits not only patient stratification but also longitudinal evaluation of treatment response, when incorporated into clinical trials. This review summarizes the results from recent and ongoing circulating tumor DNA (ctDNA)-based biomarker-driven clinical trials, with respect to ctDNA analysis' predictive role, both in adjuvant, neo-adjuvant, and metastatic setting. Furthermore, current challenges in ctDNA analysis applications are critically discussed, including pre-analytical and analytical issues, and future perspectives in this field, through the conduct of well-designed, multicenter, randomized, large-scale, biomarker-stratified trials, with robust statistical methods. Despite in its infancy, ctDNA analysis holds great promise as a minimally invasive tool regarding tailored, personalized treatment guidance for breast cancer patients.

Keywords: Breast cancer; circulating tumor DNA (ctDNA); clinical trials; predictive biomarkers

Submitted Feb 02, 2020. Accepted for publication Sep 06, 2020.

doi: 10.21037/atm-20-1175

View this article at: <http://dx.doi.org/10.21037/atm-20-1175>

Introduction

Breast cancer is the most prevalent cancer and the second leading cause of cancer mortality in women (1,2). Next-generation sequencing (NGS)-based diagnostics have identified around 40 genomic alterations, shedding light into the heterogeneity of this disease (3,4). Currently, only

a few of these somatic alterations have been validated as therapeutic targets, whereas, there are multiple targeted therapies, effective as signalling blockade, in the adjuvant, neo-adjuvant, and metastatic settings.

In particular, trastuzumab (5-7), pertuzumab (8-10), ado-trastuzumab emtansine (11), lapatinib (12) and neratinib (13)

[^] ORCID: Oraianthi Fiste: 0000-0002-7106-4162; Michael Liontos: 0000-0001-7192-4575; Konstantinos Koutsoukos: 0000-0002-1710-125X; Evangelos Terpos: 0000-0001-5133-1422; Meletios Athanasios Dimopoulos: 0000-0001-8990-3254; Flora Zagouri: 0000-0001-7153-6556.

are human epidermal growth factor receptor 2 (HER2) inhibitors for the treatment of HER2+ disease; palbociclib (14), ribociclib (15) and abemaciclib (16) are cyclin-dependent kinase 4 and 6 (CDK4/6) inhibitors for the treatment of hormone-receptor (HR)+, HER2- disease, in combination with hormonal treatment, like aromatase inhibitors (AIs), tamoxifen or fulvestrant; everolimus (17) is a mammalian target of rapamycin (mTOR) inhibitor, also, for the treatment of HR+, HER2- disease, in combination with hormonal treatment; olaparib (18) and talazoparib (19) are poly adenosine diphosphate (ADP) ribose polymerase (PARP) inhibitors for the treatment of BReast CAncer gene (BRCA)+ disease; while alpelisib (20) is a phosphoinositide-3-kinase (PI3K) inhibitor for the treatment of phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha (PIK3CA)+ disease.

Despite the rapid advance in personalized medicine strategies, metastatic breast cancer remains an incurable disease, with a 5-year survival rate of approximately 25% (21,22). Breast cancer's plasticity, over time and under treatment pressure, represents the greatest challenge in its therapeutics, due to disease recurrence and drug resistance (23,24). Thus, both American and European guidelines recommend reassessment of biomarkers, like HR and HER2 status, if feasible, in the metastatic setting (25).

Unfortunately, tissue biopsies are fraught with several caveats; they are invasive, patient-unfriendly procedures, not always feasible either because of patient's condition and comorbidities or because of tumor's accessibility, and they don't permit longitudinal monitoring of tumor (26-30). Thus, the ideal approach to address the diverse molecular profile of breast tumors would be a minimally invasive method that could capture the entire genetic make-up of the tumor, in 'real-time', during the course of treatment. Currently, analysis of circulating blood biomarkers, like circulating tumor DNA (ctDNA), under the umbrella-term of 'liquid biopsies', offers an attractive approach to evaluate patient's entire tumor burden, in a non-invasive, convenient, repetitive, dynamic, and cost-effective way (27,31-36).

Several studies have evaluated the emerging role of ctDNA in monitoring treatment response or resistance and in predicting early relapse (37-48). Nevertheless, studies investigating the potential capacity of serial ctDNA monitoring for treatment guidance are still scarce, small-scale, and lack a strict clinically-centered protocol. To the best of our knowledge, there is no other review focused on the incorporation of ctDNA-based predictive biomarkers in breast cancer patients enrolled in clinical trials, therefore,

we performed this review of the published literature, to assess the potential of ctDNA in optimizing disease management.

We present the following article in accordance with the Narrative Review reporting checklist (available at <http://dx.doi.org/10.21037/atm-20-1175>).

Methods

A review of published literature was conducted to assess the predictive value of ctDNA analysis in the setting of clinical trials in breast cancer patients. All eligible studies were identified by a search in www.clinicaltrials.gov, MEDLINE/PubMed database and Cochrane Database of Systematic Reviews (CDSR) for the period up to August 31, 2019. Clinical Trials incorporating ctDNA analysis, as source of potential predictive biomarkers, in patients with breast cancer were considered for inclusion. To create a search strategy, medical subject heading (MeSH) terms [breast, cancer, neoplasm, carcinoma, clinical trial, ctDNA, cell free DNA (cfDNA), predictive, biomarker] were used in addition with Boolean search terms (AND, OR).

Eligible for inclusion were considered all randomised and non-randomised clinical trials carried out in adult patients (≥ 18 years old), irrespective of gender, with breast cancer, reporting results of ctDNA analysis and its correlation with treatment efficacy. Abstracts presented in conferences were also included.

Language restrictions were applied (only articles published in English were considered eligible). Animal studies, book chapters, observational study designs, commentaries, case reports, reviews, meta-analyses and studies not in cancer patients were also excluded.

The following data were extracted from each clinical trial: clinical trial name and Identification number (ID number), status, first author, year of publication, setting (primary or advanced breast cancer), line of therapy (neo-adjuvant, adjuvant, and 1st or 2nd line for metastatic setting, etc.), pathological subtype/hormonal status, allocation of study (randomized, non-randomized), intervention model (sequential-, parallel-, single group-assignment), masking, phase, treatment modalities (intervention and control arm regimens), number of patients enrolled in biomarker sub-study, primary endpoint, ctDNA sequencing technique, results.

Results

Our search strategy retrieved initially 64 clinical trials,

which were screened at title and abstract (if it was available) using the study inclusion criteria. In 43 ongoing clinical trials no data were mature, one was observational study, thus 20 clinical trials, containing data on 5,890 patients with evaluable ctDNA analyses, were finally eligible for this review. Characteristics of studies are presented in *Table 1*.

Preliminary results from Palbociclib and Circulating Tumor DNA for estrogen receptor-1 gene (ESR1) Mutation Detection (PADA-1) trial demonstrated that ESR1mut detection is uncommon in untreated AI-sensitive, ER+, HER2- metastatic breast cancer patients (detection rate of 2.1% at baseline) and is related to prior AI exposure in the adjuvant setting (4.9% with AI use *vs.* 0% without AI use, Yates Chi2: P=0.009). Remarkably, 1-month use of AI and palbociclib, the first CDK4/6 inhibitor approved as an anticancer regimen, led to undetectable ESR1mut in 13 among the 17 patients with ESR1mut detected at baseline (49).

In the PALOMA-3 study, which compared the combination of palbociclib plus fulvestrant to placebo plus fulvestrant, in patients with HR+, HER2- advanced breast cancer, progressing on prior endocrine therapy, changes in PIK3CA ctDNA dynamics upon 15 days treatment predicted response to targeted therapy in combination with fulvestrant (HR 3.94, 95% CI, 1.61–9.64, log-rank P=0.0013), while ESR1 ctDNA levels change was less predictive on progression free survival (PFS) on palbociclib plus fulvestrant (14,43). Detection of PIK3CA and estrogen receptor-2 gene (ESR2) mutations in plasma ctDNA samples, compared with their detection in archived tissue samples, has been associated with significantly improved PFS and response to abemaciclib (another selective CDK4/6 inhibitor) plus fulvestrant, in postmenopausal women with HR+, HER2- advanced breast cancer, progressing on prior endocrine therapy (62,63).

On the contrary, ctDNA sequencing from 494 patients enrolled in the randomized MONALEESA-2 trial of letrozole ± ribociclib, showed a consistent PFS benefit for the combination of endocrine therapy plus CDK4/6 inhibitor, regardless of the baseline status of ctDNA biomarkers [PIK3CA, tumor protein 53 (TP53), Zinc finger protein 703 (ZNF703)/fibroblast growth factor receptor 1 (FGFR1), ESR1] (15,51). Consistent treatment benefit was observed for fulvestrant and ribociclib, irrespective of baseline ctDNA alteration status [PIK3CA, ESR1, TP53, CDC20 homolog 1 (CDH1), FGFR1/ZNF703/Wolf-Hirschhorn syndrome candidate 1-like 1 (WHSC1L1)] in Phase III MONALEESA-3 study (52,53).

In BELLE-2, which evaluated the combination of the

panPI3 kinase inhibitor buparlisib with fulvestrant in patients with refractory to AI, HR+, HER2- advanced breast cancer, the presence of PIK3CA mutations in ctDNA corresponded to improved PFS in the buparlisib arm (7.0 *vs.* 3.2 months; HR =0.58; 95% CI, 0.41–0.82; 1-sided nominal P=0.001) (54,55). Clinical benefit of the addition of buparlisib to fulvestrant in HR+, HER2- advanced breast cancer patients, with prior use of mTOR inhibitors, has also been observed in the randomized Phase III BELLE-3 trial, even if this benefit was irrespective of PIK3CA status in ctDNA (56). Both, BELLE-2 and BELLE-3 highlighted the potential of PIK3CA mutational status in plasma ctDNA as predictive biomarker for benefit of buparlisib treatment, in this subset of breast cancer patients; whereas the discordance in PIK3CA status between tumor tissue and ctDNA samples (76.7% in BELLE-2 *vs.* 84.8% in BELLE-3) underline the need for an optimal standardized assay.

In a single group assignment, Phase I/II trial the combination of alpelisib and nab-paclitaxel resulted in increased PFS in HER2- advanced breast cancer patients, harbouring ctDNA PIK3CA mutations (66).

A subsidiary analysis of the BOLERO-2 trial on 550 ER+ advanced breast cancer patients, demonstrated that the addition of everolimus to exemestane prolonged PFS, irrespective of ctDNA PIK3CA mutation status (HR =0.43 and 0.37 respectively) (17,69).

Furthermore, the ongoing POSEIDON trial and Neratinib HER Mutation Basket Study (SUMMIT) support the predictive value of early evaluation of ctDNA changes, before radiologic treatment response (58,59).

The translational sub-study of the ongoing I-SPY 2 trial demonstrated the significance of serial monitoring of ctDNA in predicting response to neo-adjuvant treatment (61). ctDNA analysis of the NeoALTTO trial demonstrated that the detection of PIK3CA and/or TP53 mutations, in the baseline (before neo-adjuvant therapy) plasma sample was correlated with lower rates of pathological complete response, whereas persistent ctDNA detection both at baseline and after 14 days of neo-adjuvant therapy was significantly associated with the lowest rate of pathological complete response (71,72).

In open-label WJOG6110B/ELTOP trial, whereas patients with HER2+ advanced breast cancer, were randomized to receive either lapatinib and capecitabine or trastuzumab and capecitabine, PIK3CA mutations in both tissue and ctDNA samples associated with shorter PFS, regardless of the treatment arm (57). The presence

Table 1 Characteristics of clinical trials incorporating ctDNA-based predictive biomarkers

Clinical trial (name/ID number)	Status	Design	Intervention model	Setting	Population characteristics	Intervention vs. Control arm	Enrollment (biomarker analysis)	Patients (%) with detectable ctDNA	Endpoints	ctDNA sequencing technique	Concordance of tissue and plasma samples	Results	References
PADA-1/ NCT03079011	Active, not recruiting	Open label, randomized, phase III	Sequential assignment	1st line (metastatic setting)	ER+, HER2-, postmenopausal female, ECOG PS: 0-2	Palbociclib + AI vs. palbociclib + fulvestrant	803	17/803 (2.1%)	Safety, efficacy	ddPCR-based assay		76.47% of patients had undetectable ESR1m after 1 month of palbociclib + AI therapy	(49)
SOLAR-1/ NCT02437318	Active, not recruiting	Triple blind, randomized (1:1), phase III	Parallel assignment	2nd line (metastatic setting)	PIK3CAm, HR+, HER2-, male or postmenopausal female, 1 prior line of endocrine therapy, ECOG PS: 0-1	Alpelisib + fulvestrant vs. placebo + fulvestrant	549	186/549 (33.87%)	PFS	Assay developed by Qiagen	94.7%	PFS of 3.7 months for tissue PIK3CAm and of 10.9 months for ctDNA PIK3CAm. Treatment benefit, with the combination of alpelisib and fulvestrant, in PFS for patients with ctDNA PIK3CAm, irrespective of prior treatment for advanced breast cancer and/or prior CDK4/6 inhibitors use	(20,50)
MONALEESA-2/ NCT01958021	Active, not recruiting	Double blind, randomized (1:1), phase III	Parallel assignment	1st line (metastatic setting)	HR+, HER2-, postmenopausal female, ECOG PS: 0-1	Ribociclib + letrozole vs. placebo + letrozole	494	427/494 (86%)	PFS	NGS		≥1 ctDNA genomic alteration: PIK3CA (33%), TP53 (12%), ZNF703/FGFR1 (5%), ESR1 (4%), and in genes involved in RTK signaling (12%). Treatment benefit, with the combination of ribociclib and letrozole, irrespective of ctDNA genetic alterations at baseline	(15,51)
MONALEESA-3/ NCT02422615	Active, not recruiting	Double blind, randomized (2:1), phase III	Parallel assignment	≤2nd line (metastatic setting)	HR+, HER2-, postmenopausal female, ≤1 prior line of endocrine therapy ECOG PS: 0-1	Ribociclib + fulvestrant vs. placebo + fulvestrant	600	124/600 (20.66%) for PIK3CAm	PFS	NGS		ctDNA genomic alterations: PIK3CA (35%), ESR1 (14%), TP53 (18%), CDH1 (12%), FGFR1/ZNF703/WHSC1L1 (11%), CCC genes (16%), genes involved in RTK signaling (20%) and genes involved in the MAPK pathway (10%). Treatment benefit, with the combination of ribociclib and fulvestrant, irrespective of ctDNA genetic alterations; shorter PFS was correlated with altered genetic status	(52,53)
BELLE-2/ NCT01610284	Completed	Double blind, randomized (1:1), phase III	Parallel assignment	2nd line (metastatic setting)	HR+, HER2-, postmenopausal female, AI-refractory disease	Buparlisib + fulvestrant vs. placebo + fulvestrant	587	200/587 (34%)	PFS	Sanger sequencing	77%	64 of 307 (21%) patients with PIK3CAwt tumour tissue had PIK3CAm ctDNA, indicating evolution between initial diagnosis and the present time. ctDNA PIK3CAm corresponded to improved median PFS in the buparlisib arm (7.0 vs. 3.2 months; HR =0.58; 95% CI, 0.41-0.82; 1-sided nominal P=0.001)	(54,55)
BELLE-3/ NCT01633060	Terminated	Double blind, randomized (2:1), phase III	Parallel assignment	≥2nd line (metastatic setting)	HR+, HER2-, postmenopausal female, prior treatment with AI, progression to the combination of mTORi and endocrine therapy, ECOG PS: 0-2	Buparlisib + fulvestrant vs. placebo + fulvestrant	348	135/348 (39%)	PFS	Inostics BEAMing assay	83%	Treatment benefit, with the combination of buparlisib and fulvestrant, irrespective of ctDNA PIK3CA mutational status (PFS of 4.2 vs. 1.6 months; HR =0.46; 95% CI 0.29-0.73; P=0.00031 for PIK3CAm and 3.9 vs. 2.7 months; HR =0.73; 95% CI, 0.53-1.00; P=0.026 for PIK3CAwt)	(56)

Table 1 (continued)

Table 1 (continued)

Clinical trial (name/ID number)	Status	Design	Intervention model	Setting	Population characteristics	Intervention vs. Control arm	Enrollment (biomarker analysis)	Patients (%) with detectable ctDNA	Endpoints	ctDNA sequencing technique	Concordance of tissue and plasma samples	Results	References
PALOMA-3/ NCT01942135	Active, not recruiting	Double blind, randomized (2:1), phase III	Parallel assignment	2nd line (metastatic setting)	HR+, HER2-, female of any menopausal status, progression to prior adjuvant or metastatic endocrine therapy, ECOG PS: 0-1	Palbociclib + fulvestrant vs. placebo + fulvestrant	455	100/455 (22%) for PIK3CAm and 114/445 (25.6%) for ESR1m	PFS	ddPCR-based assay		Both PIK3CA mutant copies and wild-type allele and ESR1 mutant copies and wild-type allele were significantly lower in the Palbociclib treatment group (Wilcoxon signed-rank test, P<0.0001). Early ctDNA PIK3CA dynamics (after 2 weeks of therapy) were predictive on response to palbociclib and fulvestrant	(14,43)
WJOG6110B/ ELTOP/ UMIN000005219	Completed	Open label, randomized (1:1), phase II	Parallel assignment	≥1st line (metastatic setting)	HER2+, female, prior use of taxanes, progression on trastuzumab-containing regimens, ECOG PS: 0-2	Lapatinib + capecitabine vs. trastuzumab + capecitabine	35	8/35 (23%)	PFS	ddPCR-based assay	85%	PIK3CAm in both tissue and plasma samples correlated with shorter PFS, irrespective of the treatment arm. Especially, for ctDNA PIK3CAwt PFS was 8.2 and 4.9 months for the lapatinib arm and for the trastuzumab arm, respectively (HR =0.38; 95% CI, 0.16-0.93; P=0.035), whereas for ctDNA PIK3CAm PFS was 4.1 and 6.1 months for the lapatinib arm and for the trastuzumab arm, respectively (HR =0.60; 95% CI, 0.11-3.13; P=0.54)	(57)
POSEIDON/ NCT02285179	Recruiting	Double blind, randomized (1:1), phase Ib (3+3 design)	Parallel assignment	≥2nd line (metastatic setting)	HR+, HER2-, female of any menopausal status, prior endocrine therapy, ≤5 chemotherapy lines in the metastatic setting	Taselisib + tamoxifen vs. placebo + tamoxifen	22		PFS	dPCR/tagged amplicon deep-sequencing		ctDNA PIK3CA dynamics were predictive on response to taselisib and tamoxifen, before radiologic treatment response	(58)
SUMMIT/ NCT03433274	Recruiting	Open label, non-randomized, phase II	Single group assignment	BASKET trial: colon, lung, breast, bladder cancer, fibromellar carcinoma, any line of therapy	HER2+ or EGFR+ or HER3+	Neratinib	14	11/14 (78.57%)	Clinical benefit rate	70-gene digital sequencing assay	93.5%	Early ctDNA HER2 dynamics were predictive on response to neratinib; ctDNA HER2mut frequency decreased in 9 of 11 paired samples, at week 4, followed by an increase upon radiographical disease progression at week 8	(59)
BEECH/ NCT01625286	Active, not recruiting	Double blind, randomized (1:1), phase I/II	Parallel assignment	1st line (metastatic setting)	ER+, HER2-, WHO PS: 0-1	Capivasertib + paclitaxel vs. placebo + paclitaxel	148		Dose-limiting toxicity events, PFS	ddPCR-based assay for ctDNA quantification. Roche cobas PIK3CA assay for PIK3CAmut identification		Early ctDNA dynamics were predictive on PFS irrespective of treatment arm (median PFS was 11.1 months in patients with decreased ctDNA levels at week 4, and 6.4 months in patients with higher ctDNA levels; HR =0.20; 95% CI, 0.083-0.50; P<0.0001)	(60)
I-SPY 2/ NCT01042379	Recruiting	Open label, randomized, phase II (adaptive design)	Parallel assignment	Locally advanced breast cancer (stage II, III), neoadjuvant setting	Any tumor ER/PR/HER2 status, female, no prior cytotoxic regimens, ECOG PS: 0-1	AMG 386 ± trastuzumab/ AMG 479 + metformin/MK-2206 ± trastuzumab/T-DM1 + pertuzumab/ganetespib/ ABT-888/neratinib/PLX3397/ pembrolizumab/talazoparib + irinotecan/patritumab ± trastuzumab/SGN-LIV1A/ durvalumab + olaparib/SD-101 + pembrolizumab/tucatinib vs. standard therapy/pertuzumab + trastuzumab	84		pCR after the use of experimental agents	Mutational profiles derived from pretreatment tumor biopsy and germline DNA whole exome sequencing were used to design personalized assays		Early ctDNA dynamics were predictive on response to neoadjuvant treatment	(61)

Table 1 (continued)

Table 1 (continued)

Clinical trial (name/ID number)	Status	Design	Intervention model	Setting	Population characteristics	Intervention vs. Control arm	Enrollment (biomarker analysis)	Patients (%) with detectable ctDNA	Endpoints	ctDNA sequencing technique	Concordance of tissue and plasma samples	Results	References
MONARCH 2/ NCT02107703	Recruiting	Double blind, randomized (2:1), phase III	Parallel assignment	2nd line (metastatic setting)	HR+, HER2-, postmenopausal female, ECOG PS: 0-1	Abemaciclib + fulvestrant vs. placebo + fulvestrant	334	96/238 (40.3%) for PIK3CAm and 190/295 (64.4%) for ESR1m	PFS	ddPCR-based assay	62.8% for PIK3CAm and 37.1% for ESR1m	ctDNA mutational status associates with improved PFS and response to abemaciclib and fulvestrant arm. For ctDNA PIK3CAm PFS was 15 and 5.7 months for the abemaciclib arm and for the control arm, respectively (HR =0.46; 95% CI, 0.27-0.78), whereas for ctDNA ESR1m PFS was 21.9 and 10.3 months for the abemaciclib arm and for the control arm, respectively (HR =0.49; 95% CI, 0.33-0.73)	(62,63)
LOTUS/ NCT02162719	Active, not recruiting	Double blind, randomized (1:1), phase II	Parallel assignment	1st line (metastatic setting)	TNBC, female of any menopausal status, ECOG PS: 0-1	lpatasertib + paclitaxel vs. placebo + paclitaxel	88		PFS	FoundationACT assay (plasma samples) and FoundationOne genomic profiling (tumor tissue samples)		ctDNA dynamics were predictive on PFS and objective response irrespective of treatment arm	(64,65)
NCT02379247	Active, not recruiting	Open label, non-randomized, phase I/II	Single group assignment	≥2nd line (metastatic setting)	HER2-, female, prior chemotherapy for metastatic disease, ECOG PS ≥2	Alpelisib + nab-paclitaxel	42	17/42 (40%)	Recommended phase II dose, objective response rate, PFS	NGS	70%	PFS of 13 months for ctDNA PIK3CAm and 7 months for ctDNA PIK3CAwt (HR =0.39; P=0.03)	(66)
INSPIRE/ NCT02644369	Active, not recruiting	Open label, non-randomized, phase II	Single group assignment	BASKET trial: squamous cell Ca of the head and neck, TNBC, high-grade serous ovarian cancer, melanoma, mixed advanced solid tumors, any line of therapy	Triple negative, male or female, ECOG PS: 0-1	Pembrolizumab	20 (TNBC)		Changes in genomic and immune biomarkers that will be measured in blood and tumor pre-treatment, on-treatment and at progression	Single cell suspensions were pooled for exome/ RNA sequencing, flow cytometry for immunophenotyping		Early ctDNA dynamics were predictive on PFS, OS and overall clinical RR	(67,68)
BOLERO-2/ NCT00863655	Completed	Double blind, randomized (2:1), phase III	Parallel assignment	2nd line (metastatic setting)	ER+, postmenopausal female, disease refractory to NSAI, recurrence or progression on or after the last systemic therapy	Everolimus + exemestane vs. placebo + exemestane	550	238/550 (43.3%)	PFS	ddPCR-based assay	70.4%	Treatment benefit, with the combination of everolimus and exemestane, irrespective of ctDNA PIK3CA status (HR =0.43 for PIK3CAwt tumors and 0.37 for PIK3CAm tumors)	(17,69)
BLTN-1c/ NCT02361112	Completed	Open label, non-randomized, phase I	Single group assignment	2nd line (metastatic setting)	HER2+, male or female of any menopausal status, no previous treatment of capecitabine during the past 1 year, ECOG PS: 0-1	Pyrotinib + capecitabine	28		MTD			Median PFS of 15.8 months for ≥2 ctDNA genetic alterations of HER2, PI3K/AKT/mTOR pathway and TP53 and of 26.2 months for ≤1 ctDNA genetic alteration (P=0.006)	(70)
Neo ALLTO/ NCT00553358	Active, not recruiting	Open label, randomized (2:2:2), phase III	Parallel assignment	Primary invasive breast cancer, neoadjuvant setting	HER2+, female, invasive breast cancer >2cm diameter, ECOG PS: 0-1	Lapatinib + paclitaxel + trastuzumab vs. paclitaxel + trastuzumab	124	94/124 (75.8%) with PIK3CAm and 30/124 (24.2%) with TP53m	Number of participants with pCR at the time of surgery	NGS		ctDNA PIK3CAm and/or TP53m detection at baseline and at serial plasma samples was predictive of low rates of pathological response	(71,72)
MONALEESA-7/ NCT02278120	Active, not recruiting	Double blind, randomized (1:1), phase III	Parallel assignment	1st line (metastatic setting)	ER+ and/or PR+, HER2-, premenopausal or perimenopausal female, ECOG PS: ≤1	Ribociclib + tamoxifen/letrozole/ anastrozole + goserelin vs. placebo + tamoxifen/letrozole/ anastrozole + goserelin	565	489/565 (86.54%)	PFS	NGS		Treatment benefit, with the combination of ribociclib and NSAI or tamoxifen and goserelin, irrespective of ctDNA mutational status at baseline	(73,74)

AI, aromatase inhibitor; AKT, protein kinase B; BEAMing, beads, emulsion, amplification, magnetics; CCC, cell cycle-related; CDH1, CDC20 homolog 1; CDK4/6, cyclin-dependent kinase 4 and 6; CI, confidence interval; ctDNA, circulating tumor DNA; ddPCR, droplet digital polymerase chain reaction; dPCR, digital polymerase chain reaction; ECOG PS, Eastern Cooperative Oncology Group Performance Status; EGFR, epidermal growth factor receptor; ESR1, estrogen receptor-1 gene; FGFR1, fibroblast growth factor receptor 1; HER2, human epidermal growth factor receptor 2; HER3, human epidermal growth factor receptor 3; HR =, hazard ratio; HR, hormone receptors; m, mutated; MAPK, mitogen-activated protein kinase; mTOR, mammalian target of rapamycin; NGS, next generation sequencing; NSAI, non-steroidal aromatase inhibitor; OS, overall survival; pCR, pathologic complete response; PFS, progression free survival; PI3K, phosphoinositide-3-kinase; PIK3CA, phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha; PR, progesterone receptors; RR, response rate; RTK, receptor tyrosine kinase; TNBC, triple negative breast cancer; TP53, tumor protein 53; WHSC1L1, Wolf-Hirschhorn syndrome candidate 1-like 1; wt, wild type; ZNF703, zinc finger protein 703.

of concomitant genetic alterations of HER2, PI3K/protein kinase B (AKT)/mTOR pathway and TP53 in ctDNA analysis was significantly correlated with worse PFS, compared to ≤ 1 genetic alteration, in the open-label, Phase I BLTN-Ic trial, of the combination of pyrotinib plus capecitabine in HER2+ advanced breast cancer patients (70).

Dynamic ctDNA analysis of plasma samples from Phase I/II trial BEECH, whereas patients with ER+ metastatic breast cancer randomized to either paclitaxel plus AKT inhibitor capivasertib or paclitaxel plus placebo, predicted long-term outcome (PFS of 11.1 months in patients with suppressed ctDNA at 21 days *vs.* 6.4 months in patients with high levels of ctDNA, HR =0.20; 95% CI, 0.083–0.50; $P < 0.0001$), thus serving as a surrogate for PFS (60).

The double-blind, Phase II LOTUS trial, comparing the combination of ipatasertib plus paclitaxel with paclitaxel monotherapy in triple negative advanced breast cancer patients, demonstrated the predictive value of dynamic evaluation of ctDNA in evaluating both objective response and PFS, consistently in both arms (64,65).

As part of the Phase II, INSPIRE basket trial, a secondary analysis of ctDNA at baseline and before the initiation of 3rd cycle of the single-agent immune checkpoint inhibitor pembrolizumab in 10 triple negative metastatic breast cancer patients, strongly correlated with PFS, overall survival (OS) and overall clinical response rate (ORR) (67,68).

In the 1st comprehensive genomic analysis of ctDNA of premenopausal patients with ER+ and/or progesterone receptors (PR)+, HER2- advanced breast cancer, the combination of the CDK 4/6 inhibitor ribociclib and non-steroidal aromatase inhibitor (NSAI) or tamoxifen and goserelin resulted in PFS benefit, irrespective of the baseline genetic landscape status (73,74).

Based on the results of SOLAR-1, Food and Drug Administration (FDA) approved, on May 24, 2019, the use of PIK3CA selective inhibitor alpelisib in combination with fulvestrant for the treatment of men and postmenopausal women, with HR+, HER2-, PIK3CA-mutated advanced breast cancer, following disease progression on or after an endocrine-based regimen. In particular, the combination of alpelisib and fulvestrant resulted in significant prolongation of PFS (HR 0.55; 95% CI, 0.39–0.79; $n=186$) in patients with ctDNA PIK3CA mutant status. Concurrently, FDA also approved the companion diagnostic test PIK3CA Rotor-Gene Q real-time polymerase chain reaction (RGQ PCR) kit to detect the PIK3CA mutation in a tissue and/or a liquid biopsy. Thus, the assessment of PIK3CA mutations

in ctDNA became the first liquid biopsy to be used in the clinical setting for breast cancer patients (20,50).

Discussion

Research into understanding breast cancer's complexity, both at cellular and molecular level, and development of targeted therapies underline the urgent need of conducting novel biomarker-driven clinical trials, with the ultimate goal of optimizing disease management. The traditional process of drug research and development, where investigational drugs were evaluated for safety and optimal dosing scheme in Phase I, for early signs of efficacy in Phase II, and for confirmation of efficacy, effectiveness and safety in Phase III, gradually fades out. Over the last decade, novel clinical trial designs have found their way into clinical research, in order not only to streamline but also to expedite drug development (75).

Master Protocol (MAPs) use a single, biomarker-driven, trial design and protocol to concurrently evaluate multiple drugs and/or diseases, and include (I) basket trials, which enrol patients based on the presence of a specific biomarker (e.g., mutation), regardless of histology, to identify efficacy of a biomarker-specific, thus targeted, therapy, and (II) umbrella and (III) adaptive platform trials, where patients who share the same cancer histology are allocated to different arms, based on their biomarker status (e.g., mutation), in order to evaluate new investigational agents matches to biomarker-derived cohorts (75). The main difference between umbrella and platform trials is that the last incorporate more adaptations, during the trial, based on efficacy results of interim analyses, by permitting in a flexible way the addition or exclusion of new treatment modalities (75).

Establishing biomarker-stratified clinical-trial design frameworks in the context of spatial and temporal heterogeneity is challenging because the traditional use of archival tissue samples may not be reflective of the dynamic genomic status of the tumor, especially in the metastatic setting (26,30). Such hurdle could potentially be overcome through the incorporation of ctDNA analyses, for the longitudinal evaluation of predictive biomarkers. Overall, results emerged from the clinical trials presented in this review highlight the importance of dynamic ctDNA monitoring in the era of precision medicine; measurement of ctDNA provides representative data of spatiotemporal tracking of mutational landscape of both primary tumour and metastases, thus serving as a sensitive biomarker

for both monitoring tumor progression and evaluating treatment response (37,76).

ctDNA dynamics could serve as a predictive biomarker independent of the histology. Indeed, the I-SPY 2 trial reported that early ctDNA dynamics could predict response to neo-adjuvant treatment (61), whereas in the basket trial SUMMIT a decrease in ctDNA HER2 mutation variant allele frequency during treatment with the pan-HER inhibitor neratinib was followed by an increase upon radiographically proven progression (59).

Moreover, in the SUMMIT trial a number of genetic aberrations were also identified co-occurring with HER2 mutations, which highlighted the acquisition of secondary resistance to targeted therapy due to clonal evolution (59). Also, in the PALOMA-3 trial, which enrolled patients with ER+, HER2- advanced, previously progressed on endocrine treatment, breast cancer, early ctDNA dynamics of truncal mutations in PIK3CA predicted sensitivity to the CDK4/6 inhibitor palbociclib; on the contrary, serial ctDNA monitoring of the, commonly sub clonal, ESR1 mutations failed to predict clinical outcome (43). Taken together, these results address the importance of assessing tumor's genetic heterogeneity and clonal evolution in real time, with minimally invasive ways, like ctDNA analyses, in order not only to predict response, but also to rapidly identify acquired resistance to targeted therapies in breast cancer.

Regarding breast cancer detection, at present mammography remains the gold standard screening method, whereas gene expression profiling tests are used to stratify patients regarding recurrence (77). The potential of ctDNA analysis both as a consistent detection biomarker and as an accurate predictor of breast cancer recurrence risk needs to be further investigated, given the data scarcity and the lack of standardized analytical methods.

Nowadays, digital PCR (dPCR)- and next generation sequencing (NGS)-based methods are most frequently used to detect ctDNA in a background of wildtype DNA. Despite the wide variety in the number of available technologies for ctDNA analysis, only 2 companion diagnostic kits are FDA-approved: cobas EGFR Mutations Test v2 for detection of epidermal growth factor receptor (EGFR) mutations in non-small cell lung cancer (NSCLC), and theascreen® PIK3CA RGQ PCR Kit for detection of PIK3CA mutations in advanced or metastatic breast cancer (78).

Standardization challenges for integration of ctDNA analysis into routine clinical practice include: (I) biological variability (thus tumor heterogeneity), (II) pre-analytical variability (e.g., specialized collecting tubes to prevent

leukocyte lysis, optimal time period between blood-draw and sample processing, centrifugation conditions, quantification methods), and (III) analytical variability (an ideal technology should be accurate, highly sensitive and specific, robust, and cost-effective) (76). To accelerate the development and establishment of liquid biopsies in clinical practice, consortium of researchers from academia, industry, regulatory agencies and public, both in United States (BloodPAC) (79), and Europe (Cancer-ID) (80) have been developed.

Conclusions

In conclusion, it can be said that the majority of published results from both recent and ongoing biomarker-driven clinical trials in breast cancer patients seem to concur that ctDNA profiling may significantly correlate with response to targeted therapies, thus indicating its potential as a non-invasive predictive biomarker, both in adjuvant, neo-adjuvant, and metastatic setting.

The incorporation of ctDNA analysis into sophisticated, biomarker-driven clinical trials, with adequate statistical power and sufficient sample sizes, remains the most reliable way to demonstrate not only the analytical and clinical validity, but also the clinical utility of ctDNA as liquid biopsy, in tailoring decision-making in breast cancer patients.

Acknowledgments

Funding: None.

Footnote

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <http://dx.doi.org/10.21037/atm-20-1175>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/atm-20-1175>). ML reports personal fees from Roche, Astra Zeneca, Astellas, MSD, Ipsen, Jansen, BMS, GSK, outside the submitted work; Spouse is Pfizer employee. KK reports personal fees from Roche, BMS, MSD, IPSEN, outside the submitted work. ET reports grants, personal fees and non-financial support from JANSSEN, personal fees from CELGENE, grants, personal fees and non-financial support from TAKEDA, grants,

personal fees and non-financial support from AMGEN, grants, personal fees and non-financial support from GENESIS PHARMA, personal fees from BMS, outside the submitted work. MAD reports personal fees from Janssen, Celgene, Takeda, Amgen, Genesis Pharma, BMS, outside the submitted work. FZ reports personal fees from Astra Zeneca, Daiichi, Eli-Lilly, Merck, Novartis, Pfizer, Roche, outside the submitted work. OF has no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-E386.
2. Bertucci F, Houlgatte R, Benziane A, et al. Gene expression profiling of primary breast carcinomas using arrays of candidate genes. *Hum Mol Genet* 2000;9:2981-91.
3. Stephens PJ, Tarpey PS, Davies H, et al. The landscape of cancer genes and mutational processes in breast cancer. *Nature* 2012;486:400-4.
4. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* 2012;490:61-70.
5. Cameron D, Piccart-Gebhart MJ, Gelber RD, et al. 11 years' follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive early breast cancer: final analysis of the HERceptin Adjuvant (HERA) trial. *Lancet* 2017;389:1195-205.
6. Gianni L, Eiermann W, Semiglazov V, et al. Neoadjuvant and adjuvant trastuzumab in patients with HER2-positive locally advanced breast cancer (NOAH): follow-up of a randomised controlled superiority trial with a parallel HER2-negative cohort. *Lancet Oncol* 2014;15:640-7.
7. Slamon DJ, Leyland-Jones B, Shak S, et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 2001;344:783-92.
8. von Minckwitz G, Procter M, de Azambuja E, et al. Adjuvant Pertuzumab and Trastuzumab in Early HER2-Positive Breast Cancer. *N Engl J Med* 2017;377:122-31.
9. Gianni L, Pienkowski T, Im YH, et al. 5-year analysis of neoadjuvant pertuzumab and trastuzumab in patients with locally advanced, inflammatory, or early-stage HER2-positive breast cancer (NeoSphere): a multicentre, open-label, phase 2 randomised trial. *Lancet Oncol* 2016;17:791-800.
10. Swain SM, Kim SB, Cortés J, et al. Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA study): overall survival results from a randomised, double-blind, placebo-controlled, phase 3 study. *Lancet Oncol* 2013;14:461-71.
11. Krop IE, Kim SB, Martin AG, et al. Trastuzumab emtansine versus treatment of physician's choice in patients with previously treated HER2-positive metastatic breast cancer (TH3RESA): final overall survival results from a randomised open-label phase 3 trial. *Lancet Oncol* 2017;18:743-54.
12. Geyer CE, Forster J, Lindquist D, et al. Lapatinib plus capecitabine for HER2-positive advanced breast cancer. *N Engl J Med* 2006;355:2733-43.
13. Martin M, Holmes FA, Ejlertsen B, et al. Neratinib after trastuzumab-based adjuvant therapy in HER2-positive breast cancer (ExteNET): 5-year analysis of a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2017;18:1688-700.
14. Cristofanilli M, Turner NC, Bondarenko I, et al. Fulvestrant plus palbociclib versus fulvestrant plus placebo for treatment of hormone-receptor-positive, HER2-negative metastatic breast cancer that progressed on previous endocrine therapy (PALOMA-3): final analysis of the multicentre, double-blind, phase 3 randomised controlled trial. *Lancet Oncol* 2016;17:425-39.
15. Hortobagyi GN, Stemmer SM, Burris HA, et al. Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer. *N Engl J Med* 2016;375:1738-48.
16. Goetz MP, Toi M, Campone M, et al. MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer. *J Clin Oncol* 2017;35:3638-46.
17. Baselga J, Campone M, Piccart M, et al. Everolimus in postmenopausal hormone-receptor-positive advanced

- breast cancer. *N Engl J Med* 2012;366:520-9.
18. Robson M, Im SA, Senkus E, et al. Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation. *N Engl J Med* 2017;377:523-33.
 19. Litton JK, Rugo HS, Ettl J, et al. Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation. *N Engl J Med* 2018;379:753-63.
 20. André F, Ciruelos E, Rubovszky G, et al. Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J Med* 2019;380:1929-40.
 21. Cardoso F, Spence D, Mertz S, et al. Global analysis of advanced/metastatic breast cancer: Decade report (2005-2015). *Breast* 2018;39:131-8.
 22. Howlader N, Noone AM, Krapcho M, et al. editors. SEER Cancer Statistics Review, 1975-2016, National Cancer Institute. Bethesda, MD, based on November 2018 SEER data submission, posted to the SEER web site, April 2019.
 23. Wang R, Li X, Zhang H, et al. Cell-free circulating tumor DNA analysis for breast cancer and its clinical utilization as a biomarker. *Oncotarget* 2017;8:75742-55.
 24. National Comprehensive Cancer Network, Inc. 2018. Available online: <https://www.nccn.org/>
 25. Cardoso F, Harbeck N, Fallowfield L, et al. Locally recurrent or metastatic breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2012;23:vii11-vii19.
 26. Gerlinger M, Rowan AJ, Horswell S, et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med* 2012;366:883-92.
 27. Yates LR, Gerstung M, Knappskog S, et al. Subclonal diversification of primary breast cancer revealed by multiregion sequencing. *Nat Med* 2015;21:751-9.
 28. Nik-Zainal S, Van Loo P, Wedge DC, et al. The life history of 21 breast cancers. *Cell* 2012;149:994-1007.
 29. Gundem G, Van Loo P, Kremeyer B, et al. The evolutionary history of lethal metastatic prostate cancer. *Nature* 2015;520:353-7.
 30. Lipinski KA, Barber LJ, Davies MN, et al. Cancer Evolution and the Limits of Predictability in Precision Cancer Medicine. *Trends Cancer* 2016;2:49-63.
 31. Ziegler A, Zangemeister-Wittke U, Stahel RA. Circulating DNA: a new diagnostic gold mine? *Cancer Treat Rev* 2002;28:255-71.
 32. Han SX, Zhou X, Sui X, et al. Serum dickkopf-1 is a novel serological biomarker for the diagnosis and prognosis of pancreatic cancer. *Oncotarget* 2015;6:19907-17.
 33. Kosaka N, Iguchi H, Ochiya T. Circulating microRNA in body fluid: a new potential biomarker for cancer diagnosis and prognosis. *Cancer Sci* 2010;101:2087-92.
 34. Shah SP, Morin RD, Khattra J, et al. Mutational evolution in a lobular breast tumour profiled at single nucleotide resolution. *Nature* 2009;461:809-13.
 35. Ellsworth RE, Blackburn HL, Shriver CD, et al. Molecular heterogeneity in breast cancer: State of the science and implications for patient care. *Semin Cell Dev Biol* 2017;64:65-72.
 36. Zardavas D, Irrthum A, Swanton C, et al. Clinical management of breast cancer heterogeneity. *Nat Rev Clin Oncol* 2015;12:381-94.
 37. Dawson SJ, Tsui DW, Murtaza M, et al. Analysis of circulating tumor DNA to monitor metastatic breast cancer. *N Engl J Med* 2013;368:1199-209.
 38. Schiavon G, Hrebien S, Garcia-Murillas I, et al. Analysis of ESR1 mutation in circulating tumor DNA demonstrates evolution during therapy for metastatic breast cancer. *Sci Transl Med* 2015;7:313ra182.
 39. Fribbens C, Garcia Murillas I, Beaney M, et al. Tracking evolution of aromatase inhibitor resistance with circulating tumour DNA analysis in metastatic breast cancer. *Ann Oncol* 2018;29:145-53.
 40. Wang P, Bahreini A, Gyanchandani R, et al. Sensitive Detection of Mono- and Polyclonal ESR1 Mutations in Primary Tumors, Metastatic Lesions, and Cell-Free DNA of Breast Cancer Patients. *Clin Cancer Res* 2016;22:1130-7.
 41. Beijs N, Sieuwerts AM, Kraan J, et al. Estrogen receptor mutations and splice variants determined in liquid biopsies from metastatic breast cancer patients. *Mol Oncol* 2018;12:48-57.
 42. Murtaza M, Dawson SJ, Pogrebniak K, et al. Multifocal clonal evolution characterized using circulating tumour DNA in a case of metastatic breast cancer. *Nat Commun* 2015;6:8760.
 43. O'Leary B, Hrebien S, Morden JP, et al. Early circulating tumor DNA dynamics and clonal selection with palbociclib and fulvestrant for breast cancer. *Nat Commun* 2018;9:896.
 44. Chung JH, Pavlick D, Hartmaier R, et al. Hybrid capture-based genomic profiling of circulating tumor DNA from patients with estrogen receptor-positive metastatic breast cancer. *Ann Oncol* 2017;28:2866-73.
 45. De Mattos-Arruda L, Weigelt B, Cortes J, et al. Capturing intra-tumor genetic heterogeneity by de novo mutation profiling of circulating cell-free tumor DNA: a proof-of-principle. *Ann Oncol* 2014;25:1729-35.
 46. Ma F, Zhu W, Guan Y, et al. ctDNA dynamics: a novel indicator to track resistance in metastatic breast

- cancer treated with anti-HER2 therapy. *Oncotarget* 2016;7:66020-31.
47. Guttery DS, Page K, Hills A, et al. Noninvasive detection of activating estrogen receptor 1 (ESR1) mutations in estrogen receptor-positive metastatic breast cancer. *Clin Chem* 2015;61:974-82.
 48. Jansen MP, Martens JW, HelmiJR JC, et al. Cell-free DNA mutations as biomarkers in breast cancer patients receiving tamoxifen. *Oncotarget* 2016;7:43412-8.
 49. Bidard FC, Pistilli B, Dalenc F, et al. Circulating ESR1 mutation detection rate and early decrease under first line aromatase inhibitor and palbociclib in the PADA-1 trial (UCBG-GINECO) [abstract]. In: Proceedings of the 2018 San Antonio Breast Cancer Symposium; 2018 Dec 4-8; San Antonio, TX. Philadelphia (PA): AACR; *Cancer Res* 2019;79:Abstract nr PD2-06.
 50. Juric D, Ciruelos E, Rubovszky G, et al. Alpelisib + fulvestrant for advanced breast cancer: Subgroup analyses from the phase III SOLAR-1 trial [abstract]. In: Proceedings of the 2018 San Antonio Breast Cancer Symposium; 2018 Dec 4-8; San Antonio, TX. Philadelphia (PA): AACR; *Cancer Res* 2019;79:Abstract nr GS3-08.
 51. Hortobagyi GN, Stemmer S, Campone M, et al. First-line ribociclib + letrozole in hormone receptor-positive, HER2-negative advanced breast cancer: Efficacy by baseline circulating tumor DNA alterations in MONALEESA-2 [abstract]. In: Proceedings of the 2017 San Antonio Breast Cancer Symposium; 2017 Dec 5-9; San Antonio, TX. Philadelphia (PA): AACR; *Cancer Res* 2018;78:Abstract nr PD4-06.
 52. Slamon DJ, Neven P, Chia S, et al. Phase III Randomized Study of Ribociclib and Fulvestrant in Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer: MONALEESA-3. *J Clin Oncol* 2018;36:2465-72.
 53. Neven P, Petrakova K, Val Bianchi G, et al. Biomarker analysis by baseline circulating tumor DNA alterations in the MONALEESA-3 study [abstract]. In: Proceedings of the 2018 San Antonio Breast Cancer Symposium; 2018 Dec 4-8; San Antonio, TX. Philadelphia (PA): AACR; *Cancer Res* 2019;79:Abstract nr PD2-05.
 54. Baselga J, Im SA, Iwata H, et al. Buparlisib plus fulvestrant versus placebo plus fulvestrant in postmenopausal, hormone receptor-positive, HER2-negative, advanced breast cancer (BELLE-2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2017;18:904-16.
 55. Campone M, Im SA, Iwata H, et al. Buparlisib plus fulvestrant versus placebo plus fulvestrant for postmenopausal, hormone receptor-positive, human epidermal growth factor receptor 2-negative, advanced breast cancer: Overall survival results from BELLE-2. *Eur J Cancer* 2018;103:147-54.
 56. Di Leo A, Johnston S, Lee KS, et al. Buparlisib plus fulvestrant in postmenopausal women with hormone-receptor-positive, HER2-negative, advanced breast cancer progressing on or after mTOR inhibition (BELLE-3): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2018;19:87-100.
 57. Takano T, Tsurutani J, Takahashi M, et al. A randomized phase II trial of trastuzumab plus capecitabine versus lapatinib plus capecitabine in patients with HER2-positive metastatic breast cancer previously treated with trastuzumab and taxanes: WJOG6110B/ELTOP. *Breast* 2018;40:67-75.
 58. Baird RD, Van Rossum A, Oliveira M, et al. POSEIDON trial phase 1b results: Safety and preliminary efficacy of the isoform selective PI3K inhibitor taselisib (GDC-0032) combined with tamoxifen in hormone receptor (HR) positive, HER2-negative metastatic breast cancer (MBC) patients (pts) - including response monitoring by plasma circulating tumor (ct) DNA. *J Clin Oncol* 2016;34:2520.
 59. Ma CX, Bose R, Gao F, et al. Neratinib Efficacy and Circulating Tumor DNA Detection of HER2 Mutations in HER2 Nonamplified Metastatic Breast Cancer. *Clin Cancer Res* 2017;23:5687-95.
 60. Hrebien S, Citi V, Garcia-Murillas I, et al. Early ctDNA dynamics as a surrogate for progression-free survival in advanced breast cancer in the BEECH trial. *Ann Oncol* 2019;30:945-52.
 61. Magbanua MJM, Brown-Swigart L, Hirst GL, et al. Personalized serial circulating tumor DNA (ctDNA) analysis in high-risk early stage breast cancer patients to monitor and predict response to neoadjuvant therapy and outcome in the I-SPY 2 TRIAL [abstract]. In: Proceedings of the 2018 San Antonio Breast Cancer Symposium; 2018 Dec 4-8; San Antonio, TX. Philadelphia (PA): AACR; *Cancer Res* 2019;79:Abstract nr PD2-01.
 62. Sledge GW Jr, Toi M, Neven P, et al. MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy. *J Clin Oncol* 2017;35:2875-84.
 63. Tolaney SM, Toi M, Neven P, et al. Clinical significance of PIK3CA and ESR1 mutations in ctDNA and FFPE samples from the MONARCH 2 study of abemaciclib plus

- fulvestrant [abstract]. In: Proceedings of the American Association for Cancer Research Annual Meeting 2019; 2019 Mar 29-Apr 3; Atlanta, GA. Philadelphia (PA): AACR; Cancer Res 2019;79:Abstract nr 4458.
64. Kim SB, Dent R, Im SA, et al. Ipatasertib plus paclitaxel versus placebo plus paclitaxel as first-line therapy for metastatic triple-negative breast cancer (LOTUS): a multicentre, randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Oncol* 2017;18:1360-72.
 65. Wongchenko MJ, Lipson D, Clark T, et al. On-treatment changes in circulating tumor DNA (ctDNA) level as an early predictor of clinical outcome in the LOTUS randomized phase 2 trial of 1st-line ipatasertib (IPAT) + paclitaxel (PAC) for metastatic triple-negative breast cancer (mTNBC) [abstract]. In: Proceedings of the American Association for Cancer Research Annual Meeting 2018; 2018 Apr 14-18; Chicago, IL. Philadelphia (PA): AACR; Cancer Res 2018;78:Abstract nr 2964.
 66. Sharma P, Abramson VG, O'Dea A. et al. Clinical and biomarker results from phase I/II study of PI3K inhibitor BYL 719 (alpelisib) plus nab-paclitaxel in HER2-negative metastatic breast cancer. *J Clin Oncol* 2018;36:1018.
 67. Clouthier DL, Lien SC, Yang SYC, et al. An interim report on the investigator-initiated phase 2 study of pembrolizumab immunological response evaluation (INSPIRE). *J Immunother Cancer* 2019;7:72.
 68. James Iafolla MA, Yang C, Dashner S, et al. Bespoke circulating tumor DNA (ctDNA) analysis as a predictive biomarker in solid tumor patients (pts) treated with single-agent pembrolizumab (P). *J Clin Oncol* 2019;37:2542.
 69. Moynahan ME, Chen D, He W, et al. Correlation between PIK3CA mutations in cell-free DNA and everolimus efficacy in HR+, HER2- advanced breast cancer: results from BOLERO-2. *Br J Cancer* 2017;116:726-30.
 70. Ma F, Li Q, Guan X, et al. Safety, efficacy, and biomarker analysis of pyrotinib in combination with capecitabine in HER2-positive metastatic breast cancer patients: a phase I clinical trial [abstract]. *J Clin Oncol* 2019;37:1035.
 71. de Azambuja E, Holmes AP, Piccart-Gebhart M, et al. Lapatinib with trastuzumab for HER2-positive early breast cancer (NeoALTTO): survival outcomes of a randomised, open-label, multicentre, phase 3 trial and their association with pathological complete response. *Lancet Oncol* 2014;15:1137-46.
 72. Rothé F, Silva MJ, Venet D, et al. Circulating Tumor DNA in HER2-Amplified Breast Cancer: A Translational Research Substudy of the NeoALTTO Phase III Trial. *Clin Cancer Res* 2019;25:3581-8.
 73. Tripathy D, Im SA, Colleoni M, et al. Ribociclib plus endocrine therapy for premenopausal women with hormone-receptor-positive, advanced breast cancer (MONALEESA-7): a randomised phase 3 trial. *Lancet Oncol* 2018;19:904-15.
 74. Bardia A, Su F, Solovieff N, et al. Genetic landscape of premenopausal HR+/HER2- advanced breast cancer (ABC) based on comprehensive circulating tumor DNA analysis and association with clinical outcomes in the Phase III MONALEESA-7 trial [abstract]. In: Proceedings of the American Association for Cancer Research Annual Meeting 2019; 2019 Mar 29-Apr 3; Atlanta, GA. Philadelphia (PA): AACR; Cancer Res 2019;79:Abstract nr CT141.
 75. Janiaud P, Serghiou S, Ioannidis JPA. New clinical trial designs in the era of precision medicine: An overview of definitions, strengths, weaknesses, and current use in oncology. *Cancer Treat Rev* 2019;73:20-30.
 76. Sellami D, Dharan B, Wilke C, et al. Circulating tumor DNA as a novel tool to shape clinical trial designs with the potential to impact outcomes: a focus on PI3K inhibitors. *Ann Oncol* 2017;28:2882-7.
 77. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2019;30:1674.
 78. Available online: www.fda.gov/medical-devices/vitro-diagnostics/list-cleared-or-approved-companion-diagnostic-devices-vitro-and-imaging-tools
 79. The Blood Profiling Atlas in Cancer (BloodPAC) Consortium. Available online: <https://www.bloodpac.org/>
 80. Cancer – ID Consortium. Available online: <https://www.cancer-id.eu/>

Cite this article as: Fiste O, Lontos M, Koutsoukos K, Terpos E, Dimopoulos MA, Zagouri F. Circulating tumor DNA-based predictive biomarkers in breast cancer clinical trials: a narrative review. *Ann Transl Med* 2020;8(23):1603. doi: 10.21037/atm-20-1175