

Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory

Gustav Uhre Jakobsen,^{1,2,*} Gustav Mogull,^{1,2,†} Jan Plefka,^{1,‡} and Jan Steinhoff^{2,§}

¹*Institut für Physik und IRIS Adlershof, Humboldt-Universität zu Berlin,
Zum Großen Windkanal 2, 12489 Berlin, Germany*

²*Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, 14476 Potsdam, Germany*

Using the recently established formalism of a worldline quantum field theory (WQFT) description of the classical scattering of two spinless black holes, we compute the far-field time-domain waveform of the gravitational waves produced in the encounter at leading order in the post-Minkowskian (weak field, but generic velocity) expansion. We reproduce previous results of Kovacs and Thorne in a highly economic way. Then using the waveform we extract the leading-order total radiated angular momentum, reproducing a recent result of Damour. Our work may enable crucial improvements of gravitational-wave predictions in the regime of large relative velocities.

When two compact objects (black holes, neutron stars or stars) fly past each other their gravitational interactions not only deflect their trajectories but they also produce gravitational radiation, or gravitational Bremsstrahlung in analogy to the electromagnetic case. The resulting waveform in the far field at leading order in Newton's constant G has been constructed (in the spinless case) in a series of papers by Kovacs, Thorne, and Crowley in the 1970s [1–4] — see Refs. [5] for recent work on slow-motion sources. Today's gravitational wave (GW) observatories routinely detect quasi-circular *inspirals and mergers* of binary black holes and neutron stars [6]. Yet Bremsstrahlung events currently appear to be out of reach as the signal is not periodic and typically less intensive [7]. Still, they represent interesting targets for GW searches, calling for accurate waveform models.

Indeed, the experimental success of GW astronomy brings up the need for high-precision theoretical predictions for the classical relativistic two-body problem [8]. A number of complementary classical perturbative approaches have been established over the years [9]. Yet quantum-field-theory based techniques founded in a perturbative Feynman-diagrammatic expansion of the path integral in the classical limit have proven to be highly efficient. These come in two alternative approaches.

The first approach, the effective field theory (EFT) formalism [10], models the compact objects as point-like massive particles coupled to the gravitational field. It has mostly been applied to a nonrelativistic post-Newtonian (PN) scenario for *bound* orbits, in which an expansion in powers of Newton's constant G implies an expansion in velocities ($\frac{Gm}{c^2 r} \sim \frac{v^2}{c^2}$). Recently it has also been extended to the post-Minkowskian (PM) expansion for *unbound* orbits [11, 12] relevant for this work, an expansion in G for arbitrary velocities. In these EFT settings the graviton field $h_{\mu\nu}(x)$ is integrated out successively (from small to large length scales) in the path integral, while

the worldline trajectories of the black holes $x_i^\mu(\tau_i)$ are kept as classical background sources — see Refs. [13] for reviews.

The second now blossoming approach starts out from scattering amplitudes of massive scalars — avatars of spinless black holes — minimally coupled to general relativity [14], thereby putting the younger innovations in on-shell techniques for scattering amplitudes (e.g. generalized unitarity [15] or the double copy [16]) to work. In order to obtain the conservative gravitational potential one performs a subtle classical limit of the scattering amplitudes [17] in order to match to a non-relativistic EFT for scalar particles with the desired potential [18] (see also Refs. [14, 18]), which is known to 3PM order [14] (complemented by certain radiation-reaction effects [17, 19, 20]). Very recently the 4PM conservative potential was also reported [21]. The so-obtained effective potential is then used to compute observables such as the scattering angle or the (PM-resummed) periastron advance in the bound system [11, 21]. Further recent PM results exist for non-spinning particles [22], for spin effects [23], tidal effects [24], and radiation effects [25].

In a recent work of three of the present authors the synthesis of these two quantum-field-theory based approaches to classical relativity was provided in the form of a worldline quantum field theory (WQFT) [26]: quantizing *both* the graviton field $h_{\mu\nu}$ and the fluctuations about the bodies' worldline trajectories z_i^μ were shown to yield an efficient approach yielding only the relevant classical contributions. In essence the WQFT formalism provides an efficient diagrammatic framework for solving the equations of motion of gravity-matter systems perturbatively.

In this Letter we employ this novel formalism to compute the time-domain gravitational waveform of a Bremsstrahlung event at leading order in G , demonstrating its effectiveness. To our knowledge the seminal result of Kovacs and Thorne [4] has not been verified in its entirety to date. As we shall see, our approach is far more efficient than the one employed back then, paving the way for calculations of higher orders. We stress that we are able to determine the *far-field waveforms* which are of direct relevance for GW observatories. As a check on

* gustav.uhre.jakobsen@physik.hu-berlin.de

† gustav.mogull@aei.mpg.de

‡ jan.plefka@hu-berlin.de

§ jan.steinhoff@aei.mpg.de




FIG. 1. The three diagrams contributing to the Bremsstrahlung at 2PM order, where $\omega_i = k \cdot v_i$ by energy conservation at the worldline vertices. All three diagrams have the integral measure in Eq. (16); in the rest frame of black hole 1 diagram (a) does not contribute as soon as the outgoing graviton is contracted with a purely spatial polarization tensor.

these waveforms we furthermore reproduce Damour's recent result for the *total radiated angular momentum* [19] at 2PM order. Our results also complement the recent result of the *total radiated momentum* at leading order in G (3PM) established with amplitude techniques [27]. We comment on how to achieve this result from our methods.

Worldline Quantum Field Theory. — The classical gravitational scattering of two massive objects m_i moving on trajectories $x_i^\mu(\tau_i) = b_i^\mu + v_i^\mu \tau_i + z_i^\mu(\tau_i)$ is described by the worldline quantum field theory (WQFT) with partition function [26]

$$\mathcal{Z}_{\text{WQFT}} := \text{const} \times \int D[h_{\mu\nu}] \int \prod_{i=1}^2 D[z_i] e^{i(S_{\text{EH}} + S_{\text{gf}})} \quad (1)$$

$$\exp \left[-i \sum_{i=1}^2 \int_{-\infty}^{\infty} d\tau_i \frac{m_i}{2} [\eta_{\mu\nu} + \kappa h_{\mu\nu}(x)] \dot{x}_i^\mu \dot{x}_i^\nu \right],$$

Diagram (a) of Fig. 1 then takes the form¹

$$k^2 \langle h_{\mu\nu}(k) \rangle_{\text{WQFT}} \Big|_{(a)} = -\frac{m_1 m_2 \kappa^3}{8} \int_{q_1, q_2} \mu_{1,2}(k) \frac{(2\omega_1 v_1^{(\mu} \delta^{\nu)}) - v_1^\mu v_1^\nu k_\rho)(2\omega_1 v_1^{(\sigma} \eta^{\lambda)\rho} - v_1^\sigma v_1^\lambda q_2^\rho)}{(\omega_1 + i\epsilon)^2} \frac{P_{\sigma\lambda; \alpha\beta}}{[(q_2^0 + i\epsilon)^2 - \mathbf{q}_2^2]} v_2^\alpha v_2^\beta, \quad (6)$$

¹ In principle we should also contract with $P_{\mu\nu; \rho\sigma}$ for an outgoing graviton line; however as the polarization tensors $e_{+,\times}^{\mu\nu}$ are

where $S_{\text{EH}} + S_{\text{gf}}$ is the gauge-fixed Einstein-Hilbert action

$$S_{\text{EH}} + S_{\text{gf}} = \int d^4x \left(-\frac{2}{\kappa^2} \sqrt{-g} R + (\partial_\nu h^{\mu\nu} - \frac{1}{2} \partial^\mu h^\nu{}_\nu)^2 \right), \quad (2)$$

with $\kappa^2 = 32\pi G$ the gravitational coupling; we have suppressed the ghost contributions in Eq. (1) as they are irrelevant in the classical setting. We work in mostly minus signature, $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$.

Correlation functions in the WQFT $\langle \mathcal{O}(h, \{x_i\}) \rangle_{\text{WQFT}}$ result from an insertion of the operator \mathcal{O} in the path integral and dividing by $\mathcal{Z}_{\text{WQFT}}$. Moving to momentum space for the graviton $h_{\mu\nu}(k)$ and energy space for the fluctuations $z^\mu(\omega)$ we have the *retarded* propagators

$$\begin{array}{c} \mu\nu \\ \bullet \text{---} \text{---} \text{---} \bullet \\ \quad \quad \quad k \\ \rho\sigma \end{array} = i \frac{P_{\mu\nu; \rho\sigma}}{(k^0 + i\epsilon)^2 - \mathbf{k}^2}, \quad (3a)$$

$$\begin{array}{c} \mu \\ \bullet \text{---} \text{---} \bullet \\ \quad \quad \quad \omega \\ \nu \end{array} = -i \frac{\eta^{\mu\nu}}{m(\omega + i\epsilon)^2}, \quad (3b)$$

with $P_{\mu\nu; \rho\sigma} := \eta_{\mu(\rho} \eta_{\sigma)\nu} - \frac{1}{2} \eta_{\mu\nu} \eta_{\rho\sigma}$. The relevant vertices for the emission of a graviton off the worldline read

$$\begin{array}{c} \bullet \\ \downarrow \\ \bullet \end{array} = -i \frac{m\kappa}{2} e^{ik \cdot b} \bar{\delta}(k \cdot v) v^\mu v^\nu, \quad (4)$$

$$h_{\mu\nu}(k)$$

with k outgoing, $\bar{\delta}(\omega) := (2\pi)\delta(\omega)$ and

$$\begin{array}{c} \bullet \\ \downarrow \\ \bullet \end{array} z^\rho(\omega) = \frac{m\kappa}{2} e^{ik \cdot b} \bar{\delta}(k \cdot v + \omega) \times \left(2\omega v^{(\mu} \delta^{\nu)\rho} + v^\mu v^\nu k_\rho \right). \quad (5)$$

$$h_{\mu\nu}(k)$$

The energy ω is also taken as outgoing. One also has the standard bulk graviton vertices, of which we shall need only the three-graviton vertex — see e.g. Ref. [28].

To determine the Bremsstrahlung of two traversing black holes we compute the expectation value $k^2 \langle h^{\mu\nu}(k) \rangle_{\text{WQFT}}$. At leading (2PM) order there are three diagrams contributing, cp. Fig. 1. We integrate over the momenta or energies of internal gravitons or fluctuations respectively; lack of three-momentum conservation at the worldline vertices leaves unresolved integrals for the tree-level diagrams.

traceless we find it unnecessary.

where $\omega_1 = k \cdot v_1$, $\int_{q_i} := \int \frac{d^4 q_i}{(2\pi)^4}$ and the integral measure is

$$\mu_{1,2}(k) = e^{i(q_1 \cdot b_1 + q_2 \cdot b_2)} \delta(q_1 \cdot v_1) \delta(q_2 \cdot v_2) \delta(k - q_1 - q_2), \quad (7)$$

with $\delta(k) := (2\pi)^4 \delta^{(4)}(k)$. The diagram (b) is naturally obtained by swapping $1 \leftrightarrow 2$. Diagram (c) includes the three-graviton vertex $V_3^{(\mu\nu)(\rho\sigma)(\lambda\tau)}(k, -q_1, -q_2)$:

$$k^2 \langle h_{\mu\nu}(k) \rangle_{\text{WQFT}} \Big|_{(c)} = -\frac{m_1 m_2 k^3}{8} \int_{q_1, q_2} \mu_{1,2}(k) V_3^{(\mu\nu)(\rho\sigma)(\lambda\tau)} \frac{P_{\rho\sigma;\alpha\beta}}{[(q_1^0 + i\epsilon)^2 - \mathbf{q}_1^2]} \frac{P_{\lambda\tau;\gamma\delta}}{[(q_2^0 + i\epsilon)^2 - \mathbf{q}_2^2]} v_1^\alpha v_1^\beta v_2^\gamma v_2^\delta. \quad (8)$$

These integrands were already given in Ref. [26]. The sum of the three integrands also agrees with a previous amplitudes-based result [14] and is gauge-invariant.

The waveform in spacetime in the *wave zone* is obtained from $\langle h^{\mu\nu}(k) \rangle_{\text{WQFT}}$ as follows: we may identify

$$k^2 \langle h_{\mu\nu}(k) \rangle_{\text{WQFT}} = \frac{\kappa}{2} S_{\mu\nu}(k), \quad (9)$$

where $S_{\mu\nu} = \tau_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} \tau^\lambda{}_\lambda$ and $\tau_{\mu\nu}$ is the combined energy-momentum pseudo-tensor of matter and the gravitational field. Consider $S_{\mu\nu}(k)$ for a fixed GW frequency $k^0 = \Omega$. In the wave zone ($r \gg \{|b_i|, \Omega^{-1}, \Omega|b_i|^2\}$) the metric perturbation $h_{\mu\nu}(\mathbf{x}, t)$ takes the form of a plane wave (see e.g. Chapter 10.4 of Weinberg [29]):

$$\kappa h_{\mu\nu}(\mathbf{x}, t) = \frac{4G}{r} S_{\mu\nu}(\Omega, \mathbf{k} = \Omega \hat{\mathbf{x}}) e^{-ik_\mu x^\mu} + c.c., \quad (10)$$

with the wave vector $k^\mu = \Omega(1, \hat{\mathbf{x}})$; $\hat{\mathbf{x}} = \mathbf{x}/r$ is the unit vector pointing in the direction of the observation point (hence $k^2 = 0$).

The total gauge-invariant frequency-domain waveform can be read off as $4G S_{ij}^{\text{TT}}(\Omega, \mathbf{k} = \Omega \hat{\mathbf{x}})$, where TT denotes the transverse-traceless projection. The corresponding time-domain waveform $f_{ij}(u, \theta, \phi)$ is essentially its Fourier transform in Ω :

$$\kappa h_{ij}^{\text{TT}} = \frac{f_{ij}}{r} = \frac{4G}{r} \int_{\Omega} e^{-ik \cdot x} S_{ij}^{\text{TT}}(k) \Big|_{k^\mu = \Omega(1, \hat{\mathbf{x}})}, \quad (11)$$

where $\int_{\Omega} := \int_{-\infty}^{\infty} \frac{d\Omega}{2\pi}$. Note that $k \cdot x = \Omega(t - r)$ yields the retarded time $u = t - r$. Our task now is to perform the integrals; in a PM expansion $f_{ij} = \sum_n G^n f_{ij}^{(n)}$, and we seek the 2PM component $f_{ij}^{(2)}$. By focusing on the time-domain instead of the frequency-domain waveform we considerably simplify the integration step — as we shall see, the integration over frequency Ω of the outgoing radiation coincides neatly with energy conservation along each worldline.

Kinematics. — We describe the waveform in a Cartesian coordinate system (t, x, y, z) where black hole 1 is initially at rest $v_1^\mu = (1, 0, 0, 0)$ and located at the spatial origin, i.e. we set $b_1^\mu = 0$. The orbit of black hole 2 we put in the x - y plane with initial velocity $v_2^\mu = (\gamma, \gamma v, 0, 0)$ in the x -direction; the impact parameter $b_2^\mu = (0, 0, b, 0) =: b^\mu$ points in the y -direction. Introducing the polar angles θ and ϕ we may write the

unit (spatial) vector \hat{x}^μ pointing from black hole 1 to our observation point as

$$\hat{x}^\mu = \hat{e}_1^\mu \cos \theta + \sin \theta (\hat{e}_2^\mu \cos \phi + \hat{e}_3^\mu \sin \phi), \quad (12)$$

where $\hat{e}_i^\mu = (0, \hat{\mathbf{e}}_i)$ are spatial unit vectors. Also, we put $\rho^\mu = v_1^\mu + \hat{x}^\mu$.

The two additional unit spatial vectors orthogonal to \hat{x}^μ are

$$\hat{\theta}^\mu = \partial_\theta \hat{x}^\mu = (0, \hat{\boldsymbol{\theta}}), \quad \hat{\phi}^\mu = \frac{1}{\sin \theta} \partial_\phi \hat{x}^\mu = (0, \hat{\boldsymbol{\phi}}). \quad (13)$$

Together with \hat{x}^μ they form a right-handed spatial coordinate system. GWs travel in the direction of \hat{x}^μ and we use $\hat{\theta}^\mu$ and $\hat{\phi}^\mu$ to define our polarization tensors in a linear basis:

$$e_{+}^{\mu\nu} = \hat{\theta}^\mu \hat{\theta}^\nu - \hat{\phi}^\mu \hat{\phi}^\nu, \quad e_{\times}^{\mu\nu} = \hat{\theta}^\mu \hat{\phi}^\nu + \hat{\phi}^\mu \hat{\theta}^\nu. \quad (14)$$

The waveform $f_{ij}(u, \theta, \phi)$ is thus decomposed as

$$f_{ij} = f_{+}(e_{+})_{ij} + f_{\times}(e_{\times})_{ij} \quad (15)$$

with $f_{+, \times} = \frac{1}{2}(e_{+, \times})_{ij} f_{ij}$.

The polarization tensors have zero time components, which conveniently implies the vanishing of diagram (a) in Fig. 1 once contracted with them. This observation follows directly from the expression for vertex (5): in the case of diagram (a) the instance of this vertex that contracts with the outgoing graviton line carries an overall factor of $v_1^\mu = (1, 0, 0, 0)$, which is orthogonal to the spatial polarization tensors above.

Integration. — The two non-zero diagrams in Fig. 1 share the integration measure $\mu_{1,2}(k)$ (7). Including also the integration with respect to Ω in Eq. (11) the full measure becomes

$$\int_{\Omega, q_1, q_2} \mu_{1,2}(k) e^{-ik \cdot x} = \frac{1}{\rho \cdot v_2} \int_{\mathbf{q}} e^{i\mathbf{q} \cdot \tilde{\mathbf{b}}}, \quad (16)$$

where we recall that $k^\mu = \Omega \rho^\mu$; using the delta function constraints in $\mu_{1,2}(k)$ we can now identify

$$q_2 = k - q_1, \quad q_1 = (0, \mathbf{q}), \quad \Omega = -\frac{v\gamma}{\rho \cdot v_2} \mathbf{q} \cdot \hat{\mathbf{e}}_1. \quad (17)$$

We are left with a three-dimensional Euclidean integral involving the shifted τ -dependent impact parameter:

$$\tilde{\mathbf{b}}(\tau) = \mathbf{b} + \tau \hat{\mathbf{e}}_1, \quad \tau = \frac{v\gamma}{\rho \cdot v_2} (u + \mathbf{b} \cdot \hat{\mathbf{x}}), \quad (18)$$

noting that $\rho \cdot v_2 = \gamma(1 - v \cos \theta)$. The polarizations of the waveform from Eq. (11) now take the schematic form (also using Eq. (15))

$$\begin{aligned} & \frac{f_{+, \times}^{(2)}}{m_1 m_2} \\ &= 4\pi \int_{\mathbf{q}} e^{i\mathbf{q} \cdot \tilde{\mathbf{b}}} \left(\frac{\mathcal{N}_{+, \times}^i \mathbf{q}^i}{\mathbf{q}^2 (\mathbf{q} \cdot \hat{\mathbf{e}}_1 - i\epsilon)} + \frac{\mathcal{M}_{+, \times}^{ij} \mathbf{q}^i \mathbf{q}^j}{\mathbf{q}^2 (\mathbf{q}^2 + \mathbf{q} \cdot L \cdot \mathbf{q})} \right), \end{aligned} \quad (19)$$

with the two terms corresponding to the non-zero diagrams (b) and (c) in Fig. 1 respectively. The rank-2 matrix L introduced here is

$$L^{ij} = 2 \frac{v\gamma}{\rho \cdot v_2} \hat{\mathbf{e}}_1^{(i} \hat{\mathbf{x}}^{j)}. \quad (20)$$

Finally the vector and matrix insertions are explicitly given as the real and imaginary parts of²

$$\begin{aligned} \mathcal{N}^i &= 2 \frac{\gamma^2 \sin^2 \theta}{\rho \cdot v_2} \left(\frac{\gamma(1 - 3v^2)}{\rho \cdot v_2} + (1 + v^2) \right) \hat{\mathbf{e}}_1^i \\ &+ 2 \frac{\gamma(1 + v^2) \sin \theta}{\rho \cdot v_2} \left(\frac{(\rho \cdot v_2)^2 - 1}{v(\rho \cdot v_2)} \cos \phi + 2i\gamma \sin \phi \right) \hat{\mathbf{e}}_2^i, \end{aligned} \quad (21a)$$

$$\begin{aligned} \mathcal{M}^{ij} &= 8 \frac{\gamma^4 v^4 \sin^2 \theta}{(\rho \cdot v_2)^3} \hat{\mathbf{e}}_1^i \hat{\mathbf{e}}_1^j + 16 \frac{\gamma^3 v^2 \sin \theta}{(\rho \cdot v_2)^2} \hat{\mathbf{e}}_1^{(i} (\hat{\boldsymbol{\theta}} + i\hat{\boldsymbol{\phi}})^{j)} \\ &+ 4 \frac{\gamma^2 (1 + v^2)}{\rho \cdot v_2} (\hat{\boldsymbol{\theta}} + i\hat{\boldsymbol{\phi}})^i (\hat{\boldsymbol{\theta}} + i\hat{\boldsymbol{\phi}})^j, \end{aligned} \quad (21b)$$

where $\mathcal{N}^i = \mathcal{N}_+^i + i\mathcal{N}_\times^i$ and $\mathcal{M}^{ij} = \mathcal{M}_+^{ij} + i\mathcal{M}_\times^{ij}$. The insertions \mathcal{N}^i and \mathcal{M}^{ij} correspond to a helicity basis in which they have a particular simple expression. We integrate the two diagrams separately.

Integration of the first diagram is achieved using the simple result (true regardless of the vector $\tilde{\mathbf{b}}$)

$$\begin{aligned} & \int_{\mathbf{q}} e^{i\mathbf{q} \cdot \tilde{\mathbf{b}}} \frac{\mathbf{q}^i}{\mathbf{q}^2 (\mathbf{q} \cdot \hat{\mathbf{e}}_1 - i\epsilon)} \\ &= \frac{1}{4\pi} \left(\frac{\hat{\mathbf{e}}_1^i}{|\tilde{\mathbf{b}}|} - \frac{\tilde{\mathbf{b}}^i - (\tilde{\mathbf{b}} \cdot \hat{\mathbf{e}}_1) \hat{\mathbf{e}}_1^i}{\tilde{\mathbf{b}}^2 - (\tilde{\mathbf{b}} \cdot \hat{\mathbf{e}}_1)^2} \left(1 + \frac{\tilde{\mathbf{b}} \cdot \hat{\mathbf{e}}_1}{|\tilde{\mathbf{b}}|} \right) \right), \end{aligned} \quad (22)$$

which we prove in the Appendix. The other integral required corresponding to diagram (c) is somewhat more involved. The denominator of this integral is composed of an isotropic propagator together with an anisotropic one. The physical interpretation is a convolution between the potentials of the two black holes, where the potential of black hole 2 is boosted and leads to the anisotropic

propagator. One compact representation is

$$\begin{aligned} & \int_{\mathbf{q}} e^{i\mathbf{q} \cdot \tilde{\mathbf{b}}} \frac{\mathbf{q}^i \mathbf{q}^j}{\mathbf{q}^2 (\mathbf{q}^2 + \mathbf{q} \cdot L \cdot \mathbf{q})} \\ &= \frac{1}{2\pi \Delta(G)} \left[\frac{(G_0 + \alpha G_1) A^{ij} - (G_1 + \alpha G_2) B^{ij}}{\sqrt{G(\alpha)}} \right]_{\alpha=0}^{\alpha=1}, \end{aligned} \quad (23)$$

where we have introduced the quadratic polynomial

$$\begin{aligned} G(\alpha) &= G_0 + 2\alpha G_1 + \alpha^2 G_2, \\ G_0 &= \tilde{\mathbf{b}}^2, G_1 = \tilde{\mathbf{b}}^i \tilde{\mathbf{b}}^j \frac{\delta^{ij} L^{kk} - L^{ij}}{2}, G_2 = (\tilde{\mathbf{b}} \cdot \hat{\boldsymbol{\phi}})^2 \text{Det}_2 L, \end{aligned} \quad (24)$$

and $\Delta(G) = 4(G_1^2 - G_0 G_2)$ is the polynomial discriminant. We have also introduced the two matrices

$$\begin{aligned} A^{ij} &= \text{Det}_2(L) \left(-2(\tilde{\mathbf{b}} \cdot \hat{\boldsymbol{\phi}}) (L^{-1} \cdot \tilde{\mathbf{b}})^i \hat{\boldsymbol{\phi}}^j \right. \\ &\quad \left. + (\tilde{\mathbf{b}} \cdot \hat{\boldsymbol{\phi}})^2 (L^{-1})^{ij} + (\tilde{\mathbf{b}} \cdot L^{-1} \cdot \tilde{\mathbf{b}}) \hat{\boldsymbol{\phi}}^i \hat{\boldsymbol{\phi}}^j \right), \end{aligned} \quad (25a)$$

$$B^{ij} = \tilde{\mathbf{b}}^2 \delta^{ij} - \tilde{\mathbf{b}}^i \tilde{\mathbf{b}}^j. \quad (25b)$$

$\text{Det}_2(L)$ and L^{-1} are computed in the 2-dimensional subspace spanned by L , while $\hat{\boldsymbol{\phi}}$ is the unit vector orthogonal to this subspace. This integral is also discussed in the Appendix; again, both of the integrals (22) and (23) are solved for arbitrary $\tilde{\mathbf{b}}^i$ and L^{ij} (with the assumption that L^{ij} is rank 2). In the present case where L^{ij} is given by Eq. (20) we find that

$$\text{Det}_2(L) = - \left(\frac{\gamma v \sin \theta}{\rho \cdot v_2} \right)^2, \quad (26a)$$

$$(L^{-1})^{ij} = \frac{\rho \cdot v_2}{2\gamma v} \sum_{\pm} \pm \frac{(\hat{\mathbf{x}} \pm \hat{\mathbf{e}}_1)^i (\hat{\mathbf{x}} \pm \hat{\mathbf{e}}_1)^j}{(1 \pm \cos \theta)^2}, \quad (26b)$$

summing over the two signs in the latter case.

Leading-Order Waveform. — By combining Eq. (19) with the insertions $\mathcal{N}_{+, \times}^i$ and $\mathcal{M}_{+, \times}^{ij}$ and the integrals above we get the full 2PM waveform:

$$\begin{aligned} & \frac{f_{+, \times}^{(2)}}{m_1 m_2} = \frac{\hat{\mathbf{e}}_1^i \mathcal{N}_{+, \times}^i}{\sqrt{\mathbf{b}^2 + \tau^2}} - \frac{\mathbf{b}^i \mathcal{N}_{+, \times}^i}{\mathbf{b}^2} \left(1 + \frac{\tau}{\sqrt{\mathbf{b}^2 + \tau^2}} \right) \\ &+ \frac{2\mathcal{M}_{+, \times}^{ij}}{\Delta(G)} \left[\frac{(G_0 + \alpha G_1) A^{ij} - (G_1 + \alpha G_2) B^{ij}}{\sqrt{G(\alpha)}} \right]_{\alpha=0}^{\alpha=1}. \end{aligned} \quad (27)$$

This is a rather compact representation of the gravitational Bremsstrahlung waveform, which we have confirmed agrees with the (rather lengthy) result of Kovacs and Thorne [4]. The two values of α in the second line correspond to contributions from the two black holes. Note that there is also a leading (and non-radiating) 1PM contribution to the waveform which is independent of the retarded time $u = t - r$:

$$f_+^{(1)} = \frac{2m_2 \gamma v^2 \sin^2 \theta}{1 - v \cos \theta}, \quad f_\times^{(1)} = 0. \quad (28)$$

² To compactify these results we have used the generalized gauge invariance $\mathcal{N}_{+, \times} \rightarrow \mathcal{N}_{+, \times} + X \hat{\mathbf{e}}_1$, $\mathcal{M}_{+, \times} \rightarrow \mathcal{M}_{+, \times} - X(\mathbb{1} + L)$ for an arbitrary function X of external kinematics. We have also dropped a term from \mathcal{N}_\pm in the $\hat{\mathbf{e}}_3$ direction which does not contribute to the final integrated result (27).




FIG. 2. Plots of the wave memories $\Delta f_{+,x}$ for $v = 0.2$. For a visualisation of the complete waveforms as they evolve with retarded time u see $f_+(u, \theta, \phi)|_{v=0.2}$ and $f_-(u, \theta, \phi)|_{v=0.2}$.

Diagrammatically this consists only of the vertex (4) with emission from worldline 2; the contribution from worldline 1 again vanishes in our frame due to $(v_1 \cdot e_{\pm} \cdot v_1) = 0$.

To illustrate this result in Fig. 2 we present the *gravitational wave memories* $\Delta f_{+,x} := [f_{+,x}]_{u=-\infty}^{u=+\infty}$. The beauty of our result (27) lies in the fact that the memories only receive contributions from the second term, and read

$$\Delta f_{+,x} = -2G^2 m_1 m_2 \frac{\mathbf{b}^i \mathcal{N}_{+,x}^i}{\mathbf{b}^2} + \mathcal{O}(G^3). \quad (29)$$

Diagrammatically they exclusively emerge from diagram (b) of Fig. 1. So they are manifestly insensitive to gravitational self-interactions — this was also pointed out in Ref. [19].

Radiated Angular Momentum. — One may now use our result for the waveform (27) to compute the total radiated momentum and angular momentum. Expressions for these quantities in terms of the asymptotic waveform are given in Refs. [19, 30]:

$$P_{\text{rad}}^{\mu} = \frac{1}{32\pi G} \int \text{d}u \text{d}\sigma [\dot{f}_{ij}]^2 \rho^{\mu}, \quad (30)$$

$$J_{ij}^{\text{rad}} = \frac{1}{8\pi G} \int \text{d}u \text{d}\sigma \left(f_{k[i} \dot{f}_{j]k} - \frac{1}{2} x_{[i} \partial_{j]} f_{kl} \dot{f}_{kl} \right), \quad (31)$$

where $\dot{f}_{ij} := \partial_u f_{ij}$ and $\text{d}\sigma = \sin\theta \text{d}\theta \text{d}\phi$ is the unit sphere measure. Here we concentrate on J_{ij}^{rad} as it contributes at leading order $\mathcal{O}(G^2)$ due to $\dot{f}_{ij}^{(1)} = 0$ and was recently obtained in the center-of-mass frame [19].

At leading order the static nature of $f_{ij}^{(1)}$ (28) allows one to trivially perform the u -integration and express the radiated angular momentum in terms of the wave memories $\Delta f_{+,x}$. Inserting the basis of polarization tensors (15) (and using $f_x^{(1)} = 0$) gives

$$J_{xy}^{\text{rad}} = \frac{1}{8\pi} \int \text{d}\sigma \left[\frac{\sin\phi}{\sin\theta} f_+^{(1)} \Delta f_x - \frac{1}{2} \cos\phi \partial_{\theta} f_+^{(1)} \Delta f_+ \right] + \mathcal{O}(G^3). \quad (32)$$

The spherical integral is elementary and yields

$$\frac{J_{xy}^{\text{rad}}}{J_{xy}^{\text{init}}} = \frac{4G^2 m_1 m_2 (2\gamma^2 - 1)}{b^2 \sqrt{\gamma^2 - 1}} \mathcal{I}(v) + \mathcal{O}(G^3), \quad (33a)$$

$$\mathcal{I}(v) = -\frac{8}{3} + \frac{1}{v^2} + \frac{(3v^2 - 1)}{v^3} \text{arctanh}(v), \quad (33b)$$

where we have normalized our result with respect to the initial angular momentum in the rest frame of black hole 1: $J_{xy}^{\text{init}} = m_2 |\mathbf{v}_2| |\mathbf{b}| = m_2 \gamma v b$. Compared with Damour's center-of-mass frame result [19] we find perfect agreement.³ Similarly evaluating Eq. (30) should reproduce the recent result of Ref. [27] for the radiated momentum.

Conclusions and Outlook. — Searching for GWs from scattering events over the full range of impact velocities requires precision predictions in the PM approximation. Furthermore, PM results may even improve GW predictions for bound systems observed by current GW detectors. Indeed, while the potential and radiation of bound systems was calculated to high PN order [31] (see Refs. [32] for spinning bodies), a resummation of PN results in the strong-field and fast-motion regimes is essential for building accurate waveform models and averting the imminent domination of systematic errors in observations [8]. The PM resummation is one promising recent attempt [18, 33, 34].

We believe our results provide a stepping stone for higher-order calculations, where a repertoire of advanced integration techniques can be put to use [14, 21, 27, 35]. This work provides an explicit link, via the WQFT [26], between quantum scattering amplitudes and classical waveforms ultimately used by GW observatories. An extension to spin and finite-size effects within the WQFT appears quite possible, and would involve integrals similar to the non-spinning case. Deriving an analytic expression for frequency-domain PM waveforms would also be very useful; in fact the WQFT readily leads to one-dimensional integrals involving Bessel functions. Yet, the class of special functions describing these waveforms remains to be identified.

Finally, our eventual aim is an extension to *bound* orbits. Recent work has shown that mappings between bound and unbound orbits exist for both conservative and dissipative observables [21, 36]; finding a similar mapping for the waveform would be of great utility.

Acknowledgments. — We would like to thank A. Buonanno and J. Vines for helpful discussions. We are also grateful for use of G. Kälin's C++ graph library. GUJ's and GM's research is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Projektnummer 417533893/GRK2575 "Rethinking Quantum Field Theory".

³ As the two frames are related by a boost in the x direction this implies that $J_{0y}^{\text{rad}} = 0$ in both frames.

INTEGRALS

We begin with the simpler integral in Eq. (22), corresponding to diagram (b) in Fig. 1. Working in Cartesian components with $\mathbf{b} = (b_1, b_2, b_3)$ (in the main text we replace $\mathbf{b} \rightarrow \tilde{\mathbf{b}}$) and $\mathbf{q} = (q_1, q_2, q_3)$ it is sufficient to show

$$\int_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{b}} \frac{q_2}{\mathbf{q}^2(q_1 - i\epsilon)} = -\frac{b_2}{4\pi(\mathbf{b}^2 - b_1^2)} \left(1 + \frac{b_1}{|\mathbf{b}|}\right). \quad (34)$$

The corresponding result with numerator q_3 is related by symmetry, and the one with q_1 is trivially given by

$$\int_{\mathbf{q}} \frac{e^{i\mathbf{q}\cdot\mathbf{b}}}{\mathbf{q}^2} = \frac{1}{4\pi|\mathbf{b}|}. \quad (35)$$

We make convenient use of the fact that

$$\int_{\omega} e^{i\omega\tau} \frac{f(\omega)}{\omega - i\epsilon} = i \int_{-\infty}^{\tau} d\tau' \int_{\omega} e^{i\omega\tau'} f(\omega), \quad (36)$$

the $i\epsilon$ prescription implying that our integrals are boundary fixed at $\tau \rightarrow -\infty$. This is to be expected given our use of retarded propagators. The original integral can therefore be re-written as

$$\int_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{b}} \frac{q_2}{\mathbf{q}^2(q_1 - i\epsilon)} = \frac{\partial}{\partial b_2} \int_{-\infty}^{b_1} db'_1 \int_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{b}'} \frac{1}{\mathbf{q}^2}, \quad (37)$$

where $\mathbf{b}' = (b'_1, b_2, b_3)$. From here using Eq. (35) it is a simple exercise to reproduce Eq. (34).

Next, let us consider the anisotropic integral (23) corresponding to the gravitational self-interaction of the two black hole potentials. The numerator can be obtained by differentiation with respect to the impact parameter; also introducing a Feynman parameter α the integral is re-expressed as

$$\begin{aligned} \mathcal{I}^{ij} &= \int_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{b}} \frac{\mathbf{q}^i \mathbf{q}^j}{\mathbf{q}^2(\mathbf{q}^2 + \mathbf{q} \cdot L \cdot \mathbf{q})} \\ &= -\frac{\partial}{\partial b^i} \frac{\partial}{\partial b^j} \int_0^1 d\alpha \int_{\mathbf{q}} \frac{e^{i\mathbf{q}\cdot\mathbf{b}}}{(\mathbf{q}^2 + \alpha \mathbf{q} \cdot L \cdot \mathbf{q})^2}. \end{aligned} \quad (38)$$

The \mathbf{q} -integration and \mathbf{b} derivatives are straightforward; we are left with the Feynman parameter integral

$$\mathcal{I}^{ij} = \int_0^1 d\alpha \frac{\det(M) \mathbf{b}^k \mathbf{b}^l (M^{-1})^{k[l} (M^{-1})^{i]j}}{4\pi G(\alpha)^{3/2}}, \quad (39)$$

where the quadratic function $G(\alpha) = \det(M) \mathbf{b} \cdot M^{-1} \cdot \mathbf{b}$ was given in Eq. (24) and the α -dependent matrix M^{ij} is defined as simply

$$M^{ij} = \delta^{ij} + \alpha L^{ij}. \quad (40)$$

The Feynman parameter integral is solved by recognizing that the numerator in Eq. (39) is a linear function in α : $B^{ij} + \alpha A^{ij}$, where all dependence on α is explicit. When the numerator in Eq. (39) is rewritten in this way the Feynman parameter integral can easily be solved to give Eq. (23).

-
- [1] K. S. Thorne and S. J. Kovacs, “The generation of gravitational waves. I. Weak-field sources.” *Astrophys. J.* **200**, 245–262 (1975).
- [2] R. J. Crowley and K. S. Thorne, “The generation of gravitational waves. II. The postlinear formation revisited.” *Astrophys. J.* **215**, 624–635 (1977).
- [3] S.J. Kovacs and K.S. Thorne, “The Generation of Gravitational Waves. 3. Derivation of Bremsstrahlung Formulas,” *Astrophys. J.* **217**, 252–280 (1977).
- [4] S.J. Kovacs and K.S. Thorne, “The Generation of Gravitational Waves. 4. Bremsstrahlung,” *Astrophys. J.* **224**, 62–85 (1978).
- [5] Lorenzo De Vittori, Philippe Jetzer, and Antoine Klein, “Gravitational wave energy spectrum of hyperbolic encounters,” *Phys. Rev. D* **86**, 044017 (2012), [arXiv:1207.5359 \[gr-qc\]](#); Gröbner, Matthias and Jetzer, Philippe and Haney, Maria and Tiwari, Shubhanshu and Ishibashi, Wako, “A note on the gravitational wave energy spectrum of parabolic and hyperbolic encounters,” *Class. Quant. Grav.* **37**, 067002 (2020), [arXiv:2001.05187 \[gr-qc\]](#); Salvatore Capozziello and Mariafelicia De Laurentis, “Gravitational waves from stellar encounters,” *Astropart. Phys.* **30**, 105–112 (2008), [arXiv:0806.4117 \[astro-ph\]](#).
- [6] B.P. Abbott *et al.* (LIGO Scientific, Virgo), “Observation of Gravitational Waves from a Binary Black Hole Merger,” *Phys. Rev. Lett.* **116**, 061102 (2016), [arXiv:1602.03837 \[gr-qc\]](#); “GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral,” *Phys. Rev. Lett.* **119**, 161101 (2017), [arXiv:1710.05832 \[gr-qc\]](#); “GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs,” *Phys. Rev. X* **9**, 031040 (2019), [arXiv:1811.12907 \[astro-ph.HE\]](#); R. Abbott *et al.* (LIGO Scientific, Virgo), “GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run,” (2020), [arXiv:2010.14527 \[gr-qc\]](#).
- [7] Bence Kocsis, Merse E. Gaspar, and Szabolcs Marka, “Detection rate estimates of gravity-waves emitted during parabolic encounters of stellar black holes in globular clusters,” *Astrophys. J.* **648**, 411–429 (2006), [arXiv:astro-ph/0603441](#); Sajal Mukherjee, Sanjit Mitra, and Sourav Chatterjee, “Detectability of hyperbolic encounters of compact stars with ground-based gravitational waves detectors,” (2020), [arXiv:2010.00916 \[gr-qc\]](#); Michael Zevin, Johan Samsing, Carl Rodriguez, Carl-Johan Haster, and Enrico Ramirez-Ruiz, “Eccentric Black Hole Mergers in Dense Star Clusters: The Role of Binary–Binary Encounters,” *Astrophys. J.* **871**, 91 (2019), [arXiv:1810.00901 \[astro-ph.HE\]](#).

- [8] Pürrer, Michael and Haster, Carl-Johan, “Gravitational waveform accuracy requirements for future ground-based detectors,” *Phys. Rev. Res.* **2**, 023151 (2020), [arXiv:1912.10055 \[gr-qc\]](#).
- [9] Luc Blanchet, “Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries,” *Living Rev. Rel.* **17**, 2 (2014), [arXiv:1310.1528 \[gr-qc\]](#); Schäfer, Gerhard and Jaranowski, Piotr, “Hamiltonian formulation of general relativity and post-Newtonian dynamics of compact binaries,” *Living Rev. Rel.* **21**, 7 (2018), [arXiv:1805.07240 \[gr-qc\]](#); Toshifumi Futamase and Yousuke Itoh, “The post-Newtonian approximation for relativistic compact binaries,” *Living Rev. Rel.* **10**, 2 (2007); Michael E. Pati and Clifford M. Will, “Post-Newtonian gravitational radiation and equations of motion via direct integration of the relaxed Einstein equations. 1. Foundations,” *Phys. Rev. D* **62**, 124015 (2000), [arXiv:gr-qc/0007087](#); LLuis Bel, T. Damour, N. Deruelle, J. Ibanez, and J. Martin, “Poincaré-invariant gravitational field and equations of motion of two pointlike objects: The postlinear approximation of general relativity,” *Gen. Rel. Grav.* **13**, 963–1004 (1981); Konradin Westpfahl, “High-Speed Scattering of Charged and Uncharged Particles in General Relativity,” *Fortsch. Phys.* **33**, 417–493 (1985); Tomas Ledvinka, Gerhard Schäfer, and Jiri Bicak, “Relativistic Closed-Form Hamiltonian for Many-Body Gravitating Systems in the Post-Minkowskian Approximation,” *Phys. Rev. Lett.* **100**, 251101 (2008), [arXiv:0807.0214 \[gr-qc\]](#).
- [10] Walter D. Goldberger and Ira Z. Rothstein, “An Effective field theory of gravity for extended objects,” *Phys. Rev. D* **73**, 104029 (2006), [arXiv:hep-th/0409156](#); “Towers of Gravitational Theories,” *Gen. Rel. Grav.* **38**, 1537–1546 (2006), [arXiv:hep-th/0605238](#); Walter D. Goldberger and Andreas Ross, “Gravitational radiative corrections from effective field theory,” *Phys. Rev. D* **81**, 124015 (2010), [arXiv:0912.4254 \[gr-qc\]](#).
- [11] Kälin, Gregor and Porto, Rafael A., “Post-Minkowskian Effective Field Theory for Conservative Binary Dynamics,” *JHEP* **11**, 106 (2020), [arXiv:2006.01184 \[hep-th\]](#).
- [12] Kälin, Gregor and Liu, Zhengwen and Porto, Rafael A., “Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach,” *Phys. Rev. Lett.* **125**, 261103 (2020), [arXiv:2007.04977 \[hep-th\]](#).
- [13] Walter D. Goldberger, “Les Houches lectures on effective field theories and gravitational radiation,” in *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime* (2007) [arXiv:hep-ph/0701129](#); Stefano Foffa and Riccardo Sturani, “Effective field theory methods to model compact binaries,” *Class. Quant. Grav.* **31**, 043001 (2014), [arXiv:1309.3474 \[gr-qc\]](#); Ira Z. Rothstein, “Progress in effective field theory approach to the binary inspiral problem,” *Gen. Rel. Grav.* **46**, 1726 (2014); Rafael A. Porto, “The effective field theorist’s approach to gravitational dynamics,” *Phys. Rept.* **633**, 1–104 (2016), [arXiv:1601.04914 \[hep-th\]](#); Michèle Levi, “Effective Field Theories of Post-Newtonian Gravity: A comprehensive review,” *Rept. Prog. Phys.* **83**, 075901 (2020), [arXiv:1807.01699 \[hep-th\]](#).
- [14] Duff Neill and Ira Z. Rothstein, “Classical Space-Times from the S Matrix,” *Nucl. Phys. B* **877**, 177–189 (2013), [arXiv:1304.7263 \[hep-th\]](#); N.E.J. Bjerrum-Bohr, John F. Donoghue, and Pierre Vanhove, “On-shell Techniques and Universal Results in Quantum Gravity,” *JHEP* **02**, 111 (2014), [arXiv:1309.0804 \[hep-th\]](#); Andrés Luna, Isobel Nicholson, Donal O’Connell, and Chris D. White, “Inelastic Black Hole Scattering from Charged Scalar Amplitudes,” *JHEP* **03**, 044 (2018), [arXiv:1711.03901 \[hep-th\]](#); N.E. J. Bjerrum-Bohr, Poul H. Damgaard, Guido Festuccia, Ludovic Planté, and Pierre Vanhove, “General Relativity from Scattering Amplitudes,” *Phys. Rev. Lett.* **121**, 171601 (2018), [arXiv:1806.04920 \[hep-th\]](#); Zvi Bern, Clifford Cheung, Radu Roiban, Chia-Hsien Shen, Mikhail P. Solon, and Mao Zeng, “Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order,” *Phys. Rev. Lett.* **122**, 201603 (2019), [arXiv:1901.04424 \[hep-th\]](#); “Black Hole Binary Dynamics from the Double Copy and Effective Theory,” *JHEP* **10**, 206 (2019), [arXiv:1908.01493 \[hep-th\]](#); Clifford Cheung and Mikhail P. Solon, “Classical gravitational scattering at $\mathcal{O}(G^3)$ from Feynman diagrams,” *JHEP* **06**, 144 (2020), [arXiv:2003.08351 \[hep-th\]](#).
- [15] Zvi Bern, Lance J. Dixon, David C. Dunbar, and David A. Kosower, “One loop n point gauge theory amplitudes, unitarity and collinear limits,” *Nucl. Phys. B* **425**, 217–260 (1994), [arXiv:hep-ph/9403226 \[hep-ph\]](#); “Fusing gauge theory tree amplitudes into loop amplitudes,” *Nucl. Phys. B* **435**, 59–101 (1995), [arXiv:hep-ph/9409265 \[hep-ph\]](#); Ruth Britto, Freddy Cachazo, and Bo Feng, “Generalized unitarity and one-loop amplitudes in N=4 super-Yang-Mills,” *Nucl. Phys. B* **725**, 275–305 (2005), [arXiv:hep-th/0412103 \[hep-th\]](#).
- [16] Z. Bern, J. J. M. Carrasco, and Henrik Johansson, “New Relations for Gauge-Theory Amplitudes,” *Phys. Rev. D* **78**, 085011 (2008), [arXiv:0805.3993 \[hep-ph\]](#); Zvi Bern, John Joseph M. Carrasco, and Henrik Johansson, “Perturbative Quantum Gravity as a Double Copy of Gauge Theory,” *Phys. Rev. Lett.* **105**, 061602 (2010), [arXiv:1004.0476 \[hep-th\]](#); Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson, and R. Roiban, “Simplifying Multiloop Integrands and Ultraviolet Divergences of Gauge Theory and Gravity Amplitudes,” *Phys. Rev. D* **85**, 105014 (2012), [arXiv:1201.5366 \[hep-th\]](#); Zvi Bern, John Joseph M. Carrasco, Wei-Ming Chen, Henrik Johansson, Radu Roiban, and Mao Zeng, “Five-loop four-point integrand of $N = 8$ supergravity as a generalized double copy,” *Phys. Rev. D* **96**, 126012 (2017), [arXiv:1708.06807 \[hep-th\]](#); Zvi Bern, John Joseph Carrasco, Wei-Ming Chen, Alex Edison, Henrik Johansson, Julio Parra-Martinez, Radu Roiban, and Mao Zeng, “Ultraviolet Properties of $\mathcal{N} = 8$ Supergravity at Five Loops,” *Phys. Rev. D* **98**, 086021 (2018), [arXiv:1804.09311 \[hep-th\]](#); Zvi Bern, John Joseph Carrasco, Marco Chiodaroli, Henrik Johansson, and Radu Roiban, “The Duality Between Color and Kinematics and its Applications,” (2019), [arXiv:1909.01358 \[hep-th\]](#).
- [17] David A. Kosower, Ben Maybee, and Donal O’Connell, “Amplitudes, Observables, and Classical Scattering,” *JHEP* **02**, 137 (2019), [arXiv:1811.10950 \[hep-th\]](#); Ben Maybee, Donal O’Connell, and Justin Vines, “Observables and amplitudes for spinning particles and black holes,” *JHEP* **12**, 156 (2019), [arXiv:1906.09260 \[hep-th\]](#); Thibault Damour, “Classical and quantum scattering in post-Minkowskian gravity,” *Phys. Rev. D* **102**, 024060 (2020), [arXiv:1912.02139 \[gr-qc\]](#).

- [18] Clifford Cheung, Ira Z. Rothstein, and Mikhail P. Solon, “From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion,” *Phys. Rev. Lett.* **121**, 251101 (2018), [arXiv:1808.02489 \[hep-th\]](#); Varun Vaidya, “Gravitational spin Hamiltonians from the S matrix,” *Phys. Rev.* **D91**, 024017 (2015), [arXiv:1410.5348 \[hep-th\]](#); Thibault Damour, “High-energy gravitational scattering and the general relativistic two-body problem,” *Phys. Rev.* **D97**, 044038 (2018), [arXiv:1710.10599 \[gr-qc\]](#).
- [19] Thibault Damour, “Radiative contribution to classical gravitational scattering at the third order in G ,” *Phys. Rev. D* **102**, 124008 (2020), [arXiv:2010.01641 \[gr-qc\]](#).
- [20] Paolo Di Vecchia, Carlo Heissenberg, Rodolfo Russo, and Gabriele Veneziano, “Universality of ultra-relativistic gravitational scattering,” *Phys. Lett. B* **811**, 135924 (2020), [arXiv:2008.12743 \[hep-th\]](#).
- [21] Zvi Bern, Julio Parra-Martinez, Radu Roiban, Michael S. Ruf, Chia-Hsien Shen, Mikhail P. Solon, and Mao Zeng, “Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$,” (2021), [arXiv:2101.07254 \[hep-th\]](#); Kälín, Gregor and Porto, Rafael A., “From Boundary Data to Bound States,” *JHEP* **01**, 072 (2020), [arXiv:1910.03008 \[hep-th\]](#); “From boundary data to bound states. Part II. Scattering angle to dynamical invariants (with twist),” *JHEP* **02**, 120 (2020), [arXiv:1911.09130 \[hep-th\]](#).
- [22] Luc Blanchet and Athanassios S. Fokas, “Equations of motion of self-gravitating N -body systems in the first post-Minkowskian approximation,” *Phys. Rev. D* **98**, 084005 (2018), [arXiv:1806.08347 \[gr-qc\]](#); Andrea Cristofoli, N. E. J. Bjerrum-Bohr, Poul H. Damgaard, and Pierre Vanhove, “Post-Minkowskian Hamiltonians in general relativity,” *Phys. Rev.* **D100**, 084040 (2019), [arXiv:1906.01579 \[hep-th\]](#); Andrea Cristofoli, Poul H. Damgaard, Paolo Di Vecchia, and Carlo Heissenberg, “Second-order Post-Minkowskian scattering in arbitrary dimensions,” *JHEP* **07**, 122 (2020), [arXiv:2003.10274 \[hep-th\]](#); Donato Bini, Thibault Damour, Andrea Geralico, Stefano Laporta, and Pierpaolo Mastrolia, “Gravitational dynamics at $\mathcal{O}(G^6)$: perturbative gravitational scattering meets experimental mathematics,” (2020), [arXiv:2008.09389 \[gr-qc\]](#); “Gravitational scattering at the seventh order in G : nonlocal contribution at the sixth post-Newtonian accuracy,” (2020), [arXiv:2012.12918 \[gr-qc\]](#); Florian Loebbert, Jan Plefka, Caxin Shi, and Tianheng Wang, “Three-Body Effective Potential in General Relativity at 2PM and Resulting PN Contributions,” (2020), [arXiv:2012.14224 \[hep-th\]](#).
- [23] Justin Vines, “Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings,” *Class. Quant. Grav.* **35**, 084002 (2018), [arXiv:1709.06016 \[gr-qc\]](#); Donato Bini and Thibault Damour, “Gravitational spin-orbit coupling in binary systems, post-Minkowskian approximation and effective one-body theory,” *Phys. Rev.* **D96**, 104038 (2017), [arXiv:1709.00590 \[gr-qc\]](#); “Gravitational spin-orbit coupling in binary systems at the second post-Minkowskian approximation,” *Phys. Rev.* **D98**, 044036 (2018), [arXiv:1805.10809 \[gr-qc\]](#); Alfredo Guevara, “Holomorphic Classical Limit for Spin Effects in Gravitational and Electromagnetic Scattering,” *JHEP* **04**, 033 (2019), [arXiv:1706.02314 \[hep-th\]](#); Justin Vines, Jan Steinhoff, and Alessandra Buonanno, “Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order,” *Phys. Rev.* **D99**, 064054 (2019), [arXiv:1812.00956 \[gr-qc\]](#); Alfredo Guevara, Alexander Ochirov, and Justin Vines, “Scattering of Spinning Black Holes from Exponentiated Soft Factors,” *JHEP* **09**, 056 (2019), [arXiv:1812.06895 \[hep-th\]](#); Ming-Zhi Chung, Yu-Tin Huang, Jung-Wook Kim, and Sangmin Lee, “The simplest massive S-matrix: from minimal coupling to Black Holes,” *JHEP* **04**, 156 (2019), [arXiv:1812.08752 \[hep-th\]](#); Alfredo Guevara, Alexander Ochirov, and Justin Vines, “Black-hole scattering with general spin directions from minimal-coupling amplitudes,” *Phys. Rev.* **D100**, 104024 (2019), [arXiv:1906.10071 \[hep-th\]](#); Ming-Zhi Chung, Yu-Tin Huang, and Jung-Wook Kim, “Classical potential for general spinning bodies,” *JHEP* **09**, 074 (2020), [arXiv:1908.08463 \[hep-th\]](#); Poul H. Damgaard, Kays Haddad, and Andreas Helset, “Heavy Black Hole Effective Theory,” *JHEP* **11**, 070 (2019), [arXiv:1908.10308 \[hep-ph\]](#); Rafael Aoude, Kays Haddad, and Andreas Helset, “On-shell heavy particle effective theories,” *JHEP* **05**, 051 (2020), [arXiv:2001.09164 \[hep-th\]](#); Zvi Bern, Andres Luna, Radu Roiban, Chia-Hsien Shen, and Mao Zeng, “Spinning Black Hole Binary Dynamics, Scattering Amplitudes and Effective Field Theory,” (2020), [arXiv:2005.03071 \[hep-th\]](#); Alfredo Guevara, Ben Maybee, Alexander Ochirov, Donal O’Connell, and Justin Vines, “A worldsheet for Kerr,” (2020), [arXiv:2012.11570 \[hep-th\]](#).
- [24] Donato Bini, Thibault Damour, and Andrea Geralico, “Scattering of tidally interacting bodies in post-Minkowskian gravity,” *Phys. Rev.* **D101**, 044039 (2020), [arXiv:2001.00352 \[gr-qc\]](#); Clifford Cheung and Mikhail P. Solon, “Tidal Effects in the Post-Minkowskian Expansion,” *Phys. Rev. Lett.* **125**, 191601 (2020), [arXiv:2006.06665 \[hep-th\]](#); Kays Haddad and Andreas Helset, “Tidal effects in quantum field theory,” *JHEP* **12**, 024 (2020), [arXiv:2008.04920 \[hep-th\]](#); Kälín, Gregor and Liu, Zhengwen and Porto, Rafael A., “Conservative Tidal Effects in Compact Binary Systems to Next-to-Leading Post-Minkowskian Order,” *Phys. Rev.* **D102**, 124025 (2020), [arXiv:2008.06047 \[hep-th\]](#); Andreas Brandhuber and Gabriele Travaglini, “On higher-derivative effects on the gravitational potential and particle bending,” *JHEP* **01**, 010 (2020), [arXiv:1905.05657 \[hep-th\]](#); Manuel Accettulli Huber, Andreas Brandhuber, Stefano De Angelis, and Gabriele Travaglini, “Note on the absence of R^2 corrections to Newton’s potential,” *Phys. Rev.* **D101**, 046011 (2020), [arXiv:1911.10108 \[hep-th\]](#); “Eikonal phase matrix, deflection angle and time delay in effective field theories of gravity,” *Phys. Rev.* **D102**, 046014 (2020), [arXiv:2006.02375 \[hep-th\]](#); Zvi Bern, Julio Parra-Martinez, Radu Roiban, Eric Sawyer, and Chia-Hsien Shen, “Leading Nonlinear Tidal Effects and Scattering Amplitudes,” (2020), [arXiv:2010.08559 \[hep-th\]](#); Clifford Cheung, Nabha Shah, and Mikhail P. Solon, “Mining the Geodesic Equation for Scattering Data,” *Phys. Rev.* **D103**, 024030 (2021), [arXiv:2010.08568 \[hep-th\]](#); Rafael Aoude, Kays Haddad, and Andreas Helset, “Tidal effects for spinning particles,” (2020), [arXiv:2012.05256 \[hep-th\]](#).
- [25] D. Amati, M. Ciafaloni, and G. Veneziano, “Higher Order Gravitational Deflection and Soft Bremsstrahlung in Planckian Energy Superstring Collisions,” *Nucl. Phys.*

- B347**, 550–580 (1990); Paolo Di Vecchia, Andrés Luna, Stephen G. Naculich, Rodolfo Russo, Gabriele Veneziano, and Chris D. White, “A tale of two exponentiations in $\mathcal{N} = 8$ supergravity,” *Phys. Lett.* **B798**, 134927 (2019), [arXiv:1908.05603 \[hep-th\]](#); Paolo Di Vecchia, Stephen G. Naculich, Rodolfo Russo, Gabriele Veneziano, and Chris D. White, “A tale of two exponentiations in $\mathcal{N} = 8$ supergravity at subleading level,” *JHEP* **03**, 173 (2020), [arXiv:1911.11716 \[hep-th\]](#); Zvi Bern, Harald Ita, Julio Parra-Martinez, and Michael S. Ruf, “Universality in the classical limit of massless gravitational scattering,” *Phys. Rev. Lett.* **125**, 031601 (2020), [arXiv:2002.02459 \[hep-th\]](#); Manuel Accettulli Huber, Andreas Brandhuber, Stefano De Angelis, and Gabriele Travaglini, “From amplitudes to gravitational radiation with cubic interactions and tidal effects,” (2020), [arXiv:2012.06548 \[hep-th\]](#); Paolo Di Vecchia, Carlo Heissenberg, Rodolfo Russo, and Gabriele Veneziano, “Radiation Reaction from Soft Theorems,” (2021), [arXiv:2101.05772 \[hep-th\]](#); Yilber Fabian Bautista and Alfredo Guevara, “From Scattering Amplitudes to Classical Physics: Universality, Double Copy and Soft Theorems,” (2019), [arXiv:1903.12419 \[hep-th\]](#); Alok Laddha and Ashoke Sen, “Gravity Waves from Soft Theorem in General Dimensions,” *JHEP* **09**, 105 (2018), [arXiv:1801.07719 \[hep-th\]](#); “Logarithmic Terms in the Soft Expansion in Four Dimensions,” *JHEP* **10**, 056 (2018), [arXiv:1804.09193 \[hep-th\]](#); Biswajit Sahoo and Ashoke Sen, “Classical and Quantum Results on Logarithmic Terms in the Soft Theorem in Four Dimensions,” *JHEP* **02**, 086 (2019), [arXiv:1808.03288 \[hep-th\]](#); Alok Laddha and Ashoke Sen, “Classical proof of the classical soft graviton theorem in $D > 4$,” *Phys. Rev.* **D101**, 084011 (2020), [arXiv:1906.08288 \[gr-qc\]](#); Arnab Priya Saha, Biswajit Sahoo, and Ashoke Sen, “Proof of the classical soft graviton theorem in $D = 4$,” *JHEP* **06**, 153 (2020), [arXiv:1912.06413 \[hep-th\]](#); Manu A, Debidirna Ghosh, Alok Laddha, and P. V. Athira, “Soft Radiation from Scattering Amplitudes Revisited,” (2020), [arXiv:2007.02077 \[hep-th\]](#); Biswajit Sahoo, “Classical Sub-subleading Soft Photon and Soft Graviton Theorems in Four Spacetime Dimensions,” *JHEP* **12**, 070 (2020), [arXiv:2008.04376 \[hep-th\]](#).
- [26] Gustav Mogull, Jan Plefka, and Jan Steinhoff, “Classical black hole scattering from a worldline quantum field theory,” (2020), [arXiv:2010.02865 \[hep-th\]](#).
- [27] Enrico Herrmann, Julio Parra-Martinez, Michael S. Ruf, and Mao Zeng, “Gravitational Bremsstrahlung from Reverse Unitarity,” (2021), [arXiv:2101.07255 \[hep-th\]](#).
- [28] Sigurd Sannan, “Gravity as the Limit of the Type {II} Superstring Theory,” *Phys. Rev. D* **34**, 1749 (1986).
- [29] Steven Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (John Wiley and Sons, New York, 1972).
- [30] Béatrice Bonga and Eric Poisson, “Coulombic contribution to angular momentum flux in general relativity,” *Phys. Rev. D* **99**, 064024 (2019), [arXiv:1808.01288 \[gr-qc\]](#).
- [31] Damour, Thibault and Jaranowski, Piotr and Schäfer, Gerhard, “Nonlocal-in-time action for the fourth post-Newtonian conservative dynamics of two-body systems,” *Phys. Rev.* **D89**, 064058 (2014), [arXiv:1401.4548 \[gr-qc\]](#); “Conservative dynamics of two-body systems at the fourth post-Newtonian approximation of general relativity,” *Phys. Rev.* **D93**, 084014 (2016), [arXiv:1601.01283 \[gr-qc\]](#); Laura Bernard, Luc Blanchet, Alejandro Bohé, Guillaume Faye, and Sylvain Marsat, “Energy and periastron advance of compact binaries on circular orbits at the fourth post-Newtonian order,” *Phys. Rev.* **D95**, 044026 (2017), [arXiv:1610.07934 \[gr-qc\]](#); Stefano Foffa, Pierpaolo Mastrolia, Riccardo Sturani, and Christian Sturm, “Effective field theory approach to the gravitational two-body dynamics, at fourth post-Newtonian order and quintic in the Newton constant,” *Phys. Rev.* **D95**, 104009 (2017), [arXiv:1612.00482 \[gr-qc\]](#); Thibault Damour and Piotr Jaranowski, “Four-loop static contribution to the gravitational interaction potential of two point masses,” *Phys. Rev.* **D95**, 084005 (2017), [arXiv:1701.02645 \[gr-qc\]](#); Stefano Foffa and Riccardo Sturani, “Conservative dynamics of binary systems to fourth Post-Newtonian order in the EFT approach I: Regularized Lagrangian,” *Phys. Rev.* **D100**, 024047 (2019), [arXiv:1903.05113 \[gr-qc\]](#); Stefano Foffa, Rafael A. Porto, Ira Rothstein, and Riccardo Sturani, “Conservative dynamics of binary systems to fourth Post-Newtonian order in the EFT approach II: Renormalized Lagrangian,” *Phys. Rev.* **D100**, 024048 (2019), [arXiv:1903.05118 \[gr-qc\]](#); Blümlein, J. and Maier, A. and Marquard, P. and Schäfer, G., “Fourth post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach,” *Nucl. Phys. B* **955**, 115041 (2020), [arXiv:2003.01692 \[gr-qc\]](#); Rafael A. Porto and Ira Z. Rothstein, “Apparent ambiguities in the post-Newtonian expansion for binary systems,” *Phys. Rev.* **D96**, 024062 (2017), [arXiv:1703.06433 \[gr-qc\]](#); Tanguy Marchand, Laura Bernard, Luc Blanchet, and Guillaume Faye, “Ambiguity-Free Completion of the Equations of Motion of Compact Binary Systems at the Fourth Post-Newtonian Order,” *Phys. Rev.* **D97**, 044023 (2018), [arXiv:1707.09289 \[gr-qc\]](#); Chad R. Galley, Adam K. Leibovich, Rafael A. Porto, and Andreas Ross, “Tail effect in gravitational radiation reaction: Time nonlocality and renormalization group evolution,” *Phys. Rev.* **D93**, 124010 (2016), [arXiv:1511.07379 \[gr-qc\]](#); Stefano Foffa, Pierpaolo Mastrolia, Riccardo Sturani, Christian Sturm, and William J. Torres Bobadilla, “Static two-body potential at fifth post-Newtonian order,” *Phys. Rev. Lett.* **122**, 241605 (2019), [arXiv:1902.10571 \[gr-qc\]](#); Blümlein, J. and Maier, A. and Marquard, P., “Five-Loop Static Contribution to the Gravitational Interaction Potential of Two Point Masses,” *Phys. Lett.* **B800**, 135100 (2020), [arXiv:1902.11180 \[gr-qc\]](#); Donato Bini, Thibault Damour, and Andrea Geralico, “Novel approach to binary dynamics: application to the fifth post-Newtonian level,” *Phys. Rev. Lett.* **123**, 231104 (2019), [arXiv:1909.02375 \[gr-qc\]](#); Blümlein, J. and Maier, A. and Marquard, P. and Schäfer, G., “The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach: potential contributions,” (2020), [arXiv:2010.13672 \[gr-qc\]](#); “Testing binary dynamics in gravity at the sixth post-Newtonian level,” *Phys. Lett.* **B807**, 135496 (2020), [arXiv:2003.07145 \[gr-qc\]](#); Donato Bini, Thibault Damour, and Andrea Geralico, “Sixth post-Newtonian local-in-time dynamics of binary systems,” *Phys. Rev.* **D102**, 024061 (2020), [arXiv:2004.05407 \[gr-qc\]](#); “Binary dynamics at the fifth and fifth-and-a-half post-Newtonian orders,”

- Phys. Rev. **D102**, 024062 (2020), [arXiv:2003.11891 \[gr-qc\]](#); “Sixth post-Newtonian nonlocal-in-time dynamics of binary systems,” Phys. Rev. **D102**, 084047 (2020), [arXiv:2007.11239 \[gr-qc\]](#); Blümlein, J. and Maier, A. and Marquard, P. and Schäfer, G., “The 6th Post-Newtonian Potential Terms at $O(G_N^4)$,” (2021), [arXiv:2101.08630 \[gr-qc\]](#); Luc Blanchet, Bala R. Iyer, and Benoit Joguet, “Gravitational waves from inspiralling compact binaries: Energy flux to third post-Newtonian order,” Phys. Rev. **D65**, 064005 (2002), [Erratum: Phys. Rev. **D71**, 129903(2005)], [arXiv:gr-qc/0105098 \[gr-qc\]](#); Luc Blanchet, Thibault Damour, Gilles Esposito-Farese, and Bala R. Iyer, “Gravitational radiation from inspiralling compact binaries completed at the third post-Newtonian order,” Phys. Rev. Lett. **93**, 091101 (2004), [arXiv:gr-qc/0406012 \[gr-qc\]](#); Luc Blanchet, Guillaume Faye, Bala R. Iyer, and Siddhartha Sinha, “The Third post-Newtonian gravitational wave polarisations and associated spherical harmonic modes for inspiralling compact binaries in quasi-circular orbits,” Class. Quant. Grav. **25**, 165003 (2008), [Erratum: Class. Quant. Grav. **29**, 239501(2012)], [arXiv:0802.1249 \[gr-qc\]](#).
- [32] Michèle Levi, Andrew J. Mcleod, and Matthew Von Hippel, “ N^3 LO gravitational spin-orbit coupling at order G^4 ,” (2020), [arXiv:2003.02827 \[hep-th\]](#); Andrea Antonelli, Chris Kavanagh, Mohammed Khalil, Jan Steinhoff, and Justin Vines, “Gravitational spin-orbit coupling through third-subleading post-Newtonian order: from first-order self-force to arbitrary mass ratios,” Phys. Rev. Lett. **125**, 011103 (2020), [arXiv:2003.11391 \[gr-qc\]](#); Michele Levi and Jan Steinhoff, “Complete conservative dynamics for inspiralling compact binaries with spins at fourth post-Newtonian order,” (2016), [arXiv:1607.04252 \[gr-qc\]](#); Michèle Levi, Andrew J. Mcleod, and Matthew Von Hippel, “NNNLO gravitational quadratic-in-spin interactions at the quartic order in G ,” (2020), [arXiv:2003.07890 \[hep-th\]](#); Michèle Levi, Stavros Mougiakakos, and Mariana Vieira, “Gravitational cubic-in-spin interaction at the next-to-leading post-Newtonian order,” JHEP **01**, 036 (2021), [arXiv:1912.06276 \[hep-th\]](#); Michèle Levi and Fei Teng, “NLO gravitational quartic-in-spin interaction,” JHEP **01**, 066 (2021), [arXiv:2008.12280 \[hep-th\]](#); Adam K. Leibovich, Natália T. Maia, Ira Z. Rothstein, and Zixin Yang, “Second post-Newtonian order radiative dynamics of inspiralling compact binaries in the Effective Field Theory approach,” Phys. Rev. **D101**, 084058 (2020), [arXiv:1912.12546 \[gr-qc\]](#); Chandra Kant Mishra, Aditya Kela, K. G. Arun, and Guillaume Faye, “Ready-to-use post-Newtonian gravitational waveforms for binary black holes with nonprecessing spins: An update,” Phys. Rev. **D93**, 084054 (2016), [arXiv:1601.05588 \[gr-qc\]](#); Alessandra Buonanno, Guillaume Faye, and Tanja Hinderer, “Spin effects on gravitational waves from inspiraling compact binaries at second post-Newtonian order,” Phys. Rev. **D87**, 044009 (2013), [arXiv:1209.6349 \[gr-qc\]](#); Rafael A. Porto, Andreas Ross, and Ira Z. Rothstein, “Spin induced multipole moments for the gravitational wave flux from binary inspirals to third Post-Newtonian order,” JCAP **1103**, 009 (2011), [arXiv:1007.1312 \[gr-qc\]](#); “Spin induced multipole moments for the gravitational wave amplitude from binary inspirals to 2.5 Post-Newtonian order,” JCAP **1209**, 028 (2012), [arXiv:1203.2962 \[gr-qc\]](#); Natalia T. Maia, Chad R. Galley, Adam K. Leibovich, and Rafael A. Porto, “Radiation reaction for spinning bodies in effective field theory I: Spin-orbit effects,” Phys. Rev. **D96**, 084064 (2017), [arXiv:1705.07934 \[gr-qc\]](#); “Radiation reaction for spinning bodies in effective field theory II: Spin-spin effects,” Phys. Rev. **D96**, 084065 (2017), [arXiv:1705.07938 \[gr-qc\]](#).
- [33] Thibault Damour, “Gravitational scattering, post-Minkowskian approximation and Effective One-Body theory,” Phys. Rev. **D94**, 104015 (2016), [arXiv:1609.00354 \[gr-qc\]](#).
- [34] Andrea Antonelli, Alessandra Buonanno, Jan Steinhoff, Maarten van de Meent, and Justin Vines, “Energetics of two-body Hamiltonians in post-Minkowskian gravity,” Phys. Rev. **D99**, 104004 (2019), [arXiv:1901.07102 \[gr-qc\]](#).
- [35] Julio Parra-Martinez, Michael S. Ruf, and Mao Zeng, “Extremal black hole scattering at $\mathcal{O}(G^3)$: graviton dominance, eikonal exponentiation, and differential equations,” JHEP **11**, 023 (2020), [arXiv:2005.04236 \[hep-th\]](#).
- [36] Donato Bini and Thibault Damour, “Gravitational radiation reaction along general orbits in the effective one-body formalism,” Phys. Rev. **D86**, 124012 (2012), [arXiv:1210.2834 \[gr-qc\]](#).