

## Clay mineralogy and its relationship with potassium forms in some paddy and non-paddy soils of northern Iran

Alireza Raheb\*, Ahmad Heidari

Soil Science Department, Faculty of Agricultural Engineering and Technology, College of Agriculture & Natural Resources, University of Tehran, Karaj, Iran

\*Corresponding author: araheb@ut.ac.ir

### Abstract

Forms of potassium in soils have been a key focus in soil-fertility studies. Clay minerals are the main source of plant nutrients in soil, as their specific surface characteristics determine the release pattern of some important nutrients, such as potassium. Three forms of K consist of unavailable, slowly available and readily available exist in equilibrium in the soil system. Relationships between soil potassium forms within the clay mineralogical suite were determined in twenty four soil samples in three pedons from paddy soils, two from kiwifruit and one from citrus land in the north of Iran. The results showed that, the characteristics of different forms of potassium in adjacent pedons under different land uses depended on their specific clay mineralogy. Mineralogical analyses showed that smectite, illite and vermiculite were the most abundant clay minerals in the studied soils. Mean soluble and exchangeable potassium in paddy soils were rather low compared to non-paddy soils due to potassium fertilization in non-paddy soils. Non-exchangeable potassium in the samples, which was dominated by smectite, was lower than that in samples dominated by vermiculite, hydroxy-interlayer vermiculite (HIV) and illite; this was related to the inability of smectite in potassium fixation. Therefore, taking into account both exchangeable and HNO<sub>3</sub>-extractable K gives a better indication for K potential and soil-quality management.

**Keywords:** Exchangeable and non-exchangeable K; Kiwi fruit; Mineralogy; Smectite.

**Abbreviations:** Illi- Illite; HI- Hydroxy Interlayer Vermiculite; Smc- Smectite; Kao- Kaolinite; Ver- Vermiculite; Cl- Chlorite; Mix- Mixed mineral.

### Introduction

Southern and southeastern Asia contains the world's most important paddy soils, comprising about 126 million ha of the worldwide total of 135 million ha (IRRI, 2002). Climate and landforms are the two determinants for this strongly biased distribution of paddy lands and their fertility (Kyuma, 2004). Rice is the staple food for billions of people worldwide (Akbar et al, 2011) and is one of the most important grains in the world (Rahimi Petroudi et al, 2011). Rice cultivation in paddy field is of great importance in the north of Iran, especially in Mazandaran and Guilan provinces with totally 78% of cultivation area and rice yield (Rahimi Petroudi et al, 2011). Potassium is an essential and major nutrient for crop production in paddy soils (Sparks and Huang, 1985). The exact function of K in plant growth has not been clearly defined, although its association with the movement of water, nutrients and carbohydrates in plant tissue has repeatedly been reported. Soils commonly contain over 20,000 ppm of total potassium (McLean and Watson, 1985). Nearly all soil K is in a structural form, and thus not available for plant growth. Plants can use only the exchangeable potassium on the surface of soil particles and potassium dissolved in the soil water (these often amount to less than 100 ppm) (Rehm and Schmitt, 2002). Soil potassium is classified in three forms according to availability: unavailable, slowly available and readily available or exchangeable (Sharpley, 1989; Bhonsle, 1992). Crystalline and insoluble soil K, which is in an unavailable

form, comprises approximately 90-98% of total soil K (Sparks, 1987; Martin and Sparks, 1985). Slowly available potassium, which is fixed and non-exchangeable, is the form trapped between the layers or sheets of certain kinds of clay minerals; plants can use only very little of it during a single growing season (Sparks, 1987). The major sources of non-exchangeable K in soils are K-rich 2:1 clay minerals such as micas and vermiculite (Sparks, 1987; Moritsuka et al. 2004). Mica (illite) clays also fix K between their layers when they become dry, but do not release all of the fixed K when wet. This fixation without release causes problems in the paddy-soil regions for management of potash fertilizers in crop production. K release from the interlayer of these minerals is very slow, depending on their weathering stage (Rehm and Schmitt, 2002). The release of K from clay minerals is influenced by particle size and chemical composition (Huang, 2005). It is generally accepted that trioctahedral micas, such as biotite and phlogopite, release K more readily than dioctahedral ones, such as muscovite (Fanning et al., 1989). Readily available potassium is a dissolved form of K (water-soluble) or held on the surface of clay particles (exchangeable K). Different articles have reported that plants can take up both exchangeable and non-exchangeable forms of potassium. Bhonsel, et al. (1992) suggested that fixed or non-exchangeable forms of K can be the main source of potassium for plants. Minerals' K release to soluble and exchangeable forms and its adsorption by exchange sites

**Table 1.** Land use, soil classification and some physico-chemical properties of the studied pedons


| Horizon                           | Depth<br>cm | Clay<br>% | Tex. | pH   | EC<br>$\mu\text{S m}^{-1}$ | OC<br>% | CEC<br>$\text{cmol}^+ \text{kg}^{-1}$ | $\text{ppm}$     |                 |                    |
|-----------------------------------|-------------|-----------|------|------|----------------------------|---------|---------------------------------------|------------------|-----------------|--------------------|
|                                   |             |           |      |      |                            |         |                                       | $K_{\text{sol}}$ | $K_{\text{ex}}$ | $K_{\text{nonex}}$ |
| (P1)Citrus: Oxiaquic Udipsamments |             |           |      |      |                            |         |                                       |                  |                 |                    |
| Ap                                | 0-13        | 15.6      | SL   | 7.85 | 1022                       | 1.97    | 15.18                                 | 17.94            | 120.9           | 641.9              |
| AC                                | 13-23       | 9.6       | LS   | 8.08 | 557                        | 0.4     | 7.91                                  | 4.29             | 35.1            | 335.4              |
| C1                                | 23-61       | 5.6       | S    | 8.13 | 347                        | 0.2     | 4.75                                  | 2.11             | 29.64           | 259.6              |
| C2                                | 61-100      | 3.6       | S    | 8.38 | 336                        | 0.2     | 4.9                                   | 2.53             | 26.13           | 284.3              |
| (P2) Paddy: Typic Endoaqualfs     |             |           |      |      |                            |         |                                       |                  |                 |                    |
| Apg                               | 0-12        | 49.9      | C    | 7.85 | 1030                       | 3.75    | 21.56                                 | 1.95             | 58.11           | 253.1              |
| Bg                                | 12-25       | 43.6      | C    | 7.89 | 1022                       | 3.75    | 25.15                                 | 1.95             | 65.52           | 276.9              |
| Btg1                              | 25-60       | 45.6      | SiC  | 8.18 | 563                        | 1.97    | 21.35                                 | 1.56             | 51.48           | 696.9              |
| Btg2                              | 60-100      | 43.6      | SiC  | 8.12 | 524                        | 0.6     | 18.03                                 | 1.56             | 52.26           | 188.6              |
| (P3) Paddy: Typic Endoaquepts     |             |           |      |      |                            |         |                                       |                  |                 |                    |
| Apg                               | 0-20        | 43.6      | C    | 7.79 | 1095                       | 0.79    | 21.04                                 | 2.73             | 54.6            | 189.9              |
| 2Bg1                              | 20-40       | 15.6      | SL   | 7.92 | 1119                       | 0.2     | 9.65                                  | 1.95             | 24.57           | 198.3              |
| 2Bg2                              | 40-75       | 27.6      | CL   | 7.93 | 866                        | 0.6     | 12.33                                 | 2.34             | 50.7            | 139.8              |
| (P4) Paddy: Typic Endoaqualfs     |             |           |      |      |                            |         |                                       |                  |                 |                    |
| Apg                               | 0-14        | 52.16     | C    | 7.81 | 1055                       | 2.57    | 24.83                                 | 2.34             | 82.29           | 299                |
| Btg1                              | 14-35       | 53.6      | C    | 8.09 | 716                        | 1.38    | 24.05                                 | 1.56             | 69.03           | 313.1              |
| Btg2                              | 35-50       | 43.6      | C    | 8.22 | 501                        | 0.99    | 18.81                                 | 2.73             | 59.67           | 229                |
| 2Bg1                              | 50-75       | 17.6      | SL   | 7.99 | 917                        | 0.39    | 9.65                                  | 2.11             | 29.64           | 180.1              |
| 3Bg2                              | 75-110      | 27.6      | L    | 7.98 | 841                        | 0.99    | 12.02                                 | 2.34             | 44.85           | 209                |
| (P5) kiwi: Typic Endoaquepts      |             |           |      |      |                            |         |                                       |                  |                 |                    |
| Ap                                | 0-12        | 42.16     | C    | 7.73 | 1005                       | 6.71    | 30.36                                 | 6.63             | 110.8           | 281                |
| Bg1                               | 12-27       | 35.6      | CL   | 7.56 | 882                        | 5.33    | 32.59                                 | 3.12             | 56.55           | 247.5              |
| Bg2                               | 27-45       | 29.6      | CL   | 7.29 | 1459                       | 5.92    | 36.22                                 | 3.12             | 41.34           | 175.5              |
| 2Cg                               | 45-100      | 19.6      | SL   | 7.28 | 2130                       | 2.76    | 17.22                                 | 3.51             | 30.03           | 207.3              |
| (P6) kiwi: Oxiaquic Hapludalfs    |             |           |      |      |                            |         |                                       |                  |                 |                    |
| Apg                               | 0-20        | 62.5      | C    | 7.56 | 1066                       | 5.33    | 34.01                                 | 10.53            | 164.9           | 268.5              |
| Bg                                | 20-38       | 43.6      | C    | 8.13 | 587                        | 0.99    | 24.21                                 | 1.95             | 62.01           | 201.6              |
| Btg1                              | 38-45       | 43.6      | SiC  | 8.28 | 494                        | 0.99    | 23.25                                 | 1.56             | 49.92           | 218.5              |
| Btg2                              | 45-70       | 31.6      | CL   | 8.26 | 578                        | 0.6     | 18.19                                 | 2.34             | 54.99           | 216.7              |

depends on the equilibrium between different phases of soil K (McLean and Watson, 1985), which may be affected by such factors as root uptake, fertilizer K applied, soil moisture, soil pH and soil temperature (Sparks, 1987). The relationship between clay mineralogy composition and potassium forms has been demonstrated by several studies (Sharply, 1989; Bhonsle et al., 1992; Liu et al., 1997; Ghosh and Singh, 2001; Surapaneni et al., 2002; Srinivasarao et al., 2006). Knowledge of clay mineralogy is crucial to understanding soils' nutritional status and nutrient-supplying power. Mineralogical composition of soils may have a considerable influence on K dynamics (Surapaneni et al., 2002). Relationships between clay mineralogy and potassium forms can be used in evaluating potential soil K fertility, prediction of K cycling and plant uptake (Sharply, 1989). Information on the  $\text{NH}_4\text{OAc-K}$  form of potassium along with knowledge of clays' mineralogical composition can provide insights into the equilibrium and release of non-exchangeable K to plants and the need for K fertilizers (Bhonsle, 1992). Soils with high 2:1 clay minerals (micas, vermiculite and high-layer-charge smectite) contain larger amounts of  $\text{HNO}_3$ -extractable K than kaolinite and other siliceous minerals (Martin and Sparks, 1985; Sharply, 1989; Ghosh and Singh, 2001). Bhonsle et al. (1992) stated that kaolinitic soils had low levels of  $\text{NH}_4\text{OAc-K}$ , mixed and illitic soils had medium levels and smectitic soils had high levels; their estimates for  $\text{HNO}_3\text{-K}$ , showed that kaolinitic, mixed and smectitic soils had low levels, and illitic soils had high. Their study set out the proportion of  $\text{NH}_4\text{OAc-K}$  to  $\text{HNO}_3\text{-K}$  in the sequence smectitic > kaolinitic > mixed > illitic. Sharpley (1989) stated

that determining both exchangeable and  $\text{HNO}_3\text{-K}$  could give a better indication of soil's potential K-supplying power. There is no report on the clay mineralogy and potassium status of the land used for different agricultural purposes in Iran (although in 1998 Towfighi estimated that available potassium in more than 100,000 ha of Iranian paddy soils varies between lower-than-critical to moderate). Therefore, the objectives of present study are: (1) to compare different forms of potassium (soluble K, exchangeable K and non-exchangeable K) in paddy soils and non-paddy soils, and (2) to compare the relationships between forms of K and clay mineralogy in paddy soils, and land uses under citrus and kiwi fruit cultivation.

## Results and discussion

The paddy soils showed very distinct morphological, physical and chemical properties from those of non-paddy soils. Table 1 shows some physicochemical characteristics of the studied pedons, including pH, EC, CEC, OC, PSD and ammonium acetate and the forms of potassium that are extractable using nitric acid. pH values of saturated paste in the paddy and non-paddy pedons varied between 7.79 and 8.22 for paddy soils, and between 7.28 and 8.38 for non-paddy soils (Table 1). Electrical conductivity of saturated extracts (ECe) ranged between 336 and 2130  $\mu\text{S m}^{-1}$ . Relatively strong differences (0.2 to 6.71 %) were observed in the horizontal (between pedons) and vertical (between horizons) distribution of soils' organic carbon. The largest amount of organic carbon was observed in the surface layer



**Fig 1.** Map of studied region in Mazandaran province, Iran.

of the kiwi fruit soil (pedons 5 and 6) due to higher decay of the surface litter and rotting of roots caused by very poor drainage. Cation exchange capacities (CEC) varied between 4.75 (pedon 1) to 36.22 (pedon 5)  $\text{cmol}^+ \text{kg}^{-1}$ . Kyuma (1984) also found an average CEC for some paddy soils from tropical Asia of 18.6  $\text{cmol}^+ \text{kg}^{-1}$ . The lowest CEC was obtained in the deeper horizons of the citrus pedon, due to its lower clay and organic-matter contents. In all studied pedons, the highest CEC values were found in the surface layers as a result of their higher organic carbon contents. The paddy soils had the highest clay content among the studied pedons. The highest water-soluble K content was observed in the citrus pedons, followed by kiwifruit and paddy. In contrast, the highest  $\text{NH}_4\text{OAc-K}$  was observed in the kiwi fruit pedons, followed by citrus and paddy. For  $\text{HNO}_3\text{-K}$ , the citrus pedon had the highest value, followed by paddy and kiwi fruit (Table 1). The mineral composition of studied soils shows by the X-ray diffractograms of the clay fraction (Fig.2 and Table 2). Clay mineralogical composition was determined using the intensity and position of the X-ray diffractogram peaks, considering clay apparent CEC (CEC/Clay %). The high intensity of 1.43 and 0.71 nm peaks in Mg-saturated and Mg-saturated, glycerol-solvated treatments, and the collapse of the peaks in K-saturated and K550 treatments demonstrate the presence of vermiculite (Fig.2A). Also, 1.0, 0.5 and 0.33 nm peaks in Mg-saturated diffractograms indicate the presence of illite. Kaolinite was detected based on the presence of 0.71 and 3.56 nm peaks in Mg-saturated treatment that collapsed in K550 treatment. A shifting 1.43 nm peak in the Mg-saturated treatment toward 1.8 nm in Mg saturated-glycerol solvated treatment also indicated the presence of some expandable clay minerals (smectites). A shifting 1.43 nm peak toward 1.1-1.3 nm in K550 treatment showed the presence of hydroxy-interlayered minerals. Vermiculite, smectite-illite and vermiculite, hydroxy-interlayered vermiculite were the most prevailing


clay minerals in citrus, paddy and kiwi fruit pedons, respectively (Table 2). Differences in clay mineralogical suites were most probably related to land management, different parent materials and soil-drainage conditions. Figure 3 shows the relationship between dominant clay minerals with exchangeable and non-exchangeable potassium in different horizons for some soil pedons. The exchangeable and non-exchangeable potassium in the surface layers of the citrus pedons were higher than that of the subsurface layers due to more suitable thermal and moisture status preconditioned for mica and feldspar weathering (Fig.3A). Potassium does not leach out from clayey and silty soils, but the sandy soil used for citrus is not able to hold high amounts of exchangeable and non-exchangeable potassium. Sinha and Biswas (2003) stated that water soluble, available and nonexchangeable potassium showed a positive and significant correlation with clay content and cation exchange capacity for soils of West Bengal, India. Further, Nabiollahy et al. (2006) reported that mineral K,  $\text{HNO}_3$ -extractable K and the clay content of the soils containing lesser illite (10-30%) were significantly different from those with more illite (30-50%). Besides the clay content, K-bearing minerals and clay mineralogy may play a more important role in determining a soil's extractable K status. For example, the clay contents in pedon 1 were relatively low, and their dominant clay mineral was vermiculite, whereas the measured exchangeable and non-exchangeable K was medium and high, respectively (Table 1). Wedge zones or specific K-adsorption sites in vermiculite may have caused these results (Rich, 1972; Sparks and Liebhart, 1982). Plant-available potassium (soluble K) and exchangeable potassium in the studied soils were very low (Table 1). Mean soluble and exchangeable potassium in paddy soil samples were lower than in kiwi fruit and citrus soil samples. Smectite was the most prevalent mineral in the paddy soils; this may be the main reason for their lower K forms compared to other land

**Table 2.** Clay mineralogical suite of the studied soil samples

| Horizon    | Mineralogy                        |
|------------|-----------------------------------|
| (P2) Paddy |                                   |
| Apg        | Ill. » HIV. » Smc. » Kao. » Ver.  |
| Bg         | Smc. » Ver. » HIV. » Illi. » Kao. |
| Btg1       | Ill. » HIV. » Ver. » Kao. » Smc.  |
| Btg2       | Ill. » HIV. » Ver. » Kao. » Smc.  |
| (P3) Paddy |                                   |
| Apg        | Smc. » Illi. » Kao. » HIV. » Ver. |
| 2Bg1       | Smc. » Ver. » Illi. » Kao. » HIV. |
| 2Bg2       | Smc. » Illi. » Kao. » HIV. » Ver. |
| (P4) Paddy |                                   |
| Apg        | Ver. » Illi. » HIV. » Kao. » Smc. |
| Btg1       | HIV. » Kao. » Illi. » Ver. » Smc. |
| Btg2       | Smc. » HIV. » Ver. » Illi. » Kao. |
| 2Bg1       | Smc. » HIV. » Ver. » Illi. » Kao. |
| 3Bg2       | Smc. » Illi. » Ver. » Kao. » HIV. |

| Horizon    | Mineralogy                               |
|------------|--|
| (P5) kiwi  |  |
| Ap         | Smc. » Ver. » Illi. » Kao. » HIV.        |
| Bg1        | HIV. » Smc. » Ver. » Illi. » Kao.        |
| Bg2        | HIV. » Smc. » Ver. » Illi. » Kao.        |
| 2Cg        | HIV. » Smc. » Ver. » Illi. » Kao.        |
| (P6) kiwi  |  |
| Apg        | Smc. » HIV. » Ver. » Illi. » Kao.        |
| Bg         | Ver. » HIV-Cl. » Kao. » Illi. » Smc.     |
| Btg1       | Ver. » HIV-Cl. » Kao. » Illi. » Smc.     |
| Btg2       | Ver. » Smc. » HIV-Cl. » Kao. » Illi.     |
| (P1)Citrus |  |
| Ap         | Ver. » Smc. » Mix. » HIV. » Illi. » Kao. |
| AC         | Ver. » Smc. » HIV. » Kao. » Illi.        |
| C1         | Ver. » Illi. » HIV. » Kao. » Smc.        |
| C2         | Ver. » Illi. » HIV. » Kao. » Smc.        |

\* Illi.: Illite, HI.: Hydroxy-Interlayer Vermiculite, Smc.: Smectite, Kao.: Kaolinite, Ver.: Vermiculite, Cl.: Chlorite, Mix.: Mixed mineral.

**Fig 2.** Representative X-ray diffractograms: A: Btg1 horizon, pedon no. 2 and B: Bg1 horizon, pedon no. 5.


uses. Suitable soil aeration and high oxygen levels are necessary for K uptake by plants. K uptake decreases as soil moisture content increases to saturation. Shahbazi and Towfighi (2006) showed that exchangeable K decreases with increasing soil-saturation time. Results in this study showed that in soil samples in which smectite predominated, non-exchangeable potassium was about 200-300 ppm, while in vermiculitic and hydroxy-interlayered vermiculite and illite-dominated soil samples, non-exchangeable potassium was about 300-600 ppm (Fig.3). Illitic soils, due to the higher fixed potassium in their interlayer positions, and vermiculitic soils, due to their more-negative layer charge, are among the most important factors in controlling soils' potassium status. Sparks (1987) and Moritsuka et al. (2004) reported that the major sources of non-exchangeable K in soils are K-rich 2:1 clay minerals such as micas and vermiculite. In general, paddy soils' mean non-exchangeable potassium, extracted by boiling nitric acid, was between moderately low and low. This may be attributed to the prevalence of smectite, which can cause these soils to be rich in ammonium acetate

extractable potassium, yet moderately poor in  $\text{HNO}_3$ -K concentration. Years of cultivation without the use of potassium fertilizers have led to a drastic decrease in exchangeable potassium, followed by a decrease in non-exchangeable potassium, in paddy soils. So, for better understanding of the relationship between clay mineralogy and different forms of potassium, studying the processes and mechanisms of potassium absorption and release is essential to achieve sustainable agriculture and increase the efficiency of crop production.

## Materials and methods

### Study Area

This study examined paddy and non-paddy soils (about 10 ha) obtained from the Rice Research Institute of Iran ( $36^\circ 51' 53''$  to  $36^\circ 51' 48''$  northern latitudes and  $50^\circ 46' 56''$  to  $50^\circ 46' 44''$  eastern longitudes, 20 m below sea level) (Fig.1). Mean annual precipitation in this region is 1253 mm,



**Fig 3.** Relationship between dominant clay minerals with exchangeable and non-exchangeable potassium in different horizons, A: pedon no.1, B: pedon no.3, C: pedon no.4, D: pedon no.6.

and mean annual air temperature is 15.8 °C. Soil moisture and temperature regimes, calculated according to the Newhall Simulation Model (Newhall and Berdanier, 1996), were udic and thermic, respectively.

#### Field Sampling

Three adjacent land uses, including paddy, kiwifruit and citrus, were selected. Ten pedons were dug, described and sampled according to the field book for describing and sampling soils, and were classified according to the Soil Taxonomy (Soil Survey Staff, 2010). From the pedon samples, six were included for comprehensive analysis in this study (three pedons from paddy soils, two pedons from kiwifruit and one pedon from citrus).

#### Physico-Chemical Analysis

All analyses were performed on air-dried soil samples passed through a 2 mm sieve (Pansu and Gautheyrou, 2006). The samples' physicochemical and mineralogical characteristics were analyzed according to standard methods (Kunze and Dixon, 1986).  $pH_{sp}$  was determined using a pH meter applied to soil saturated to a paste using deionised water. Electrical conductivity ( $EC_{se}$ ) was also measured in the saturated extracts (Carter and Gregorich, 2008). Particle-size distribution was measured by the hydrometer method

(Carter and Gregorich, 2008). Organic carbon was determined using wet oxidation (Pansu and Gautheyrou, 2006). Cation exchange capacity (CEC) was measured by the ammonium acetate method (pH=7) (Carter and Gregorich, 2008). Exchangeable potassium was determined using shaking of 2.5 g air-dried soil with 50 ml of 1 M  $NH_4OAc$  solution (pH=7) (Knudsen et al., 1996). Non-exchangeable potassium was determined by heating 10 g of soil with 25 ml of 1 M  $HNO_3$  solution in an oil bath (113 °C) for 25 minutes (Pratt, 1965). The extracts' K concentrations were determined by flame photometry.

#### Mineralogical Analysis

Clay mineralogical studies were performed by removing soluble salts and gypsum by washing out (Konse and Rich, 1959), carbonates by neutralization with sodium acetate (pH=5) (Grossman and Millet, 1961), organic materials by oxidation with  $H_2O_2$  (Konse and Rich, 1959) and ferrous oxides by citrate dithionate bicarbonate (CDB) (Mehra and Jackson, 1960). Clay fraction was separated by sedimentation and saturated with  $Mg^{2+}$  and  $K^+$  ions using 1 N  $MgCl_2$  and KCl solutions. Mineralogical composition was determined by X-ray diffraction using a Siemens D5000 diffractometer via  $CuK\alpha$  ( $\lambda=1.5409 \text{ \AA}$ ) and 30 kV voltage and 30 mA (Fig. 2, Table 2).

## Conclusion

NH<sub>4</sub>OAc-K is used as an indicator for the availability of potassium in soils, but it does not give similar trends in all mineralogical suites. A combination of both NH<sub>4</sub>OAc-K and HNO<sub>3</sub>-K gives a more applicable index for smectitic, illitic and vermiculitic soils. This study found that while smectitic soils are rich in NH<sub>4</sub>OAc-K, they have modest amounts of HNO<sub>3</sub>-K, with moderate K release rates. On the other hand, illitic soils have only modest amounts of NH<sub>4</sub>OAc-K with high amounts of HNO<sub>3</sub>-K, and high rates of K release. Accordingly, shallow smectitic soils could experience K exhaustion and the need for K fertilization more than illitic soils. Non-exchangeable K plays a significant role in supplying available K, particularly in soils containing K-bearing minerals. Therefore determination of both exchangeable and HNO<sub>3</sub>-extractable K gives a better indication of the K potential and management for rice production in paddy soils. Hence, clay mineralogy should be considered as a main determinant in paddy soils' nutrient status. To improve rice production and potassium reserves in these paddy soils requires precise potassium management.

## Acknowledgement

The financial support provided by the University of Tehran, Iran, is gratefully acknowledged.

## References

- Akbar N, Ehsanullah, Jabran K, Amjad Ali M (2011) Weed management improves yield and quality of direct seeded rice. *Aust J Crop Sci* 5(6):688-694.
- Bhonsle NS, Pal SK, Sekhon GS (1992) Relationship of K forms and release characteristics with clay mineralogy. *Geoderma* 54: 285-293.
- Carter MR, Gregorich EG (2008) *Soil Sampling and Methods of Analysis*. 2<sup>nd</sup> (ed). Can Soc Soil Sci, p 1224.
- Fanning DS, Keramidas VZ, El-Desoky MA (1989) Micas. In: Dixon JB, Weed SB (ed) *Minerals in Soil Environments*. Soil Sci Soc Am J, Madison, Wisconsin, USA. pp551-634.
- Ghosh BN, Singh RD (2001) Potassium release characteristics of some soils of Uttar Pradesh hills varying in altitude and their relationship with forms of soil K and clay mineralogy. *Geoderma* 104: 135-144.
- Grossman RB, Millet JL (1961) Carbonate removal from soils by a modification of the acetate buffer method. *Soil Sci Soc Am J* 25: 325-326.
- Huang PM (2005) Chemistry of potassium in soils. In: Tabatabai MA, Sparks DL (ed) *Chemical Processes in Soils*. Soil Sci Soc Am J, Madison, WI, USA, pp227-292.
- IRRI (International Rice Research Institute) (2002) *World rice statistics*, Los Baños, Philippines.
- Knudsen D, Peterson GA, Pratt PF (1996) Lithium, Sodium and Potassium. In: Page AL (ed) *Methods of Soil Analysis*. Part 2, Second edition, Am Soc Agron, Madison, WI, pp403-429.
- Kunze GW, Rich CI (1959) Mineralogical methods. In: Rich CI, Satz LF, Kunze GW (ed) *Certain properties of selected southeastern United States soils and mineralogical procedures for the study*. Southern Coop, Series Bul, pp135-146.
- Kunze GW, Dixon JB (1986) *Method of soil Analysis, Part 1. Physical and Mineralogical Methods*. Am Soc Agron.
- Kyuma K (1984) Fundamental characteristics of wetland soils. In: *Proceedings of workshop on wetland soils: Characterization, Classification and Utilization*, IRRI, LosBonos, pp191-206.
- Kyuma K (2004) *Paddy soil science*. Kyoto, Japan, Kyoto University Press, p 280.
- Liu G, Nielsen NE, Hansen HCB, Borggaard OK (1997) Mineral changes in a danish alfisol caused by 30 years of potassium depletion in the field. *Soil Plan Sci* 47:1-6.
- Martin HW, Sparks DL (1985) On the behavior of non-exchangeable potassium in soils. *Soil Sci Plan Anal* 16: 133-162.
- McLean EO, Watson ME (1985) Soil measurements of plant available potassium, In: Munson RD (ed) *Potassium in Agriculture*. Soil Sci Soc Am J, Madinson, USA, pp277-308.
- Mehra GW, Jackson ML (1960) Iron oxide removal from soils and clays by a dithionite-citrate system buffered with sodium bicarbonate. In: *Clay Clay Miner, Proceeding, 7<sup>th</sup> Conference*, pp317-327.
- Moritsuka N, Yanai J, Kosaki T (2004) Possible processes releasing nonexchangeable potassium from the rhizosphere of maize. *Plan Soil* 258: 261-268.
- Nabiollahy K, Khormali F, Bazargan K, Ayoubi Sh (2006) Forms of K as a function of clay mineralogy and soil development. *Clay Clay Miner* 41: 739-749.
- Newhall F, Berdanier CR (1996) Calculation of soil moisture regimes from the climatic record. NRCS, Soil Survey Investigation Report, No. 46.
- Pansu M, Gautheyrou J (2006) *Handbook of Soil Analysis: Mineralogical, Organic and Inorganic Methods*. Springer, p 995.
- Pratt PF (1965) *Method of soil Analysis*. Am Soc Agron, Madison, WI.
- Rahimi Petroudi E, Noormohammadi Gh, Mirhadi MJ, Madani H, Mobasser HR (2011) Effects of nitrogen fertilization and rice harvest height on agronomic yield indices of ratoon rice-berseem clover intercropping system. *Aust J Crop Sci* 5(5):566-574.
- Rehm G, Schmitt M (2002) Potassium for crop production. University of Minnesota Extension. <http://www.extension.umn.edu/distribution/cropsystems/DC6794.html>.
- Rich CI (1972) Potassium in minerals. *Proceedings of the 9<sup>th</sup> Colloquium of the International Potash Inst* 9: 15-31.
- Shahbazi K, Towfighi H (2006) Effects of different moisture regimes on soil exchangeable potassium. *Iran J Agric Sci* 38: 161-171.

- Sharpley N (1989) Relationship between Soil Potassium Forms and Mineralogy. *Soil Sci Soc Am J* 52: 1023-1028.
- Sinha AK, Biswas S (2003) Distribution of different forms of potassium in surface and subsurface horizons of some well established soils of West Bengal under the order Inceptisols. *J Interacademia* 7 (3): 286-291.
- Soil Survey Staff. (2010) *Keys to Soil Taxonomy*, United States Department of Agriculture. 11<sup>nd</sup> (ed), NRCS.
- Sparks DL, Liebhardt WC (1982) Temperature effects on potassium exchange and selectivity in Delaware soils. *Soil Sci* 133: 10-17.
- Sparks DL, Huang PM (1985) Physical chemistry of soil potassium. In: Munson RD (ed) *Potassium in Agriculture*. *Soil Sci Soc Am J*, Madison, WI, USA, pp201-276.
- Sparks DL (1987) Potassium dynamics in soils. *Adv Soil Sci* 6: 1-63.
- Srinivasarao Ch, Rupab TR, Subba Raoc A, Ramesha G, Bansald SK (2006) Release kinetics of non-exchangeable potassium by different extractants from soils of varying mineralogy and depth, *Soil Sci Plan Anal* 37(3): 473-491.
- Surapaneni A, Palmer AS, Tillman RW, Kirkman JH, Gregg PEH (2002) The mineralogy and potassium supplying power of some loessial and related soils of New Zealand. *Geoderma* 110: 191-204.
- Towfighi H (1998) Study of rice response to potassium fertilizer in paddy soils of northern Iran. *Iran J Agric Sci* 29: 869-883.