

## Climate change and Arctic ecosystems:

### 1. Vegetation changes north of 55°N between the last glacial maximum, mid-Holocene, and present

Nancy H. Bigelow,<sup>1</sup> Linda B. Brubaker,<sup>2</sup> Mary E. Edwards,<sup>3,4,5</sup> Sandy P. Harrison,<sup>6,7</sup> I. Colin Prentice,<sup>6</sup> Patricia M. Anderson,<sup>8</sup> Andrei A. Andreev,<sup>9</sup> Patrick J. Bartlein,<sup>10</sup> Torben R. Christensen,<sup>11,12</sup> Wolfgang Cramer,<sup>13</sup> Jed O. Kaplan,<sup>6,14,15</sup> Anatoly V. Lozhkin,<sup>16</sup> Nadja V. Matveyeva,<sup>17</sup> David F. Murray,<sup>18</sup> A. David McGuire,<sup>19,20</sup> Volodya Y. Razzhivin,<sup>17</sup> James C. Ritchie,<sup>21</sup> Benjamin Smith,<sup>6,11,12</sup> Donald A. Walker,<sup>3,22</sup> Konrad Gajewski,<sup>23</sup> Victoria Wolf,<sup>24</sup> Björn H. Holmqvist,<sup>25</sup> Yaeko Igarashi,<sup>26</sup> Konstantin Kremenetskii,<sup>27</sup> Aage Paus,<sup>28</sup> Michael F. J. Pisaric,<sup>29</sup> and Valentina S. Volkova<sup>30</sup>

Received 23 May 2002; revised 1 October 2002; accepted 18 December 2002; published 8 October 2003.

[1] A unified scheme to assign pollen samples to vegetation types was used to reconstruct vegetation patterns north of 55°N at the last glacial maximum (LGM) and mid-Holocene (6000 years B.P.). The pollen data set assembled for this purpose represents a comprehensive compilation based on the work of many projects and research groups. Five tundra types (cushion forb tundra, graminoid and forb tundra, prostrate dwarf-shrub tundra, erect dwarf-shrub tundra, and low- and high-shrub tundra) were distinguished and mapped on the basis of modern pollen surface samples. The tundra-forest boundary and the distributions of boreal and temperate forest types today were realistically reconstructed. During the mid-Holocene the tundra-forest boundary was north of its present position in some regions, but the pattern of this shift was strongly asymmetrical around the pole, with the largest northward shift in central Siberia (~200 km), little change in Beringia, and a southward shift in Keewatin and Labrador (~200 km). Low- and high-shrub tundra extended farther north than today. At the LGM, forests were absent from high latitudes. Graminoid and forb tundra abutted on temperate steppe in northwestern Eurasia while prostrate dwarf-shrub, erect dwarf-shrub, and graminoid and forb tundra formed a mosaic in Beringia. Graminoid and forb tundra is restricted today and

<sup>1</sup>Alaska Quaternary Center, College of Science, Engineering and Mathematics, University of Alaska, Fairbanks, Alaska, USA.

<sup>2</sup>College of Forest Resources, University of Washington, Seattle, Washington, USA.

<sup>3</sup>Institute of Arctic Biology, University of Alaska, Fairbanks, Alaska, USA.

<sup>4</sup>Department of Geography, Norges Teknisk-Naturvitenskapelige Universitet, Trondheim, Norway.

<sup>5</sup>Now at Department of Geography, University of Southampton, Southampton, UK.

<sup>6</sup>Max Planck Institute for Biogeochemistry, Jena, Germany.

<sup>7</sup>Dynamic Palaeoclimatology, Lund University, Lund, Sweden.

<sup>8</sup>Quaternary Research Center, University of Washington, Seattle, Washington, USA.

<sup>9</sup>Alfred-Wegener-Institut für Polar- und Meerforschung, Potsdam, Germany.

<sup>10</sup>Department of Geography, University of Oregon, Eugene, Oregon, USA.

<sup>11</sup>Climate Impacts Group, Department of Ecology, Lund University, Lund, Sweden.

<sup>12</sup>Now at Department of Physical Geography and Ecosystems Analysis, Lund University, Lund, Sweden.

<sup>13</sup>Potsdam Institut für Klimafolgenforschung, Potsdam, Germany.

<sup>14</sup>Plant Ecology, Department of Ecology, Lund University, Lund, Sweden.

<sup>15</sup>Now at Canadian Centre for Climate Modeling and Analysis, Victoria, British Columbia, Canada.

<sup>16</sup>Northeast Interdisciplinary Scientific Research Institute, Far East Branch, Russian Academy of Sciences, Magadan, Russia.

<sup>17</sup>Department of Vegetation of the Far North, Komarov Botanical Institute, St. Petersburg, Russia.

<sup>18</sup>University of Alaska Museum, Fairbanks, Alaska, USA.

<sup>19</sup>Department of Biology and Wildlife, University of Alaska, Fairbanks, Alaska, USA.

<sup>20</sup>Now at U.S. Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska, Fairbanks, Alaska, USA.

<sup>21</sup>Pebbledash Cottage, Corfe, Taunton, UK.

<sup>22</sup>Institute of Arctic and Alpine Research, University of Colorado, Boulder, Colorado, USA.

<sup>23</sup>Department of Geography, University of Ottawa, Ottawa, Ontario, Canada.

<sup>24</sup>Alaska SAR Facility, Geophysical Institute, University of Alaska, Fairbanks, Alaska, USA.

<sup>25</sup>Department of Geology, Quaternary Geology, Lund University, Lund, Sweden.

<sup>26</sup>Earthscience Co. Ltd., Sapporo, Japan.

<sup>27</sup>Institute of Geography, Russian Academy of Sciences, Moscow, Russia.

<sup>28</sup>Botanisk Institutt, University of Bergen, Bergen, Norway.

<sup>29</sup>Big Sky Institute, Montana State University, Bozeman, Montana, USA.

<sup>30</sup>Joint Institute for Geology, Geophysics and Mineralogy, Siberian Branch, Russian Academy of Sciences, Novosibirsk, Russia.

does not form a large continuous biome, but the pollen data show that it was far more extensive at the LGM, while low- and high-shrub tundra were greatly reduced, illustrating the potential for climate change to dramatically alter the relative areas occupied by different vegetation types. *INDEX TERMS:* 1615 Global Change: Biogeochemical processes (4805); 1620 Global Change: Climate dynamics (3309); 1851 Hydrology: Plant ecology; 3344 Meteorology and Atmospheric Dynamics: Paleoclimatology; *KEYWORDS:* vegetation maps, mid-Holocene, last glacial maximum, Arctic ecosystems, palaeoclimate, biomization

**Citation:** Bigelow, N. H., et al., Climate change and Arctic ecosystems: 1. Vegetation changes north of 55°N between the last glacial maximum, mid-Holocene, and present, *J. Geophys. Res.*, 108(D19), 8170, doi:10.1029/2002JD002558, 2003.

## 1. Introduction


[2] Arctic climate and ecosystems are expected to be highly sensitive to anthropogenic changes in atmospheric composition and hence radiative forcing. Early simulations of the global climatic response to CO<sub>2</sub> doubling indicated increases in temperature by up to 12 K in the northern polar regions [Manabe and Wetherald, 1975]. More recent simulations that take into account the cooling effect of sulphate aerosols show less extreme changes [Cubasch et al., 2001], but it remains a robust generalization that the simulated high-latitude temperature response to increased greenhouse gas concentrations is stronger than the responses of the tropics and midlatitudes. This generalization has also been derived independently of models, by examining inferred patterns of climate change for various geological epochs [Hoffert and Covey, 1992]. The particular sensitivity of high-latitude climates can be explained by the operation of two powerful (and synergistic) positive feedbacks: changes in the extent and duration of sea-ice cover in the Arctic Ocean [Ganopolski et al., 1998; Braconnot et al., 1999; Vavrus, 1999], and changes in the albedo of the land surface as a consequence of changes in snow cover [Bonan et al., 1992; Foley et al., 1994; Berger, 2001]. Model experiments indicate that changes in land-surface albedo resulting from climatically induced shifts in the tundra-forest boundary play a role in the initiation of glaciations [Gallée et al., 1992; Gallimore and Kutzbach, 1996; de Noblet et al., 1996], amplification of high-latitude cooling during glaciations [Levis et al., 2000], and enhancement of the direct effects of higher than present northern summer insolation during interglacial periods [Foley et al., 1994; TEMPO Members, 1996; Texier et al., 1997].

[3] The palaeorecord provides an opportunity to test the simulations of past climates and hence to evaluate the climatic and ecological sensitivity of the Arctic as indicated by models. The Palaeovegetation Mapping Project (BIOME 6000; Prentice and Webb [1998]) produced the first maps of vegetation distribution across the northern high latitudes at 6000 <sup>14</sup>C years B.P. (6 ka) and 18,000 <sup>14</sup>C years B.P. (18 ka) using a formal procedure (biomization; Prentice et al. [1996]) based on plant functional types (PFTs). The biomization procedure was applied in BIOME 6000 region by region, where the regions were defined pragmatically so as to build on pre-existing collaborative projects. The Arctic was thus fragmented among four regions, with slight overlaps between them: Beringia (Alaska and part of NW Canada plus the Russian Far East; Edwards et al. [2000]), Canada and Eastern North America [Williams et al., 2000], Europe [Prentice et al., 1996; Tarasov et al., 2000], and the Former Soviet Union, excluding Beringia [Tarasov et al.,

1998, 2000]. Slightly different allocations of pollen taxa to PFTs, and of PFTs to biomes, were adopted by the different regional working groups. There are no apparent discontinuities at regional boundaries in the BIOME 6000 maps reconstructed from modern surface samples [see Prentice et al., 2000]. This finding suggests that the differences in the regional biome schemes adopted in BIOME 6000 are not crucial, at least for the broad-scale patterns that BIOME 6000 was intended to reconstruct. However, the degree to which the reconstructed positions of transitions between steppe and tundra biomes at 18 ka have been influenced by differences in the regional biomization schemes has been questioned [Edwards et al., 2000; Tarasov et al., 2000; Elenga et al., 2000]. Thus a re-examination of the Arctic data using a consistent biomization scheme is timely.

[4] Several additional factors motivate a reconsideration of the distribution of Arctic biomes at key times in the past. The regional biomizations made in BIOME 6000 concentrated on differentiating forest types, and forest from tundra. There was no attempt to distinguish different types of tundra. Yet tundra is highly differentiated, structurally and floristically, and the differences are closely linked to climatic gradients. Knowledge of the distribution of these different tundra vegetation types in the past could improve our understanding of regional climate changes in the high latitudes. The distinctions among tundra types are also potentially important for feedbacks involving changing water and carbon exchanges, as these vegetation types differ greatly in their biogeochemical and biophysical characteristics [Chapin et al., 2000a, 2000b; Epstein et al., 2001] including the height of perennial biomass and the annual net primary production (NPP). The height of perennial biomass ranges from <5 cm in cushion-forb and prostrate dwarf-shrub tundra, to nearly 2 m in high-shrub tundra near treeline. The taller shrub-dominated vegetation types present a many times rougher surface and can greatly reduce surface albedo in the presence of snow. NPP ranges from <50 g C m<sup>-2</sup> yr<sup>-1</sup> in cushion-forb tundra to >300 g C m<sup>-2</sup> yr<sup>-1</sup> in high-shrub tundra [e.g., Christensen et al., 2000] implying a large range in carbon storage potential which is reinforced by slow decomposition rates in some shrub tundras. The lumping of all treeless Arctic vegetation into a single “tundra” biome for vegetation mapping and modeling purposes thus discards information about the land surface that is both ecologically and geophysically important.

[5] The international Pan-Arctic Initiative (PAIN) was set up with the goal of improving the ability of the ecological and geophysical communities to model Arctic vegetation types and their responses to environmental changes. The first part of the work carried out by PAIN was the produc-



**Figure 1.** Types of sites providing (a) modern pollen data, (b) 6 ka pollen data, and (c) 18 ka pollen data. In cases where two or more 6 ka or 18 ka sites lie very close to one another, they have been slightly shifted in these plots to permit each site to be individually resolved.

tion of pollen-based biome reconstructions compatible with a new global biogeography-biogeochemistry model, BIOME4, which was also developed in part within the PAIN project [Kaplan *et al.*, 2003]. This paper documents the data sets and the biomization procedure developed by PAIN. The work enabled us to make a spatially detailed and methodologically consistent reconstruction of vegetation distribution in the high northern latitudes. Our primary aims were (a) to delimit the boundary between forest and tundra now and in the past, and (b) to reconstruct changes in the distribution of tundra types. We address some specific outstanding research questions: (a) the location of the Arctic treeline in the mid-Holocene, and in particular whether its late Holocene shifts have been zonally uniform; and (b) the nature of the vegetation of northern Eurasia and Beringia during the last glacial maximum (LGM), and in particular the interface between steppe and tundra vegetation types. The steppe-tundra interface at the LGM has been controversial because of the apparent lack of convincing modern analogues for this vegetation [Vartanyan *et al.*, 1993; Goetcheus and Birks, 2001; Guthrie, 2001; Walker *et al.*, 2001; Yurtsev, 2001]. It is important to understand the nature of the LGM vegetation because it provided the habitat for megaherbivore populations and hence support

for human populations, including those that colonized the Americas (by way of the Bering land bridge) toward the end of the last glacial period.

## 2. Methods

### 2.1. Study Region

[6] The southernmost occurrence of tundra at low elevations in the Northern Hemisphere today is at about 55°N, along the southern coast of Hudson Bay in Canada. By confining our study area to the circumpolar region north of 55°N, we could encompass the full latitudinal range of the modern lowland tundra. This latitudinal limit also enabled us to process a great deal of information concerning the past and present distribution of boreal and cool-temperate forest types that occur north of 55°N (and in Fennoscandia, even north of 70°N).

### 2.2. Choice of Time Slices

[7] We assembled a comprehensive pollen data set for 0 ka (i.e., core tops and surface sediment samples), 6 ka (<sup>14</sup>C timescale) and 18 ka (<sup>14</sup>C timescale) (Figure 1). This choice of time slices allowed comparison with previous attempts to construct pollen-based biome maps covering

different regions of the Arctic as generated by the BIOME 6000 project. These specific times were selected in BIOME 6000 in part to allow comparison with climate model simulations carried out within the framework of the Palaeoclimate Modeling Intercomparison Project (PMIP; *Joussaume and Taylor [1995]*) for 6 ka and 21 ka BP (astronomical timescale). Within the time resolution that can actually be achieved by correlating independently dated sediment cores, 6 ka  $^{14}\text{C}$ -years is effectively the same as 6 ka astronomical years BP, and 18 ka  $^{14}\text{C}$ -years is effectively equivalent to 21 ka astronomical years BP. The 6 ka (mid-Holocene) time period represents an interval when the total annual insolation received in the northern high latitudes was larger than today and the annual cycle of insolation was enhanced (Northern Hemisphere summer insolation was greater and winter insolation less than today). Other boundary conditions, most importantly ice sheet distributions, sea level and atmospheric greenhouse gas concentrations ( $\text{CO}_2$ ,  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ ), were not materially different from those of the pre-industrial Holocene. The 21 ka (last glacial maximum, LGM) time period represents a contrasting interval when insolation was quite similar to today, but the continental ice sheets were at a maximum and greenhouse gas concentrations were at a minimum.

### 2.3. Pollen Data

[8] We generally used the single pollen sample that was closest in estimated  $^{14}\text{C}$  age to the designated time, provided it fell within a certain time range. However, because of the larger dating uncertainties and/or higher inter-sample variability characteristic of some 18 ka records, we used multiple samples from these sites. Each of these samples was independently allocated to a biome, and the reconstructed biome at the site was assumed to correspond to the biome given by the majority of the samples. Following the BIOME 6000 convention, we allowed samples that fell within 0 to 500 years for the modern sample data set, within  $\pm 500$  years of 6 ka, and within  $\pm 1000$  years of 18 ka. A few samples that fell just outside these designated time ranges were allowed in exceptional cases. At 0 ka, eight samples that were dated in the range 500 to 850 years B.P. were included, for comparison with fossil pollen samples from the same cores. Four mid-Holocene samples that fell beyond the designated time range but within  $\pm 700$  years of 6 ka BP were included. We included one sample in the 18 ka data set that was dated to 20.7 ka because it provided critical information from the Bering Land Bridge [*Elias et al., 1996*] and because data from other parts of Beringia suggest relatively stable vegetation and climate during the interval 21 to 18 ka BP.

[9] Pollen data were available to us either as full pollen counts, or as pollen percentages derived from summary tables or scanned from published pollen diagrams. Typically, published pollen percentage summary tables and diagrams contain only major tree, shrub and herb taxa. Digitized pollen percentage data were used extensively in earlier studies in order to ensure a reasonably dense spatial coverage. The modern pollen-based maps of Europe made by *Prentice et al. [1996]*, for example, are almost entirely based on digitized pollen percentage data, and about 50% of the sites used to construct the corresponding maps for Russia were digitized [*Tarasov et al., 1998*]. These studies,

and similar studies from other regions, suggest that pollen percentage data of major taxa are generally adequate to discriminate between forest types but are less good at discriminating non-forest biomes. Since herbaceous taxa are an important diagnostic of different tundra vegetation types, we used original pollen counts to the greatest possible extent. Digitized pollen percentage data were used only from those regions (e.g., some parts of northern Fennoscandia and European Russia) from which no other data were available to us.

[10] The modern pollen data set consists of 2098 samples (Table 1), 382 of which are associated with a fossil sample at 6 ka and/or 18 ka (e.g., they are lacustrine or peat core tops, or terrestrial sections). The original pollen counts were available for most of the samples (1809). However, 289 samples (13%) were only available as pollen percentages. The samples are from a variety of site types, including lacustrine core tops (185), lacustrine surface samples (755), peat core tops (151), peat samples (19), moss polsters (596), terrestrial section tops (16), terrestrial soils and/or surface sediments (269), and air pollen collectors (7). Site information is lacking for 100 samples, 22 of which are associated with a fossil sample. The mid-Holocene data set includes 493 samples (Table 2). Original pollen counts were available for most of the samples (404). However, 89 samples (18%) were only available as pollen percentages. The samples are from lacustrine cores (230), peat cores (158) and terrestrial sections (33). Site information is lacking for 72 samples. The LGM data set includes 67 samples from 39 sites (Table 2). Original pollen counts were available for most of the samples (58). Nine samples (13%) were available only as pollen percentages. The samples are from lacustrine cores (28), peat cores (2) and terrestrial sections (37).

[11] Some sites that were used in the BIOME 6000 publications have been excluded from the PAIN data set. For example, some modern sites from Russia were excluded because they appeared to duplicate samples represented in the data for Europe (30 sites). A further 466 modern sites were excluded for one or more of the following reasons: (a) they had poor dating control, (b) they had pollen counts of  $< 100$ , (c) they were based on digitized records in cases where full pollen counts have subsequently become available, or (d) key background information (e.g., about the source of the samples) was lacking. A number of fossil sites that were included in the BIOME 6000 compilation for 6 ka (117 samples) and for 18 ka (6 samples) were excluded from the current data set for the same reasons. We applied the same quality-control criteria in selecting new sites to include in the compilation.

### 2.4. Assignment of Pollen Samples to Biomes

[12] The “biomization” method [*Prentice et al., 1996*] is based on the recognition that plant taxa can be grouped into functional types (PFTs) that have both identifiable sets of traits and distinctive climatic requirements. The functional characteristics are expressed in life form (e.g., tree, erect dwarf-shrub, forb), leaf morphology (e.g., needle-leaved, broad-leaved), phenology (e.g., evergreen, cold-deciduous) and mechanism of extreme cold tolerance (e.g., tropical, i.e., frost intolerant, temperate, boreal). These functional characteristics are an expression of the mechanisms whereby

**Table 1 (Representative Sample).** Characteristics of the Modern Pollen Samples<sup>a</sup> [The full Table 1 is available in the HTML version of the article.]

Sample Code	Site Name	Sector	Latitude, deg	Longitude, deg	Elevation, m	Local Vegetation	Vegetation Formation	Sample Type	Data Type	Record Length, years	Number of <sup>14</sup> C Dates	References
NI148_0k	Tämet	Western Europe	69.67	30.10	n/a	undisturbed		peat core top	count	ca. 0 – > 5000	2 + top	Vorren [1983]
S030	Juvjalampi	Western Europe	62.63	33.67	150	taiga dominated by <i>Pinus/betris</i>	taiga	moss polster	unknown %	0	0	Tarasov <i>et al.</i> [1998]
S302	10 Mile Lake	Eastern Beringia (Alaska)	63.05	-145.42	n/a		alpine tundra	lacustrine surface sample	count	0	0	M. E. Edwards (unpublished data, 2003)
S15-1	15/1	Labrador	55.09	-75.25	244		boreal forest	lacustrine surface sample	count	0	0	Gajewski [1991]
S15-2	15/2	Labrador	55.83	-75.02	305		forest tundra/boreal forest transition	lacustrine surface sample	count	0	0	Gajewski [1991]
S16-1	16/1	Labrador	55.41	-75.07	285		boreal forest	lacustrine surface sample	count	0	0	Gajewski [1991]
S17-1	17/1	Labrador	55.07	-75.70	235		boreal forest	lacustrine surface sample	count	0	0	Gajewski [1991]
S17-2	17/2	Labrador	55.25	-74.93	335		boreal forest	lacustrine surface sample	count	0	0	Gajewski [1991]
S17-3	17/3	Labrador	55.12	-75.95	255		boreal forest	lacustrine surface sample	count	0	0	Gajewski [1991]
S143	6 Asseyes Lake	Eastern Beringia (Alaska)	62.65	-141.07	549	<i>Picea glauca</i> and sedge meadows	black spruce muskeg-sedge meadows	lacustrine surface sample	count	0	0	P. M. Anderson and L. B. Brubaker (unpublished data, 2003)
TA020	70 Mile Lake, Richardson Hwy	Eastern Beringia (Alaska)	61.52	-145.23	564		boreal forest	lacustrine surface sample	count	0	0	T. A. Ager (unpublished data, 2003)
S303	Ace Lake	Eastern Beringia (Alaska)	64.52	-147.56	n/a		boreal forest	lacustrine surface sample	count	0	0	M. E. Edwards and A. P. Krumhardt (unpublished data, 2003)
iep38	Adamsvatnet	Western Europe	70.53	29.45	322		tundra	lacustrine surface sample	digitized %	0	0	Prentice [1978]
AGE	Ageröds Mosse	Western Europe	55.83	13.42	58	pasture and arable land	memoral forest	peat core top	count	0 – 9660	21 + top	Nilsson [1964]
S039	Ahaliorak Lake	Eastern Beringia (Alaska)	68.92	-151.32	329	<i>Betula</i> shrub tussock tundra, <i>Salix</i> present (no <i>Alnus</i> )	tundra	lacustrine surface sample	count	0	0	P. M. Anderson and L. B. Brubaker (unpublished data, 2003)
S273	Ahaliorak Lake	Eastern Beringia (Alaska)	68.91	-151.32	n/a		shrub tussock tundra	lacustrine surface sample	count	0	0	P. M. Anderson and L. B. Brubaker (unpublished data, 2003)
iep32	Ahvenjärvi	Western Europe	68.93	26.97	155.5		pine forest	lacustrine surface sample	digitized %	0	0	Prentice [1978]

<sup>a</sup>Latitude and longitude are given in decimal degrees, where N and E are conventionally positive and S and W are negative.

**Table 2 (Representative Sample).** Characteristics of the 6 ka and 18 ka Pollen Samples<sup>a</sup> [The full Table 2 is available in the HTML version of this article.]

Site Name	Latitude, deg	Longitude, deg	Elevation, m	Sample Type	Data Type	Record Length, years	Number of <sup>14</sup> C Dates	6 ka Dating Control	18 ka Dating Control	Database Source	References
Adycha Section	67.57	134.42	130	peat core	count	1000–8820	5 + top	1C	n/a	PALE	A. V. Lozhkin (unpublished data, 2003)
Ageröds Mosse	55.83	13.42	58	peat core	count	0–9660	21 + top	1C	n/a	EPD	Nilsson [1964]
Aholammi	61.88	25.22	114	lacustrine core	count	1480–11,552	3	2C	n/a	EPD	Koivula [1987]
Aiatskoe	57.00	60.08	229	peat core	digitized %	0–> = 9000	7 + top	1C	n/a		Peterson [1993]
Ailsa so	75.32	–19.67	88	lacustrine core	count	6000–11,500	5 (1 rejected)	1D	n/a	EPD	Björck and Persson [1981]
Akerhultagöl [Torntabaken]	57.48	14.47	303	peat core	count	6387–12,967	14	6D	n/a		Bjelm [1976], Björck and Hakansson [1982]
Akuvaara	69.13	27.68	170	lacustrine core	count	670–9434	5	1C	n/a	EPD	Hyvärinen [1975]
Alazeya	68.50	154.00	40	peat core	count	4000–10,000	5	2C	n/a		Kaplina and Lozhkin [1982]
Altersvatn	60.80	4.93	4	lacustrine core	count	ca. 2000–9000	1 (1 rejected) + pollen	7	n/a	Nordmap	Kaland [1984]
Alut Lake	60.30	152.31	480	lacustrine core	count	0–53,576	24 (11 rejected) + top + 1 tephra	1C	6C	PALE	P. M. Anderson and A. V. Lozhkin (unpublished data, 2003)
Andy Lake	64.65	–128.08	1360	lacustrine core	count	0–11,310	5 (1 rejected) + top	1C	n/a	NAPD	Szeicz et al. [1995]
Angal Lake	67.13	–153.88	853	lacustrine core	count	0–14,130	3 + top	3D	n/a	NAPD	Brubaker et al. [1983]
Antifreeze Pond	62.35	–140.83	706	lacustrine core	digitized %	0–>13,000	8 (1 rejected) + top + 1 tephra	2C	7C		Rampton [1971]
Ántu sinijarv	59.13	26.33	95	lacustrine core	count	1987–12,742	9	2C	n/a	EPD	L. Saarse (unpublished data, 2003), Saarse and Liva [1995]
Arkad'ovo	56.50	84.00	70	unknown	unknown %		1	2D	n/a		N. A. Berezhina and P. S. Liss (personal communication to P. E. Tarasov, 1997)
Åsen	59.67	9.17	668	unknown	count	0–6000	3 + top	1D	n/a	Nordmap	Hafsten et al. [1979]
Ayakli	69.25	89.00	125	terrestrial section	count	ca. 11,000–20,000	2	n/a	6C		Kind [1974]
B. Kuropatochya	71.07	156.50	77	unknown	digitized %	6000–>/= 9000	4	1	n/a		Peterson [1993]
Babozero	66.37	37.52	138	lacustrine core	count	0–11,800	3 + top	3C	n/a		Kremenetski and Palyk-Kara [1997]
Baidara	68.85	66.90	30	unknown	count	3995–11,066	10 (3 rejected)	1C	n/a	EPD	Andreev et al. [1998]
Baie du Diana	60.78	–69.83	50	lacustrine core	count	250–6330	1 + top + deglaciation	2D	n/a	NAPD	Richard [1977]
Baird Inlet	78.49	–76.78	295	lacustrine core	count	0–9000	3 + top	1C	n/a		Hyvärinen [1985a]
“Rock Basin Lake”											
Bakkemyra	69.20	17.55	140	peat core	count	0–8800	5 + top	1C	n/a	Nordmap	Vorren [1979]
Bankfjörn	60.12	5.47	21	lacustrine core	count	ca 2500–9000	0 + pollen	7	n/a	Nordmap	Somsteigaard and Mangerud [1977]
Barsebäcksmossen	55.77	12.88	1	lacustrine core	count	0–10,000	15 + top	1C	n/a	Nordmap	Digerfeldt [1975]
Begljanskii Riam	55.50	81.57	77	peat core	digitized %	0–6000	0 + 8 poll or strat	7	n/a		Peterson [1993]
Belkachi	59.15	131.98	458	unknown	digitized %	6000	0	7	n/a		Peterson [1993]
Bell's Lake	65.02	–127.48	580	lacustrine core	count	0–11,450	5 + top	2C	n/a	NAPD	Szeicz et al. [1995]
Bereyekh Section	70.58	145.00	20	terrestrial section	count	9468–31,542	4	n/a	7D	PALE	A. V. Lozhkin (unpublished data, 2003)
Bering Land Bridge [1995]	65.23	–167.42	0	peat core	digitized %	ca. 13,000–20,700	4 (+1 rejected)	n/a	7D		Elias et al. [1996]

<sup>a</sup>Latitude and longitude are given in decimal degrees, where N and E are conventionally positive and S and W are negative. The dating control (DC) follows the COHMAP scheme, as described by Yu and Harrison [1995].

<sup>b</sup>For mapping purposes, these sites, which are very close to one another, have been displaced slightly.

<sup>c</sup>These sites have multiple samples within the 18 ka time window; all samples have been used in the biomization procedure.

plants maximize productivity while surviving environmental stress. Plant functional types are expected to occupy a contiguous area in environmental space, even though their distribution in geographic space is typically discontinuous.

[13] In some cases, the physiological adaptations to climate stress are well established from experimental work (e.g., adaptations to extreme cold, budburst, adaptations to drought). In other situations, the mechanism is unclear although the geographic distribution of PFTs makes it abundantly clear that there are climatic limits operating. In practice, the definition of PFT distributions in climate space has relied on a mixture of known mechanistic limits and correlations of biogeographical distribution limits and climate variables. The first step in biomization is the derivation of a conceptual framework in which PFTs are arranged in climate space and biomes are identified as combinations of PFTs with particular climatic locations. ("Climate" here is interpreted in a broad sense, incorporating the idea that plants are sensitive to aspects of the environment that are modified by soil and topography.) In this study, this step has been made explicit and used a basis for model development [Kaplan *et al.*, 2003] as well as for the definitions of biomes based on taxa assigned to PFTs.

[14] The definitions of biomes adopted here are outlined in the companion paper [Kaplan *et al.*, 2003]. We aimed to distinguish as biomes those vegetation types that are recognizable throughout the region and distinctive in terms of land-surface parameters, carbon storage and climatic controls. We required that the biomes be recognisable on the modern landscape (although not necessarily occupying large continuous areas) and further that they should be able to be discriminated on the basis of modern pollen data, thus allowing their distribution to be reconstructed from the fossil record. Using these criteria, we arrived at a five-fold classification of tundra (cushion forb tundra; graminoid and forb tundra; prostrate dwarf-shrub tundra; erect dwarf-shrub tundra; low- and high-shrub tundra). For the non-tundra types we adopted the same classification as in the BIOME4 model [Kaplan, 2001; Kaplan *et al.*, 2003], except that we did not attempt to separate temperate deciduous broadleaf savanna from temperate deciduous broad-leaved forest (we lumped both as "forest"), nor did we attempt to separate temperate xerophytic shrubland from temperate grassland (we lumped both as "steppe"). In all, twelve biomes were recognized as occurring north of 55°N today.

[15] As the biomization method requires the maximum use of floristic information in pollen records, it is usually necessary to define a larger number of PFTs for this purpose than can practically be used in a modeling context. Here we have defined 29 PFTs (Table 3). We adopted the classification for arboreal PFTs that has been used in previous biomizations [e.g., Prentice *et al.*, 1996; Tarasov *et al.*, 1998; Edwards *et al.*, 2000], retaining just those that occur in the northern high latitudes. Thus we recognised 11 arboreal PFTs. We modified their names in order to make the classification according to bioclimate, phenology, leaf morphology and life form more explicit.

[16] Non-arboreal PFTs have been treated in a highly simplified way in previous biomizations. We devised a new functional classification, paying special attention to the diversity of shrub life forms that characterizes the vegetation of the Arctic. We divided shrubs into three primary

categories according to stature: prostrate dwarf-shrubs are less than 5 cm tall; erect dwarf-shrubs are between 5 and 25 cm tall; and low- and high-shrubs are greater than 25 cm, but less than 200 cm tall. The shrubs are then further divided according to differences in bioclimatic range, leaf morphology and phenology. In this way, we define 11 shrub PFTs. Among forbs, we distinguished three PFTs: rosette or cushion forb, arctic forb, and drought-tolerant boreal or temperate forb. The characteristic shape of rosette and cushion forbs enables them to trap air close to the leaf surface and thus to survive extremely low air temperatures through thermal enhancement [Sonesson and Callaghan, 1991]. Typical examples are *Draba* spp. and *Silene acaulis*. Arctic forbs do not have a unique growth form but characteristically survive extreme winter air temperatures by being snow covered and have a rapid growth cycle once the snow melts [Bliss, 1962]. Typical examples include *Polemonium* spp., *Pedicularis* spp., *Gentiana* spp. and *Oxyria digyna*. The drought-tolerant boreal or temperate forbs show no characteristic adaptations to temperature stress but rather have adaptations that enable them to survive drought conditions, such as succulence and early senescence. Characteristic taxa include *Sarcobatus* spp. and *Atriplex* spp. Two additional forb PFTs (boreal forbs and temperate forbs) were recognised but not used in previous biomizations because they appear to lack diagnostic value [e.g., Edwards *et al.*, 2000]. Most of the taxa that could be allocated to boreal or temperate forbs can tolerate a wide range of bioclimatic conditions. The relatively few forbs that occur uniquely in boreal or temperate environments are encountered only rarely in the pollen data. We therefore followed previous work in excluding boreal or temperate forbs from our analysis. The PFTs grass (Poaceae) and sedge (Cyperaceae) were retained, and bog moss (*Sphagnum* spp.) and rushes (Juncaceae) were further identified as distinct PFTs because of their potential discriminatory power in the Arctic. Following previous work, we excluded ferns and aquatics from our analysis.

[17] On the basis of the definitions of the biomes as presented in Table 4, Table 5 indicates the characterization of each biome in terms of constituent PFTs. Table 6 gives the assignments of pollen taxa to PFTs. The information in Tables 5 and 6 allows each pollen sample to be assigned uniquely to the most likely biome according to its numerical affinity score, following the biomization method of Prentice *et al.* [1996]. Pollen taxa that are present but constitute less than 0.5% of the pollen sum do not contribute to the calculation of the affinity score. We applied a weighting ( $\times 15$ ) to the pollen percentages of *Larix* prior to calculating the affinity score (including the square root transformation of pollen percentages used in this calculation) in order to compensate for the known low pollen production characteristic of this taxon. This weighting was chosen because it resulted in the most realistic modern distribution of cold deciduous forest in Siberia. In the case when equal affinity scores were obtained for more than one biome, biomes were assigned in the order shown in Table 5. We varied the standard biomization method in one respect: namely, to perform the biomization in two steps. At the first step, a composite biome "tundra" was distinguished. The composite biome was defined as the union of all of the tundra

**Table 3.** Definition of Plant Functional Types Used in the Biomization Procedure<sup>a</sup>

Biome 6000 Code	New Code	PFT Name	Characteristic Taxa
sp	m	bog moss	<i>Sphagnum</i>
g	g	grass graminoid	Poaceae
s	s	sedge graminoid	<i>Carex</i>
r	r	rush graminoid	<i>Juncus</i>
crc	rcfb	rosette or cushion forb	<i>Draba</i> , <i>Silene aucaulis</i>
af	ar.fb	arctic forb	<i>Polemonium</i> , <i>Pedicularis</i>
sf	bo/te-dt.fb	boreal or temperate drought-tolerant forb	<i>Sarcobatus</i> , <i>Atriplex</i>
dpms	ar.cd.mb.pds	arctic cold-deciduous malacophyll broad-leaved prostrate dwarf shrub	<i>Salix arctica</i>
eps	ar.e.mb.pds	arctic evergreen malacophyll broad-leaved prostrate dwarf shrub	<i>Diapensia</i> , <i>Dryas</i>
juni	ab.e.n.pds	arcto-boreal evergreen needle-leaved prostrate dwarf shrub	<i>Juniperus</i>
dds	ab.cd.mb.eds	arcto-boreal cold-deciduous malacophyll broad-leaved erect dwarf shrub	<i>Betula nana</i>
eds	ab.e.mb.eds	arcto-boreal evergreen malacophyll broad-leaved erect dwarf shrub	<i>Empetrum</i> , <i>Cassiope</i> , <i>Vaccinium vitis-idaea</i>
dlhs	ab.cd.mb.lhs	arcto-boreal cold-deciduous malacophyll broad-leaved low or high shrub	<i>Alnus crispa</i>
elhs	ab.e.mb.lhs	arcto-boreal evergreen malacophyll broad-leaved low or high shrub	<i>Calluna</i> , <i>Ledum</i>
cbc	ab.e.n.lhs	arcto-boreal evergreen needle-leaved low or high shrub	<i>Pinus pumila</i>
bss	bo.cd.mb.lhs	boreal cold-deciduous malacophyll broad-leaved low or high shrub	<i>Alnus incana</i> , <i>Myrica</i>
bes	bo.e.mb.lhs	boreal evergreen malacophyll broad-leaved low or high shrub	<i>Chamaedaphne</i>
ss	bo/te-dt.cd/e.mb.lhs	boreal or temperate drought-tolerant cold-deciduous or evergreen malacophyll broad-leaved low or high shrub	<i>Euphorbia</i>
bec	bo.e.n.t	boreal evergreen needle-leaved tree	<i>Picea abies</i>
cdc	bo.cd.n.t	boreal cold-deciduous needle-leaved tree	<i>Larix</i>
bst	bo.cd.mb.t	boreal cold-deciduous malacophyll broad-leaved tree	<i>Populus</i>
ec	eu.e.n.t	eurythermic evergreen needle-leaved tree	<i>Pinus sylvestris</i>
ctc	te.e.n.t	temperate evergreen needle-leaved tree	<i>Tsuga canadensis</i> , <i>Tsuga heterophylla</i> , <i>Thuja</i>
ctc1	ma.e.n.t	maritime evergreen needle-leaved tree	<i>Taxus</i>
ctc2	c-te.e.n.t	cool-temperate evergreen needle-leaved tree	<i>Tsuga mertensiana</i>
ts	te-fa.cd.mb.t	temperate (spring-frost avoiding) cold-deciduous malacophyll broad-leaved tree	<i>Quercus</i> (deciduous), <i>Acer</i>
ts1	te-ft.cd.mb.t	temperate (spring-frost tolerant) cold-deciduous malacophyll broad-leaved tree	<i>Tilia cordata</i> , <i>Fagus grandifolia</i>
ts2	te-fi.cd.mb.t	temperate (spring-frost intolerant) cold-deciduous malacophyll broad-leaved tree	<i>Juglans</i> , <i>Castanea</i>
wte1	te.e.mb.wp	temperate evergreen malacophyll broad-leaved woody plants	<i>Ilex</i> , <i>Hedera</i>

<sup>a</sup>We have adopted PFT names and codes that make the relationship to bioclimate, phenology, leaf morphology, and life form explicit. We include the PFT codes used in the BIOME 6000 project for information and to facilitate cross-comparison with earlier biomization schemes.

biomes; that is, it was defined by the presence of PFTs assigned to any tundra biome. The separate tundra biomes were identified in a second step. This two-step procedure was adopted because it was found to allow a slightly more accurate reconstruction of the modern tundra-forest boundary.

### 2.5. Modern Vegetation at the Surface Pollen Sites

[18] In order to assess the accuracy of present-day vegetation reconstructions based on pollen data, we required a way to assign each pollen sample to a modern biome independently, based on the present-day distribution of the biomes. In principle this can be done in two ways, which do not necessarily agree. Both are approximations of what the pollen samples actually “see.”

[19] 1. On the one hand, we had access to information from field notes about the vegetation surrounding most of the surface pollen samples in Beringia, and a proportion of the samples from other regions. Field notes indicate the local (i.e., within 1–2 km) vegetation immediately around the sampling site, which can be influenced by local soil, orography and drainage patterns.

[20] 2. On the other hand, we could estimate the regional vegetation around each site from a contemporary biome map. We constructed this map using the observation-based gridded (0.5°) map of potential natural vegetation map of *Haxeltine and Prentice* [1996] as a starting point (see Table 7). This map itself was derived from a variety of existing regional and global maps. We overlaid the map of *Walker* [2000] digitized at the same (0.5°) resolution to provide further information on regional tundra types. Data from *Haxeltine and Prentice* [1996] were retained for all grid cells for which *Walker* [2000] does not provide a classification including the whole of the forested area. The regional vegetation around each pollen sample was assumed to be the potential natural vegetation of the grid cell in which the sample is located. In cases where a sample fell in a grid cell not covered by the map (for example some coastal sites), the most prevalent biome in immediately adjacent grid cells was used. The biomes assigned in this way form the basis for the actual vegetation assignments plotted in Figure 2b, and used in the construction of Table 8.

[21] This procedure does not assign any modern sites to graminoid and forb tundra, because this tundra type does



Table 4. Definition of the Biomes Used in the Biomization Procedure

Biome Code	Biome Name in BIOME4	Definition	Characteristic Species or Assemblages in NH Arctic	Equivalents
STEP	temperate grassland	Treeless vegetation dominated by drought-tolerant forbs and grasses	Grasses, Chenopodiaceae, Asteraceae, Liliaceae	prairie (North America), steppe (Ukraine, North China)
STEP	temperate xerophytic shrubland	Treeless vegetation dominated by drought-tolerant, and locally salt-tolerant, shrubs with grasses	<i>Artemisia</i> , <i>Purshia</i> , <i>Chrysothamnus</i> , <i>Hippophae</i> , and grasses	sagebrush steppe of North American Great Basin, cold semi-desert in central Asia
CUSH	cushion-forb tundra	Discontinuous treeless vegetation characterised by high-arctic rosette or cushion forbs, grasses, mosses, lichens	Saxifragaceae, Caryophyllaceae, <i>Papaver</i> , <i>Draba</i>	polar desert, cold semi-desert
DRYT	graminoid and forb tundra	Treeless and predominantly herbaceous vegetation dominated by arctic forbs, graminoids, true mosses, and lichens	<i>Artemisia</i> , <i>Kobresia</i> , Brassicaceae, Asteraceae, Caryophyllaceae, grass, true mosses	<i>Yurtsev's</i> [2001] cryoxeric vegetation, <i>Kobresia</i> meadows
PROS	prostrate dwarf-shrub tundra	Treeless vegetation dominated by arcto-boreal prostrate dwarf shrubs, grasses, arctic forbs, true mosses, lichens	<i>Salix</i> , <i>Dryas</i> , <i>Pedicularis</i> , Asteraceae, Brassicaceae, grass and sedge	northern Arctic tundra
DWAR	erect dwarf-shrub tundra	Treeless vegetation dominated by arcto-boreal dwarf shrubs, with graminoids, true mosses, and lichens	<i>Betula</i> , <i>Salix</i> , <i>Vaccinium</i> , <i>Empetrum</i> , <i>Cassiope</i> , grass and sedge	low Arctic tundra, tussock tundra
SHRU	low- and high-shrub tundra	Treeless vegetation dominated by arcto-boreal deciduous or evergreen low and high shrubs, sometimes with tussock-forming graminoids and true mosses, bog mosses, and lichens	<i>Alnus</i> , <i>Betula</i> , <i>Salix</i> , <i>Pinus pumila</i> , <i>Eriophorum</i> , <i>Sphagnum</i>	shrub tundra
CLDE	cold deciduous forest	Forest dominated by boreal broad-leaved or needle-leaved deciduous trees, sometimes with significant component of arcto-boreal evergreen needle-leaved shrubs	<i>Larix</i> , <i>Betula</i> or <i>Populus</i> subg. <i>tremula/tremuloides</i> , with <i>Pinus pumila</i>	E. Siberian larch forests; aspen parkland (North America); maritime birch forests
TAIG	cold evergreen needle-leaved forest	Forest dominated by boreal evergreen needle-leaved trees, with boreal deciduous or needle-leaved trees	<i>Picea</i> and <i>Abies</i> , with some <i>Pinus</i> , <i>Betula</i> , <i>Populus</i> or <i>Larix</i>	dark taiga (Russia); boreal forest (North America)
COCO	cool evergreen needle-leaved forest	Forest dominated by boreal and temperate evergreen needle-leaved trees	<i>Pinus strobus</i> , <i>Pinus resinosa</i> , <i>Picea glauca</i> , <i>Pinus banksiana</i> , <i>Abies balsamea</i> , <i>Acer saccharum</i>	southern boreal forest (Europe); cool conifer forest (south-central Ontario-Quebec)
CLMX	cool-temperate evergreen needle-leaved forest	Forest lacking boreal evergreen needle-leaved trees, with boreal deciduous broad-leaved trees and evergreen needle-leaved trees	<i>Tsuga</i> and <i>Picea sitchensis</i> , <i>Pinus</i> with <i>Betula</i> , <i>Pinus</i> with <i>Betula</i> and <i>Populus</i>	<i>Pinus-Populus-Betula</i> forest (W. Norway & Scotland); (Central Asia)
TEDE	temperate deciduous forest	Forest dominated by temperate deciduous broad-leaved trees	<i>Quercus</i> , <i>Fagus</i> , <i>Carya</i> , <i>Ulmus</i> , <i>Castanea</i>	temperate deciduous forest (eastern North America);
COMX	cool mixed forest	Forest with boreal and temperate evergreen needle-leaved trees and deciduous broad-leaved trees	<i>Picea</i> , <i>Pinus</i> , <i>Quercus</i> , <i>Acer</i> , <i>Ulmus</i>	memoral forest (western Europe) North American northern hardwoods; Scandinavian boreo-memoral forest



**Table 6.** Assignment of Pollen Taxa to Plant Functional Types

PFT Name	PFT Code	Taxa
bog moss	m	<i>Sphagnum</i>
grass graminoid	g	Gramineae, Poaceae
sedge graminoid	s	<i>Carex</i> -type, Cyperaceae
rush graminoid	r	Juncaceae undiff.
rosette or cushion forb	rcfb	Caryophyllaceae undiff., <i>Draba</i> , <i>Saxifraga hirculus</i> -type, Saxifragaceae undiff., <i>Saxifraga caesia</i> , <i>Saxifraga cernua</i> -type, <i>Saxifraga hieracifolia</i> -type, <i>Saxifraga nivalis</i> -type, <i>Saxifraga nelsoniana</i> , <i>Saxifraga oppositifolia</i> , <i>Saxifraga tricuspidata</i> -type, <i>Saxifraga</i> , <i>Silene</i> -type
arctic forb	ar.fb	<i>Achillea</i> -type, <i>Achillea</i> , <i>Aconitum</i> , Alliaceae, <i>Allium</i> , <i>Androsace</i> , <i>Anemone</i> -type, <i>Anemone</i> , <i>Angelica</i> -type, <i>Angelica</i> , <i>Antennaria</i> -type, <i>Apiaceae</i> , <i>Arnica</i> , <i>Artemisia</i> , Asteraceae subf. Asteroideae, Asteraceae subf. Asteroideae (thick spines), Asteraceae subf. Carduoideae, <i>Aster</i> , Asteraceae undiff., <i>Astragalus</i> -type, Brassicaceae, Brassicaceae-type undiff., <i>Caltha</i> , Campanulaceae, <i>Campanula</i> -type, <i>Cardamine</i> , <i>Cerastium</i> -type, Asteraceae subf. Chichorioideae, <i>Claytonia acutifolia</i> , <i>Claytonia</i> , <i>Claytoniella vassilievii</i> , Asteraceae (high spine), <i>Corydalis</i> , Crassulaceae, Caryophyllaceae undiff., <i>Dodecatheon</i> -type, <i>Epilobium angustifolium</i> , <i>Epilobium</i> , Fabaceae, Fabaceae undiff., <i>Galium</i> , <i>Galium</i> -type, Gentianaceae, <i>Gentiana</i> undiff., <i>Geum</i> , <i>Hedysarum</i> -type, <i>Koenigia</i> , <i>Koenigia islandica</i> , Liliaceae, <i>Lloydia</i> , <i>Lychnis dianthus</i> -type, Onagraceae, <i>Oxyria digyna</i> , <i>Oxyria</i> , <i>Oxytropis</i> , Papaveraceae, <i>Papaver</i> , <i>Parnassia palustris</i> , <i>Parnassia</i> , <i>Pedicularis</i> , <i>Pedicularis langsdorfii</i> , <i>Pedicularis lanceolata</i> , <i>Pedicularis verticillata</i> , <i>Phlox</i> , <i>Phlox sibirica</i> , <i>Plantago</i> undiff., <i>Plantago canescens</i> -type, Plantaginaceae, Polygonaceae, Polemoniaceae undiff., <i>Polemonium</i> , <i>Polygonum</i> , <i>Polygonum bistorta</i> -type, <i>Polygonum viviparum</i> , <i>Polygonum aviculare</i> , <i>Polygonum aviculare</i> -type, Polygonaceae, <i>Polygonum sect. Bistorta</i> , Portulacaceae, <i>Potentilla</i> , <i>Potentilla</i> -type, Primulaceae, <i>Pyrola</i> , Pyrolaceae, <i>Ranunculus</i> undiff., Ranunculaceae undiff., <i>Ranunculus acris</i> -type, Ranunculaceae type 1, Ranunculaceae type 2, Rosaceae undiff., Rubiaceae undiff., <i>Rubus</i> , <i>Rubus arcticus</i> , <i>Rumex</i> Subgen. <i>Acetosa</i> /R. Subgen. <i>Acetosella</i> , <i>Rumex</i> , <i>Rumex acetosa</i> , <i>Rumex/Oxyria digyna</i> , <i>Rumex/Oxyria</i> , <i>Rumex</i> -type, <i>Sagina</i> , <i>Sanguisorba</i> , <i>Sanguisorba officinalis</i> , <i>Saussurea alpina</i> , <i>Saussurea</i> , <i>Saxifraga hirculus</i> -type, Saxifragaceae undiff., <i>Saxifraga caesia</i> , <i>Saxifraga cernua</i> -type, <i>Saxifraga hieracifolia</i> -type, <i>Saxifraga nivalis</i> -type, <i>Saxifraga nelsoniana</i> , <i>Saxifraga oppositifolia</i> , <i>Saxifraga tricuspidata</i> -type, <i>Saxifraga</i> , Scrophulariaceae, <i>Senecio</i> -type, <i>Sedum</i> , <i>Silene</i> -type, <i>Solidago</i> -type, <i>Spiraea</i> , <i>Stellaria</i> , <i>Taraxacum</i> , <i>Taraxacum</i> -type, <i>Thalictrum</i> , <i>Tofieldia</i> , <i>Trientalis europaea</i> , <i>Trollius europaeus</i> , Umbelliferae undiff., <i>Valeriana</i> , Valerianaceae, <i>Veratrum</i> , Violaceae
boreal or temperate drought-tolerant forb	bo/te-dt.fb	<i>Achillea</i> -type, <i>Achillea</i> , Alliaceae, <i>Allium</i> , <i>Amarantha</i> , <i>Ambrosia</i> -type, <i>Antennaria</i> -type, <i>Anthemis</i> -type, <i>Apiaceae</i> , <i>Artemisia</i> , Asteraceae subf. Asteroideae, Carduoideae, Asteraceae undiff., <i>Astragalus</i> -type, <i>Atriplex nudicaulis</i> , Boraginaceae, Brassicaceae, Brassicaceae-type undiff., <i>Bupleurum</i> , Campanulaceae, <i>Campanula</i> -type, <i>Centaurea</i> , <i>Centaurea cyanus</i> , <i>Centaurea cyanus</i> -type, <i>Centaurea jacea</i> , <i>Cerastium</i> -type, Chenopodiaceae, Chenopodiaceae/Amaranthaceae, Asteraceae subf. Chichorioideae, Asteraceae (high spine), Crassulaceae, Cruciferae, Caryophyllaceae undiff., Dipsacaceae, <i>Euphorbia</i> , Fabaceae undiff., <i>Galium</i> , <i>Galium</i> -type, Gentianaceae, <i>Hedysarum</i> -type, Iridaceae, <i>Iris</i> , <i>Kochia laniflora</i> , Labiatae undiff., Lamiaceae, Liguliflorae, Liliaceae, Onagraceae, <i>Oxytropis</i> , Papaveraceae, <i>Papaver</i> , <i>Phlox</i> , <i>Plantago</i> undiff., <i>Plantago canescens</i> -type, Plantaginaceae, Plumbaginaceae, <i>Polygonum</i> , Polygonaceae, Portulacaceae, <i>Potentilla</i> , <i>Potentilla</i> -type, Primulaceae, <i>Ranunculus</i> undiff., Ranunculaceae undiff., <i>Ranunculus acris</i> -type, Ranunculaceae type 1, Ranunculaceae type 2, Ranunculaceae, Rosaceae undiff., Rubiaceae undiff., <i>Rumex</i> , <i>Rumex/Oxyria digyna</i> , <i>Rumex/Oxyria</i> , <i>Rumex</i> -type, <i>Sanguisorba</i> , <i>Sanguisorba officinalis</i> , <i>Sarcobatus vermiculatus</i> , <i>Sarcobatus vermiculatus</i> , Scrophulariaceae, <i>Senecio</i> -type, <i>Sedum</i> , <i>Silene</i> -type, <i>Solidago</i> -type, <i>Spergula arvensis</i> , <i>Sphaeralcea</i> , <i>Spiraea</i> , <i>Stellaria</i> , <i>Taraxacum</i> , <i>Taraxacum</i> -type, <i>Thalictrum</i> , <i>Thymus</i> , Umbelliferae undiff., <i>Valeriana</i> , Valerianaceae

Table 6. (continued)

PFT Name	PFT Code	Taxa
arctic cold-deciduous malacophyll broad-leaved prostrate dwarf shrub	ar.cd.mb.pds	<i>Arctostaphylos</i> , <i>Salix</i> , <i>Salix herbacea</i> -type, <i>Salix cf. herbacea</i> , <i>Salix vestita</i> -type
arctic evergreen malacophyll broad-leaved prostrate dwarf shrub	ar.e.mb.pds	<i>Arctostaphylos</i> , <i>Diapensia</i> , <i>Dryas</i> , <i>Dryas</i> -type
arcto-boreal evergreen needle-leaved prostrate dwarf shrub	ab.e.n.pds	Cupressaceae, <i>Juniperus</i> -type, <i>Juniperus communis</i>
arcto-boreal cold-deciduous malacophyll broad-leaved erect dwarf shrub	ab.cd.mb.eds	<i>Arctostaphylos</i> , <i>Betula sect. nanae</i> , <i>Betula nana</i> -type, <i>Betula</i> undiff., <i>Betula</i> corroded, <i>Betula exilis</i> , <i>Betula</i> small, Ericaceae undiff., <i>Ericales</i> , <i>Salix</i> , <i>Salix herbacea</i> -type, <i>Salix cf. herbacea</i> , <i>Salix vestita</i> -type
arcto-boreal evergreen malacophyll broad-leaved erect dwarf shrub	ab.e.mb.eds	<i>Arctostaphylos</i> , <i>Cassiope</i> , <i>Empetrum</i> , Ericaceae undiff., Ericaceae undiff., <i>Ericales</i> , <i>Vaccinium</i> -type, <i>Vaccinium vitis-idaea</i>
arcto-boreal cold-deciduous malacophyll broad-leaved low or high shrub	ab.cd.mb.lhs	<i>Alnus crispa</i> , <i>Alnus viridis</i> ssp. <i>crispa</i> -type, <i>Alnus viridis</i> ssp. <i>fruticosa</i> , <i>Duschekia fruticosa</i> , <i>Alnus viridis</i> ssp. <i>sinuata</i> -type, <i>Alnus viridis</i> , <i>Alnus viridis</i> -type, <i>Alnaster</i> , <i>Alnus fruticosa</i> -type, <i>Alnaster</i> , <i>Alnus</i> undiff., <i>Betula sect. nanae</i> , <i>Betula nana</i> -type, <i>Betula</i> undiff., <i>Betula</i> corroded, <i>Betula exilis</i> , <i>Betula fruticosa</i> , <i>Betula humilis</i> , <i>Betula</i> small, Betulaceae, Ericaceae undiff., <i>Ericales</i> , <i>Salix</i> , <i>Salix vestita</i> -type, <i>Vaccinium</i> -type, <i>Vaccinium uliginosum</i> -type
arcto-boreal evergreen malacophyll broad-leaved low or high shrub	ab.e.mb.lhs	<i>Calluna vulgaris</i> , <i>Calluna</i> , Ericaceae undiff., <i>Ericales</i> , <i>Ledum palustre</i> , <i>Ledum</i> -type, <i>Rhododendron</i> , <i>Rubus chamaemorus</i>
arcto-boreal evergreen needle-leaved low or high shrub	ab.e.n.lhs	<i>Pinus pumila</i>
boreal cold-deciduous malacophyll broad-leaved low or high shrub	bo.cd.mb.lhs	<i>Alnus crispa</i> , <i>Alnus incana</i> , <i>Alnus incana</i> -type, <i>Alnus viridis</i> ssp. <i>crispa</i> -type, <i>Alnus viridis</i> ssp. <i>fruticosa</i> , <i>Duschekia fruticosa</i> , <i>Alnus viridis</i> ssp. <i>sinuata</i> -type, <i>Alnus viridis</i> , <i>Alnus viridis</i> -type, <i>Alnus fruticosa</i> -type, <i>Alnaster</i> , <i>Alnus</i> undiff., <i>Betula</i> undiff., <i>Betula</i> corroded, Betulaceae, Caprifoliaceae undiff., Cornaceae, <i>Cornus</i> , <i>Cornus alba</i> , <i>Cornus canadensis</i> , <i>Cornus mas/C. svecica</i> , <i>Cornus sericea</i> , <i>Cornus suecica</i> -type, <i>Linnea</i> , <i>Lonicera</i> , <i>Myrica</i> , <i>Myrica gale</i> , <i>Ribes</i> , <i>Salix</i> , <i>Sambucus</i> , <i>Shepherdia</i> , <i>Shepherdia canadensis</i> , <i>Viburnum</i> , <i>Viburnum opulus</i>
boreal evergreen malacophyll broad-leaved low or high shrub	bo.e.mb.lhs	<i>Chamaedaphne calyculata</i>
boreal or temperate drought-tolerant cold-deciduous or evergreen malacophyll broad-leaved low or high shrub	bo/te-dt.cd/e.mb.lhs	<i>Euphorbia</i> , Fabaceae undiff.
boreal evergreen needle-leaved tree	bo.e.n.t	<i>Abies</i> , <i>Abies sibirica</i> , bi-saccate pollen, <i>Picea abies</i> ssp. <i>obovata</i> , <i>Picea abies</i> , <i>Picea</i> abnormal, <i>Picea eupicea</i> , <i>Picea glauca</i> , <i>Picea mariana</i> , <i>Picea</i>
boreal cold-deciduous needle-leaved tree	bo.cd.n.t	<i>Larix</i> , <i>Larix gmelinii</i> , <i>Larix laricina</i> , <i>Larix siberica</i> , <i>Larix dahurica</i>
boreal cold-deciduous malacophyll broad-leaved tree	bo.cd.mb.t	<i>Alnus glutinosa</i> , <i>Alnus incana</i> , <i>Alnus incana</i> -type, <i>Alnus viridis</i> ssp. <i>sinuata</i> -type, <i>Alnus</i> undiff., <i>Alnus hirsuta</i> , <i>Betula arbor s. albae</i> , <i>Betula albae</i> , <i>Betula</i> undiff., <i>Betula arbor.</i> , <i>Betula</i> corroded, <i>Betula pendula</i> , <i>Betula platyphylla</i> , <i>Betula pubescens</i> , <i>Betula tortuosa</i> , Betulaceae, <i>Chosenia</i> , <i>Corylus</i> , <i>Corylus</i> -type, <i>Populus</i> , <i>Populus balsamifera</i> , <i>Populus tremuloides</i> , <i>Populus tremula</i> , <i>Salix</i> , <i>Salix vestita</i> -type, <i>Sorbus</i>
eurythermic evergreen needle-leaved tree	eu.e.n.t	Bi-saccate pollen, Cupressaceae, <i>Juniperus</i> -type, <i>Juniperus communis</i> , <i>Pinus diploxylon</i> -type, <i>Pinus haploxylon</i> , <i>Pinus</i> , <i>Pinus banksiana</i> -type, <i>Pinus contorta</i> , <i>Pinus subg. Pinus</i> , <i>Pinus sect.</i> , <i>Pinus sibirica</i> , <i>Pinus subg. Strobos</i> , <i>Pinus</i> undiff., <i>Pinus sylvestris</i>
temperate evergreen needle-leaved tree	te.e.n.t	<i>Abies</i> , <i>Picea</i> abnormal, <i>Picea eupicea</i> , <i>Picea omorica</i> , <i>Picea</i> , <i>Thuja</i> , <i>Tsuga</i> , <i>Tsuga canadensis</i> , <i>Tsuga diversifolia</i> , <i>Tsuga heterophylla</i>
maritime evergreen needle-leaved tree	ma.e.n.t	<i>Taxus</i>
cool-temperate evergreen needle-leaved tree	c-te.e.n.t	<i>Tsuga mertensiana</i>
temperate (spring-frost avoiding) cold-deciduous malacophyll broad-leaved tree	te-fa.cd.mb.t	<i>Acer</i> , <i>Acer saccharum</i> , <i>Acer spicatum</i> , <i>Alnus glutinosa</i> , <i>Alnus</i> undiff., Caprifoliaceae undiff., <i>Carya</i> , <i>Carya ovata</i> , Cornaceae, <i>Cornus</i> , <i>Cornus alba</i> , Fabaceae undiff., <i>Fraxinus americana</i> -type, <i>Fraxinus nigra</i> -type, <i>Fraxinus pennsylvanica</i> -type, <i>Lonicera</i> , <i>Populus</i> , <i>Prunus pennsylvanica</i> , <i>Quercus</i> (deciduous), <i>Quercus</i> , <i>Salix</i> , <i>Salix vestita</i> -type, <i>Sambucus</i> , <i>Sorbus</i> , <i>Viburnum</i> , <i>Viburnum opulus</i>
temperate (spring-frost tolerant) cold-deciduous malacophyll broad-leaved tree	te-ft.cd.mb.t	<i>Carpinus betulus</i> , <i>Carpinus</i> , <i>Corylus</i> , <i>Corylus avellana</i> , <i>Corylus cornuta</i> , <i>Corylus</i> -type, <i>Fagus</i> , <i>Frangula</i> , <i>Frangula alnus</i> , <i>Fraxinus excelsior</i> -type, <i>Ostrya/Carpinus</i> , <i>Ribes</i> , <i>Tilia</i> , <i>Tilia cordata</i> , <i>Ulmus</i> , <i>Ulmus glabra</i>
temperate (spring-frost intolerant) cold-deciduous malacophyll broad-leaved tree	te-fi.cd.mb.t	<i>Carpinus</i> , <i>Castanea</i> , <i>Castanea dentata</i> , <i>Juglans</i> , <i>Ostrya</i> , <i>Ostrya</i> -type, <i>Ostrya/Carpinus</i> , <i>Platanus</i> , <i>Rhamnus</i> , <i>Ulmus</i>
temperate evergreen malacophyll broad-leaved woody plants	te.e.mb.wp	<i>Genista</i> , <i>Hedera</i> , <i>Hedera helix</i> , <i>Ilex</i> , <i>Rhododendron</i>

**Table 7.** Equivalences Between Biome Classifications Used in the BIOME4 Model, in the Pollen-Based Mapping Scheme, by *Haxeltine and Prentice* [1996], and by *Walker* [2000]

BIOME4 <sup>a</sup>	Pollen-Based Mapping Scheme <sup>a</sup>	<i>Haxeltine and Prentice</i> [1996]	<i>Walker</i> [2000]
temperate deciduous broadleaf forest	temperate deciduous broadleaf forest <sup>b</sup>	temperate deciduous forest	
cool mixed forest	cool mixed forest	temperate/boreal mixed forest	
cool evergreen needleleaf forest	cool evergreen needleleaf forest	temperate conifer forest	
cool-temperate evergreen needleleaf forest	cool-temperate evergreen needleleaf forest	boreal evergreen forest/woodland or temperate conifer forest	
cold evergreen needleleaf forest	cold evergreen needleleaf forest	boreal evergreen forest/woodland	
cold deciduous forest	cold deciduous forest	boreal deciduous forest/woodland	
temperate xerophytic shrubland	temperate grassland and xerophytic shrubland (steppe) <sup>c</sup>	tall grassland or short grassland	
temperate deciduous broadleaf savanna	temperate deciduous broadleaf forest <sup>b</sup>	moist savanna	
temperate grassland	temperate grassland and xerophytic shrubland (steppe) <sup>c</sup>	tall grassland or short grassland	
graminoid and forb tundra	graminoid and forb tundra		
low- and high-shrub tundra	low- and high-shrub tundra		oceanic evergreen low-shrub and low-shrub subzones
erect dwarf shrub tundra	erect dwarf shrub tundra		oceanic herb, dwarf-shrub and dwarf-shrub subzones
prostrate dwarf shrub tundra	prostrate dwarf shrub tundra		prostrate-shrub subzone
cushion forb lichen and moss tundra	cushion-forb tundra		lichens, moss subzone

<sup>a</sup>The BIOME4 model and pollen-based mapping scheme are given by *Kaplan et al.* [2003].

<sup>b</sup>We did not attempt to separate temperate deciduous broadleaf savanna from forest.


<sup>c</sup>We did not attempt to separate temperate shrubland from grassland.

reconstructed presence of low- and high-shrub tundra in the Canadian archipelago is incorrect and is caused by long-distance transport of pollen of *Alnus viridis* from farther south. We assume this would not be an issue at LGM, when shrub tundra was restricted.

[26] The ability to successfully reproduce the regional vegetation is dependent on the nature of the samples used. Lacustrine surface samples yield reconstructions in better agreement with the regional vegetation (61% correct allocations) than moss polsters (54%). However, when compared to descriptions of local vegetation (as given by the original investigators' field notes), moss polsters give a better representation than lacustrine samples. These differences are consistent with the general empirical finding, explained by pollen transport theory [*Prentice*, 1985, 1988], that lake sediments have a wider source region than peat or other terrestrial pollen sampling sites. Samples derived from terrestrial soils and/or surface sediments yield considerably worse reconstructions (43% correct) than either lake or peat samples. This probably reflects differential oxidation of the components of the original pollen assemblage. Samples obtained in geological contexts (lacustrine core tops, peat core tops or terrestrial section tops) are systematically less successful in reproducing the observed modern vegetation than other samples from the same type of environment. Thus predictions of the modern vegetation using lacustrine core tops are successfully in only 55% of all cases (compared to 61% for lacustrine surface samples), peat core tops in 46% of all cases (compared to 54% for moss polsters), and section core tops in 38% of all cases (compared to 43% for surface soils/sediments). In some regions (e.g., Beringia, western Europe) it would be possible to reconstruct modern vegetation patterns on the basis of only lacustrine surface samples or moss polsters; this would not be possible for central Siberia or western Beringia. Furthermore, it would appear that although the biases due to site type impact on our numerical assessment of the success of the biomization procedure, it does not affect the ability of the biomization technique to successfully reproduce the geographic patterns of vegetation. We have therefore made both modern and palaeo-vegetation reconstructions using all types of sample, and have checked to see that the conclusions drawn from comparisons of these maps are not affected by site-type (or other) biases.

### 3.2. Biome Reconstructions for 6 ka

[27] The biome reconstructions for 6 ka show systematic changes from present (Figure 2c and Table 9). Table 9 shows the frequency of transitions, based only on sites where both 6 ka and present biomes have been reconstructed from samples in one core. The reconstructed treeline was farther north than present (Table 10) in Fennoscandia (western Europe) and central Siberia (e.g., Taimyr peninsula). In contrast, Beringia shows little or no change from present, and treeline was south of its present position in Labrador and Keewatin. These results are consistent with the preliminary synthesis by *TEMPO Members* [1996], subsequent studies using biomization (as summarized by *Prentice et al.* [2000]), and with the reconstructions based on subfossil tree remains by *MacDonald et al.* [2000]. They do not support the con-



**Figure 2.** Biomes (a) reconstructed from modern pollen data, compared to (b) modern vegetation at the pollen data sites, and biomes reconstructed from fossil pollen data at (c) 6 ka and (d) 18 ka. In cases where two or more 6 ka or 18 ka sites lie very close to one another, they have been slightly shifted in these plots to permit each site to be individually resolved. The sites that have been shifted are indicated in Table 2.

ventional idea that the Arctic treeline was consistently north of its present position in a circumpolar sense [e.g., *Foley et al.*, 1994]. More extensive geographic changes are seen in the boundaries among forest types south of the Arctic treeline, including a northward displacement of

temperate deciduous forest in eastern North America (south of our study region) and especially Fennoscandia (western Europe), and a more restricted distribution of cold evergreen needle-leaved forest toward the western edges of the continents (western Europe and eastern Beringia).

**Table 8.** Observed Versus Reconstructed Biomes Using the Modern Pollen Data Set<sup>a</sup>

Observed	Reconstructed Biomes												Observed	Percent Correct	
	STEP	CUSH	DRYT	PROS	DWAR	SHRU	CLDE	TAIG	COCO	CLMX	TEDE	COMX			
STEP														0	
CUSH			1											1	0.0
DRYT														0	
PROS	1		44	<b>16</b>	46	20		3						130	12.3
DWAR	2		26	17	<b>110</b>	89	15	16						275	40.0
SHRU	1		11	9	74	<b>258</b>	18	109				1		481	53.6
CLDE	3		4	1	7	77	<b>123</b>	34						249	49.4
TAIG			2	2	19	132	25	<b>596</b>	5	7		15		803	74.2
COCO			1	1		2	4	61	<b>8</b>		2	13		92	8.7
CLMX				1		3		5	1	<b>2</b>				12	16.7
TEDE					2	1	1	2	2					8	0.0
COMX					2		2	19	2		1	<b>21</b>		47	44.7
Total	7	0	89	47	260	582	188	845	18	9	3	50		<b>2098</b>	

<sup>a</sup>The definitions of the BIOME codes are given in Table 4.

[28] North of the treeline, a few samples north of 75 °N in Greenland are assigned to low- and high-shrub tundra at 6 ka. This indicates that low- and high-shrub tundra may have been present >500 km farther north at 6 ka than today. Even if the classification reflects long-distance pollen transport of shrub pollen, the presence of this pollen at very high latitudes implies that the pollen source was substantially farther north than today. However, there is no evidence in the data for systematic shifts in the distribution of the more cold-tolerant tundra types. The boundary between erect and prostrate dwarf-shrub tundra was apparently located near its present position.

### 3.3. Biome Reconstructions for the LGM

[29] The LGM reconstruction (Figure 2d) suggests that forests were absent (or at least, highly restricted) north of 55°N. Low- and high-shrub tundra was reconstructed at a single site on the Beringian land bridge, and two sites in what is now the *Larix* forest region of eastern Siberia. Cushion forb tundra was identified on the basis of samples of LGM age from three small lakes on Andøya, northwest Norway, which lay at the very edge of the Scandinavian ice sheet at its maximum extent [Vorren, 1978]. Elsewhere, the data indicate that the landscape consisted of a mosaic of tundra biomes. In Beringia the dominant biomes were erect dwarf-shrub tundra, prostrate dwarf-shrub tundra, and graminoid and forb tundra. Farther west, in western and central

(Taimyr) Siberia, the dominant biomes were graminoid and forb tundra and temperate grassland or xerophytic shrubland (steppe), suggesting the prevalence of very dry conditions downwind of the Scandinavian ice sheet [cf. Siebert and Marsiat, 2001]. The data, although sparse, suggest that there may have been a transition in central Siberia from graminoid and forb tundra north of about 65°N, to true temperate steppe farther south. This steppe was distinguished by the occurrence of drought-tolerant forbs or shrubs, in place of arctic forbs (Table 5).

## 4. Discussion and Conclusions

[30] The comprehensive data set of mid-Holocene and LGM pollen data north of 55°N, and the biome reconstructions made using these data based on a unified circumpolar classification of plant functional types and biomes, represent a key resource for understanding and modeling changes in northern-high latitude vegetation and climate in response to changing global boundary conditions between glacial and interglacial states. Given the success of the biome reconstruction method in reconstructing the tundra-forest boundary and the distributions of major forest and tundra biomes from surface pollen data, we can now make more confident statements about the large-scale, climatically induced changes in vegetation distribution between the LGM, mid-Holocene and present.

**Table 9.** Changes in Biome Assignments Between 0 ka and 6 ka<sup>a</sup>

Reconstructed 6k Biomes	Reconstructed Modern Biomes												Total	
	STEP	CUSH	DRYT	PROS	DWAR	SHRU	CLDE	TAIG	COCO	CLMX	TEDE	COMX		
STEP								1						1
CUSH														0
DRYT				1	2	1		1						5
PROS			1	<b>2</b>										3
DWAR					7	3								10
SHRU			3		10	<b>35</b>	1	17						66
CLDE					2	5	<b>14</b>	20						41
TAIG					1	7	5	<b>64</b>					1	78
COCO					1		1	19	<b>6</b>				2	29
CLMX										<b>1</b>				1
TEDE					2	1	3	10	1			<b>1</b>	11	29
COMX					1			10	2				<b>12</b>	25
Total	0	0	4	3	26	52	24	142	9	1	1	1	26	<b>288</b>

<sup>a</sup>The definitions of the BIOME codes are given in Table 4.

**Table 10.** Approximate Changes in the Position of Arctic Treeline at 6 ka Compared to Present Shown by Pollen Data

Sector	Change
Mackenzie Delta	100 km N
Keewatin	280 km S
Labrador	170 km S
Greenland	no evidence for treeline
Atlantic	no evidence for treeline
Western Europe	70 km N
Eastern Europe	insufficient data
Western Siberia	insufficient data
Central Siberia (Taimyr)	180 km N
Central Siberia (Lena)	70 km N
Western Beringia (Eastern Siberia)	no change
Western Beringia (Chukotka)	no evidence for treeline
Eastern Beringia (Alaska)	no change

[31] Our findings confirm the conclusions of the BIOME 6000 studies [Prentice *et al.*, 1996; Tarasov *et al.*, 1998; Edwards *et al.*, 2000; Williams *et al.*, 2000] concerning the mid-Holocene to present changes in the location of the Arctic tree limit, now based on a much larger set of modern and 6 ka pollen samples and a unified biomization scheme applied across the entire circumpolar region. These results establish that the shift in the forest limit between 6 ka and present was characterized by a strong circumpolar asymmetry, with the largest poleward shifts in central Siberia, little or no change in Beringia, and treeline south of present in eastern Canada. This pattern was first documented by *TEMPO Members* [1996] and is also shown in estimates of relative warming, using a rule-based interpretation of pollen data, made by *CAPE Project Members* [2001]. A small residual ice sheet was still present in Labrador around 6 ka, and this presumably accounts for the anomalous treeline situation there [Richard, 1995; Williams *et al.*, 2000]. The treeline in this region reached its maximum northward extension after 6 ka [Payette, 1992; Richard, 1995]. In the rest of the circum-polar region, it remains a challenge to deduce how a zonally uniform pattern of increased summer and annual insolation, due to the greater-than-present tilt of the Earth's rotational axis, could produce a strongly asymmetrical response in the location of the Arctic treeline. Recent climate simulations using a model with sea-ice dynamics [Vavrus and Harrison, 2003] suggest that the explanation may lie in the modulation of the orbital forcing through sea-ice dynamics. Modern sea ice is thinnest and least compact in the eastern Arctic Ocean and thickest and most compact in the western Arctic Ocean, largely as a result of the climatological drift of ice from the Siberian to the North American coast. In a warmer climate, sea ice melts more easily in the eastern part of the basin where ice divergence occurs, but tends to persist in the western sector because of ice convergence. Simulated changes in ice concentration at 6 ka show a dipole spatial pattern, with large reductions in the east and little or no change in the west, and these spatial differences in the change in ice coverage lead in turn to larger increases in surface temperature in central Siberia than elsewhere [Vavrus and Harrison, 2003].

[32] At LGM, conditions are generally assumed to have been far colder and drier than present throughout the high latitudes. This is confirmed by the extreme southward

displacement of the forest belts [Peyron *et al.*, 2000] and supported by geomorphological evidence that indicates cold, dry conditions in the unglaciated part of Beringia [Hopkins *et al.*, 1982]. Our data also indicate a major change in the relative importance of different tundra biome types between LGM and present. Thus low- and high-shrub tundra was extremely restricted in extent and, in contrast, there was a major extension of graminoid and forb tundra and an intergradation of steppe and tundra in the north Eurasian interior. It seems likely that dryness (lack of snow cover) contributed to this change. The contrasting fates of low- and high-shrub tundra, and graminoid and forb tundra, during the LGM illustrate how an altered climate regime can cause dramatic shifts in the relative areal extent of different biomes.

[33] Investigating the mechanisms behind regional patterns of vegetational change on orbital timescales requires the use of climate and vegetation models with altered boundary conditions. In the companion paper [Kaplan *et al.*, 2003] we use the data from the present paper as a benchmark for models of the mid-Holocene and LGM environments. In doing so, we provide physically possible explanations for the major features of the 6 ka and 18 ka palaeovegetation maps.

[34] **Acknowledgments.** The Pan-Arctic Initiative was supported by a research grant from the Palaeoecology of Arctic Lakes and Estuaries (PALE) initiative of the U.S. National Science Foundation (NSF) and research resources provided by the Potsdam Institute for Climate Impact Research (PIK) and the Max Planck Institute for Biogeochemistry, Jena (MPI-BGC). Workshop funds were provided by the International Geosphere-Biosphere Programme (IGBP) through its the Task Force on Global Integration, Analysis and Modeling (GAIM) and the IGBP Data and Information System (IGBP-DIS), and by the International Arctic Science Council (IASC) through the joint IASC-IGBP activity Feedbacks and Arctic Terrestrial Ecosystems (FATE). We thank Gerhard Bönisch and Matthew Duvall for assistance with the PAIN pollen database and Silvana Schott for assistance in the construction of the observed vegetation map. We thank James Jordan for providing unpublished data from Alaska.

## References

- Ager, T. A., Surficial geology and Quaternary history of the Healy Lake area, Alaska, Master's thesis, Univ. of Alaska, Dep. of Geol., Fairbanks, 1972.
- Ager, T. A., Late Quaternary environmental history of the Tanana Valley, Alaska, *Rep. 54*, Inst. of Polar Stud., Ohio State Univ., Columbus, 1975.
- Ager, T. A., A 16,000 year pollen record from St Michael Island, Norton Sound, western Alaska, paper presented at Sixth Biennial Meeting, Am. Quat. Assoc., Orono, Maine, 18–20 Aug. 1980.
- Ager, T. A., Vegetational history of western Alaska during the Wisconsin glacial interval and the Holocene, in *Paleoecology of Beringia*, edited by D. M. Hopkins *et al.*, pp. 75–93, Academic, San Diego, Calif., 1982.
- Ager, T. A., Holocene vegetational history of Alaska, in *Late-Quaternary Environments of the United States*, vol. 2, *The Holocene*, edited by H. E. Wright Jr., pp. 128–141, Univ. of Minn. Press, Minneapolis, 1983.
- Ager, T. A., and J. P. Bradbury, Quaternary history of vegetation and climate of the Yukon Delta-Norton Sound Area, in *The United States Geological Survey in Alaska: Accomplishments During 1980*, edited by W. L. Coonrad, *U.S. Geol. Surv. Circ.*, 844, 78–80, 1982.
- Ager, T. A., and L. B. Brubaker, Quaternary palynology and vegetational history of Alaska, in *Pollen Records of Late Quaternary North American Sediments*, edited by V. M. Bryant Jr. and R. G. Holloway, pp. 353–384, Am. Assoc. of Stratigr. Palynol. Found., Dallas, Tex., 1985.
- Ager, T. A., and J. D. Sims, Postglacial pollen and tephra records from lakes in the Cook Inlet region, southern Alaska, in *The United States Geological Survey in Alaska: Accomplishments During 1981*, edited by W. L. Coonrad and R. L. Elliot, *U.S. Geol. Surv. Circ.*, 868, 103–105, 1984.
- Aleshinskaya, Z. V., and V. S. Gunova, Istoriya ozera Nero kak otrazhenie dinamiki okruzhayushego landschafta (History of Nero as reflection on the surrounding landscape dynamics), in *Problemy Paleohidrologii (Problems of Paleohydrology)*, edited by G. P. Kalinin and K. Klige, pp. 214–222, Nauka, Moscow, 1976.



- Alhonen, P., Palaeolimnological investigations of three inland lakes in south-western Finland, *Acta Bot. Fenn.*, 76, 1–55, 1967.
- Alm, T., Øvre Æräsvatn: Palynostratigraphy of a 22,000 to 10,000 BP lacustrine record on Andøya, northern Norway, *Boreas*, 22, 171–188, 1993.
- Andersen, S. T., Forests at Løvenholm, Djursland, Denmark, at present and in the past, *Biol. Skr.*, 24, 1–208, 1984.
- Andersen, S. T., B. Aaby, and B. V. Odgaard, Environment and Man, *J. Dan. Archaeol.*, 2, 184–196, 1983.
- Anderson, J. H., A palynological study of late Holocene vegetation and climate in the Healy Lake Area of Alaska, *Arctic*, 28, 62–69, 1975.
- Anderson, P. M., Late Quaternary vegetational change in the Kotzebue Sound area, northwestern Alaska, *Quat. Res.*, 24, 307–321, 1985.
- Anderson, P. M., Late Quaternary pollen records from the Kobuk and Noatak River drainages, northwestern Alaska, *Quat. Res.*, 29, 263–276, 1988.
- Anderson, P. M., R. E. Reanier, and L. B. Brubaker, Late Quaternary vegetational history of the Black River region in northeastern Alaska, *Can. J. Earth Sci.*, 25, 84–94, 1988.
- Anderson, P. M., R. E. Reanier, and L. B. Brubaker, A 14,000-year pollen record from Sithylenkat Lake, north-central Alaska, *Quat. Res.*, 33, 400–404, 1990.
- Anderson, P. M., A. V. Lozhkin, W. R. Eisner, D. M. Hopkins, and L. B. Brubaker, Pollen records from Ten Mile and Wonder Lake, Alaska, *Geogr. Phys. Quat.*, 48, 131–141, 1994.
- Anderson, P. M., B. V. Belaya, O. Yu. Glushkova, and A. V. Lozhkin, Novye dannye o istorii rastitel'nosti severnogo Priokhot'ya v pozdnem pleistotsene i golotsene (New data about vegetation history in northern Priokhot'e in Late Pleistocene and Holocene), in *Pozdnyy pleistotsen i golotsen Beringii (Late Pleistocene and Holocene of Beringia)*, edited by M. Kh. Gagiev, pp. 33–54, Northeast Interdisciplinary Res. Inst., Far East Branch, Russ. Acad. of Sci., Magadan, 1997a.
- Anderson, P. M., B. V. Belaya, O. Yu. Glushkova, A. V. Lozhkin, and L. B. Brubaker, A lacustrine pollen record from near altitudinal forest limit, upper Kolyma region, northeastern Siberia, *Holocene*, 7, 331–335, 1997b.
- Andreev, A. A., and V. A. Klimanov, Istoriya rastitel'nosti i klimata Tsentral'noi Yakutii v golotsene i pozdnem pleistotsene (Vegetation and climate history of Central Yakutia during the Holocene and Late Pleistocene), in *Formirovaniye otlozhenii i rossypeli na severo-vostoke SSSR (Formation of Deposits and Placers on North-East of the USSR)*, pp. 26–51, Northeast Interdisciplinary Res. Inst., Far East Branch, Russ. Acad. of Sci., Magadan, 1989.
- Andreev, A. A., and V. A. Klimanov, Izmeneniya rastitel'nosti i klimata mezhdurech'ya rek Ungra i Yakokit (Yuzhnaya Yakutia) v golotsene (Vegetation and climate changes in the interfluvium of the rivers Ungra and Yakokit (the Southern Yakutia) during the Holocene), *Bot. Zh.*, 76, 334–351, 1991.
- Andreev, A. A., V. A. Klimanov, L. D. Sulerzhitsky, and N. A. Khotinsky, Khronologiya landshaftno-klimaticheskikh izmenenii Tsentral'noi Yakutii v golotsene (Chronology of landscape and climate changes in Central Yakutia during the Holocene), in *Paleoklimaty pozdnelednikov'ya i golotsena (Paleoclimates of Late Glacial and Holocene)*, edited by N. A. Khotinsky, pp. 115–121, Nauka, Moscow, 1989.
- Andreev, A. A., F. A. Romanenko, L. D. Sulerzhitsky, P. E. Tarasov, and E. I. Terekhov, History of relief and vegetation dynamics on the western coast of Baidara Gulf during late Pleistocene and Holocene, *Stratigr. Geol. Correl.*, 6(5), 520–525, 1998.
- Andreev, A. A., V. A. Klimanov, and L. D. Sulerzhitsky, Vegetation and climate history of the Yana River lowland, Russia, during the last 6400 yr, *Quat. Sci. Rev.*, 20, 259–266, 2001.
- Andrews, J. T., and H. Nichols, Modern pollen deposition and Holocene paleotemperature reconstructions, central northern Canada, *Arct. Alp. Res.*, 13, 387–408, 1981.
- Arkipov, S. A., and V. I. Astakhov, *Paleogeografiya Zapadno-Sibirskoi ravniny v pozdnezryanskom lednikovom maksimume (Paleogeography of the Western Siberian Plain During the Late Zyryan Glacial Maximum)*, 130 pp., Nauka, Moscow, 1980.
- Arkipov, S. A., and M. R. Votakh, Palinologicheskaya kharakteristika i absolyutnyi vozrast torfyanka bliz ust'ya reki Tom' (Palynological characteristics and the absolute age of peat near the mouth of the Tom' River), in *Palinologiya Sibiri (Palynology of Siberia)*, edited by V. N. Saks, pp. 112–118, Nauka, Moscow, 1980.
- Arkipov, S. A., T. P. Levina, and V. A. Panychev, Palynological characteristics of two Holocene peats from the middle and lower Ob' river valley, in *Paleopalynology of Siberia*, edited by V. N. Saks, pp. 123–127, Nauka, Moscow, 1980.
- Arslanov, Kh., V. Auslender, L. A. Gromova, A. Zubkov, and V. Khomutova, Paleogeograficheskie osobennosti i absolyutnyi vozrast maksimal'noi stadii Valdaiskogo oledeneniya v raione ozera Kubenskogo (Paleogeographical peculiarities and absolute age of the maximum stage of Valday glaciation in the Lake Kubenskoe region), *Dokl. Akad. Nauk SSSR*, 195, 1395–1399, 1970.
- Arslanov, Kh., N. Davydova, D. Subetto, and V. Khomutova, Kareliiskii peresheek (Karelian Isthmus), in *Istoriya ozer Vostochno-Evropeiskoi ravniny (The Lake History of East-European Plain)*, pp. 64–77, Nauka, Moscow, 1992.
- Arslanov, Kh. A., N. A. Gey, N. N. Davydova, R. N. Dzhinoridze, B. I. Koshechkin, M. Ya. Pushenko, A. E. Rybalko, M. A. Spiridonov, D. A. Subetto, and V. I. Khomutova, Novye dannye po pleistotsenovoi i golotsenovoi istorii ozera Ladoga (New data on the Late Pleistocene and Holocene history of Lake Ladoga), *Izv. Ross. Geogr. Obshchestva*, 128/2, 12–21, 1996.
- Asplund, H., and I. Vuorela, Settlement studies in Kemiö: Archaeological problems and palynological evidence, *Fennoscandia Archaeol.*, VI, 67–79, 1989.
- Bakhareva, V. A., Palinologicheskaya kharakteristika pozdnechetvertichnykh i golotsenovykh otlozhenii v okrestnostyakh derevni Pershino na reke Irtysh (The palynological characteristics of late Quaternary and Holocene deposits in the vicinity of Pershino Village on the Irtysh River), in *Palinostratigrafiya Mezozoya i Zenozya Sibiri (The Palinostratigraphy of Mesozoic and Cenozoic of Siberia)*, edited by V. S. Volkova and A. F. Khlonova, pp. 115–120, Nauka, Moscow, 1985.
- Belorusova, Zh. M., N. V. Lovelius, and V. V. Ukraintseva, Paleogeografiya pozdnego pleistotsena i golotsena v raione nakhodki selirikaniskoi loshadi (Paleogeography of the Late Pleistocene and Holocene in the area of Selirikan Horse discovery), in *Fauna i flora antropogena Severo-Vostoka Sibiri (Fauna and Flora of North-Eastern Siberia During the Anthropogene)*, pp. 265–276, Nauka, Moscow, 1977.
- Bender, M. M., R. A. Bryson, and D. A. Baerreis, University of Wisconsin radiocarbon dates I, *Radiocarbon*, 7, 399–407, 1965.
- Bender, M. M., R. A. Bryson, and D. A. Baerreis, University of Wisconsin radiocarbon dates II, *Radiocarbon*, 8, 522–533, 1966.
- Bender, M. M., R. A. Bryson, and D. A. Baerreis, University of Wisconsin radiocarbon dates III, *Radiocarbon*, 9, 530–544, 1967.
- Bender, M. M., R. A. Bryson, and D. A. Baerreis, University of Wisconsin radiocarbon dates IV, *Radiocarbon*, 10, 161–168, 1968.
- Bender, M. M., R. A. Bryson, and D. A. Baerreis, University of Wisconsin radiocarbon dates XV, *Radiocarbon*, 20, 157–167, 1978.
- Bennett, K. D., S. Boreham, M. J. Sharp, and V. R. Switsur, Holocene history of environment, vegetation and human settlement on Catta Ness, Lunnasting, Shetland, *J. Ecol.*, 80, 241–273, 1992.
- Berdovskaya, G. N., K paleogeografii ozera Chany (About paleogeography of Lake Chany), in *Pul'siruyushchee ozero Chany (Pulsating Lake Chany)*, edited by N. P. Smirnova and A. V. Shnitnikov, pp. 33–40, Nauka, Moscow, 1982.
- Berger, A., The role of CO<sub>2</sub>, sea level and vegetation during the Milankovitch-forced glacial-interglacial cycles, in *Geosphere-Biosphere Interactions and Climate*, edited by L. O. Bengtsson and C. U. Hammer, pp. 119–146, Cambridge Univ. Press, New York, 2001.
- Berglund, B. E., The Post-glacial shore displacement in eastern Blekinge, southeastern Sweden, *Sver. Geol. Unders.*, Ser. C, 58(5), 1–47, 1964.
- Berglund, B. E., Late-Quaternary vegetation in eastern Blekinge, southeastern Sweden, A pollen-analytical study: II. Post-Glacial time, *Opera Bot.*, 12, 1–190, 1966.
- Bick, H., A postglacial pollen diagram from Angmagssalik, east Greenland, *Medd. Groenl.*, 204, 5–22, 1978.
- Bigelow, N. H., Late Quaternary vegetation and lake level changes in central Alaska, Dissertation, Univ. of Alaska, Fairbanks, 1997.
- Bigelow, N. H., and M. E. Edwards, A 14,000-yr paleoenvironmental record from Windmill Lake, central Alaska: Late-Glacial and Holocene vegetation in the Alaska Range, *Quat. Sci. Rev.*, 20, 203–215, 2001.
- Birks, H. J. B., *Past and Present Vegetation of the Isle of Skye: A Palaeoecological Study*, 415 pp., Cambridge Univ. Press, New York, 1973.
- Birks, H. J. B., and B. J. Madsen, Flandrian vegetational history of Little Loch Roag, Isle of Lewis, Scotland, *J. Ecol.*, 67, 825–842, 1979.
- Bjelm, L., Deglaciation of the Smalard Highland, with special reference to deglaciation, icethickness and chronology, *Lundqua theses*, 2, 1–78, 1976.
- Bjerck, L. B., Del 2 vegetasjonshistorie, *Arkeol. Rapp. Bergen Mus.*, 5, 133–175, 1983.
- Björck, S., and S. Hakansson, Radiocarbon dates from Late Weichselian lake sediments in south Sweden as a basis for chronostratigraphic subdivision, *Boreas*, 11, 141–150, 1982.
- Björck, S., and T. Persson, Late Weichselian and Flandrian biostratigraphy and chronology from Hochstetter Forland, Northeast Greenland, *Medd. Groenl. Geosci.*, 5, 3–19, 1981.
- Bliss, L. C., Adaptations of arctic and alpine plants to environmental conditions, *Arctic*, 15, 117–144, 1962.
- Blyakharchuk, T. A., Sporovo-pyl'tsevaya kharakteristika bolot Verhneketskogo raiona (Spore and pollen characteristics of peatlands of the

- Verhneketskii region during the Holocene), in *Molodye uchoeny narodnomy khozyaystvy Ekonomicheskaya ozenka landshafta Tomskoi oblast (Abstracts of the Conference "Young Scientist for Peoples Economy: Economical Evaluation of the Tomsk Region")*, pp. 24–25, Tomsk Univ. Press, Tomsk, Russia, 1980.
- Blyakharchuk, T. A., Sopryazhenost' sporovo-pyl'zevogo i botanicheskogo analizov torfa pri izuchenii lesno-bolotnykh landshaftov yuga Zapadnoi Sibiri (Palynological and botanical analyses of peat in the investigation of forest-mire landscapes of the southern West Siberia), in *Tezisy konferentsii: Problemy Golotsena (Abstracts of the Conference "Problems of the Holocene")*, pp. 16–18, Akad. Nauk Gruzinskoy SSR, T'bilisi, Georgian Republic, 1988.
- Blyakharchuk, T. A., Istoriya rastitel'nosti yugo-vostoka Zapadnoi Sibiri v goltsene po dannym botanicheskogo i sporovo-pyl'tsevogo analiza torfa (The Holocene history of vegetation of south-eastern West Siberia by botanical and pollen analyses of peat deposits), Ph.D. thesis, 248 pp., Tomsk State Univ., Tomsk, Russia, 1989a.
- Blyakharchuk, T. A., Detal'noe raschlenenie rannego i srednego golotsena po dannym issledovaniya azonal'nogo merzlogo torfyanika na yuge Zapadnoi Sibiri (Middle and early Holocene by study of unzonal pals), in *Tezisy konferentsii "Palinologiya i poleznye iskopaemye" (Abstracts of the Conference "Palynology and Mineral Sources")*, pp. 1105–1145, Akad. Nauk BSSR, Minsk, 1989b.
- Blyakharchuk, T. A., Suktsessii bolotnoi rastitel'nosti i klimata po dannym issledovaniya dvukh torfyanikov na yuge Zapadnoi Sibiri (Succession of the mire vegetation and climate according to investigation of two peat-bogs in the southern West Siberia), in *Struktura i razvitiye bolotnykh ekosistem i rekonstruktsii paleogeograficheskikh uslovii, (Structure and Development of Peatland Ecosystems and Reconstructions of Paleogeographical Conditions)*, edited by M. A. Ilomets, pp. 45–49, Akad. Nauk Est. SSR, Tallinn, 1989c.
- Bogdel', I. I., Razvitiye prirody Belorussii v goltsene (Holocene environmental changes of Byelorussia during the Holocene), Ph.D. thesis, 192 pp., Minsk State Univ., Minsk, 1984.
- Bonan, G. B., D. Pollard, and S. L. Thompson, Effects of boreal forest vegetation on global climate, *Nature*, 359, 716–718, 1992.
- Boyarskaya, T. D., and T. N. Kaplina, Novye dannye o razvitiit rastitel'nosti severnoi Yakutii v goltsene (New data about vegetation development in northern Yakutia during the Holocene), *Vestn. Mosk. Univ., Ser. 5: Geogr.*, 5, 70–75, 1979.
- Braconnot, P., S. Joussaume, O. Marti, and N. Noblet, Synergistic feedbacks from ocean and vegetation on the African monsoon response to mid-Holocene insolation, *Geophys. Res. Lett.*, 26, 2481–2484, 1999.
- Brubaker, L. B., H. L. Garfinkel, and M. E. Edwards, A late Wisconsin and Holocene vegetation history from the central Brooks Range: Implications for Alaskan paleocology, *Quat. Res.*, 20, 194–214, 1983.
- Brubaker, L. B., P. M. Anderson, B. M. Murray, and D. Koon, A palynological investigation of true-moss (Bryidae) spores: Morphology and occurrence in modern and late Quaternary lake sediments of Alaska, *Can. J. Bot.*, 76, 2145–2157, 1998.
- CAPE Project Members, Holocene paleoclimate data from the Arctic: Testing models of global climate change, *Quat. Sci. Rev.*, 20, 1275–1287, 2001.
- Chapin, F. S., III, W. Eugster, J. P. McFadden, A. H. Lynch, and D. A. Walker, Summer differences among Arctic ecosystems in regional climate forcing, *J. Clim.*, 13, 2002–2010, 2000a.
- Chapin, F. S., III, et al., Arctic and boreal ecosystems of western North America as components of the climate system, *Global Change Biol.*, 6, 211–223, 2000b.
- Christensen, T. R., T. Friberg, M. Sommerkorn, J. Kaplan, L. Illeris, H. Soegaard, C. Nordstroem, and S. Jonasson, Trace gas exchange in a high-arctic valley: 1. Variations in CO<sub>2</sub> and CH<sub>4</sub> flux between tundra vegetation types, *Global Biogeochem. Cycles*, 14, 701–713, 2000.
- Clayden, S. L., L. C. Cwynar, and G. M. MacDonald, Stomate and pollen content of lake surface sediments from across the tree line on the Taimyr Peninsula, Siberia, *Can. J. Bot.*, 74, 1009–1015, 1996.
- Clayden, S. L., L. C. Cwynar, G. M. MacDonald, and A. A. Velichko, Holocene pollen and stomates from a forest-tundra site on the Taimyr Peninsula, Siberia, *Arct. Alp. Res.*, 29, 327–333, 1997.
- Cole, H., Objective reconstruction of the paleoclimate record through application of eigenvectors of present-day pollen spectra and climate to the late Quaternary pollen stratigraphy, Dissertation, Univ. of Madison, Madison, Wisc., 1969.
- Cubasch, U., G. A. Meehl, G. J. Boer, R. J. Stouffer, M. Dix, A. Noda, C. A. Senior, S. Raper, and K. S. Yap, Projection of future climate change, in *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J. T. Houghton et al., pp. 525–582, Cambridge Univ. Press, New York, 2001.
- Cwynar, L. C., A Late-Quaternary vegetation history from Hanging Lake, northern Yukon, Dissertation, Univ. of Toronto, Toronto, Ont., Canada, 1980.
- Cwynar, L. C., A Late-Quaternary vegetation history from Hanging Lake, northern Yukon, *Ecol. Monogr.*, 52, 1–24, 1982.
- Cwynar, L. C., A late Quaternary vegetation history from Lily Lake, Chilkat Peninsula, southeast Alaska, *Can. J. Bot.*, 68, 1106–1112, 1990.
- Cwynar, L. C., Paleovegetation and paleoclimatic changes in the Yukon at 6 ka BP, *Geogr. Phys. Quat.*, 49, 29–35, 1995.
- Cwynar, L. C., and R. W. Spear, Reversion of forest to tundra in the central Yukon, *Ecology*, 72, 202–212, 1991.
- Danielsen, A., Pollen-analytical late Quaternary studies in the Ra district of Østfold, southeast Norway, *Arbok Univ. Bergen Mat. Naturvitensk. Ser.*, 14, 1–143, 1970.
- Davidovich, T. D., Sovremennyye sporovo-pyl'tsevye spektry vostochnogo i yuzhnogo poberezh'ya Chukotskogo poluostrova (Recent pollen spectra of the eastern and southern coast of the Chukcha Peninsula, in *Palinologicheskii issledovaniya na Severo-Vostoke SSSR (Palynological Investigations on the Northeastern USSR)*, pp. 74–80, Northeast Interdisciplinary Res. Inst., Far East Branch, USSR Acad. of Sci., Vladivostok, 1978.
- Davydova, N., Diatomovyi analiz (Diatom analysis), in *Obshchie zakonomernosti vozniknoveniya i razvitiya ozer: Metody izucheniya istorii ozer, Istoriya ozer SSSR*, vol. 1, edited by D. D. Kvasov, 253 pp., Nauka, Leningrad, 1986.
- Davydova, N., Diatomovyi analiz donnykh otlozhenii (Diatom analyses of bottom deposits), in *Ratsional'noe ispol'zovanie prirodnykh resursov v Belorussii (Rational Utilization of the Natural Resources in Byelorussia)*, pp. 19–25, Nauka, Moscow, 1988.
- Davydova, N., D. Subetto, and V. Khomutova, Paleolimnologiya ozer Vishnevskoe i Michurinskoe, Karel'skii peresheek (Paleolimnology of Vishnevskoe and Michurinskoe Lakes (Karelian Isthmus), in *Antropogeniye izmeneniya sistem malyykh ozer (Anthropological Changes of Small Lake Systems)*, pp. 195–198, Gidrometizdat, St. Petersburg, 1991.
- Davydova, N., D. Subetto, and V. Khomutova, Ozero Il'men' (Lake Ilmen), in *Istoriya ozer Vostochno-Evropейskoi ravniny (The Lake History of East-European Plain)*, edited by N. N. Davydova, pp. 101–117, Nauka, Moscow, 1992.
- Dennegård, B., Pollenanalys av en Lagerföljd från Tranemosjön, Västergötland, *Publ. B 142*, 13 pp., Geol. Inst. Chalmers Tek. Högskola, Göteborg, Sweden, 1980.
- Dennegård, B., Pollenanalys av en Lagerföljd från Dalstropasjön, Västergötland, *Publ. B 168*, 15 pp., Geol. Inst. Chalmers Tek. Högskola, Göteborg, Sweden, 1981.
- de Noblet, N. I., I. C. Prentice, S. Joussaume, D. Texier, A. Botta, and A. Haxeltine, Possible role of atmosphere-biosphere interactions in triggering the last glaciation, *Geophys. Res. Lett.*, 23, 3191–3194, 1996.
- Dickson, J. H., D. A. Stewart, R. Thompson, G. Turner, M. S. Baxter, N. D. Drndarsky, and J. Rose, Palynology, palaeomagnetism, radiometric dating of Flandrian marine and freshwater sediments of Loch Lomond, *Nature*, 274, 548–553, 1978.
- Digerfeldt, G., The post-glacial development of Lake Trummen. Regional vegetation history, water level changes and palaeolimnology, *Folia Limnol. Scand.*, 16, 1–96, 1972.
- Digerfeldt, G., The post-glacial development of the bay Ranviken, Lake Immeln, *Lundqua Rep.*, 1, 1–59, 1973.
- Digerfeldt, G., A standard profile for Littorina transgressions in western Skåne, south Sweden, *Boreas*, 4, 125–142, 1975.
- Digerfeldt, G., The Flandrian development of Lake Flarken: Regional vegetation history and palaeolimnology, *Lundqua Rep.*, 13, 1–101, 1977.
- Digerfeldt, G., The Holocene development of Lake Sambosjön, 1. The regional vegetation history, *Lundqua Rep.*, 23, 1–24, 1982.
- Digerfeldt, G., and S. Welinder, Settlement development and human impact in the Hallsjön area, Västergötland, W. Sweden, *Lundqua Rep.*, 15, 1–30, 1978.
- Donner, J. J., Pollen frequencies in the Flandrian sediments of Lake Vakojärvi, south Finland, *Comment. Biol.*, 53, 3–19, 1972.
- Donner, J. J., P. Alhonen, M. Eronen, H. Jungner, and I. Vuorela, Biostratigraphy and radiocarbon dating of the Holocene lake sediments of Työtjärvi and the peats in the adjoining bog Varrassuo west of Lahti in southern Finland, *Ann. Bot. Fenn.*, 15, 258–280, 1978.
- Ebel, T., M. Melles, and F. Niessen, Laminated sediments from Levinson-Lessing lake, northern central Siberia: A 30,000 year record of environmental history?, in *Land-Ocean Systems in the Siberian Arctic: Dynamics and History*, edited by H. Kassens et al., pp. 425–435, Springer-Verlag, New York, 1999.
- Edwards, M. E., and L. B. Brubaker, Late Quaternary vegetation history of the Fishhook Bend area, Porcupine River, Alaska, *Can. J. Earth Sci.*, 23, 1765–1773, 1986.

- Edwards, M. E., P. M. Anderson, H. L. Garfinkel, and L. B. Brubaker, Late Wisconsin and Holocene vegetational history of the Upper Koyukuk region, Brooks Range, AK, *Can. J. Bot.*, 63, 616–626, 1985.
- Edwards, M. E., et al., Pollen-based biomes for Beringia 18,000, 6000 and 0 <sup>14</sup>C yr BP, *J. Biogeogr.*, 27, 521–554, 2000.
- Eide, F., Vegetasjonshistoriske undersøkelser på Valborgmyr, Kårstø, Tysvær i Rogaland, in *Vegetasjonshistoriske Undersøkelser på Kårstø, Tysvær i Rogaland, Rapp. 23*, edited by F. G. Eide and A. Paus, pp. 1–45, Bot. Inst., Univ. Bergen, Bergen, Norway, 1982.
- Elenka, H., et al., Pollen-based biome reconstruction for southern Europe and Africa 18,000 yr BP, *J. Biogeogr.*, 27, 621–634, 2000.
- Elias, S. A., S. K. Short, C. H. Nelson, and H. H. Birks, Life and times of the Bering land bridge, *Nature*, 382, 60–63, 1996.
- Eliina, G. A., Printsipy i metody rekonstruktsii i kartirovaniya rastitel'nosti golotsena (*Principles and Methods for Reconstruction and Mapping of Holocene Vegetation*), 159 pp., Nauka, Moscow, 1981.
- Eliina, G. A., and L. V. Filimonova, Late-Glacial vegetation on the territory of Karelia, in *Palaeohydrology of the Temperate Zone III: Mires and Lakes*, edited by A. Raukas and L. Saarse, pp. 53–69, Valgus, Tallinn, 1987.
- Eliina, G. A., and L. V. Filimonova, Russian Karelia, in *Palaeoecological Events During the Last 15,000 Years: Regional Syntheses of Palaeoecological Studies of Lakes and Mires in Europe*, edited by B. E. Berglund et al., pp. 353–366, John Wiley, New York, 1996.
- Eliina, G. A., and V. I. Khomutova, Correlation of Holocene sequences of bottom sediments from Onega Lake and its old bays in terms of palynological data, in *Methods for the Investigation of Lake Deposits: Palaeoecological and Paleoclimatological Aspects, Proceedings of the International Symposium, Vilnius, September 15–19, 1986*, edited by M. Kabailene, pp. 193–203, V. Kapsukas Univ., Vilnius, Lithuania, 1987.
- Eliina, G. A., and T. K. Yurkovskaya, Bolotnye ekosistemy nizkogorii severnoi taigi (Bog ecosystems of northern taiga in the low elevation mountains), in *Bolotnye ekosistemy Evropeiskogo Severa (Bog Ecosystems of European North)*, edited by L. V. Danilovich, pp. 5–24, Inst. Biol., Karel. Fil., Akad. Nauk SSSR, Petrozavodsk, Russia, 1988.
- Eliina, G. A., O. L. Kuznetsov, E. I. Devyatova, R. I. Lebedeva, A. I. Maksimov, and N. V. Stoikina, Sovremennaya i golotsenovaya rastitel'nost' natsional'nogo parka Paanayarvi (severo-zapadnaya Kareliya) (Modern and Holocene vegetation of the Paanayarvi National Park (North-Western Karelia)), *Bot. Zh.*, 79, 13–31, 1994.
- Eliina, G. A., Kh. A. Arslanov, V. A. Klimanov, and L. I. Usova, Paleorastitel'nost' i klimatostratigrafiya golotsena Lovozerskoi ravniny Kol'skogo poluostrova (Paleovegetation and climatic stratigraphy of the Lovozero Plain (Kola Peninsula)), *Bot. Zh.*, 80, 3–16, 1995.
- Elliot-Fisk, D. L., J. T. Andrews, S. K. Short, and W. N. Mode, Isopoll maps and analysis of the distribution of the modern pollen rain, eastern and central northern Canada, *Geogr. Phys. Quat.*, 36, 91–108, 1982.
- Elovicheva, Y. K., and I. I. Bogdel', Novye razrezy golotsena Belarusii (New Holocene sections in Byelorussia), in *Geologicheskoe stroenie osadochnoi tolshchi Belorussii (Geological Composition of Sedimentary Sequence of Byelorussia)*, edited by V. A. Kuznetsov, V. F. Ropot, and Ia. K. Elovicheva, pp. 141–169, Nauka i Tekhnika, Minsk, 1985.
- Engelmark, R., The vegetational history of the Umeå area during the past 4000 years, in *Palaeo-ecological Investigation in Coastal Västerbotten, N. Sweden, Early Norrland*, vol. 9, edited by R. Engelmark et al., pp. 75–111, K. Vitterhets Hist. och Antikvitets Akad., Almqvist and Wiksell Int., Stockholm, 1976.
- Engelmark, R., The comparative vegetational history of inland and coastal sites in Medelpad, N. Sweden, during the Iron Age, in *Archaeological and Palaeoecological Studies in Medelpad, N. Sweden, Early Norrland*, vol. 11, edited by E. Baudou et al., pp. 25–62, K. Vitterhets Hist. och Antikvitets Akad., Almqvist and Wiksell Int., Stockholm, 1978.
- Engelmark, R., The Paleoenvironment, in *Coastal Resources and Settlement Stability, Aun*, vol. 3, edited by N. Broadbent, pp. 158–173, Inst. of North Eur. Archaeol., Uppsala Univ., Uppsala, Sweden, 1979.
- Engelmark, R., and J.-E. Wallin, Pollen analytical evidence for Iron Age agriculture in Hälsingland, central Sweden, *Archaeol. Environ.*, 4, 353–366, 1985.
- Epstein, H. E., F. S. Chapin, M. D. Walker, and A. M. Starfield, Analyzing the functional type concept in arctic plants using a dynamic vegetation model, *Oikos*, 95, 239–252, 2001.
- Eronen, M., and H. Hyvärinen, Subfossil pine dates and pollen diagrams from northern Fennoscandia, *Geol. Foeren. Stockholm Foerh.*, 103, 437–455, 1982.
- Filimonova, L. V., K palinologicheskomu izucheniyu bolot morennykh ravnin srednei Karelii (Palynological studies of the peatlands on the moraine plains in middle Karelia), in *Zaprosy ekologii rastenii bolot, bolotnykh mestoobitaniy i torfyanykh zalezhei (Questions of Ecology of Plant of Bogs, Bog Habitats and Peatlands)*, pp. 122–132, Inst. Biol., Karel. Fil., Akad. Nauk SSSR, Petrozavodsk, Russia, 1995.
- Filimonova, L. V., and Ya. K. Elovicheva, Osnovnye stadii razvitiya lesnoi i bolotnoi rastitel'nosti territorii zapovednika Kivach (Main stages of the development of forest and mire vegetation on the territory of the Kivach Nature Reserve), in *Bolotnye ekosistemy evropeiskogo Severa (Bog Ecosystems of European North)*, edited by L. V. Danilovich, pp. 94–109, Inst. Biol., Karel. Fil., Akad. Nauk SSSR, Petrozavodsk, Russia, 1988.
- Firsov, L. V., T. P. Levina, and S. L. Troitskii, Holocene climatic changes in the Northern Siberia, in *Climatic Changes in Arctic Areas During the Last Ten Thousand Years: A Symposium Held at Oulanka and Kevo, 4–10 October, 1971, Acta Univ. Ouluensis, Ser. A*, vol. 2, edited by Y. Vasari, H. Hyvärinen, and S. Hicks, pp. 341–349, Univ. of Oula, Oula, Finland, 1972.
- Firsov, L. V., S. L. Troitskii, T. P. Levina, V. P. Nikitin, and V. A. Panychev, Absolutnyi vozrast i pervaya dlya severa Sibiri standartnaya pyl'tsevaya diagramma golotsenogo torfyanyka (Absolute age and first standard pollen diagram of Holocene peatland on northern Siberia), *Biul. Komissii Izuch. Chev. Perioda*, 41, 121–127, 1974.
- Firsov, L. V., V. S. Volkova, T. P. Levina, I. V. Nikolaeva, L. A. Orlova, V. A. Panychev, and I. A. Volkov, Stratigrafiya, geokhronologiya i standartnaya sporovo-pyl'tsevaya diagramma golotsenovogo torfyanyka, Gladkoye boloto, Novosibirsk (Stratigraphy, geochronology, and standard spore-pollen diagram for Holocene peat, Gladkoye Bog, Novosibirsk), in *Problemy stratigrafii i paleogeografii pleistotsena Sibiri: K XI Kongressu INQUA v SSSR (Problems of Stratigraphy and Paleogeography of Pleistocene of Siberia: For XI Congress INQUA in USSR)*, edited by S. A. Arkhipov, V. S. Volkova, and T. S. Troitskaya, *Tr. Inst. Geol. Geofiz. Akad. Nauk SSSR, Sib. Otd.*, no. 521, 96–107, 1982.
- Firsov, L. V., V. A. Panychev, and L. A. Orlova, *Catalog of Radiocarbon Dates USSR Academy of Sciences*, 87 pp., Geol. Inst. of Siberia Branch, Novosibirsk, Russia, 1985.
- Florin, M. B., Late-glacial and pre-boreal vegetation in southern central Sweden, II. Pollen, spore and diatom analyses, *Striae*, 5, 3–60, 1977.
- Foley, J. A., J. E. Kutzbach, M. T. Coe, and S. Levis, Feedbacks between climate and boreal forests during the Holocene epoch, *Nature*, 371, 52–54, 1994.
- Fredskild, B., Postglacial plant succession and climatic changes in a west Greenland Bog, *Rev. Palaeobot. Palynol.*, 4, 113–127, 1967.
- Fredskild, B., Studies in the vegetational history of Greenland: Palaeobotanical investigations of some holocene lake and bog deposits, *Medd. Groenl.*, 198, 1–245, 1973.
- Fredskild, B., The Holocene vegetational development of the Godthåbsfjord area, West Greenland, *Medd. Groenl. Geosci.*, 10, 3–28, 1983.
- Fredskild, B., The Holocene vegetational development of Tugtutligssuaq and Qeqertat, northwest Greenland, *Medd. Groenl. Geosci.*, 14, 3–20, 1985.
- Fries, M., Studies of the sediments and the vegetational history in the Ösbynsjö basin, north of Stockholm, *Oikos*, 13, 76–96, 1962.
- Fromm, E., Kvartära bildningar, in *Beskrivning till geologiska kartbladet Örebro SV*, edited by E. Fromm, pp. 30–98, *Sver. Geol. Unders., Ser. Ae*, 5, 1972.
- Funder, S., Holocene stratigraphy and vegetation history in the Scoresby Sund area, East Greenland, *Bull. Groenl. Geol. Unders.*, 129, 1–66, 1978.
- Funder, S., and N. Abrahamsen, Palynology in a polar desert, eastern North Greenland, *Boreas*, 17, 195–207, 1988.
- Gaillard, M.-J., A palaeohydrological study of Krageholmssjön (Scania, south Sweden), Regional vegetation history and water-level changes, *Lundqua Rep.*, 25, 1–40, 1984.
- Gajewski, K., Modern pollen assemblages at the tree-line in New-Quebec, *Can. J. Earth Sci.*, 28, 643–648, 1991.
- Gajewski, K., Modern and Holocene pollen assemblages from some small Arctic lakes on Somerset Island, NWT, Canada, *Quat. Res.*, 44, 228–236, 1995.
- Gajewski, K., and S. Garralla, Holocene vegetation histories from 3 sites in the tundra of northwestern Quebec, Canada, *Arct. Alp. Res.*, 24, 329–336, 1992.
- Gajewski, K., S. Payette, and J. C. Ritchie, Holocene vegetation history at the boreal-forest: Shrub-tundra transition in north-western Québec, *J. Ecol.*, 81, 433–443, 1993.
- Gallée, H., J. P. van Ypersele, T. Fichefet, I. Marsiat, C. Tricot, and A. Berger, Simulation of the last glacial cycle by a coupled, sectorially averaged climate-ice sheet model: 2. Response to insolation and CO<sub>2</sub> variations, *J. Geophys. Res.*, 97, 15,713–15,740, 1992.
- Gallimore, R. G., and J. E. Kutzbach, Role of orbitally induced changes in tundra area in the onset of glaciation, *Nature*, 381, 503–505, 1996.
- Ganopolski, A., C. Kubatzki, M. Claussen, V. Brovkin, and V. Petoukhov, The influence of vegetation-atmosphere-ocean interaction on climate during the mid-Holocene, *Science*, 280, 1916–1919, 1998.

- Glebov, F. Z., L. V. Karpenko, V. A. Klimanov, and T. N. Mindeeva, Paleoeokologicheskaya kharakteristika golotsena mezhdurech'ya Obi i Vasyugana (Palynological characteristics of the Holocene in the interfluvium of the rivers Ob' and Vasyugan), *Ekologiya*, 28(6), 364–370, 1997.
- Goetcheus, V. G., and H. H. Birks, Full-glacial upland tundra vegetation preserved under tephra in the Beringia National Park, Seward Peninsula, Alaska, *Quat. Sci. Rev.*, 20, 135–147, 2001.
- Göransson, H., The Flandrian vegetational history of southern Östergötland, thesis, Univ. of Lund, Lund, Sweden, 1977.
- Gunova, V. S., Palynological characteristic of Holocene Nero lake sediments, *Vestn. Mosk. Univ., Ser. 5: Geogr.*, 3, 117–119, 1972.
- Gunova, V. S., Istoriya ozera Nero po palinologicheskim dannym (The history of Lake Nero by pollen record), Can. Sci. Dissertation, Moscow State Univ., 1975.
- Gunova, V. S., and A. R. Sirin, Paleogeographical condition of bog development in Zapadnaya Dvina lowland during Holocene, in *Palynology in Russia, Part 2*, pp. 27–36, Nauka, Moscow, 1995.
- Guthrie, R. D., Origin and causes of the mammoth steppe: A story of cloud cover, woolly mammoth tooth pits, buckles, and inside-out Beringia, *Quat. Sci. Rev.*, 20, 549–574, 2001.
- Hafsten, U., Pollen-analytic investigation on the late Quaternary development in the inner Oslofjord area, *Univ. Bergen Arbok Naturvitensk. Rekke*, 8, 161 pp., 1956.
- Hafsten, U., The Norwegian *Cladium mariscus* communities and their post-glacial history, *Arbok Univ. Bergen Mat.-Naturvitensk. Ser.*, 4, 1–55, 1965.
- Hafsten, U., and T. Solem, Age, origin, and palaeo-ecological evidence of blanket bogs in Nord-Trøndelag, Norway, *Boreas*, 5, 119–141, 1976.
- Hafsten, U., K. E. Henningsmoen, and H. I. Høeg, Innvandringen av gran til Norge, in *Fortiden i Søkelyset: 14C datering gjennom 25 år*, edited by R. Nydal et al., pp. 171–198, Lab. for Radiol. Datering, Trondheim, Norway, 1979.
- Hahne, J., and M. Melles, Late- and post-glacial vegetation and climate history of the south-western Taymyr Peninsula, central Siberia, as revealed by pollen analysis of a core from Lake Lama, *Veg. Hist. Archaeobot.*, 6, 1–8, 1997.
- Hahne, J., and M. Melles, Climate and vegetation history of the Taymyr Peninsula since middle Weichselian time: Palynological evidence from Lake sediments, in *Land-Ocean Systems in the Siberian Arctic: Dynamics and History*, edited by H. Kassens et al., pp. 407–423, Springer-Verlag, New York, 1999.
- Hallsdóttir, M., On the pre-settlement history of Icelandic vegetation, *Icelandic Agric. Sci.*, 9, 17–29, 1996.
- Hallsdóttir, M., A. Geirsdóttir, B. G. Robertsdóttir, G. Larsen, H. Norddahl, J. Eiríksson, and J. Hardardóttir, Grodurfar á Sudurlandi á sidari hluta nutima: í ljósi frjögreiningar á seti Vestra Gíslholtsvatns í Holtum, in *Abstracts: Geological Society of Iceland Spring Meeting, Reykjavík, April 3, 1996*, pp. 48–50, Geol. Soc. of Iceland, Reykjavík, 1996.
- Hammar, T., The development of the cultural landscape in the Hyndevad area, SW of Eskilstuna, southern middle Sweden, *Striae*, 24, 172–176, 1986.
- Haxeltine, A., and I. C. Prentice, BIOME3: An equilibrium terrestrial biosphere model based on ecophysiological constraints, resource availability, and competition among plant functional types, *Global Biogeochem. Cycles*, 10, 693–709, 1996.
- Hicks, S., A method of using modern pollen rain values to provide a time-scale for pollen diagrams from peat deposits, *Memo. Soc. Fauna Flora Fenn.*, 49, 21–33, 1974.
- Hicks, S., Variations in pollen frequency in a bog at Kanjerjoki, NE Finland during the Flandrian, *Comment. Biol.*, 80, 4–28, 1975a.
- Hicks, S., New evidence for the presence of prehistoric man in the Kuusamo area, *Nordia*, 1, 1–16, 1975b.
- Hicks, S., Pollen analysis and archaeology in Kuusamo, N. E. Finland, an area of marginal human interference, *Inst. Brit. Geogr. Trans., New Ser.*, 1, 361–384, 1976.
- Hicks, S., Problems and possibilities in correlating historical/archaeological and pollen-analytical evidence in a northern boreal environment: An example from Kuusamo Finland, *Fennoscandia Archaeol.*, 2, 51–84, 1985.
- Høeg, I., En pollenanalytisk undersøkelse i slidreåsen i vestre Slidre, *Viking*, 37, 137–165, 1974.
- Høeg, H. I., En pollenanalytisk undersøkelse i Storgamafeltet i Nissedal, interne rapport IR 57/80, Sur nedbørs virkning på skog og fisk (SNSF), Ås, Norway, 1980.
- Høeg, H. I., Introduksjon av jordbruk i Øst-Norge, in *Introduksjonen av jordbruk i Norden*, edited by T. Sjøvold, pp. 143–151, Universitetsforlaget, Oslo, 1982.
- Hoffert, M. I., and C. Covey, Deriving global climate sensitivity from paleoclimate reconstructions, *Nature*, 360, 573–576, 1992.
- Hopkins, D. M., J. V. Matthews Jr., C. E. Schweger, and S. B. Young, *Paleoecology of Beringia*, 489 pp., Academic, San Diego, Calif., 1982.
- Hu, F. S., L. B. Brubaker, and P. M. Anderson, A 12000 year record of vegetation change and soil development from Wien Lake, central Alaska, *Can. J. Bot.*, 71, 1133–1142, 1993.
- Hu, F. S., L. B. Brubaker, and P. M. Anderson, Postglacial vegetation and climate change in the northern Bristol Bay region, southwestern Alaska, *Quat. Res.*, 43, 382–392, 1995.
- Hu, F. S., L. B. Brubaker, and P. M. Anderson, Boreal ecosystem development in the northwestern Alaska range since 11,000 yr BP, *Quat. Res.*, 45, 188–201, 1996.
- Huntley, B., The past and present vegetation of Morrone Birkwoods and Caenlochan National Nature Reserves, Ph.D. thesis, vol. VI, 259 pp., vol. VII, 118 pp., Cambridge Univ., Cambridge, England, 1976.
- Huntley, B., and H. J. B. Birks, Past and present vegetation of Morrone Birkwoods National Nature Reserve, Scotland, I. Primary phytosociological survey, *J. Ecol.*, 67, 417–446, 1979.
- Huttunen, A., Vegetation and palaeoecology of a bog complex in southern Finland, *Aquilo, Ser. Bot.*, 28, 27–37, 1990.
- Huttunen, P., and M. Tolonen, Pollen-analytical studies of prehistoric agriculture in northern Angermanland, in *Early Norrland I: Palaeo-ecological Investigations in Northern Sweden*, edited by P. Huttunen et al., pp. 9–34, K. Vitterhets Hist. och Antikvitets Akad., Stockholm, 1972.
- Hyvärinen, H., Absolute and relative pollen diagrams from northernmost Fennoscandia, *Fennia*, 142, 1–23, 1975.
- Hyvärinen, H., Flandrian pollen deposition rates and tree-line history in northern Fennoscandia, *Boreas*, 5, 163–175, 1976.
- Hyvärinen, H., Holocene pollen stratigraphy of Baird Inlet, east-central Ellesmere Island, arctic Canada, *Boreas*, 14, 19–32, 1985a.
- Hyvärinen, H., Holocene pollen history of the Alta area, an isolated pine forest north of the general pine forest region in Fennoscandia, *Ecol. Mediter.*, 11, 69–71, 1985b.
- Hyvärinen, H., Holocene pine and birch limits near Kilpisjärvi, western Finnish Lapland: Pollen stratigraphical evidence, *Palaeoklimaforschung*, 9, 19–27, 1993.
- Igarashi, Y., M. Fukuda, K. Saijo, N. Sento, and D. Nagaoka, Holocene vegetation around alasses, northeast Siberia, in *Proceedings of the Sixth Symposium on the Joint Siberian Permafrost Studies Between Japan and Russia in 1997*, edited by S. Mori et al., pp. 83–91, Natl. Inst. for Environ. Stud., Tsukuba, Japan, 1998.
- Il'ina, N. S., E. I. Lapshina, and N. I. Lavrenko, *Rastitel'nyi pokrov Zapadnosibirskogo kontinental'nogo plato (Vegetation Cover of the Western Siberian Continental Plateau)*, 251 pp., Nauka, Moscow, 1985.
- Ivanov, V. F., *Chetvertichnye otlozheniya poberezh'ya Vostochnoi Chukotki (Quaternary Deposits of Eastern Chukcha Peninsula Coast)*, 144 pp., Northeast Interdisciplinary Res. Inst., Far East Branch, USSR Acad. of Sci., Vladivostok, 1986.
- Ivanov, V. F., A. V. Lozhkin, S. S. Kol'nichenko, A. I. Kyschtmov, V. E. Narkhinova, and V. E. Terakhova, Pozdnii pleistotsen i golotsen Chukotskogo poluostrova i Severnoi Kamchatki (Late Pleistocene and Holocene of Chukcha Peninsula and Northern Kamchatka), in *Geologiya i mineral'nye resursy severovostochnoi Azii (Geology and Mineral Sources of North-Eastern Asia)*, edited by V. I. Goucharov, pp. 33–42, Far-East. Sci. Cent., USSR Acad. of Sci., Vladivostok, 1984.
- Jevne, O. E., *Vegetasjons-, klima- og jordbrukshistorie i Beitstad, Nord-Trøndelag*, Hovedoppgave, Dep. of Bot., Univ. of Trondheim, Trondheim, Norway, 1982.
- Johansen, J., Vegetational development in the Faroes from 10 Kyr BP to the present, *Dan. Geol. Unders. Arbog*, 1981, 111–136, 1982.
- Joussaume, S., and K. E. Taylor, Status of the Paleoclimate Modeling Intercomparison Project (PMIP), in *The First International AMIP Scientific Conference, 15–19 May 1995, WMO/TD-732*, edited by W. L. Gates, pp. 425–430, World Climate Res. Programme, Monterey, Calif., 1995.
- Kaland, P. E., Ble lyngheiene skapt av fimbulvinter eller ved mennskeverk?, *Forskningssnytt*, 19, 7–14, 1974.
- Kaland, P. E., Holocene shore displacement and shorelines in Hordaland, western Norway, *Boreas*, 13, 203–242, 1984.
- Kaplan, J. O., Geophysical applications of vegetation modeling, Doctoral dissertation, Lund Univ., Lund, Sweden, 2001.
- Kaplan, J. O., et al., Climate change and Arctic ecosystems: 2. Modeling, paleodata-model comparisons, and future projections, *J. Geophys. Res.*, 108, doi:10.1029/2002JD002559, in press, 2003.
- Kaplina, T. N., and A. V. Lozhkin, Istoriya rastitel'nosti Primorskikh nizmenostei Yakutii v golotsene (Vegetation history of Primorskikh Lowlands of Yakutia during the Holocene), in *Evolutsiya prirody territorii SSSR v pozdnem pleistotsene i golotsene (Environmental Evolution of USSR Territory During Late Pleistocene and Holocene)*, edited by A. A. Velichko, I. I. Spasskaya, and N. A. Khotinskiy, pp. 207–220, Nauka, Moscow, 1982.

- Karpenko, L. V., Dinamika rastitel'nogo pokrova, torfonakopleniya i ugleroda v Tugulanskoj kotlovine (Dynamics of vegetation cover, peat accumulation, and carbon in Tugulan Basin), *Geogr. Prirodnye Resursy*, 3, 74–81, 1966.
- Kartashova, G. G., Sporovo-pyl'tsevyje spektry sovremennykh otlozhenij v bassejne reki Oly (severnoe poberezh'e Okhotskogo morya) (Palynological spectra of recent deposits in River Oly basin (northern coast of Okhotsk Sea)), in *Sporovo-pyl'tsevoi analiz v geomorfologicheskikh issledovaniyakh (Palynological Analysis in Geomorphological Studies)*, edited by S. S. Voskresenskii and M. P. Grichuk, pp. 90–105, Moscow State Univ., 1971.
- Kay, P. A., Multivariate statistical estimates of Holocene vegetation and climate change, forest-tundra transition zone, NWT, Canada, *Quat. Res.*, 11, 125–140, 1979.
- Khomutova, V. I., Geokhronologiya donnykh otlozhenij po palinologicheskim dannym (Geochronology of bottom sediments by palynological data), in *Palinologiya Onezhskogo ozera (Palynology of Onega Lake)*, pp. 45–67, Nauka, Moscow, 1976.
- Khomutova, V. I., Paleogeografiya i biostratigrafiya ozernykh otlozhenij iz lesnoj zony evropejskoj chasti SSSR po sporovo-pyl'tsevym dannym (Paleogeography and biostratigraphy of lake sediments from forest zone of European part of USSR by spore-pollen data), Doctor of Sci. thesis, 491 pp., Leningrad State Univ., St. Petersburg, 1989.
- Khomutova, V. I., and G. A. Elina, Stratigrafiya ozernykh otlozhenij po palinologicheskim dannym (Stratigraphy of lake sediments by palynological data), in *Istoriya Ladozhskogo, Onezhskogo, Pskovsko-Chudskogo ozer, Baikala i Khanki (The history of Ladoga, Onega, Pskovsko-Chudskoe, Baikal and Khanka lakes)*, edited by D. D. Kvasov, G. G. Martinson, and A. Raukas, pp. 92–96, Nauka, Moscow, 1990.
- Khomutova, V., and M. Pushenko, Evolution of lake ecosystem of Southern Ural (Russia) from palynological data, paper presented at XIVème Symposium de l'Association des Palynologues de langue française "Palynologie & changements globaux," Paris, 1995.
- Khotinskiy, N. A., *Golotsen Severnoj Evrazii*, Nauka, Moscow, 1977.
- Kimmel, K., R. Rajamäe, and M. Sakson, The Holocene development of Tondi Mire, North Estonia: Pollen, diatom and chronological studies, *PACT*, 51, 85–101, 1996.
- Kind, N. V., Pozdne- i poslednikov'e Sibiri (novye materialy po absoljutyj khronologii) (Late- and post Glacial of Siberia (new data about absolute chronology)), in *Golotsen (Holocene)*, edited by M. I. Neishtadt, pp. 195–201, Nauka, Moscow, 1969.
- Kind, N. V., *Geokhronologiya pozdnego antropogena po izotopnym dannym (Geochronology of the Late Anthropogene by Isotopic Data)*, Nauka, Moscow, 1974.
- Kjemperud, A., A shoreline displacement investigation from Frosta in Trondheimsfjorden, Nord-Trøndelag, Norway, *Nor. Geol. Tidsskr.*, 61, 1–15, 1981.
- Kjemperud, A., Late Weichselian and Holocene shoreline displacement in parts of Trøndelag, central Norway, Doctor of Sci. thesis, 191 pp., Dep. of Geol., Univ. of Oslo, 1982.
- Klimanov, V. A., and A. A. Andreev, Korrelyatsionnyj analiz sovremennykh sporovopyl'tsevnykh spektrov Yakutii (Correlation analysis of recent pollen spectra from Yakutia), *Izv. Akad. Nauk SSSR, Ser. Geogr.*, 5, 83–93, 1992.
- Koivula, L., Keski-Suomen viljelyhistoriaa sittepoilytitukimus viidestä Keski-Suomen kunnasta, Licentiat Philos. Thesis, Dep. of Bot., Univ. of Jyväskylä, Jyväskylä, Finland, 1987.
- Königsson, L. K., The Holocene history of the Great Alvar of Öland, *Acta Pytogeogr. Suecica*, 55, 1–72, 1968.
- Königsson, L. K., Vegetationsgeschichte und Kultureinflüsse in der Landschaftsentwicklung der südschwedischen Gebirge in Härjedalen, in *Disserationes Botanicae 72 (Festschrift Max Welten)*, edited by G. Lang, pp. 177–189, J. Cramer, Vaduz, 1984.
- Königsson, L. K., The Fjällnäs project: Natural and cultural components in landscape formation, *Striae*, 24, 177–186, 1986.
- Kremenetski, C. V., and N. G. Patyk-Kara, Holocene vegetation dynamics of the southeast Kola peninsula, Russia, *Holocene*, 7, 473–479, 1997.
- Kremenetski, C., T. Vaschalova, S. Goriachkin, A. Cherkinsky, and L. Sulerzhitsky, Holocene pollen stratigraphy and bog development in the western part of the Kola Peninsula, Russia, *Boreas*, 26, 91–102, 1997.
- Kremenetski, C., T. Vaschalova, and L. Sulerzhitsky, The Holocene vegetation history of the Khibiny Mountains: Implications for the post-glacial expansion of spruce and alder on the Kola Peninsula, northwestern Russia, *J. Quat. Sci.*, 14, 29–43, 1999.
- Kuprina, N. P., Stratigrafiya i istoriya osadkonakopleniya pleistosenovykh otlozhenij Tsentral'noj Kamchatki (Stratigraphy and history of sedimentation during Pleistocene in Central Kamchatka), *Tr. Geol. Inst. Akad. Nauk SSSR*, 216, 148 pp., Moscow, 1970.
- Küttel, M., Vuolep Allakasjaure: Eine pollenanalytische Studie zur Vegetationsgeschichte der Tundra in Nordschweden, in *Festschrift für Max Welten, Diss. Bot.*, vol. 72, edited by G. Lang, pp. 191–212, J. Cramer, Vaduz, 1984.
- Küttel, M., Biostratigraphische und paläoökologische Untersuchungen in SE-Småland, Schweden, *Striae*, 21, 3–34, 1985.
- Kvamme, M., Del II vegetasjonshistorie, *Arkeol. Rapp. Bergen Mus.*, 3, 89–140, 1982.
- Kvamme, M., Vegetasjonshistoriske undersøkelser i Etnefjellene 1983/1984, *Arkeol. Rapp. Bot. Inst. Univ. Bergen*, 8, 112–142, 1985.
- Lamb, H. F., Late Quaternary vegetational history of the forest-tundra ecotone in north-central Labrador, Dissertation, Univ. of Cambridge, Cambridge, England, 1982.
- Lamb, H. F., Modern pollen spectra from Labrador and their use in reconstructing Holocene vegetational history, *J. Ecol.*, 72, 37–59, 1984.
- Lamb, H. F., Palynological evidence for postglacial change in the position of tree limit in Labrador, *Ecol. Monogr.*, 55, 241–258, 1985.
- Lamb, H. F., and M. E. Edwards, The Arctic, in *Vegetation History*, edited by B. Huntley and T. Webb III, pp. 519–555, Kluwer Acad., Norwell, Mass., 1988.
- Levina, T. P., Palynologicheskaya kharakteristika otlozhenij pozdnechetvertichnoj mezhdnikovoi epokhi v doline srednei Obi (Palynological characteristics of sediments of late Quaternary interglacial epoch in the middle Ob' valley), in *Stratigrafiya i palinologiya Mezozoja i Zenozoya Sibiri (Stratigraphy and Palynology of Mesozoic and Cenozoic of Siberia)*, edited by V. S. Volkova, pp. 74–97, Nauka, Moscow, 1979.
- Levina, T. P., and V. P. Nikitin, Paleobotanicheskaya kharakteristika golotsenovogo torfyanka v raione mysy Kargin'skogo na Yenisee (Paleobotanical characteristics of Holocene peat in the Kargin'skii Cape region of the Yenisei River), in *Pleistotsen Sibiri i smezhnykh oblastei (Pleistocene of Siberia and Bordering Regions)*, edited by V. N. Saks et al., pp. 80–85, Nauka, Moscow, 1973.
- Levina, T. P., L. A. Orlova, V. A. Panychev, and E. A. Ponomareva, Radiochronometry and pollen stratigraphy of Holocene peat of Kayakskoye Zaimitschye (Barabinskaya forest-steppe), in *Regional Geochronology of Siberia and Far East*, edited by I. V. Nikolaeva, pp. 136–143, Nauka, Moscow, 1987.
- Levis, S., J. A. Foley, and D. Pollard, Large-scale vegetation feedbacks on a doubled CO<sub>2</sub> climate, *J. Clim.*, 13, 1313–1325, 2000.
- Levkovskaya, G., *Priroda i chelovek na Lubanskikh nizmenostyakh v srednem golotsene (Nature and Man in the Mid-Holocene on the Lubanas Lowland)*, 93 pp., Zinatne, Riga, 1987.
- Levkovskaya, G. M., N. V. Kind, F. S. Zavel'skii, and V. S. Forova, Absoljutyj vozrast torfyanikov iz raiona Igarki i raschlenenie golotsena Zapadnoj Sibiri (Absolute age of peatlands from the Igarka region and Holocene stratigraphy of western Siberia), *Biul. Komissii Izuch. Chev.*, 37, 94–101, 1970.
- Lichti-Federovich, S., and J. C. Ritchie, Recent pollen assemblages from the western interior of Canada, *Rev. Palaeobot. Palynol.*, 7, 297–344, 1968.
- Liljegren, R., Paleoekology och strandförskjutning i en Littorinavik vid Spjälk i mellersta Blekinge, thesis, Univ. of Lund, Lund, Sweden, 1982.
- Lillealter, J., *Vegetasjons-, klima- og jordbruks historie på Frosta, Nord-Trøndelag*, Hovedoppgave, Dep. of Botany, Univ. of Trondheim, Trondheim, Norway, 1972.
- Lozhkin, A. V., and P. M. Anderson, Pollen data on upper Quaternary lake sediments in northeast Siberia, in *Proceedings of the International Conference on Arctic Margins (Magadan, Russia, September 1994)*, pp. 71–74, Far East Branch, Russ. Acad. of Sci., Magadan, 1995.
- Lozhkin, A. V., and O. Yu. Glushkova, Boreal'nye torfa v bassoon verkhnei Kolymy (Boreal peats in Upper Kolyma basin), in *Pozdnyj pleistotsen i golotsen Beringii (Late Pleistocene and Holocene of Beringia)*, edited by M. Kh. Gagiev, pp. 55–62, Northeast Interdisciplinary Res. Inst., Far East Branch, Russ. Acad. of Sci., Magadan, 1997.
- Lozhkin, A. V., and T. P. Prokhorova, Subfossil'nye sporovo-pyl'tsevyje spektry basseina r. Bol. Kuropatoch'ya (Kolym'skaya nizmennost') (Recent pollen spectra from Bol'shaya Kuropatoch'ya River basin (Kolyma lowlands)), in *Palynologicheskie metody v paleogeografii stratigrafii (Palynological Methods in Paleogeography and Stratigraphy)*, pp. 65–70, Northeast Interdisciplinary Res. Inst., Far East Branch, Russ. Acad. of Sci., Magadan, 1982.
- Lozhkin, A. V., P. M. Anderson, W. R. Eisner, L. G. Ravako, D. M. Hopkins, L. B. Brubaker, P. A. Colinvaux, and M. C. Miller, Late Quaternary lacustrine pollen records from southwestern Beringia, *Quat. Res.*, 39, 314–324, 1993.
- Lozhkin, A. V., P. M. Anderson, and B. V. Belaya, Radiouglerodnye daty i pyl'tsevyje zony iz ozernykh otlozhenij v rayone Kolymo-Okhotskogo vodorazdela, *Dokl. Akad. Nauk Rossii*, 343, 396–399, 1995.

- Lozhkin, A. V., P. M. Anderson, W. R. Eisner, D. M. Hopkins, and L. B. Brubaker, Changes of vegetation cover of Western Alaska during the last 18000 years, in *Quaternary Climates and Vegetation of Beringia*, edited by Y. M. Bychkov, pp. 31–42, Northeast Interdisciplinary Res. Inst., Far East Branch, Russ. Acad. of Sci., Magadan, 1996.
- Lundqvist, J., Beskrivning till jordartskarta över Värmlands län, 228 pp., *Sver. Geol. Unders., Ser. Ca*, 38, 1958.
- Lundqvist, J., Beskrivning till jordartskarta över Jämtlands län, 418 pp., *Sver. Geol. Unders., Ser. Ca*, 45, 1969.
- MacDonald, G. M., Holocene vegetation history of the upper Natla River area, Northwest Territories, Canada, *Arct. Alp. Res.*, 15, 169–180, 1983.
- MacDonald, G. M., Postglacial plant migration and vegetation development in the western Canadian boreal forest, Dissertation, Univ. of Toronto, Toronto, Ont., Canada, 1984.
- MacDonald, G. M., Postglacial development of the subalpine-boreal transition forest of western Canada, *J. Ecol.*, 75, 303–320, 1987.
- MacDonald, G. M., and L. C. Cwynar, A fossil pollen based reconstruction of the Late Quaternary history of Lodgepole Pine (*Pinus contorta* ssp. *latifolia*) in the western interior of Canada, *Can. J. For. Res.*, 15, 1039–1044, 1985.
- MacDonald, G. M., and L. C. Cwynar, Post-glacial population growth rates of *Pinus contorta* ssp. *latifolia* in western Canada, *J. Ecol.*, 79, 417–429, 1991.
- MacDonald, G. M., and J. C. Ritchie, Modern pollen spectra from the western interior of Canada and the interpretation of late Quaternary vegetation development, *New Phytol.*, 103, 245–268, 1986.
- MacDonald, G. M., T. W. D. Edwards, K. A. Moser, R. Pienitz, and J. P. Smol, Rapid response of treeline vegetation and lakes to past climate warming, *Nature*, 361, 243–246, 1993.
- MacDonald, G. M., et al., Holocene treeline history and climate change across northern Eurasia, *Quat. Res.*, 53, 302–311, 2000.
- MacKay, J. R., and J. Terasmae, Pollen diagrams in the Mackenzie delta area, N. W. T., *Arctic*, 16, 228–238, 1963.
- Magnusson, E., Beskrivning till geologiska kartbladet Örebro NV, 103 pp., *Sver. Geol. Unders., Ser. Ae*, 6, 1970.
- Magnusson, E., Kvartära bildningar, in *Beskrivning till geologiska kartbladet Örebro NO*, edited by E. Magnusson and R. Gorbatshev, *Sver. Geol. Unders., Ser. Ae*, 7, 25–74, 1972a.
- Magnusson, E., Kvartära bildningar, in *Beskrivning till jordartskartan Örebro SO*, edited by E. Magnusson and P. Ludegårdh, *Sver. Geol. Unders., Ser. Ae*, 8, 32–94, 1972b.
- Manabe, S., and R. T. Wetherald, The effects of doubling CO<sub>2</sub> concentration on climate of a general circulation model, *J. Atmos. Sci.*, 32, 3–15, 1975.
- Matveev, A. V., E. A. Krutous, and V. P. Zernitskaya, Geochronology of the Holocene of the Byelorussian Polessie, *Radiocarbon*, 35, 435–439, 1993.
- McAndrews, J. H., and G. Samson, Analyse pollinique et implications archéologiques et géomorphologiques, lac de la Hutte Sauvage (Mushuau Nipi), Nouveau-Québec, *Geogr. Phys. Quat.*, 31, 177–183, 1977.
- Mikkelsen, E., and H. I. Høeg, A reconsideration of Neolithic agriculture in eastern Norway, *Norw. Archaeol. Rev.*, 12, 33–47, 1979.
- Miller, U., and A.-M. Robertsson, Biostratigraphical investigations in the Anundsjö region, Ånermanland, northern Sweden, in *Early Norrland 12: Geological Investigations in the Anundsjö Region, Northern Sweden*, edited by U. Miller, S. Modig, and A.-M. Robertsson, pp. 1–76, K. Vitterhets Hist. och Antikvitets Akad., Uppsala, Sweden, 1979.
- Moe, D., Studies in the Holocene vegetation development on Hardangervidda, southern Norway. I. The occurrence and origin of pollen of plants favoured by man's activity, *Norw. Archaeol. Rev.*, 6, 67–73, 1973.
- Moser, K. A., and G. M. MacDonald, Holocene vegetation change at tree-line north of Yellowknife, Northwest Territories, Canada, *Quat. Res.*, 34, 227–239, 1990.
- Mott, R. J., Palynological studies in Central Saskatchewan: Contemporary pollen spectra from surface samples, *Pap. Geol. Surv. Can.*, 69–32, 13 pp., 1969.
- Mott, R. J., Palynological studies in central Saskatchewan: Pollen stratigraphy from lake sediment sequences, *Pap. Geol. Surv. Can.*, 72–49, 18 pp., 1973.
- Nakao, K., J. La Perriere, and T. A. Ager, Climatic changes in interior Alaska, in *Climatic Changes in Interior Alaska*, edited by K. Nakao, pp. 16–23, Dep. of Geophys., Hokkaido Univ., Sapporo, Japan, 1980.
- Neishtadt, M. I., Golotsenovye protsessy v Zapadnoi Sibiri i svyazannye s etim problemy (Holocene processes in western Siberia and associated problems), in *Izuchenie i okruzhayushei sredy (Studying and Mastering the Environment)*, edited by M. I. Neishtadt, pp. 90–99, Inst. of Geogr., USSR Acad. of Sci., Moscow, 1976a.
- Neishtadt, M. I., Regional'nye zakonomernosti istorii fitotsenozov SSSR v golotsene po palinologicheskim dannym (Regional regularities of phytocoenoses history in the USSR during the Holocene by palynological data), in *Istoriya Biogeotsenozov SSSR v golotsene (History of Biogeocoenoses in the USSR During the Holocene)*, edited by L. G. Dinesman, pp. 79–91, Nauka, Moscow, 1976b.
- Neustadt, M. I., and E. M. Zelikson, Neue Angaben zur Stratigraphie der Torfmoore Westsibiriens, *Acta Agralia Fenn.*, 123, 27–32, 1985.
- Nichols, H., Pollen diagrams from sub-arctic Central Canada, *Science*, 155, 1665–1668, 1967a.
- Nichols, H., Central Canadian palynology and its relevance to northwestern Europe in the late Quaternary period, *Rev. Palaeobot. Palynol.*, 2, 231–243, 1967b.
- Nichols, H., The post-glacial history of vegetation and climate at Ennadai Lake, Keewatin, and Lynn Lake, Manitoba (Canada), *Eiszeitalter Gegenw.*, 18, 176–197, 1967c.
- Nichols, H., Arctic North American palaeoecology: The recent history of vegetation and climate deduced from pollen analysis, in *Arct. Alp. Environ.*, edited by J. D. Ives and R. G. Barry, pp. 637–668, Methuen, New York, 1974.
- Nichols, H., Palynological and paleoclimatic study of the late Quaternary displacements of the boreal forest-tundra ecotone in Keewatin and Mackenzie, N. W. T., Canada, *Occas. Pap.* 15, Inst. of Arct. and Alp. Res., Univ. of Colo., Boulder, 1975.
- Nilsson, T., Standard pollen diagramme und <sup>14</sup>C datierungen aus dem Ageröds mosse in mittleren schonen, *Lunds Univ. Arsskr. Andra Avd., Medicin Mat. Naturvetensk. Aemnen*, 59(7), 52 pp., 1964.
- Odgaard, B. V., Heathland history in western Jutland, Denmark, in *The Cultural Landscape-Past, Present and Future*, edited by H. H. Birks et al., pp. 311–319, Cambridge Univ. Press, New York, 1988.
- Odland, A., S. Sivertsen, O. Nordmark, A. Botnen, and B. Brunstad, Stordalsvassdragei i Etne, *Rapp.* 35, Bot. Inst., Univ. of Bergen, Bergen, Norway, 1985.
- Olsson, I. U., <sup>14</sup>C dating of samples for botanical studies of prehistoric agriculture in northern Angermanland, in *Palaeo-ecological Investigations in Northern Sweden, Early Norrland*, vol. 1, edited by P. Huttunen et al., pp. 35–41, K. Vitterhets Hist. och Antikvitets Akad., Uppsala, Sweden, 1972.
- Olsson, I. U., A discussion of the <sup>14</sup>C datings from Prästsjön, Joningsmyren and Stormyren, in *Palaeo-ecological Investigation in Coastal Västerbotten, N. Sweden, Early Norrland*, vol. 9, edited by R. Engelman et al., pp. 161–164, K. Vitterhets Hist. och Antikvitets Akad., Uppsala, Sweden, 1976.
- Olsson, I. U., A discussion of the <sup>14</sup>C ages of samples from Medelpad, Sweden, in *Archaeological and Palaeoecological Studies in Medelpad, N. Sweden, Early Norrland*, vol. 11, edited by E. Baudou et al., pp. 93–97, K. Vitterhets Hist. och Antikvitets Akad., Uppsala, Sweden, 1978.
- Olsson, I. U., and M. B. Florin, Radiocarbon dating of dy and peat in the Getsjo Area, Kolmarden, Sweden, to determine the rational limit of picea, *Boreas*, 9, 289–305, 1980.
- Øren, I. J., Plantesosillogiske og plantegeografiske studier over Ledum Palustre (L) i Sør Norge, Hovedfagsoppgave, Can. Sci. thesis, Univ. i Oslo, 1982.
- Orlova, L. A., *Golotsen Baraby: Stratigrafiya i radiouglerodnaya khronologiya (The Holocene of Baraba: Stratigraphy and Radiocarbon Chronology)*, 125 pp., Nauka, Moscow, 1990.
- Oswald, W. W., L. B. Brubaker, and P. M. Anderson, Late Quaternary vegetational history of the Howard Pass area, northwestern Alaska, *Can. J. Bot.*, 77, 570–581, 1999.
- Ovenden, L., Vegetation history of a polygonal peatland, northern Yukon, *Boreas*, 11, 209–224, 1982.
- Panova, N. K., Novye dannye po paleoekologii i istorii rastitel'nosti yuzhno-golotsen Yamala v golotsene (New data about paleoecology and vegetation history of southern Yamal during the Holocene), in *Chetvertichnyi period: Metody issledovaniya, stratigrafiya i ekologiya. VII Vsesoyuznoe soveshanie (Quaternary: Methods, Stratigraphy and Ecology, VII All-Union conference, Abstracts)*, vol. 3, edited by A. M. Miidel and A. N. Molod'kov, pp. 45–46, Akad. Nauk Estonii, Tallinn, 1990.
- Panova, N. K., Palinologicheskoe issledovanie Karas'eozerkogo torfyarnika na srednem Urale (Palynological study of Karasieozerskii peatland on middle Ural), in *Issledovanie lesov Urala: Materialy nauchnykh chteniy posvyashchennykh pamyati B. P. Kolesnikova, (Investigations of Ural Forests: Proceedings of Scientific Reports in the Memory of B. P. Kolesnikov)*, pp. 28–31, Inst. of For., Akad. Nauk SSSR, Ekaterinburg, Russia, 1997.
- Panova, N. K., and T. G. Korotkovskaya, Palinologicheskoe issledovaniya torfyarnik i ozera Peschanoe (Pollen studies of peatland and Lake Peschanoe), in *Lesoekologicheskoe i palinologicheskoe issledovaniya bolot na srednem Urale (Forest Ecology and Pollen Studies of Peatlands on Middle Ural)*, pp. 49–55, Inst. of For., Akad. Nauk SSSR, Ekaterinburg, Russia, 1990.

- Panova, N. K., V. I. Makovsky, and N. G. Yerokhin, Golotsenovaya dinamika rastitel'nosti v raione Krasnoufimskoi stepi (Holocene dynamics of vegetation in Krasnoufimskaya forest-steppe area), in *Lesoobrazovatelnyy protsess na Urale i v Zaural'e (Foresterisation Process on Ural and Zaural'e)*, pp. 80–93, Inst. of For., Akad. Nauk SSSR, Ekaterinburg, Russia, 1996.
- Parra-Vergara, I., Analyse pollinique du bassin de Sobrestany (Girona, Catalunya): Action anthropique et changements climatiques pendant l'Holocène, Diplôme, 96 pp., Sci., terre et vie, Ecole pratique des hautes Etud., Montpellier, France, 1988.
- Paus, A., Paleoökologiske undersøkelser på Frøya, Sør-Trøndelag, thesis, 234 pp., Univ. of Trondheim, Trondheim, Norway, 1982.
- Paus, A., Interpretative problems and sources of error related to pollen-analytical studies of the Holocene on the Timan ridge, western Pechora Basin, northern Russia, *Ams-Skr.*, 16, 111–126, 2000.
- Paus, A., and O. E. Jevne, Innerdalens historie belyst ved den pollenanalytiske metoden, *Rapp. Arkeol. Ser. 1987-1*, pp. 7–89, Vitenskapsmuseet, Norg. Tek.-Naturvitensk. Univ., Trondheim, Norway, 1987.
- Payette, S., Fire as a controlling process in the North American boreal forest, in *A Systems Analysis of the Global Boreal Forest*, edited by H. H. Shugart, R. Leemans, and G. B. Bonan, pp. 145–169, Cambridge Univ. Press, New York, 1992.
- Pennington, W., Modern pollen samples from west Greenland and the interpretation of pollen data from the British Late-Glacial (Late Devensian), *New Phytol.*, 84, 171–201, 1980.
- Peteet, D. M., A. A. Andreev, W. Bardeen, and F. Mistretta, Long-term Arctic peatland dynamics, vegetation and climate history of the Pur-Taz region, western Siberia, *Boreas*, 27, 115–126, 1998.
- Peterson, G. M., Recent pollen spectra and zonal vegetation in the western USSR, *Quat. Sci. Rev.*, 2, 281–321, 1983.
- Peterson, G. M., Vegetational and climatic history of the western former Soviet Union, in *Global Climates Since the Last Glacial Maximum*, edited by H. E. Wright Jr. et al., pp. 169–193, Univ. of Minn. Press, Minneapolis, 1993.
- Peyron, O., D. Jolly, R. Bonnefille, A. Vincens, and J. Guiot, Climate of east Africa 6000 <sup>14</sup>C yr B.P. as inferred from pollen data, *Quat. Res.*, 54, 90–101, 2000.
- Pirrus, R., A. U. Rõuk, and A. Liiva, Geology and stratigraphy of the reference site of lake Raigastvere in Saardjarve dreemlin, in *Palaeohydrology of the Temperate Zone. II. Lakes*, edited by A. Raunas and L. Saarse, pp. 101–122, Valgus, Tallin, Estonia, 1987.
- Pisarcic, M. F. J., The Late-Quaternary vegetation history of the Lower Lena River regions, Siberia, Master's thesis, 115 pp., McMaster Univ., Hamilton, Ont., Canada, 1996.
- Poska, A., Three pollen diagrams from coastal Estonia, *Kvartärgeol. Avd. 170*, 40 pp., Univ. of Uppsala, Uppsala, Sweden, 1994.
- Prentice, I. C., Modern pollen spectra from lake sediments in Finland and Finnmark, north Norway, *Boreas*, 7, 131–153, 1978.
- Prentice, I. C., Pollen representation, source area, and basin size: Toward a unified theory of pollen analysis, *Quat. Res.*, 23, 76–86, 1985.
- Prentice, I. C., Records of vegetation in time and space: The principles of pollen analysis, in *Vegetation History*, edited by B. J. Huntley and T. Webb III, pp. 17–42, Kluwer Acad., Norwell, Mass., 1988.
- Prentice, I. C., and T. Webb III, BIOME 6000: Reconstructing global mid-Holocene vegetation patterns from palaeoecological records, *J. Biogeogr.*, 25, 997–1005, 1998.
- Prentice, I. C., J. Guiot, B. Huntley, D. Jolly, and R. Cheddadi, Reconstructing biomes from palaeoecological data: A general method and its application to European pollen data at 0 and 6 ka, *Clim. Dyn.*, 12, 185–194, 1996.
- Prentice, I. C., D. Jolly, and BIOME 6000 Participants, Mid-Holocene and glacial-maximum vegetation geography of the northern continents and Africa, *J. Biogeogr.*, 27, 507–519, 2000.
- Ramfjord, H., Vegetasjons- og klimahistorie gjennom de siste 9000 ar i Nærøy, Nord-Trøndelag, 108 pp., Cand. Real. thesis, Dep. of Bot., Univ. of Trondheim, Trondheim, Norway, 1979.
- Ramfjord, H., On the late Weichselian and Flandrian shoreline displacement in Nærøy, Nord-Trøndelag, Norway, *Nor. Geol. Tidsskr.*, 3, 191–205, 1982.
- Rampton, V. N., Late Quaternary vegetational and climatic history of the Snag-Klutlan area, southwestern Yukon Territory, Canada, *Geol. Soc. Am. Bull.*, 82, 959–978, 1971.
- Rankama, T., and I. Vuorela, Between inland and coast in Metal Age Finland-human impact on the primeval forests of Southern Häme during the Iron Age, *Memo. Soc. Fauna Flora Fenn.*, 64, 25–34, 1988.
- Reeburgh, W. S., and M. S. Young, Univ. of Alaska radiocarbon dates 1, *Radiocarbon*, 18, 1–15, 1976.
- Renberg, I., Palaeolimnology and varve counts of the annually laminated sediment of Lake Rudejtjärn, northern Sweden, in *Archaeological and Palaeoecological Studies in Nedelpad, N. Sweden, Early Norrland*, vol. 11, edited by E. Baudou et al., pp. 63–92, K. Vitterhets Hist. och Antikvitets Akad., Uppsala, Sweden, 1978.
- Richard, P. J. H., Histoire post-wisconsinienne de la végétation du Québec méridional par l'analyse pollinique, publications et rapports divers, Serv. de la rech., Dir. Gén. des for., Minist. des Terres et For. du Québec, Québec, Québec, Canada, 1977.
- Richard, P. J. H., *Paléophytogéographie postglaciaire en Ungava par l'analyse pollinique*, Collect. Paléo-Québec, vol. 13, 154 pp., Lab. d'archéol. De l'UQAM, Montréal, Qué., Canada, 1981.
- Richard, P. J. H., Le couvert végétal du Québec-Labrador il y a 6000 ans BP: Essai, *Geogr. Phys. Quat.*, 49, 117–140, 1995.
- Ritchie, J. C., Modern pollen assemblages near the Arctic tree line, Mackenzie Delta region, Northwest Territories, *Can. J. Bot.*, 52, 381–396, 1974.
- Ritchie, J. C., The modern and late Quaternary vegetation of the Campbell-Dolomite Uplands, near Inuvik, N. W. T. Canada, *Ecol. Monogr.*, 47, 401–423, 1977.
- Ritchie, J. C., The modern and Late-Quaternary vegetation of the Doll Creek Area, North Yukon, Canada, *New Phytol.*, 90, 563–603, 1982.
- Ritchie, J. C., A Holocene pollen record of boreal forest history from the Travaillant Lake area, lower Mackenzie River Basin, *Can. J. Bot.*, 62, 1385–1392, 1984.
- Ritchie, J. C., and L. C. Cwynar, The Late Quaternary vegetation of the North Yukon, in *Palaeoecology of Beringia*, edited by D. M. Hopkins et al., pp. 113–126, Academic, San Diego, Calif., 1982.
- Ritchie, J. C., and F. K. Hare, Late-Quaternary vegetation and climate near the arctic tree line of northwestern North America, *Quat. Res.*, 1, 331–342, 1971.
- Ritchie, J. C., and S. Lichti-Federovich, Pollen dispersal phenomena in Arctic-subarctic Canada, *Rev. Palaeobot. Palynol.*, 3, 255–266, 1967.
- Robertsson, A.-M., Pollenanalytical investigation of the Leveäniemi sediments, in *The Interglacial Deposit at the Leveäniemi Mine, Svappavaara, Swedish Lapland*, edited by J. Lundqvist, *Sver. Geol. Unders.*, Ser. C, 658, 83–97, 1971.
- Rowe, J. S., D. Spittlehouse, E. Johnson, and M. Jasieniuk, Fire studies in the Upper Mackenzie Valley and adjacent Precambrian uplands, *INA Publ. QS-8045-000-EE-A1*, Dep. of Indian and North. Affairs, Ottawa, 1975.
- Rymer, M. J., and J. D. Sims, Lake-sediment evidence for the date of deglaciation of the Hidden Lake area, Kenai Peninsula, Alaska, *Geology*, 10, 314–316, 1982.
- Saarnisto, M., Holocene emergence history and stratigraphy in the area north of the Gulf of Bothnia, *Ann. Acad. Sci. Fenn.*, Ser. A, III, *Geol.-Geogr.*, 130, 1–42, 1981.
- Saarnisto, M., Long varve series in Finland, *Boreas*, 14, 133–137, 1985.
- Saarse, L., *Bottom Deposits of Small Estonian Lakes*, Inst. of Geol., Est. Acad. of Sci., Tallinn, 1994.
- Saarse, L., and L.-K. Königsson, Holocene environmental changes on the Island of Saaremaa, Estonia, *PACT*, 37, 97–131, 1992.
- Saarse, L., and A. Liiva, Geology of the Äntu group of lakes (abstract in Estonian and Russian), *Proc. Est. Acad. Sci.*, *Geol.*, 44, 119–132, 1995.
- Saarse, L., and R. Rajamäe, Holocene vegetation and climatic change on the Haanja Heights, SE Estonia, *Proc. Est. Acad. Sci.*, *Geol.*, 46, 75–92, 1997.
- Saarse, L., S. Veski, R. Rajamäe, A. Sarv, and A. Heinsalu, *Geologiya ozero Maardu (Geology of Lake Maardu)*, 32 pp., Inst. of Geol., Est. Acad. of Sci., Tallinn, 1990a.
- Saarse, L., Y. Vishnevskaya, A. Sarv, R. Rajamäe, and E. Ilves, Evolyutsiya ozer na ostrove Saaremaa (Evolution of the lakes of Saaremaa Island), *Eesti Tead. Akad. Toim. Biol.*, 39, 34–45, 1990b.
- Saarse, L., S. Veski, A. Heinsalu, R. Rajamäe, and T. Martma, Litho- and biostratigraphy of lake Päidre, south Estonia, *Proc. Est. Acad. Sci.*, *Geol.*, 44, 45–59, 1995.
- Samson, G., Prehistorie du Mushuau Nipi, Nouveau-Québec: Etude du mode d'adaptation à l'intérieur des terres hemi-arctiques, Dissertation, Univ. of Toronto, Toronto, Ont., Canada, 1983.
- Sandvik, P. U., Paleoökologisk undersökning i Nord-Trøndelag: Med hovedvekt på innvandringa og etableringa av granskogen, Hovedoppgave, Dep. of Bot., Univ. of Trondheim, Trondheim, Norway, 1986.
- Sarmaja-Korjonen, K., Y. Vasari, and C.-A. Haeggström, Taxus baccata and influence of iron Age man on the vegetation in Åland, SW Finland, *Ann. Bot. Fenn.*, 28, 143–159, 1991.
- Sarv, A., and E. Ilves, On the age of the Holocene deposits at the mouth of Emajõgi River (based on material from Saviku), *Eesti NSV Tead. Akad. Toim.*, 24, 64–69, 1975.
- Sarv, A. A., and E. Ilves, Geokronologischeskoe podrazdelenie golotsenovykh bolotno-ozernykh otlozhenii v yugo-vostochnoi Estonii (Geochronological subdivision of Holocene bog-lacustrine deposits in south-western Estonia), in *Palinologiya v kontinental'nykh i morskikh geologicheskikh*

- issledovaniyakh (Palynology in Terrestrial and Marine Geological Studies), edited by T. D. Bartos, pp. 47–59, Riga, 1976.
- Savvinova, G. M., Sporovo-pyl'tsevye spektry sovremennoi tundry severovostoka Yakutii (Palynological spectra of modern tundra on north-east Yakutia, in *Stratigrafiya, paleontologiya i litologiya osadochnykh format-siy Yakutii (Stratigraphy, Paleontology and Lithology of Sediment Formations of Yakutia)*), pp. 165–172, Yakutian Publ. House, Yakutsk, Russia, 1975a.
- Savvinova, G. M., Pyl'tsevye spektry iz razlichnykh travyanistykh soobshestv Tsentral'noi Yakutii (Pollen spectra from different herb associations of Central Yakutia), in *Palynologicheskie materialy k stratigrafii osadochnykh otlozhenii Yakutii (Palynological Data for Sediment Stratigraphy of Yakutia)*, edited by V. I. Ivanov, pp. 98–112, Yakutian Publ. House, Yakutsk, Russia, 1975b.
- Selnes, H., Paleo-økologiske undersøkelser omkring israndavsetninger på Fosenhalvøya, Midt-Norge, Hovedoppgave, Dep. of Bot., Univ. of Trondheim, Trondheim, Norway, 1982.
- Selvik, S. F., Paleøkologiske undersøkelser i Nord-Trøndelag: Med hovedvekt på granskogens innvandring og etablering, Hovedoppgave, Dep. of Bot., Univ. of Trondheim, Trondheim, Norway, 1985.
- Sergeeva, L., V. Khomutova, and I. Trifonova, Paleogeographical stages in the development of Latgale lakes, in *Palaeohydrology of the Temperate Zone*, vol. 2, *Lakes*, edited by A. Raukas and L. Saarse, pp. 154–163, Valgus, Tallinn, 1987.
- Sergeeva, L., I. Trifonova, and V. Khomutova, Sporovo-pyl'tsevoi, litologicheskii i pigmentnyi analyzy donnykh otlozhenii (Palynological, lithological and pigment analyses of bottom deposits, in *Ratsional'noe ispol'zovanie prirodnykh resursov v Belorussii (Rational Utilization of Natural Resources in Byelorussia)*), pp. 14–19, St. Petersburg, 1988.
- Shilo, N. A., A. V. Lozhkin, E. E. Titov, and Yu. V. Schumilov, *Kirgilyakhskiy mamont: Paleogeograficheskii aspekt*, Nauka, Moscow, 1983.
- Short, S. K., and H. Nichols, Holocene pollen diagrams from subarctic Labrador-Ungava: Vegetational history and climatic change, *Arct. Alp. Res.*, 9, 265–290, 1977.
- Short, S. K., S. A. Baker, J. T. Andrews, and P. J. Webber, Pollen, vegetation, and climate relationships along the Alaskan Haul Road: The basis for Holocene paleoecological and paleoclimatic studies, final report to the National Science Foundation, Division of Polar Programs, *DPP-79-26238*, Arlington, Va., 1983.
- Short, S. K., J. T. Andrews, and P. J. Webber, Pollen, vegetation, and climate relationships along the Dalton highway, Alaska, USA: A basis for Holocene paleoecological and paleoclimatic studies, *Arct. Alp. Res.*, 18, 57–72, 1986.
- Siegert, C., A. Yu. Derevyagin, G. N. Shilova, W.-D. Hermichen, and A. Hiller, Paleoclimatic indicators from permafrost sequences in the eastern Tymyr Lowland, in *Land-Ocean Systems in the Siberian Arctic: Dynamics and History*, edited by H. Kassens et al., pp. 477–499, Springer-Verlag, New York, 1999.
- Siegert, M. J., and I. Marsiat, Numerical reconstructions of LGM climate across the Eurasian Arctic, *Quat. Sci. Rev.*, 20, 1595–1605, 2001.
- Simola, H., P. Huttenen, and J. Meriläinen, Varve-dated eutrophication history of a small lake, *Verh. Int. Ver. Theor. Angew. Limnol.*, 22, 1404–1408, 1984.
- Simonsen, A., *Vertikale variasjoner i Holocen pollensedimentasjon i Ulvik, Hardanger, Ams-Varia*, vol. 8, 75 pp., Arkeol. mus. i Stavanger, Stavanger, Norway, 1980.
- Sonesson, M., Pollen zones at Abisko, Torne Lappmark, Sweden, *Bot. Not.*, 121, 491–500, 1968.
- Sonesson, M., Late Quaternary forest development of the Torneträsk area, north Sweden. 2. Pollen analytical evidence, *Oikos*, 25, 288–307, 1974.
- Sonesson, M., and T. V. Callaghan, Strategies of survival in plants of the Fennoscandian tundra, *Arctic*, 44, 95–105, 1991.
- Sønstegaard, E., and J. Mangerud, Stratigraphy and dating of Holocene gully sediments in Os, western Norway, *Norsk Geol. Tidsskr.*, 57, 313–346, 1977.
- Sorsa, P., Pollenanalytische Untersuchungen zur Spätquartären Vegetations- und Klimaentwicklung im östlichen Nord-Finland, *Ann. Bot. Fenn.*, 2, 301–413, 1965.
- Spear, R. W., Paleoecological approaches to the study of tree-line fluctuation in the MacKenzie delta region, Northwest Territories: Preliminary results, *Nordicana*, 47, 61–72, 1983.
- Spear, R. W., The palynological record of Late-Quaternary arctic tree-line in northwest Canada, *Rev. Palaeobot. Palynol.*, 79, 99–111, 1993.
- Stravers, L. K. S., Palynology and deglaciation history of the central Labrador-Ungava Peninsula, M.Sc. thesis, 171 pp., Univ. of Colo., Boulder, 1981.
- Surova, T. G., and M. M. Chernavskaya, Paleobotanicheskoe opisaniye bolota Chistik v svyazi s klimaticheskimi izmeneniyami v golotsene (Paleobotanical description of peat bog Chistik in connection with climatic change in the Holocene), in *Geophysicheskii komitet: Dannye meteorologicheskikh issledovaniy (Geophysical Committee: Data of Meteorological Studies)*, vol. 16, pp. 10–15, Akad. Nauk SSSR, Moscow, 1997.
- Szeicz, J. M., G. M. MacDonald, and A. Duk-Rodkin, Late Quaternary vegetation history of the central Mackenzie Mountains, Northwest Territories, Canada, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 113, 351–371, 1995.
- Tarasov, P. E., et al., Lake status records from the former Soviet Union and Mongolia: Database documentation, *NOAA Paleoclimatol. Publ. Ser. Rep.* 2, 86–88, Natl. Oceanic and Atmos. Admin., Boulder, Colo., 1994.
- Tarasov, P. E., et al., Present-day and mid-Holocene biomes reconstructed from pollen and plant macrofossil data from the former Soviet Union and Mongolia, *J. Biogeogr.*, 25, 1029–1053, 1998.
- Tarasov, P. E., et al., Last glacial maximum biomes reconstructed from pollen and plant macrofossil data from northern Eurasia, *J. Biogeogr.*, 27, 609–620, 2000.
- TEMPO Members, Potential role of vegetation feedback in the climate sensitivity of high-latitude regions: A case study at 6000 years before present, *Global Biogeochem. Cycles*, 10, 727–736, 1996.
- Terasmae, J., and O. L. Hughes, Late-Wisconsinan chronology and history of vegetation in the Olgivie Mountains, Yukon Territory, Canada, *Palaeobotanist*, 15, 235–242, 1966.
- Ter-Grigoryan, E. V., Sovremennyye sporovo-pyl'tsevye spektry poberezh'ya Chukotskogo poluostrova: Palinologicheskie issledovaniya na Severo-Vostoke SSSR (Recent pollen spectra from Chukcha Peninsula coast: Palynological studies on the North-East of the USSR), in *Palynologicheskie issledovaniya na Dal'nem Vostoke: Materialy II Mezhdomstvennogo seminara po palinologicheskim issledovaniyam na Dal'nem Vostoke (Palynological Studies on Far-East: Proceedings of II Interdepartmental Seminar of Palynological Studies on Far-East)*, edited by M. P. Grichuk and A. M. Korotkii, pp. 67–73, Far East Branch, Russ. Acad. of Sci., Vladivostok, 1978.
- Texier, D., N. de Noblet, S. P. Harrison, A. Haxeltine, D. Jolly, S. Joussaume, F. Laarif, I. C. Prentice, and P. Tarasov, Quantifying the role of biosphere-atmosphere feedbacks in climate change: Coupled model simulation for 6000 years BP and comparison with paleodata for northern Eurasia and northern Africa, *Clim. Dyn.*, 13, 865–882, 1997.
- Tolonen, K., Über die Entwicklung der Moore im Finnischen Nordkarelien, *Ann. Bot. Fenn.*, 4, 219–416, 1967.
- Tolonen, M., Palaeoecological reconstruction of vegetation in a prehistoric settlement area, Salo, SW Finland, *Ann. Bot. Fenn.*, 22, 101–116, 1985.
- Tolonen, K., and R. Ruuhijärvi, Standard pollen diagrams from the Salpausselkä region of southern Finland, *Ann. Bot. Fenn.*, 13, 155–196, 1976.
- Trifonova, I. S., and N. N. Davydova, Diatoms in the plankton and sediments of two lakes of different trophic type, *Hydrobiologia*, 103, 265–268, 1983.
- Vance, R. E., Pollen stratigraphy of Eaglenest Lake, Northeastern Alberta, *Can. J. Earth Sci.*, 23, 11–20, 1986.
- Vartanyan, S. L., V. E. Garutt, and A. V. Sher, Holocene dwarf mammoths from Wrangel Island in the Siberian Arctic, *Nature*, 362, 337–340, 1993.
- Vasari, Y., Studies on the vegetational history of the Kuusamo district (north east Finland) during the late quaternary period. III. Maanselänsuo, a late-glacial site in Kuusamo, *Ann. Bot. Fenn.*, 2, 219–235, 1965.
- Vas'kovskiy, A. P., Pyl'tsevye spektry na Karin Severo-Vostoke SSSR i ikh znachenie dlya rekonstruktsii chetvertichnoi rastitel'nosti (Pollen spectra on extreme north-east of the USSR and their use for reconstruction of Quaternary vegetation), in *Mater. Geol. Polezn. Iskop. Severo-Vostoka*, 11, 130–178, 1957.
- Vavrus, S. J., The response of the coupled Arctic sea ice-atmosphere system to orbital forcing and ice motion at 6 ka and 115 ka BP, *J. Clim.*, 12, 873–896, 1999.
- Vavrus, S., and S. P. Harrison, The impact of sea ice dynamics on the Arctic climate system, *Clim. Dyn.*, 20, 741–757, 2003.
- Velichko, A. A., A. A. Andreev, and V. A. Klimanov, Dinamika rastitel'nosti i klimata v tundrovoi i lesnoi zonakh Severnoi Evrazii v pozdnelednikov'e i golotsene, in *Korotkoperiodichnye i rezkie landshafno-klimaticheskie izmeneniya za poslednie 15,000 let*, edited by A. A. Velichko, E. E. Gurtovaia, and A. N. Drenova, pp. 4–60, Inst. of Geogr., Russ. Acad. of Sci., Moscow, 1994.
- Veselova, M., Prirodnye usloviya baseina ozera Kubenskoe (Natural condition of watershed area of Lake Kubenskoe), in *Ozero Kubenskoe (Lake Kubenskoe)*, vol. 1, pp. 5–15, Nauka, Moscow, 1977.
- Volkov, I. A., and S. A. Arkhipov, *Chevertichnye otlozheniya Novosibirskoi oblasti (Quaternary Deposits in the Novosibirsk Region)*, 89 pp., Inst. of Geol. and Geophys., Siberian Branch, USSR Acad. of Sci., Novosibirsk, 1978.
- Volkov, I. A., E. E. Gurtovaya, L. V. Firsov, V. A. Panychev, and L. A. Orlova, Stroeenie, vozrast i istoriya formirovaniya golotsenovogo torfyarnika u s. Gorno-Slinkina na Irtyshye (Structure, age, and history of for-



- mation Holocene peat near Gorno-Slinkino Village), in *Pleistotsen Sibiri i smezhnykh oblastei (Pleistocene of Siberia and Bordering Regions)*, edited by V. N. Saks et al., pp. 34–39, Nauka, Moscow, 1973.
- Volkova, V. S., *Chetvertichnye otlozheniya Nizhnego Irtysha i ikh biostratigraficheskie kharakteristiki (Quaternary Deposits of the Lower Irtysh River and Their Biostratigraphic Characteristics)*, 173 pp., Nauka, Moscow, 1966.
- Vorren, K.-D., Stratigraphical investigations of a palsa bog in northern Norway, *Astarte*, 5, 39–71, 1972.
- Vorren, K.-D., Late and middle Weichselian stratigraphy of Andøya, north Norway, *Boreas*, 7, 19–38, 1978.
- Vorren, K.-D., Anthropogenic influence on the natural vegetation in coastal north Norway during the Holocene: Development of farming and pastures, *Norw. Archaeol. Rev.*, 12, 1–21, 1979.
- Vorren, K.-D., Den eldste komdryking i det nordlige Norge, in *Folk og Ressurser I Nord*, edited by J. Sandnes, A. Kjelland, and I. Østerlie, pp. 11–46, Tapir, Trondheim, Norway, 1983.
- Vorren, K.-D., Vegetasjonshistorien i gamle Helgøy herred, Troms, nord-Norge; med særlig henblik på menneskets innvirkning, *Publ. 9 fra Helgøy Prosjektet*, Univ. i Tromsø, Tromsø, Norway, 1985.
- Vorren, K.-D., and E. Nilssen, Det eldstejordbruk i Nord-Norge, en palaeoekologisk oversikt, in *Introduksjonen av jordbruk i Norden*, edited by T. Sjøvold, pp. 173–208, Universitetsforlaget, Oslo, 1982.
- Vorren, K.-D., and B. Vorren, The problem of dating a palsa: Two attempts involving pollen diagrams, determination of moss subfossils, and <sup>14</sup>C-datings, *Astarte*, 8, 73–81, 1976.
- Vorren, T. O., K.-D. Vorren, T. Alm, S. Gulliksen, and G. Løvlie, The last deglaciation (20,000 to 11,000 B. P.) on Andøya, northern Norway, *Boreas*, 17, 41–77, 1988.
- Vuorela, I., The vegetational and settlement history in Sysmae, central South Finland, interpreted on the basis of two pollen diagrams, *Bull. Geol. Soc. Finland*, 53, 47–61, 1981.
- Vuorela, I., Pollenanalytiska studier, in *Finska skären: Studier i aboländsk kulturhistoria*, edited by K. Zilliacus, pp. 115–133, Konstsfundet, Helsingfors, Finland, 1990a.
- Vuorela, I., Helsingin Vanhankaupungin siitepöly- ja makrofossilitutkimukset, *Geol. Tutkimuskeskus Arktistorap. KA43/90/3*, 31 pp., Helsinki, 1990b.
- Vuorela, I., Lounais-suomen varhaismetallikautinen asutus ja viljely siitepölyanalyysin valossa, *Karhunhammas*, 13, 2–23, 1991a.
- Vuorela, I., Turvetutkimus Suomen asutushistorian selvittämisessä, *Suo*, 42, 101–108, 1991b.
- Vuorinen, J., and K. Tolonen, Flandrian pollen deposition in Lake Pappilampi, eastern Finland, *Publ. Univ. Joensuu, Ser. B2*, 3, 1–12, 1975.
- Walker, D. A., Hierarchical subdivision of Arctic tundra based on vegetation response to climate, parent material and topography, *Global Change Biol.*, 6, 19–34, 2000.
- Walker, D. A., J. G. Bockheim, F. S. Chapin, W. Eugster, F. E. Nelson, and C. L. Ping, Calcium-rich tundra, wildlife, and the “Mammoth Steppe,” *20*, 149–163, 2001.
- Wallin, J.-E., Vegetationshistorisk och naturgeografisk undersökning vid Stalon, Vilhelmina, *Acta Bothniensia Occident.*, 8, 20–32, 1986.
- Watts, W. A., Late-Glacial pollen zones in western Ireland, *Irish Geogr.*, 4, 367–376, 1963.
- Webber, P. J., V. Komarkova, D. A. Walker, and E. Werbe, Geobotanical studies along a latitudinal gradient between the Yukon River and Prudhoe Bay, Alaska, *U.S. Army CRREL Internal Rep. 585*, 366 pp., U.S. Army Corps of Eng., Hanover, N. H., 1979.
- Williams, J. W., T. Webb III, P. H. Richard, and P. Newby, Late Quaternary biomes of Canada and the eastern United States, *J. Biogeogr.*, 27, 585–607, 2000.
- Wolf, V. G., A window to the past: Macrofossil remains from an 18,000 year-old buried surface, Seward Peninsula, Alaska, Master’s thesis, Dep. of Geol. and Geophys., Univ. of Alaska, Fairbanks, 2001.
- Yelovicheva, Ya., and I. Bogdel’, *Novyi golotsenovyi razrez iz Belorussii (New Holocene Section From Byelorussia)*, Nauka i Tekhnika, Minsk, 1985.
- Yu, G., and S. P. Harrison, Lake status records from Europe: Database documentation, *NOAA Paleoclimatol. Publ. Ser. Rep. 3*, 451 pp., Natl. Oceanic and Atmos. Admin., Boulder, Colo., 1995.
- Yurkovskaya, T. K., and G. A. Elina, Krupnomasshtabnoe kartografirovaniye paleorastitel’nosti golotsena (Large scale mapping of Holocene paleovegetation), in *Geobotanisheskoe kartografirovaniye (Geobotanical Mapping)*, pp. 3–12, Nauka, Moscow, 1991.
- Yurkovskaya, T. K., G. A. Elina, and V. A. Klimanov, Rastitel’nost’ i paleogeografiya lesnykh i bolotnykh ekosistem pravoberezh’ya r. Pinega (Arkhangel’skaya oblast’) (Vegetation and paleogeography forest and bog ecosystems right side of Pinega River (Archangelsk region)), *Bot. Zh.*, 74, 1711–1722, 1989.
- Yurtsev, B. A., The Pleistocene “Tundra-Steppe” and the productivity paradox: The landscape approach, *Quat. Sci. Rev.*, 20, 165–174, 2001.
- Zernitskaya, V. P., *Paleogeografiya Belorusskogo Poles’ya v pozdnelednikov’e i golotsene (Paleogeography of Belorussian Poles’e in Late Glacial and Holocene)*, Ph.D. thesis, Byeloruss. State Univ., Minsk, 1991.
- Zernitskaya, V. P., E. A. Krutous, and V. A. Klimanov, Izuchenie bolota dlya tselei rekonstrukzii klimaticheskikh osobennosti Belorusskogo Poles’ya (Studies of mire for the purpose of reconstruction of the climatic peculiarities of the Byelorussian Poles’e), in *Voprosy prikladnoi geomorfologii (Problem of the Practical Geomorphology)*, pp. 68–73, Nauka i Tekhnika, Minsk, 1988.

P. M. Anderson, Quaternary Research Center, University of Washington, AK-60, Box 351360, Seattle, WA 98195, USA. (pata@u.washington.edu)

A. A. Andreev, Alfred-Wegener-Institut für Polar- und Meerforschung, Telegrafenberg A43, D-14473 Potsdam, Germany. (aandreev@awi-potsdam.de)

P. J. Bartlein, Department of Geography, University of Oregon, 107 Condon Hall, Eugene, OR 97403-1251, USA. (bartlein@oregon.uoregon.edu)

N. H. Bigelow, Alaska Quaternary Center, College of Science, Engineering and Mathematics, University of Alaska, Fairbanks, P. O. Box 755940, Fairbanks, AK 99775-5940, USA. (fnhb@uaf.edu)

L. B. Brubaker, College of Forest Resources, University of Washington, AR-10, Seattle, WA 98195, USA. (lbru@u.washington.edu)

T. R. Christensen and B. Smith, Department of Physical Geography and Ecosystems Analysis, Lund University, Sölvegatan 13, S-223 62 Lund, Sweden. (torben.christensen@nateko.lu.se; benjamin.smith@nateko.lu.se)

W. Cramer, Potsdam Institut für Klimafolgenforschung (PIK), Telegrafenberg A31, Postfach 60 12 03, D-14412 Potsdam, Germany. (wolfgang.cramer@pik-potsdam.de)

M. E. Edwards, Department of Geography, University of Southampton, Southampton, SO95NH, UK. (m.e.edwards@soton.ac.uk)

K. Gajewski, Department of Geography, University of Ottawa, 60 University, PR Simard Hall, Ottawa, Ontario K1N 6N5, Canada. (gajewski@aix1.uottawa.ca)

S. P. Harrison and I. C. Prentice, Max Planck Institute for Biogeochemistry, Postfach 100164, D-07701 Jena, Germany. (sharris@bgc-jena.mpg.de; cprentic@bgc-jena.mpg.de)

B. H. Holmquist, Department of Geology, Quaternary Geology, Lund University, Tomnavägen 13, S-22363 Lund, Sweden. (bjorn.holmquist@geol.lu.se)

Y. Igarashi, Earthscience Co. Ltd., N39W3 Kitaku, Sapporo 001-0039, Japan. (vzq06055@nifty.ne.jp)

J. O. Kaplan, Canadian Centre for Climate Modeling and Analysis, P. O. Box 1700 STN CSC, Victoria BC V8W 2Y2, Canada. (jed.kaplan@ec.gc.ca)

K. Kremenetskii, Institute of Geography, Russian Academy of Sciences, Staromonetny Lane 29, 109017 Moscow, Russia. (paleo@glasnet.ru)

A. V. Lozhkin, Northeast Interdisciplinary Scientific Research Institute, Far East Branch, Russian Academy of Sciences, 68500 Magadan, Russia. (lozhkin@neisri.magadan.ru)

N. V. Matveyeva and V. Y. Razzhivin, Department of Vegetation of the Far North, Komarov Botanical Institute, Prof. Popova Street 2, 197376 St. Petersburg, Russia. (nadyam@nveget.bin.ras.spb.ru; volodyar@north.bin.ras.spb.ru)

A. D. McGuire, U.S. Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska, Fairbanks, Fairbanks, AK 99775, USA.

D. F. Murray, University of Alaska Museum, Fairbanks, AK 99775-6960, USA. (ffdfm@uaf.edu)

A. Paus, Botanisk Institutt, University of Bergen, Allégaten 41, N-5007 Bergen, Norway. (aage.paus@bot.uib.no)

M. F. J. Pisarcic, Big Sky Institute, Montana State University, 106 AJM Johnson Hall, Bozeman, MT, 59717-3490, USA.

J. C. Ritchie, Pebbledash Cottage, Corfe, Taunton, Somerset TA3 7AJ, UK. (jcr@aber.ac.uk)

V. S. Volkova, Joint Institute for Geology, Geophysics and Mineralogy, Siberian Branch, Russian Academy of Sciences, 3 Universitetskii Prospekt, Novosibirsk 90, 630090 Russia. (volkova@uigm.nsc.ru)

D. A. Walker, Institute of Arctic Biology, University of Alaska, Fairbanks, Box 757000, Fairbanks, AK 99775-7000, USA. (ffdaw@uaf.edu)

V. Wolf, Alaska SAR Facility, Geophysical Institute, University of Alaska, Fairbanks, Box 757320, Fairbanks, AK 99775-7320, USA. (vwolf@asf.alaska.edu)