

Climbing plants of Porto Ferreira State Park, southeastern Brazil

Betânia da Cunha Vargas^{*1,2}; Ana Paula Caldeira Oliveira¹; Renata Giassi Udulutsch³;
Gabriel Mendes Marcusso¹; Gabriel Pavan Sabino¹; Pablo Hendrigo Alves de Melo¹;
Roberta Marotti Martelletti Grillo¹; Vitor de Andrade Kamimura¹; Marco Antonio Assis¹

¹Universidade Estadual Paulista, Instituto de Biociências de Rio Claro, Departamento de Botânica, Rio Claro, SP, Brasil

²Universidade Estadual Paulista, Instituto de Biociências de Rio Claro, Departamento de Botânica,
Laboratório de Fenologia, Rio Claro, SP, Brasil

³Universidade Estadual Paulista, Faculdade de Ciências e Letras, Departamento de Ciências Biológicas, Assis, SP, Brasil

*Corresponding author: Betânia da Cunha Vargas, e-mail: betaniadacunhavargas@gmail.com

VARGAS, B. C.; OLIVEIRA, A. P. C.; UDULUTSCH, R. G.; MARCUSO, G. M.; SABINO, G. P.; MELO, P. H. A.; GRILLO, R. M. M.; KAMIMURA, V. A.; ASSIS, M. A. **Climbing plants of Porto Ferreira State Park, southeastern Brazil**. *Biota Neotropica*. 18(2): e20170346. <http://dx.doi.org/10.1590/1676-0611-BN-2017-0346>

Abstract: A floristic survey of climbing plants was carried out in an ecotone area of seasonal semideciduous forest (SSF) and forested savanna (CER), in Porto Ferreira State Park (PFSP), Southeastern Brazil. We sampled the reproductive specimens every month during two periods, March 2010 to September 2011 and April and July 2015. The surveys were performed by the walking method, and the sampled individuals were classified by habit, climbing mechanism and dispersal mode. Overall, 109 species, belonging to 67 genera and 29 families, were recorded; 49 species occurred in both, 29 and 31 were exclusive to SSF and CER, respectively. Bignoniaceae and Malpighiaceae were the richest families, with 17 species, followed by Sapindaceae (12 species), Asteraceae and Apocynaceae (8 species each) and Fabaceae (6). The majority of climbers were lianas, twining and anemochoric species, corresponding to 70%, 47% and 66% of all samples, respectively. In this work, we added one new family and 14 species to the Cerrado's list of climbing plants from São Paulo state, and 10 species to the Brazilian seasonal semideciduous forest's list. Therefore, we contributed to the understanding of diversity of climbing plants in vegetation types poorly studied for this plant group, mainly in the Cerradão, wherein we found new records for several species.

Key words: *Cerrado; Seasonal Semideciduous Forest; ecotone; climbing plants.*

Trepadeiras do Parque Estadual de Porto Ferreira, Brasil

Resumo: O levantamento florístico das trepadeiras foi realizado em um ecótono de Floresta Estacional Semidecidual (FES) e Cerradão (CER), no Parque Estadual de Porto Ferreira, Sudeste do Brasil. Realizamos coletas mensais dos espécimes reprodutivos ao longo de dois períodos, março 2010 a setembro 2011, abril e julho 2015. Os levantamentos foram realizados por meio do método de caminhada e os indivíduos amostrados foram classificados quanto ao hábito, mecanismo de ascensão e síndrome de dispersão. No geral, foram registradas 109 espécies, pertencentes a 67 gêneros e 29 famílias. Dentre essas, 49 espécies ocorrem nos dois tipos de vegetação, sendo que 29 e 31 espécies são exclusivas de FES e CER, respectivamente. Bignoniaceae e Malpighiaceae foram as famílias mais ricas com 17 espécies, seguidas por Sapindaceae (12 espécies), Asteraceae e Apocynaceae (8 espécies cada) e Fabaceae (6). A maioria das espécies de trepadeiras são lianas, volúveis e anemocóricas, correspondendo a 70%, 47% e 66% de toda a amostra, respectivamente. Neste trabalho, acrescentamos uma nova família e 14 espécies para a lista de trepadeiras do Cerrado paulista e 10 espécies para a lista brasileira de trepadeiras em Floresta Estacional Semidecidual. Portanto, contribuimos para o conhecimento da diversidade de trepadeiras em tipos vegetacionais pouco estudados para este grupo de planta, destacando o Cerradão, no qual encontramos novos registros para um grande número de espécies.

Palavras-chave: *Cerrado, Floresta Estacional Semidecidual; ecótono; trepadeiras.*

Introduction

Climbing plants germinate and remain on the ground, requiring external mechanical sustentation provided by others plants for access to light in the forest canopy (Darwin 1867; Putz & Windsor 1987; Gerwing et al. 2006;

Lawdig & Meiners 2010; van der Heijden et al. 2013). These plants occur in many climates and vegetation types, representing up to 25% of plant diversity in some tropical forests (Gentry 1991; Engel et al. 1998; Pérez Salicrup et al. 2001). Besides climbing plants playing an important role in biodiversity maintenance, they establish competition for resources with the

host tree (phorophyte), and can change dynamic and natural regeneration in different biomes (Laurence et al. 2001; Schnitzer & Bongers 2002).

Although floristic and structural surveys of climbing plants have increased over the past decades (e.g. Durigon & Waechter 2011; Yorke et al. 2013; Vargas et al. 2014; Ibarra-Marinquez et al. 2015), climbing plants receive less attention in contrast to studies carried out with tree communities (Phillips et al. 2002; Letcher & Chazdon 2009; Carvalho et al. 2013). In Southeastern Brazil, climbing plant studies have been conducted by floristic and phytosociological surveys, mostly in semideciduous seasonal forest (e.g. Morellato & Leitão-Filho 1998; Hora & Soares 2002; Udulutsch et al. 2004; Rezende & Ranga 2005; Tibiriçá et al. 2006; Rezende et al. 2007; Santos et al. 2009; Udulutsch et al. 2010). Nevertheless, studies on vegetation types of the Cerrado are still incipient (Weiser 2002; 2007; Carvalho et al. 2013).

Porto Ferreira State Park has been the subject of numerous floristic surveys (Bertoni & Martins 1987; Bertoni et al. 2001; Colli et al. 2003; Oliveira 2012; Osaco 2012; Sabino 2013; Marcusso et al. 2016); however, studies on climbing plants have not been performed in this area. The main objective of this study was to survey the climbing flora of an ecotone area, between seasonal semideciduous forest and forested savanna (Cerradão), in Porto Ferreira State Park, São Paulo, Southeastern Brazil.

Materials and methods

1. Study site

Porto Ferreira State Park (PFSP) is located in the municipality of Porto Ferreira, in São Paulo state, Southeastern Brazil (Figure 1). The study area is 611.55 hectares, and has altitudinal variation between 540 and 608 m.a.s.l. (São Paulo 2003). PFSP is located in a geomorphological region of peripheral depression, in the central region of the watershed of Mogi-Guaçu River (São Paulo 2003).

The climate of the region is classified as Aw, according to Köppen's classification (Bertoni & Martins 1987). The annual mean temperature is 22.3 °C and the annual mean precipitation is 1.497 mm, presenting variation between the rainy (247.9 mm) and dry seasons (26.6 mm) (CEPAGRI 2016).

PFSP has distinct vegetation types according to IBGE (2012), presenting forested savanna (Cerradão) and seasonal semideciduous forest. They are mainly differentiated in the field by their physiognomy (small average height and high density in the Cerradão), and greater average height and lower density in Semideciduous forests), ecological aspects (soil types, deciduousness) and typical and exclusive species of each vegetation type (Durigan et al. 2012). Furthermore, there are ecotonal areas among these vegetation types (Osaco 2012; Sabino 2013), in which a floristic mixture and phytophysiognomic indistinguishability occurs (IBGE 2012).




Figure 1. A. Brazilian Atlantic Forest and the Cerrado and the location of São Paulo state; B. location of Porto Ferreira State Park; C. Porto Ferreira State Park and the main vegetation types studied.

1.2 Data collection

Climbing plants sampling was performed by the walking method (Filgueiras et al. 1994), every month between March 2010 and September 2011. According to Morellato et al. (1996), transitional and final periods between climatic seasons can be related to the flowering and fruiting peaks of climbing plants, especially of the families most abundant in species. In this way, aiming to improve the sample, we performed additional floristic surveys in April and July 2015. The reproductive specimens collected in the field were processed and herborized following the protocols of Fidalgo & Bononi (1984). The vouchers were incorporated in the Herbarium Rioclarense (HRCB). Taxonomic identity was determined through specialized literature, direct comparison with identified specimens in HRCB's collection and by expert taxonomists (cited in the acknowledgements). Synonyms were verified by BFG (2015), and we adopted APG IV (2016) to classify the botanical families.

Identified plants were classified into two different habits: woody (lianas – plants with a secondary growth stem) or herbaceous (plants without a secondary growth stem) climbing plants, following the protocols of Gentry (1991) and of Villagra & Romaniuc Neto (2014). Climbing mechanisms were categorized as twining, tendrill climbing or scandent (Hergaty 1991; Villagra & Romaniuc Neto 2014). Additionally, we classified climbing plants according to their dispersal modes: anemochory, autochory or zoochory (van der Pijl 1982; Barroso et al. 1999; Peres 2016). We compared the species sampled here with the list of climbing plants for seasonal semideciduous forest (Santos et al. 2009), savanna (Weiser & Martins 2014) and Brazilian flora (BFG 2015).

Results

We recorded 109 species, distributed in 67 genera and 29 families (Table 1). We found 78 and 80 species in Seasonal Semideciduous Forest and Cerradão, respectively, of which 49 (45%) occurred in both, and 29 (26.6%) and 31 (28.4%) were exclusive for the specific vegetation types. The richest families were Bignoniaceae and Malpighiaceae (17 species each), followed by Sapindaceae (12), Asteraceae and Apocynaceae (8 species each) and Fabaceae (6), corresponding to 46.7% of overall richness (Figure 2). In contrast, 11 families (38% of total families) had only one species recorded.

Among the habits, 76 species (70%) were lianas (woody climbers). The three richest families were entirely composed by woody plants (Table 1). The twining climbing mode was the most representative strategy (51 species; 47%), followed by tendrill climbing (37 species, 37%) and scandent (18 species, 16%).

Anemochory was the most representative dispersal mechanism among the climbing plants species surveyed (72 species; 66%), followed by zoochory (23 species, 21%) and autochory (14 species, 13%). Among lianas, anemochory was the main dispersal mode (63 species, 83%), followed by zoochory (8 species, 10.5%) and autochory (5 species, 6.5%), while among herbaceous species, zoochory was the most frequent dispersal mode (15 species, 45.4%), followed by autochory and anemochory (9 species each, 27.3%).

In this study, we have registered one new family and 14 species for the Cerrado list, and 10 species for the seasonal semideciduous forest list (Table 1).

Table 1. Climbing plants sampled in Porto Ferreira State Park, Porto Ferreira, Southeastern Brazil. CER: forested savanna (Cerradão); SSF: seasonal semideciduous forest. Habit: Herb: herbaceous climbing plants; Wood: woody climbing plants (liana). Climbing mode: scandent; tendrill; twining. Dispersion: Ane: anemochory; Aut: autochory; Zoo: zoochory. Voucher: APO: Ana Paula Caldeira Oliveira; HRCB: Herbário de Rio Claro; JAL: Júlio Antonio Lombardi. * First family and/or species record for FES; ● first record of the family and/or species for the Cerrado (Santos et al. 2009; Weiser & Martins 2014; Brazilian Flora Group 2015).

Family/species	Vegetation type	Habit	Climbing mode	Dispersion	Voucher
Acanthaceae					
<i>Mendoncia puberula</i> Mart.	SSF	Herb.	Twining	Zoo.	JAL 7712
Amaranthaceae					
<i>Alternanthera brasiliana</i> (L.) Kuntze	SSF*	Herb.	Scandent	Ane.	JAL 7710
<i>Chamissoa acuminata</i> Mart.	SSF*	Herb.	Scandent	Ane.	HRCB 68052
Apocynaceae					
<i>Condylocarpon isthmicum</i> (Vell.) A.DC.	CER*, SSF	Wood	Twining	Ane.	JAL 7763
<i>Forsteronia australis</i> Müll.Arg.	CER	Wood	Twining	Ane.	HRCB 67992
<i>Forsteronia pubescens</i> A.DC.	CER, SSF	Wood	Twining	Ane.	HRCB 67993
<i>Odontadenia lutea</i> (Vell.) Markgr.	CER	Wood	Twining	Ane.	HRCB 68001
<i>Oxypetalum appendiculatum</i> Mart.	CER	Herb.	Twining	Ane.	HRCB 67994
<i>Prestonia coalita</i> (Vell.) Woodson	CER, SSF	Wood	Twining	Ane.	HRCB 67995
<i>Secondatia densiflora</i> A.DC.	CER, SSF	Wood	Twining	Ane.	HRCB 67996
<i>Temnadenia violacea</i> (Vell.) Miers	CER, SSF	Wood	Twining	Ane.	HRCB 68066
Aristolochiaceae					
<i>Aristolochia labiata</i> Willd.	CER, SSF	Wood	Twining	Aut.	HRCB 67998
Asteraceae					
<i>Bidens segetum</i> Mart. ex Colla	CER, SSF	Wood	Scandent	Ane.	HRCB 67999
<i>Cyrtocymura scorpioides</i> (Lam.) H.Rob.	SSF	Herb.	Scandent	Ane.	APO 116
<i>Dasyphyllum brasiliense</i> (Spreng.) Cabrera	SSF	Wood	Scandent	Ane.	HRCB 68000
<i>Mikania cordifolia</i> (L. f.) Willd.	CER	Herb.	Twining	Ane.	APO 7
<i>Mikania laevigata</i> Sch.Bip. ex Baker	CER	Wood	Scandent	Ane.	APO 110
<i>Mikania</i> sp. 1	SSF	Wood	Twining	Ane.	APO 117
<i>Mikania</i> sp. 2	SSF	Wood	Twining	Ane.	APO 104
<i>Mikania triangularis</i> Baker	SSF	Herb.	Twining	Ane.	APO 103

Table 1. Continued...

Family/species	Vegetation type	Habit	Climbing mode	Dispersion	Voucher
Bignoniaceae					
<i>Adenocalymma bracteatum</i> (Cham.) DC.	CER, SSF	Wood	Tendrill	Ane.	JAL 7730
<i>Adenocalymma marginatum</i> (Cham.) DC.	CER*, SSF	Wood	Tendrill	Ane.	JAL 7784
<i>Amphilophium crucigerum</i> (L.) L.G.Lohmann	CER, SSF	Wood	Tendrill	Ane.	HRCB 68002
<i>Amphilophium elongatum</i> (Vahl) L.G. Lohmann	CER, SSF	Wood	Tendrill	Ane.	HRCB 68003
<i>Anemopaegma chamberlayni</i> (Sims) Bureau & K. Schum.	CER, SSF	Wood	Tendrill	Ane.	JAL 7745
<i>Bignonia campanulata</i> Cham.	CER*, SSF	Wood	Tendrill	Ane.	HRCB 68006
<i>Cuspidaria convoluta</i> (Vell.) A.H.Gentry	CER, SSF	Wood	Tendrill	Ane.	HRCB 68007
<i>Cuspidaria pulchra</i> (Cham.) L.G.Lohmann	CER, SSF	Wood	Tendrill	Ane.	JAL 7765
<i>Dolichandra unguis-cati</i> (L.) L.G. Lohmann	SSF	Wood	Tendrill	Ane.	HRCB 68009
<i>Fridericia craterophora</i> (DC.) L.G.Lohmann	CER	Wood	Tendrill	Ane.	HRCB 68005
<i>Fridericia florida</i> (DC.) L.G.Lohmann	CER	Wood	Tendrill	Ane.	HRCB 68004
<i>Fridericia formosa</i> (Bureau) L.G. Lohmann	CER, SSF	Wood	Tendrill	Ane.	HRCB 68010
<i>Fridericia speciosa</i> Mart.	CER, SSF	Wood	Tendrill	Ane.	HRCB 68061
<i>Lundia obliqua</i> Sond.	CER*, SSF	Wood	Tendrill	Ane.	APO 62
<i>Pyrostegia venusta</i> (Ker-Gawl.) Miers	CER, SSF	Wood	Tendrill	Ane.	HRCB 68060
<i>Tanaecium selloi</i> (Spreng.) L.G. Lohmann	CER, SSF	Wood	Tendrill	Ane.	APO 40
<i>Stizophyllum perforatum</i> (Cham.) Miers	CER, SSF	Wood	Tendrill	Ane.	HRCB 68012
Boraginaceae					
<i>Myriopus rubicundus</i> (Salzm. ex DC.) Luebert	CER*	Wood	Scandent	Zoo.	HRCB 68013
<i>Varronia urticifolia</i> (Cham.) J.S.Mill.	CER*	Wood	Scandent	Zoo.	HRCB 68014
Cactaceae					
<i>Pereskia aculeata</i> Mill.	SSF	Wood	Scandent	Zoo.	JAL 8087
Cannabaceae					
<i>Celtis iguanaea</i> (Jacq.) Sarg.	SSF	Wood	Scandent	Zoo.	JAL 8532
Celastraceae					
<i>Semialarium paniculatum</i> (Mart. ex Schult.) N.Hallé	SSF	Wood	Scandent	Ane.	JAL 8534
Commelinaceae					
<i>Dichorisantra hexandra</i> (Aubl.) C.B.Clarke	SSF	Herb.	Scandent	Zoo.	JAL 7671
Convolvulaceae					
<i>Ipomoea chondrosepala</i> Hallier f.	CER, SSF*	Herb.	Twining	Aut.	JAL 7743
<i>Ipomoea saopaulista</i> O'Donell	CER, SSF	Herb.	Twining	Aut.	JAL 7744
<i>Merremia macrocalyx</i> (Ruiz & Pav.) O'Donell	CER, SSF	Herb.	Twining	Aut.	HRCB 68015
<i>Merremia umbellata</i> (L.) Hallier f.	CER, SSF*	Herb.	Twining	Aut.	APO 60
Cucurbitaceae					
<i>Gurania</i> sp.	CER*, SSF*	Herb.	Tendrill	Zoo.	APO 55
<i>Psiguria ternata</i> (M.Roem.) C.Jeffrey	CER	Herb.	Tendrill	Zoo.	HRCB 68016
Cyperaceae					
<i>Scleria latifolia</i> Sw.	SSF	Herb.	Scandent/thorn	Zoo.	HRCB 68059
Dilleniaceae					
<i>Davilla elliptica</i> A.St.-Hil.	SSF*	Wood	Twining	Zoo.	HRCB 68017
<i>Davilla rugosa</i> Poir.	CER, SSF	Wood	Twining	Zoo.	APO 46
<i>Doliocarpus dentatus</i> (Aubl.) Standl.	CER, SSF	Wood	Twining	Zoo.	APO 21
Dioscoreaceae					
<i>Dioscorea dodecaneura</i> Vell.	CER, SSF	Herb.	Twining	Ane.	JAL 7736
<i>Dioscorea offersiana</i> Klotzsch ex Griseb.	CER, SSF	Herb.	Twining	Ane.	JAL 7741
Euphorbiaceae					
<i>Dalechampia pentaphylla</i> Lam.	CER*, SSF	Herb.	Twining	Aut.	JAL 7777
<i>Dalechampia stipulacea</i> Müll.Arg.	SSF	Herb.	Twining	Aut.	JAL 7709
<i>Dalechampia triphylla</i> Lam.	SSF	Herb.	Twining	Aut.	HRCB 68019
Fabaceae					
<i>Canavalia picta</i> Mart. ex Benth.	CER	Wood	Twining	Aut.	APO 92
<i>Centrosema sagittatum</i> (Humb. & Bonpl. ex Willd.) Brandegee	SSF	Herb.	Twining	Aut.	HRCB 68020
<i>Dioeclea violacea</i> Mart. ex Benth.	SSF	Wood	Twining	Aut.	HRCB 68021
<i>Rhynchosia minima</i> (L.) DC.	CER* SSF	Herb.	Twining	Aut.	JAL 7752
<i>Rhynchosia phaseoloides</i> (Sw.) DC.	SSF	Wood	Twining	Aut.	HRCB 68022
<i>Senegalia polyphylla</i> (DC.) Britton & Rose	CER	Wood	Scandent/thorn	Aut.	JAL 8531

Table 1. Continued...

Family/species	Vegetation type	Habit	Climbing mode	Dispersion	Voucher
Lygodiaceae					
<i>Lygodium volubile</i> Sw.	CER, SSF	Herb.	Twining	Ane.	HRCB 68050
Malpighiaceae					
<i>Banisteriopsis adenopoda</i> (A.Juss.) B.Gates	CER*	Wood	Twining	Ane.	APO 91
<i>Banisteriopsis argyrophylla</i> (A.Juss.) B.Gates	CER, SSF	Wood	Twining	Ane.	JAL 7755
<i>Banisteriopsis latifolia</i> (A.Juss.) B.Gates	CER	Wood	Twining	Ane.	HRCB 68025
<i>Banisteriopsis malifolia</i> (Nees & Mart.) B.Gates var. <i>malifolia</i>	CER, SSF*	Wood	Twining	Ane.	HRCB 68026
<i>Banisteriopsis</i> cf. <i>muricata</i> (Cav.) Cuatrec.	CER, SSF	Wood	Twining	Ane.	HRCB 68031
<i>Banisteriopsis oxyclada</i> (A.Juss.) B.Gates	CER, SSF	Wood	Twining	Ane.	HRCB 68027
<i>Banisteriopsis stellaris</i> (Griseb.) B.Gates	CER	Wood	Twining	Ane.	HRCB 68028
<i>Banisteriopsis variabilis</i> B.Gates	CER	Wood	Scandent	Ane.	HRCB 68029
<i>Diplopterys pubipetala</i> (A.Juss.) W.R.Anderson & C.C.Davis	CER	Wood	Twining	Ane.	APO 114
<i>Heteropterys umbellata</i> A.Juss.	SSF*	Wood	Twining	Ane.	JAL 8089
<i>Heteropterys</i> sp.	CER	Wood	Twining	Ane.	HRCB 68030
<i>Mascagnia cordifolia</i> (A.Juss.) Griseb.	CER	Wood	Twining	Ane.	APO 115
<i>Mascagnia sepium</i> (A.Juss.) Griseb.	CER, SSF	Wood	Twining	Ane.	JAL 7732
<i>Niendenzuella lucida</i> (A.Juss.) W.R.Anderson	SSF*	Wood	Twining	Ane.	HRCB 68034
<i>Niendenzuella multiglandulosa</i> (A.Juss.) W.R.Anderson	CER, SSF	Wood	Twining	Ane.	HRCB 68032
<i>Stigmaphyllon lalandianum</i> A.Juss.	SSF	Wood	Twining	Ane.	HRCB 68033
<i>Tetrapterys</i> sp.	SSF	Wood	Twining	Ane.	HRCB 68049
Menispermaceae					
<i>Cissampelos pareira</i> L.	CER	Wood	Twining	Zoo.	JAL 8536
Passifloraceae					
<i>Passiflora alata</i> Curtis	SSF*	Herb.	Tendrill	Zoo.	JAL 7767
<i>Passiflora miersii</i> Mast.	CER	Herb.	Tendrill	Zoo.	APO 95
<i>Passiflora suberosa</i> L.	CER	Herb.	Tendrill	Zoo.	APO 2
Polygalaceae					
<i>Bredemeyera floribunda</i> Willd.	CER, SSF	Wood	Scandent	Ane.	HRCB 68035
<i>Securidaca tomentosa</i> A.St.Hil. & Moq.	CER*	Wood	Scandent	Ane.	HRCB 68065
Ranunculaceae*					
<i>Clematis dioica</i> L.	CER*	Wood	Twining	Ane.	JAL 8530
Rhamnaceae					
<i>Gouania virgata</i> Reissek	CER, SSF	Wood	Tendrill	Ane.	HRCB 68037
Rubiaceae					
<i>Galianthe laxa</i> (Cham. & Schltdl.) E.L.Cabral	CER, SSF	Herb.	Scandent	Zoo.	JAL 7738
<i>Manettia cordifolia</i> Mart.	CER, SSF	Herb.	Twining	Zoo.	HRCB 68038
Sapindaceae					
<i>Cardiospermum grandiflorum</i> Sw.	SSF	Wood	Tendrill	Ane.	HRCB 68039
<i>Paullinia rhomboidea</i> Radlk.	CER, SSF	Wood	Tendrill	Ane.	APO 102
<i>Serjania</i> cf. <i>acoma</i> Radlk.	CER	Wood	Tendrill	Ane.	HRCB 68040
<i>Serjania communis</i> Cambess.	CER*	Wood	Tendrill	Ane.	APO 3
<i>Serjania fuscifolia</i> Radlk.	CER*, SSF	Wood	Tendrill	Ane.	APO 44
<i>Serjania laruooteana</i> Cambess.	CER	Wood	Tendrill	Ane.	HRCB 68041
<i>Serjania lethalis</i> A.St.-Hil.	CER	Wood	Tendrill	Ane.	APO 107
<i>Serjania meridionalis</i> Cambess.	SSF	Wood	Tendrill	Ane.	JAL 7715
<i>Serjania paradoxa</i> Radlk.	CER	Wood	Tendrill	Ane.	APO 120
<i>Serjania pinnatifolia</i> Radlk.	CER, SSF	Wood	Tendrill	Ane.	HRCB 68055
<i>Serjania reticulata</i> Cambess.	CER, SSF	Wood	Tendrill	Ane.	APO 48
<i>Urvillea laevis</i> Radlk.	CER, SSF	Wood	Tendrill	Ane.	APO 63
Smilacaceae					
<i>Smilax elastica</i> Griseb.	CER	Herb.	Tendrill	Zoo.	HRCB 68043
<i>Smilax fluminensis</i> Steud.	CER	Herb.	Tendrill	Zoo.	HRCB 68058
Vitaceae					
<i>Cissus erosa</i> Rich.	CER, SSF	Herb.	Tendrill	Zoo.	HRCB 68045
<i>Cissus tinctoria</i> Mart.	SSF	Herb.	Tendrill	Zoo.	HRCB 68046
<i>Cissus verticillata</i> (L.) Nicolson & C.E.Jarvis	SSF	Herb.	Tendrill	Zoo.	HRCB 68047



Figure 2. A. *Bidens segetum*; B. *Odontadenia lutea*; C. *Fridericia craterophora*; D. *Centrosema sagittatum*; E. *Banisteriopsis argyrophylla*; F. *Banisteriopsis adenopoda*.
By: Marcusso, G., Sabino, G. and Udulutsch, R.G.

Discussion

The present study showed elevated richness of climbing plants in Cerradão, compared to other studies carried out in Cerrado *sensu stricto* in São Paulo state, where 41 (Mantovani & Martins 1993) and 15 species (Weiser & Godoy 2001) were recorded. On the other hand, in a Cerradão in Bauru, 52 species of climbers were recorded (Weiser 2007), corroborating our results that the Cerradão is richer in climbing plants than Cerrado *sensu stricto*. Although these studies had different sample time spans, the richness recorded in the present study can be considered high by the fact that the studies carried out in Cerrado *sensu stricto* had very similar

sample times to ours (16 to 22 months, Mantovani & Martins 1993; Weiser & Godoy 2001, respectively), while the study in Cerradão considered a much larger sample time (51 months, Weiser 2007). However, the last considered a smaller area than the present study (1 versus 169 hectares, respectively). In this way, it is likely that other Cerradão areas are as rich in this vegetation type as that recorded in the present study.

Otherwise, the number of species found in Seasonal Semideciduous Forest can be considered low compared to the average for this vegetation type (e.g. Morellato & Leitão-Filho 1998; Udulutsch et al. 2004; Rezende & Ranga 2005; Tibiriçá et al. 2006; Carneiro & Vieira 2012). However, we did not sample months with the same frequency, as we collected one more

time during transitional months and those at the end of the climatic season (rainy and dry season), and these reproductive periods are considered the peak of flowering and fruiting for climbing plants (Morellato & Leitão-Filho 1996). This was also the case in the study of Udulutsch et al. (2010), so much that these surveys showed a similar number of species (74 species) to the present study (78 species). Furthermore, floristic surveys are usually carried out for different time spans and in areas with different sizes, which compromises comparisons of diversity and the relationships between flora of different sites (e.g. Forzza et al. 2014). In this way, we suggest that floristic studies are made considering equal sample frequency during the year, considering the reproductive stages of the plants. Therefore, it may be that the Seasonal Semideciduous Forest of PFSP is so rich as those in other studies which have reported high values for richness (e.g. Morellato & Leitão-Filho 1998; Udulutsch et al. 2004; Tibiriçá et al. 2006).

In general, the reasonably substantial number of climbing plant species recorded in this study is perhaps associated with the ecotonal condition of this area, where the floristic elements of both vegetation types (seasonal semideciduous forest and Cerradão) contribute to the richness. Although for climbing plants this aspect is poorly known, this has already been documented for vascular epiphytes (e.g. Kersten 2010; Bonnet et al. 2011; Marcusso & Monteiro 2016) and trees (e.g. Pinheiro & Monteiro 2008). This elevated richness and abundance of species in an ecotone is related to the meeting of adjacent areas, and thus encompasses the diversity of the community around it (Odum 1969; Kent et al. 1997; Kark 2012). This transitional area shares environmental conditions but also establishes the particular conditions of an ecotone, allowing coexistence of species from the surrounding vegetation, as well as rare and endemic species (Kark & Rensburg 2006; Kark 2012).

The majority of climbing plants species registered belong to few families, corroborating the hypothesis described by Gentry (1991) in which the majority of species (70% in the present study) are concentrated in only 13 families. Among the richest families, Bignoniaceae, Malpighiaceae and Sapindaceae stand out (e.g. Lombardi et al. 1999; Nabe-Nielsen 2001; Hora & Soares 2002; Tibiriçá et al. 2006). These families have a widespread geographical distribution, occurring among different vegetation types (Lohmann 2015; Mamede et al. 2015; Somner et al. 2015), and certain genera are predominantly (e.g. Bignoniaceae: *Fridericia*; Sapindaceae: *Serjania*; Malpighiaceae: *Banisteriopsis*) or exclusively (e.g. Bignoniaceae: *Lundia*, *Pyrostegia*, *Tanaecium*, *Stizophyllum*; Sapindaceae: *Urvillea*; Malpighiaceae: *Mascagnia*, *Niedenzuella*) composed by climbing species (Gentry 1991; BFG 2015).

We found a prevalence of lianas in the study area. Although an equal proportion of lianas and herbaceous climbing plants is expected in tropical forests (Gentry & Dodson 1987; Gentry 1991; Durigon et al. 2009), studies of climbing plants realized in semideciduous forest found 64% (Udulutsch et al. 2004) and 60% (Vargas et al. 2013) lianas, contradicting this estimation. On the other hand, herbaceous vines can represent almost 85% of the richness in subtropical and temperate forests (Durigon et al. 2014).

The pattern of climbing mechanism registered for tropical forests was corroborated in this study, wherein twining was the predominant mechanism, followed by tendril climbing (e.g. Gentry 1991; Araujo & Alves 2010; Vivek & Parthasarathy 2015). Despite some species having specialized structures to assist in colonization of the phorophyte, such as tendrils modified with adhesive pads and hooks, improving their colonization success (Gentry 1979; Lohmann 2006), this climbing mechanism has not been the most encountered in tropical forests (Gentry 1991). In this study, we recorded *Scleria latifolia* and *Senegalia polyphylla* with a twining habit. In general, these species are not included in climbing plant floristic surveys, but in PFSP these species were observed with this growth mode; perhaps they use scabrous and thorn features to ascend in the phorophyte.

In general, anemochory (72 species) was the most commonly found dispersal mode among climbing plants (lianas + herbaceous vines) (Durigon & Waechter 2011; Gallagher et al. 2011; Vargas et al. 2013), which is

associated with environments with pronounced climatic seasonality (Dewalt et al. 2010; Schnitzer & Bongers 2011). According to Morellato & Leitão-Filho (1996), the pattern of dispersion can be considered complementary in forest phenology, wherein in periods with a smaller amount of annual precipitation, the tree community totally or partially loses its leaves (Montovani & Martins 1998), and climbing plants can use wind currents to disperse their seeds (Morellato & Leitão-Filho 1998). Although, when considering only herbaceous vines, this relation can be altered by the predominance of zoochory, thus their fruits mature during the rainy season (Morellato & Leitão-Filho 1996).

Despite floristic inventories of climbing plants increasing, they are concentrated in semideciduous forest (e.g. Hora & Soares 2002; Tibiriçá et al. 2006; Rezende et al. 2007; Udulutsch et al. 2010; Vargas et al. 2013), to the detriment of those realized in the Cerrado (Weiser & Godoy 2001; Weiser 2007; Oliveira et al. 2014). However, this study included new registers of species for both vegetation types. This is because climbing plants comprise a very diverse group, so much so that they are considered key to innovation throughout evolution, increasing angiosperm diversity (Gianoli 2015; Gianoli et al. 2016). For this reason, we highlight the importance of including climbing plants in community research (Durigon et al. 2014; Schnitzer et al. 2015), and realizing floristic inventories that include or integrate climbing plants.

The present study reveals the considerable richness of this sinusia in Porto Ferreira State Park, with a similar number of species between the vegetation types. The mainly contribution is relation to diversity, recorded species, like *Scleria latifolia* and *Senegalia polyphylla*, as climbing plants. Both to vegetation types showed new records to the climbing plants checklist. Thus, we encourage the initiatives of floristic surveys with climbing plants, with similar frequency of sample, in these vegetation types, mainly in the Cerradão, in view of the importance of its floristic richness and to filling out the remaining gaps.

Acknowledgements

We would like to thank the following experts, followed by their respective linked herbaria, for help with some identifications: Maria Cândida Henrique Mamede (SP), Rosângela Simão Bianchini (SP), Lidyanne Y. Saleme Aona (UFRB), Luiza Sumiko Kinoshita (UEC), Rubens Teixeira Queiroz (UEC), Ana Maria Goulart de Azevedo Tozzi (UEC), Ana Paula Fortuna-Perez (BOTU), Marcelo Monge Egea (UEC), João Semir (UEC) and Rafael Felipe de Almeida (HUEFS); the employees of PFSP; Capes and CNPq for the fellowship grant; and São Paulo Research Foundation (FAPESP) for fellowship grant 2015/13112-7 to P.H.A. Melo. We like to thank to J.A. Lombardi contributions with data collection.

Author Contributions

Marco Antonio Assis and Renata Giassi Udulutsch: Contribution in the concept and design of the study

Betânia da Cunha Vargas, Renata Giassi Udulutsch, Gabriel Mendes Marcusso, Gabriel Pavan Sabino, Pablo Hendrigo Alves de Melo, Roberta Marotti Martelletti Grillo, Vitor de Andrade Kamimura, Marco Antonio Assis: Contribution to data collection

Betânia da Cunha Vargas, Gabriel Mendes Marcusso, Gabriel Pavan Sabino, Pablo Hendrigo Alves de Melo, Vitor de Andrade Kamimura: Contribution to data analysis and interpretation

Betânia da Cunha Vargas, Gabriel Mendes Marcusso, Gabriel Pavan Sabino, Roberta Marotti Martelletti Grillo, Vitor de Andrade Kamimura: Contribution to manuscript preparation

Betânia da Cunha Vargas, Renata Giassi Udulutsch, Gabriel Mendes Marcusso, Gabriel Pavan Sabino, Vitor de Andrade Kamimura, Marco Antônio Assis: Contribution to critical revision, adding intellectual content

Conflicts of interest

The authors declare that they have no conflict of interest related to the publication of this manuscript.

References

- ARAUJO, D. & ALVES, M. 2010. Climbing plants of a fragmented area of lowland Atlantic Forest, Igarassu, Pernambuco (Northeastern Brazil). *Phytotaxa* 8: 1-24.
- APG. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181 (1): 1-20.
- BARROS, A.A.M.; RIBAS, L.A. & ARAUJO, D.S.D. 2009. Trepadeiras do Parque Estadual da Serra da Tiririca, Rio de Janeiro, Brasil. *Rodriguesia* 60 (3): 681-694.
- BARROSO, G.M.; MORIM, M.P.; PEIXOTO, A.L. & ICHASO, C.L.F. 1999. Frutos e sementes - morfologia aplicada à sistemática de dicotiledôneas. Viçosa: Editora UFV. 444 pp.
- BERTONI, J.E.A. & MARTINS, F.R. 1987. Composição florística de uma floresta ripária na Reserva Estadual de Porto Ferreira, SP. *Acta Botanica Brasilica* 1 (1): 17-26.
- BERTONI, J.E.A.; TOLEDO FILHO, D.V.; LEITÃO FILHO, H.F.; FRANCO, G.A.D.C. & AGUIAR, O.T. 2001. Flora arbórea e arbustiva do cerrado do Parque Estadual de Porto Ferreira (SP). *Revista do Instituto Florestal* 13 (2): 169-188.
- BFG. 2015. Growing knowledge: an overview of Seed Plant diversity in Brazil. *Rodriguesia* 66 (4): 1085-1113.
- BONNET, A.; CURCIO, G.R.; LAVORANTI, O.J. & GALVÃO, F. 2011. Flora epifítica vascular em três unidades vegetacionais do Rio Tibagi, Paraná, Brasil. *Rodriguesia* 62: 491-498.
- CARNEIRO, J.S. & VIEIRA, A.O.S. 2012. Trepadeiras: florística da Estação Ecológica do Caiuá e chave de identificação vegetativa para espécies do Norte do Estado do Paraná. *Acta Scientiarum Biological Sciences* 34 (2): 217-223.
- CARVALHO, P.G.; MELIS J.V.; MORELLATO, L.P. & GROMBONE-GUARATINI, M.T. 2013. Critérios para a amostragem de lianas: comparação e estimativa da abundância e biomassa de lianas no cerrado. *Revista Árvore* 37 (6): 1037-1043.
- CEPAGRI. Clima dos municípios paulistas. Disponível em: <http://www.cpa.unicamp.br/outras-informacoes/clima_muni_457.html>. Acesso em: 31 de janeiro de 2016.
- COLLI, A.M.T.; SOUZA, S.A. & SILVA, R.T. 2003. Pteridófitas do Parque Estadual de Porto Ferreira (SP), Brasil. *Revista do Instituto Florestal* 15 (1): 29-35.
- DARWIN, C. 1867. On the movements and habits of climbing plants. *Journal of the Linnean Society, ser. Botany*, 9: 1-118.
- DEWALT, S.J.; SCHNITZER, S.A.; CHAVE, J.; BONGER, F.; BURNHAM, R.J.; CAI, Z.; CHUYONG, G.; CLARCK, D.B.; EWAANGO, C.E.N.; GERWING, J.J.; GORTAIRE, E.; HART, T.; IBARRA-MANRÍQUES, G.; ICKES, K.; KENFACK, D.; MACÍA, M.J.; MAKANA, J.; MARTÍNEZ-RAMOS, M.; MASCARO, J.; MOSES, S.; MULLER-LANDAU, H.C.; PARREN, M.P.E.; PARTHASARATHY, N.; PÉREZ-SALICRUP, D.R.; PUTZ, F.E.; ROMERO-SALTOS, H. & THOMAS, D. 2010. Annual rainfall and seasonality predict pan-tropical patterns of liana density and basal area. *Biotropica* 42 (3): 309-317.
- DURIGAN, G.; RAMOS, V.S.; IVANAUSKAS, N. M.; FRANCO, G.A.D.C. 2012. Espécies indicadoras de fitofisionomias na transição Cerrado-Mata Atlântica no Estado de São Paulo. São Paulo: Secretaria do Meio Ambiente, 145p.
- DURIGAN, J.; CANTO-DOROW, T.S. & EISINGER, S.M. 2009. Composição florística de trepadeiras ocorrentes em bordas de fragmentos de floresta estacional, Santa Maria, Rio Grande do Sul, Brasil. *Rodriguesia* 60 (2): 415-422.
- DURIGAN, J. & WAECHTER, J.L. 2011. Floristic composition and biogeographic relations of a subtropical assemblage of climbing plants. *Biodiversity Conservation* 20 (5): 1027-1044.
- DURIGAN, J., MIOTTO, S.T.S. & GIANOLI, E. 2014. Distribution and traits of climbing plants in subtropical and temperate South America. *Journal of Vegetation Science* 25: 1484-1492.
- ENGEL, V.L.; FONSECA, R.C.B. & OLIVEIRA, R.E. 1998. Ecologia de lianas e o manejo de fragmentos florestais. *Série Técnica IPEF* 12 (32): 43-64.
- FIDALGO, O. & BONONI, V.L.R. 1984. Técnicas de coleta, preservação e herborização de material botânico. Instituto de Botânica, São Paulo. 62 p.
- FILGUEIRAS, T.S.; NOGUEIRA, P.E.; BROCHADO, A.L. & GUALALL, G.F. 1994. Caminhamento: um método expedito para levantamentos florísticos qualitativos. *Cadernos de Geociências* 12: 39- 347.
- FORZZA, R.C.; PIFANO, D.S.; OLIVEIRA-FILHO, A.R.; MEIRELES, L.D.; FARIA, P.L.; SALIMENA, F.R.; MYNSSSEN, C.M. & PRADO, J. 2014. Flora vascular da Reserva Biológica da Represa do Grama, Minas Gerais, e sua relação florística com outras florestas do sudeste brasileiro. *Rodriguesia* 65 (2): 275-292.
- GALLAGHER, R. V.; LEISHMAM, M. R. & MOLES, A. T. 2011. Traits and ecological strategies of Australian tropical and temperate climbing plants. *Journal of Biogeography* 38 (5): 828-839.
- GENTRY, A. H. 1979. Additional generic mergers in Bignoniaceae. *Annals of the Missouri Botanical Garden* 66: 778-787.
- GENTRY, A.H. & DODSON, C. 1987. Contribution of Nontrees to Species Richness of a Tropical Rain Forest. *Biotropica* 19 (2): 149-156.
- GENTRY, A.H. 1991. The distribution and evolution of climbing plants, pp. 3-49, in: Putz, F.E. and Mooney, H.A. (ed.). *The biology of vines*. Cambridge, Cambridge University Press.
- GERWING, J.J.; SCHNITZER, S.A.; BURNHAM, R.J.; BONGERS, F.; CHAVE, J.; DEWALT, S.; EWANGO, C.E.N.; FOSTER, R.; KENFACK, D.; MARTÍNEZ-RAMOS, M.; PARREN, M.; PARTHASARATHY, N.; PÉREZ-SALICRUP, D.; PUTZ, F.E. & THOMAS, D.W. 2006. A Standard Protocol for Liana Censuses. *Biotropica* 38 (2): 256-261.
- GIANOLI, E. Evolutionary implications of the climbing habit in plants. 2015. In: Schnitzer, S.A.; Bongers, F.; Burnham, R.J. & Putz, F.E. *Ecology of lianas*, Wiley Blackwell, Oxford. Pp. 239-250.
- GIANOLI, E.; TORRES-DIAZ, C.; RUIZ, E., SALGADO-LUARTE, C.; MOLINA-MONTENEGRO, M.A.; SALDAÑA, A. & RIOS, R.S. 2016. Woody climbers show greater population genetic differentiation than trees: Insights into the link between ecological traits and diversification. *Evolution* 70 (12): 2736-2745.
- HERGATY, E.E. 1991. Vine-host interactions, pp. 357-375, in: Putz, F.E. and Mooney, H.A. (ed.). *The biology of vines*. Cambridge, Cambridge University Press.
- HORA, R.C. & SOARES, J.J. 2002. Estrutura fitossociológica da comunidade de lianas em uma floresta estacional na Fazenda Cachim, São Carlos, SP. *Revista Brasileira de Botânica* 25 (3): 323-329.
- IBARRA-MANRÍQUEZ, G.; RENDÓN-SANDOVAL, F.J.; CORNEJO-TENORIO, G. & CARRILLO-REYES, P. 2013. Lianas of Mexico. *Botanical Sciences* 9 (3): 365-417.
- IBGE. Instituto Brasileiro de Geografia e Estatística. 2012. Manual técnico da vegetação brasileira. Instituto Brasileiro de Geografia e Estatística. Rio de Janeiro. 276 p.
- KARK, S. & VAN RENSBURG, B.J. 2006. Ecotones: marginal or central areas of transition? *Israel Journal of Ecology and Evolution* 52 (1): 29-53.
- KARK, S. 2012. Ecotones and Ecological Gradients, pp. 147-160, in: Meyers, R.A (ed.). *Encyclopedia of Sustainability Science and Technology*. New York: Springer Science + Business Media.
- KENT, M.; GILL, W.J.; WEAVER, R.E. & ARMITAGE, R.P. 1997. Landscape and plant community boundaries in biogeography. *Progress in Physical Geography* 21 (3): 315-353.
- KERSTEN, R.A. 2010. Epifitas vasculares - histórico, participação taxonômica e aspectos relevantes, com ênfase na Mata Atlântica. *Hoehnea* 37: 9-38.
- LADWIG, L. & MEINERS, S. 2010. Spatiotemporal dynamics of lianas during 50 years of succession to temperate forest, *Ecology* 91 (3): 671-680.
- LETCHER, S.G. & CHAZDON, R.L. 2000. Lianas and self-supporting plants during tropical forest succession. *Forest Ecology and Management* 257 (10): 2150-2156.
- LOHMANN, L. 2006. Untangling the phylogeny of Neotropical lianas (Bignoniaceae). *American Journal of Botany* 93 (4): 304-318.
- LOHMANN, L.G. 2015. Bignoniaceae in Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. Disponível em: <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB112305>. Acesso em: 31 de janeiro de 2017.
- LOMBARDI, J.A.; TEMPONI, L.G. & LEITE C.A. 1999. Mortality and diameter growth of lianas in a semideciduous forest fragmente in southeastern Brazil. *Acta Botanica Brasilica* 13 (2): 159-165.

- MAMEDE, M.C.H.; SEBASTIANI, R.; ALMEIDA, R.F.; FRANCENER, A. & AMORIM, A.M.A. 2015. Malpighiaceae in Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. Disponível em: <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB155>. Acesso em: 31 de janeiro de 2017.
- MANTOVANI, W. & MARTINS, F.R. 1993. Florística do cerrado na Reserva Biológica de Moji Guaçu, estado de São Paulo. *Acta Botânica Brasilica* 7 (1): 33-60.
- MARCUSSO, G.M.; DICKFELDT, E.P.; BERTONI, J.E.A. & MONTEIRO, R. 2016. Epífitas vasculares do Parque Estadual de Porto Ferreira, São Paulo, Brasil. *Revista do Instituto Florestal* 28 (2): 119-133.
- MARCUSSO, G.M. & MONTEIRO, R. 2016. Composição florística das epífitas vasculares em duas fisionomias vegetais no município de Botucatu, estado de São Paulo, Brasil. *Rodriguésia*, 67 (3): 553-569.
- MORELLATO, P.C. & LEITÃO FILHO, H.F. 1996. Reproductive phenology of climbers in Southeastern Brazilian forest. *Biotropica* 28 (2): 180-191.
- MORELLATO, P.C. & LEITÃO FILHO, H.F. 1998. Levantamento florístico da comunidade de trepadeiras de uma floresta semidecídua no Sudeste do Brasil. *Boletim do Museu Nacional, Nova Série Botânica* 103: 1-15.
- NABE-NIELSEN, J. 2001. Diversity and distribution of lianas in a Neotropical rain forest, Yasuni National Park, Ecuador. *Journal of Tropical Ecology* 17: 1-19.
- ODUM, E.P. 1969. *Ecologia*. Interamericana, México. 412p.
- OLIVEIRA, A.P.C. 2012. Flora vascular não-arbórea do Parque Estadual de Porto Ferreira, SP, Brasil. Dissertação de Mestrado. Universidade Estadual Paulista “Júlio Mesquita Filho”, Rio Claro.
- OLIVEIRA, E.A.; MARIMON, B.S.; FELDPAUSCH, T.R.; COLLIF, G.R.; MARIMON-JUNIOR, B.H.; LLOYD, J.; LENZA, E.; MARACAHIPES, L.; OLIVEIRA-SANTOS, C. & PHILLIPS, O.L. 2014. Diversity, abundance and distribution of lianas of the Cerrado–Amazonian forest transition, Brazil. *Plant Ecology and Diversity* 7 (1-2): 231-240.
- OSACO, M. 2012. Florística e fitossociologia do estrato arbustivo de área de transição savana-floresta no Parque Estadual de Porto Ferreira. Dissertação de Mestrado. Universidade Estadual Paulista “Júlio de Mesquita Filho”, Rio Claro.
- PERES, M.K. 2016. Estratégias de dispersão de sementes no Bioma Cerrado: considerações ecológicas e filogenéticas. Tese de Doutorado, Brasília, Universidade de Brasília.
- PÉREZ-SALICRUP, D.R.; SORK, V.L. & PUTZ, F.E. 2001. Lianas and trees in a Lianas Forest of Amazonian Bolivia. *Biotropica* 33 (1): 34-47.
- PHILLIPS, O.L.; VÁSQUEZ, M.R.; ARROYO, L.; BAKER, T.; KILLEEN, T.; LEWIS, S.L.; YANDVINDER, M.; MENDONZA, A.M.; NEIL, D.; VARGAS, P.C.; ALEXIADES, M.; CERÓN, C.; DI FIORE, A.; ERWIN, T.; JARDIM, A.; PALACIOS, W.; SALDIAS, M. & VINCETI, B. 2002. Increasing dominance of large lianas in Amazonian forests. *Nature* 418: 770-774.
- PINHEIRO, M.H.O. & MONTEIRO, R. 2008. Florística de uma Floresta Estacional Semidecidual, localizada em ecótono savânico-florestal, no município de Bauru, SP, Brasil. *Acta Botanica Brasilica*, 22 (4): 1085-1094.
- PUTZ, F.E. & CHAI, P. 1987. Ecological studies of lianas in Lambir National Park, Sarawak. *Journal of Ecology* 75 (2): 523-531.
- REZENDE, A.A. & RANGA, N.T. 2005. Lianas da Estação Ecológica do Noroeste Paulista, São José do Rio Preto/Mirassol, SP, Brasil. *Acta Botânica Brasilica* 19 (2): 273-279.
- REZENDE, A.A.; RANGA, N.T. & PEREIRA, R.A.S. 2007. Lianas de uma floresta estacional semidecidual, município de Paulo de Faria, norte do estado de São Paulo, Brasil. *Revista Brasileira de Botânica* 30 (3): 451-461.
- São Paulo. 2003. Instituto Florestal. Plano de Manejo do Parque Estadual de Porto Ferreira.
- SANTOS, K.; KINOSHITA, L.S. & REZENDE, A.A. 2009. Species composition of climbers in seasonal semideciduous forest fragments of Southeastern Brazil. *Biota Neotropica* 9: 175-188.
- SABINO, G.P. 2013. Florística e fitossociologia de uma comunidade arbórea em contato savana-floresta no Parque Estadual de Porto Ferreira, Porto Ferreira, São Paulo. Trabalho de Conclusão de Curso (Graduação em Ciências Biológicas) -Universidade Estadual Paulista “Júlio Mesquita Filhos”, Rio Claro.
- SCHNITZER, S.A. & BONGERS, F. 2002. The ecology of lianas and their role in forest. *Trends in Ecology & Evolution* 17 (5): 223-230.
- SCHNITZER, S.A. & BONGERS, F. 2011. Increasing liana abundance and biomass in tropical forests: emerging patterns and putative mechanisms. *Ecology Letters* 14 (4): 397-402.
- SCHNITZER, S.A.; VAN DER HEIJDEN, G.M.F. & POWERS, J.S. 2016. Addressing the challenges of including lianas in global vegetation models. *Proceedings of the National Academy of Sciences of the United States of America* 113 (1): 5-6.
- SOMNER, G.V.; FERRUCCI, M.S.; ACEVEDO-RODRÍGUEZ, P.; PERDIZ, R.O.; COELHO, R.L.G. & MEDEIROS, H. 2015. Sapindaceae in Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. Disponível em: <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB216>. Acesso em: 31 de janeiro de 2017.
- TIBIRIÇÁ, Y.J.A.; COELHO, L.F.M. & MOURA, L.C. 2006. Florística de lianas em um fragmento de floresta estacional semidecidual, Parque Estadual de Vassununga, Santa Rita do Passa Quatro, SP, Brasil. *Acta Botanica Brasilica* 20 (2): 339-346.
- UDULUTSCH, R.G.; ASSIS, M.A. & PICCHI, D.G. 2004. Florística de trepadeiras numa floresta estacional semidecidual, Rio Claro–Araras, estado de São Paulo, Brasil. *Revista Brasileira de Botânica* 27 (1): 125-134.
- UDULUTSCH, R.G.; SOUZA, V.C.; RODRIGUES, R.R. & DIAS, P. 2010. Composição florística e chaves de identificação para as lianas da Estação Ecológica dos Caetetus, estado de São Paulo, Brasil. *Rodriguésia* 61 (4): 715-730.
- VAN DER HEIDJEN, G.M.; SCHNITZER, S.A.; POWERS, J.S. & PHILLIPS, O.L. 2013. Liana Impacts on Carbon Cycling, Storage and Sequestration in Tropical Forests. *Biotropica* 45(6): 682-692.
- VAN DER PIJL, L. 1982. *Principles of dispersal in higher plants*. 3rd ed. Springer Verlag, New York. 139p.
- VARGAS, B.C.; ARAÚJO, G.M.; SCHIAVINI, I.; ROSA, P.O. & HATTORI, E.K.O. 2013. Florística de trepadeiras em floresta semidecidual e em Mata ciliar no vale do Rio Araguari, MG. *Bioscience Journal* 29 (1): 185-197.
- VARGAS, B.C. & ARAÚJO, G.M. 2014. Florística de trepadeiras em fragmentos de florestas semidecíduais em Uberlândia, Minas Gerais, Brasil. *Rodriguésia* 65 (1): 49-59.
- VILLAGRA, B.L.P. & ROMANIUC NETO, S.R. 2010. Florística de trepadeiras no Parque Estadual das Fontes do Ipiranga, São Paulo, SP, Brasil. *Revista Brasileira de Biociências* 8 (2): 186-200.
- VILLAGRA, B.L.P. & ROMANIUC NETO, S.R. 2014. Nomenclatura das plantas de hábito trepador, pp. 1-12, in: B.L.P., Villagra, M.M.R.F., Melo, S., Romaniuc-Neto, L.M. & Barbosa, L.M. (eds). *Diversidade e conservação de trepadeiras: contribuição para a restauração de ecossistemas brasileiros*. São Paulo: Instituto de Botânica.
- VIVEK, P. & N. PARTHASARATHY. 2015. Liana community and functional trait analysis in tropical dry evergreen forest of India. *Journal of Plant Ecology* 8 (5): 501-512.
- YORKE, S.R.; SCHNITZER, S.A.; MASCARO, J.; LECHTER, S.G. & CARSON, W.P. 2013. Increasing liana abundance and basal area in a tropical forest: the contribution of long-distance clonal colonization. *Biotropica* 45 (3): 317-324.
- WEISER, V. de L. & GODOY, S.A.P. 2001. Florística em um hectare de cerrado sensu stricto na ARIE – Cerrado Pé de Gigante, Santa Rita do Passa Quatro, SP. *Acta Botanica Brasilica* 15 (2): 201-212.
- WEISER, V. de L. 2002. *Ecologia e sistemática de lianas em um hectare de cerrado sensu stricto da ARIE- Cerrado Pé-de Gigante, Santa Rita do Passa Quatro, SP*. Dissertação de Mestrado, Ribeirão Preto, Universidade de São Paulo.
- WEISER, V. de L. 2007. *Árvores, arbustos e trepadeiras do cerrado do Jardim Botânico Municipal de Bauru, SP*. Tese de Doutorado, Campinas, Universidade de Campinas.
- WEISER, V. de L. & MARTINS, F.R. 2014. Trepadeiras do cerrado paulista, pp. 58-71, in: B.L.P., Villagra, M.M.R.F., Melo, S., Romaniuc-Neto, L.M. & Barbosa, L.M. (eds). *Diversidade e conservação de trepadeiras: contribuição para a restauração de ecossistemas brasileiros*. São Paulo: Instituto de Botânica.

Received: 28/03/2017

Revised: 17/07/2017

Accepted: 24/01/2018

Published online: 12/03/2018