



Low, K., Ashraf, T., Canham, N., Clayton-Smith, J., Deshpande, C., Donaldson, A., Fisher, R., Flinter, F., Foulds, N., Fryer, A., Gibson, K., Hayes, I., Hills, A., Holder, S., Irving, M., Joss, S., Kivuva, E., Lachlan, K., Magee, A., ... DDD Study (2016). Clinical and genetic aspects of KBG syndrome. *American Journal of Medical Genetics, Part A*, 170(11), 2835-2846. <https://doi.org/10.1002/ajmg.a.37842>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1002/ajmg.a.37842](https://doi.org/10.1002/ajmg.a.37842)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at DOI: 10.1002/ajmg.a.37842. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Clinical and Genetic Aspects of KBG Syndrome

Karen Low,^{1*} Tazeen Ashraf,² Natalie Canham,³ Jill Clayton-Smith,^{4,5} Charu Deshpande,² Alan Donaldson,¹ Richard Fisher,⁶ Frances Flinter,² Nicola Foulds,⁷ Alan Fryer,⁸ Kate Gibson,⁹ Ian Hayes,¹⁰ Alison Hills,¹¹ Susan Holder,³ Melita Irving,² Shelagh Joss,¹² Emma Kivuva,¹³ Kathryn Lachlan,⁷ Alex Magee,¹⁴ Vivienne McConnell,¹⁴ Meriel McEntagart,¹⁵ Kay Metcalfe,⁴ Tara Montgomery,¹⁶ Ruth Newbury-Ecob,¹ Fiona Stewart,¹⁴ Peter Turnpenny,¹³ Julie Vogt,¹⁷ David Fitzpatrick,¹⁸ Maggie Williams,¹¹ DDD Study,¹⁹ and Sarah Smithson¹

¹University Hospitals Bristol NHS Trust/University of Bristol, Bristol, United Kingdom

²Guy's and St Thomas' NHS Trust, London, United Kingdom

³North West Thames Regional Genetics Service, Harrow, London, United Kingdom

⁴Manchester Centre For Genomic Medicine, St Mary's Hospital Manchester, United Kingdom

⁵Institute of Human Development, University of Manchester, Manchester, United Kingdom

⁶Teesside Genetics Unit, The James Cook University Hospital, Middlesbrough, United Kingdom

⁷Wessex Clinical Genetics Service, Southampton, United Kingdom

⁸Liverpool Women's NHS Trust, Liverpool, United Kingdom

⁹Genetic Health Service NZ, Christchurch Hospital, Christchurch, New Zealand

¹⁰Genetic Health Service NZ, Auckland Hospital, Auckland, New Zealand

¹¹Bristol Genetics Laboratory, North Bristol NHS Trust, Bristol, United Kingdom

¹²West of Scotland Department of Clinical Genetics, Glasgow, United Kingdom

¹³Royal Devon and Exeter Hospital, Exeter, United Kingdom

¹⁴Northern Ireland Regional Genetics Service, Belfast City Hospital, Belfast, Ireland

¹⁵South West Thames Clinical Genetics Service, St Georges Hospital, London, United Kingdom

¹⁶Northern Genetics Service, Newcastle Upon Tyne, United Kingdom

¹⁷West Midlands Regional Genetics Service, Birmingham, United Kingdom

¹⁸MRC Human Genetics Unit, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, United Kingdom

¹⁹Wellcome Trust Sanger Institute, Cambridgeshire, United Kingdom

Manuscript Received: 3 March 2016; Manuscript Accepted: 17 June 2016

KBG syndrome is characterized by short stature, distinctive facial features, and developmental/cognitive delay and is caused by mutations in *ANKRD11*, one of the ankyrin repeat-containing cofactors. We describe 32 KBG patients aged 2–47 years from 27 families ascertained via two pathways: targeted *ANKRD11* sequencing (TS) in a group who had a clinical diagnosis of KBG and whole exome sequencing (ES) in a second group in whom the diagnosis was unknown. Speech delay and learning difficulties were almost universal and variable behavioral problems frequent. Macrodontia of permanent upper central incisors was seen in 85%. Other clinical features included short stature, conductive hearing loss, recurrent middle ear infection, palatal abnormalities, and feeding difficulties. We recognized a new feature of a wide anterior fontanelle with delayed closure in 22%. The subtle facial features of KBG syndrome were recognizable in half the patients. We identified 20 *ANKRD11* mutations (18 novel: all truncating) confirmed by Sanger sequencing in

32 patients. Comparison of the two ascertainment groups demonstrated that facial/other typical features were more subtle in

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Conflicts of interest: None.

Grant sponsor: Health Innovation Challenge Fund; Grant number: HICF-1009-003; Grant sponsor: Wellcome Trust and the Department of Health; Grant sponsor: Wellcome Trust Sanger Institute; Grant number: WT098051; Grant sponsor: National Institute for Health Research.

*Correspondence to:

Karen Low, Clinical Genetics, Level B, St Michaels Hospital, Southwell Street, Bristol, BS2 8EG, UK.

E-mail: Karen.low1@nhs.net

Article first published online in Wiley Online Library (wileyonlinelibrary.com): 26 September 2016

DOI 10.1002/ajmg.a.37842

the ES group. There were no conclusive phenotype–genotype correlations. Our findings suggest that mutation of *ANKRD11* is a common Mendelian cause of developmental delay. Affected patients may not show the characteristic KBG phenotype and the diagnosis is therefore easily missed. We propose updated diagnostic criteria/clinical recommendations for KBG syndrome and suggest that inclusion of *ANKRD11* will increase the utility of gene panels designed to investigate developmental delay.

© 2016 The Authors. *American Journal of Medical Genetics Part A* Published by Wiley Periodicals, Inc.

Key words: KBG syndrome; macrodontia; *ANKRD11*

INTRODUCTION

KBG syndrome (MIM # 148050) [Herrmann et al., 1975] combines short-stature, macrodontia of the permanent central upper incisors, distinctive facial features, learning difficulties, and neuro-behavioral problems [Herrmann et al., 1975; Smithson et al., 2000; Skjei et al., 2007; Sirmaci et al., 2011; Youngs et al., 2011]. Other recognized features include seizures, cardiac abnormalities, and hearing loss [Brancati et al., 2006; Youngs et al., 2011; Ockeloen et al., 2015]. Heterozygous mutations in *ANKRD11* (ankyrin repeat domain-containing protein 11) were shown to cause KBG syndrome [Sirmaci et al., 2011]. Deletions of 16q24 including *ANKRD11* were reported to lead to a similar phenotype [Willemsen et al., 2010; Isrie et al., 2012; Sacharow et al., 2012]. In 2006, the total number of cases reported worldwide was 45 [Brancati et al., 2006], but the advent of whole exome sequencing (ES) has increased recognition, including recent reports of over 25 new KBG patients [Ockeloen et al., 2015; Walz et al., 2015].

We established single gene analysis of *ANKRD11* (TS) as a diagnostic test at Bristol Genetics Laboratory in 2014 for patients considered to have KBG syndrome. Concurrently, the Deciphering Developmental Disorders study utilized ES to identify diagnoses for children with developmental delay and additional features [Wright et al., 2015]. Here, we describe clinical and molecular genetic findings in 31 previously unreported KBG patients and one adult who was previously reported as a child [Smithson et al., 2000], ascertained through both sources. We compare the phenotypic characteristics in each of our ascertainment groups and with patients reported by other authors. We alert clinicians to the key features of KBG syndrome including a wide fontanelle, which may be the first indication that an infant is affected. Data on older patients is presented, providing insight into the long-term outcome of the condition. Recognition of this diagnosis in young adults has important implications for genetic counselling.

METHODS

Patients were ascertained either via TS or ES [Wright et al., 2015] as above. In the TS group, a Clinical Geneticist in the UK (or New Zealand in one case) had suspected the diagnosis of KBG syndrome based on clinical features and sent a DNA sample to Bristol Genetics Laboratory for *ANKRD11* analysis. Consent for genetic testing was obtained at the time of

How to Cite this Article:

Low K, Ashraf T, Canham N, Clayton-Smith J, Deshpande C, Donaldson A, Fisher R, Flinter F, Foulds N, Fryer A, Gibson K, Hayes I, Hills A, Holder S, Irving M, Joss S, Kivuva E, Lachlan K, Magee A, McConnell V, McEntagart M, Metcalfe K, Montgomery T, Newbury-Ecob R, Stewart F, Turnpenny P, Vogt J, Fitzpatrick D, Williams M, DDD Study, Smithson S. 2016. Clinical and genetic aspects of KBG syndrome.

Am J Med Genet Part A 170A:2835–2846.

sample collection. The ES group of children had been recruited to the DDD study because no clinical diagnosis had been made (20 patients from 10 of the 23 UK Clinical Genetic Centers). These children had a history of developmental delay/intellectual disability with or without additional features. DNA samples were collected as parent/child trios. Array CGH and ES were performed looking for copy number variation (CNV) and de novo variants. Variants were interrogated on the basis of link to phenotype and using standard in silico tools [Wright et al., 2015]. A DDD-approved complementary analysis project (CAP) allowed us to access the exome data and phenotypes of all subjects identified with *ANKRD11* variants through the DDD Decipher website [Firth et al., 2009]. For both groups, clinicians responsible for the patients' care were approached. Patient consent and phenotyping was initially undertaken by the referring clinician and reviewed by KL and SS. A deep phenotyping questionnaire was devised based on current literature. Broad subheadings in the questionnaire included growth, development, learning, vision and hearing, neurobehavioral, skeletal, dental, cardiac, gastrointestinal, immune, and other features. We paid particular attention to learning difficulties and behavioral phenotype which are often significant in KBG syndrome [Lo-Castro et al., 2013].

Genomic DNA was received from an external laboratory or isolated from peripheral blood leukocytes using a Gentra Puregene cell kit (QIAGEN Ltd). Coding regions of the *ANKRD11* gene (13 exons including intron/exon boundaries extending to the branch sites) were amplified in 27 fragments using a MegaMix (Micro-Zone). Primers were designed to GenBank Reference Sequence NM_013275.5 using Primer3 software [Koressaar and Remm, 2007; Untergasser et al., 2012]. The large size of exon 9 (>6 kb of coding sequence) required the design of 17 overlapping sequencing amplicons in order to cover the whole exon. The high GC content of the *ANKRD11* gene necessitated the use of a GC-RICH (Roche) PCR system. Primer sequences are available on request.

M13 tagged bidirectional sequencing was undertaken using a BigDye Terminator v3.1 Cycle Sequencing Kit and 3730 DNA Genetic Analyzer (Applied Biosystems) with Mutation Surveyor DNA Variant Analysis Software v3.97 (Softgenetics). Alamut software v2.3.1 (Interactive Biosoftware, Rouen, France) was used to predict the effect of genetic variation. The software integrates PolyPhen-2, Align GVGD and SIFT and five splice

site prediction programs: SpliceSiteFinder, MaxEntScan, Human Splice Finder, NNSPLICE, and GeneSplicer. All DDD results were confirmed by Sanger sequencing either in our laboratory or locally.

RESULTS

We studied 32 patients from 27 families in total. Thirteen patients were ascertained through our Bristol laboratory and for 12 (11 from the UK and one New Zealand) TS was undertaken. One additional patient had a microdeletion of 16p24.3 including *ANKRD11* identified by array CGH but was excluded from the study. Four patients were from one family (a father and three daughters). Twenty patients were ascertained through DDD including one family (a mother, son, and daughter).

All but two patients consented to clinical photographs. In our analysis of the clinical findings of KBG we initially considered our cohort separately. We then amalgamated our data with those previously published which enabled us to estimate the frequency of features in 100 patients (Fig. 1i and j). The wide age range in our cohort provided some insight into the developmental outcome and prognosis in KBG syndrome.

Facial Features

The facial features of our patients with KBG syndrome are shown in Figure 2a and b (ascertained through ES above and TS below). Subjective appraisal of the images confirmed the presence of characteristic facial features in some but not all patients. These included broad triangular face, short neck, wide or bushy eyebrows, often with synophrys, hypertelorism, prominent ears or dysplastic helices, bulbous nasal tip, prominent nasal bridge, long and smooth philtrum, and thin upper lip. We found that patients in the TS group were more likely to express several of these features. A few patients in the ES group could not be identified as having KBG syndrome on the basis of their facial features.

Growth

The parameters of growth in our patients (Fig. 1a–c) are in keeping with the other reported cohorts [Kim et al., 2015; Ockeloen et al., 2015] and cumulative data showed the frequency of short stature (height <3rd centile) was 40% (Fig. 1i).

Perinatal Period

Few complications occurred in pregnancy and most 20-week anomaly scans were normal apart from single instances of a single umbilical artery, an atrio-ventricular septal defect (AVSD), placenta praevia, and decreased fetal movements. All babies reached a gestation of at least 36 weeks and the vast majority were delivered at term. Only two babies were breech presentation. Birthweights were all within normal parameters (2nd–98th centiles). Nineteen percent (6/32) of babies were reported to have significant feeding difficulties after birth with some requiring nasogastric tube-feeding (Fig. 1h).

Developmental Progress/Learning Difficulties

The ages at which milestones were reached was delayed (Fig. 1d), especially speech; most patients used their first single words between 2 and 3 years of age and one patient had only a few words aged 4. All children achieved walking and the oldest to do so was aged 3 years and 2 months. In no instances did patients lose skills or regress. Four patients attended a school for children with special needs and four went to a mainstream school but within a specified special needs unit (two within an autism unit). The rest attended mainstream school usually with a statement of educational needs and received variable one to one support within the classroom. Across the literature, 99% of patients had learning difficulties (Fig. 1i), but they could be very mild.

Neurodevelopment/Behavior

Fourteen patients (43%) had seizures although the age of onset varied from infancy to mid-teens. Seizure type was heterogeneous and included absences, grand mal, myoclonic jerks, and nocturnal seizures. Some patients without a diagnosis of epilepsy had staring episodes, infantile seizures, or febrile fits which resolved. Brain MRI scans were available in 13 children; 2 showed single abnormalities including mild periventricular leukomalacia with normal ventricles and an isolated dilated left ventricle. Two scans showed moderate enlargement of the cisterna magna with normal ventricles (one also showing atrophic changes). Only four children had a movement disorder including one with Tourette-like tics. One adult had a wide-based gait with apparent foot drop and involuntary flailing of the arms on walking. Eight children (25%) had a formal diagnosis of autistic spectrum disorder (ASD) and one more had some autistic features. Five patients had a diagnosis of attention deficit hyperactivity disorder (ADHD) and three others had mild symptoms in this spectrum.

Behavioral issues occurred in all but two of the patients; however, their nature was vary variable and in some cases mild. There were recurring themes, such as poor concentration, lack of attention span, and restless movement. Some patients had obsessions and could become fixated on a favorite object or routine or disliked change with behavior deteriorating at times of stress. Anxiety and shyness were reported in several patients associated with difficulty in understanding social situations and making friendships. In contrast, some parents noted a lack of stranger-awareness and disinhibition in their children when interacting with others. Some patients were described as having a low anger threshold and a few had severely challenging behavior such as tantrums, inconsolable upset, and occasional aggressive outbursts. One patient had a diagnosis of oppositional defiant disorder. Other problems included a sensory processing disorder, rocking and flicking of elastic bands, a very unusual phenotype of echolalia-imitation of an American accent, and autistic-type behavior. Sleep disturbances were reported in six patients and the majority were resistant to melatonin. Two further patients had severe sleep disruption as babies which resolved with age. One child had sleep apnea which persisted after adenotonsillectomy.




FIG. 1. Clinical features of KBG syndrome. [a–c] Bar charts show number of patients in each centile category for weight, head circumference, and height. [d] Box and whisker plot indicates age in months at which patients were first able to sit, walk, and speak first words [dots denote statistical outliers]. [e–h] Bar charts demonstrate numbers of patients showing specific oral, skeletal, ocular anomalies, and feeding difficulties. [i–j] Tables compare the frequency of specified features in patients our cohort compared with cumulative data [ours and previously reported cases] and the frequency of features in the two ascertainment groups. In e, f, and i the bracketed numbers denotes the total number of patients in which the feature/data could be sought. [i] Cumulative data from Ockeloen et al. [2015]; Busa et al. [2015]; and Walz et al. [2015].

i	This study (32)	Ockeloen (19)	Busa (20)	Walz (11)	Cumulative(total number)
Learning difficulties	31	19	20	10	80(82) 98%
Macrodonia	23	19	14(19)	11	67(81) 82%
Behavioural issues	27	19	4		50(71) 70%
Delayed bone age	3(5)	3	4(7)	5(5)	15(36) 42%
Hand anomalies	15	18	17		50(71) 70%
Seizures	13		3	1(8)	17(60) 28%
Hearing loss	8	6	5		19(71) 27%
Postnatal short stature	9	11	11	8	39(82) 40%
Palate abnormalities	9	2	4		15(71) 21%
Feeding issues	10				
Autism	8				
Persistent/large fontanelle	7				
Cardiac anomalies	4	3	7		14(71) 20%
ADHD	4				
Breech presentation	3				
Cryptorchidism	5(16)	6(10)			11(26) 42%
Decreased fetal movements	3				
Polyhydramnios	1				
Hernia	1	1			2(51) 4%

j	ES (DDD)	Targeted
Macrodonia (or gestalt features in child with primary dentician)	14/16 88%	9/11 82%
Learning difficulties	20/20 100%	12/12 100%
Speech delay	12/15 80%	9/12 75%
Behaviour problems	17/20 85%	11/12 92%
Height below 2nd centile	5/20 25%	4/12 58%
Gestalt	2/20 10%	11/12 92%
Recurrent otitis media and/or hearing loss	7/20 35%	7/12 58%
Hand abnormality-	7/20 35%	8/12 67%
Seizures	7/20 35%	7/12 58%
Palate abnormality	4/20 20%	2/12 17%
Feeding problems	6/20 30%	5/12 42%
Large anterior fontanelle and/or delayed closure	4/20 20%	3/12 25%
Cryptorchidism	2/9 22%	3/7 42%

Fig. 1. Continued.




FIG. 2. Facial, dental and limb features in KBG syndrome. (a) Facial features of patients with KBG syndrome during infancy, childhood, and targeted testing (TS, below). (b) Evolution of the facial phenotype and profile of 3 patients with KBG syndrome during childhood and adolescence ascertained through TS [2 patients above] and TS [1 patient below]. (c) Secondary dentition of patients with KBG showing macrodontia of central incisors, dental crowding, and extra teeth in some cases. The hands and feet show subtle brachydactyly and clinodactyly of fifth fingers. [Color figure can be viewed at wileyonlinelibrary.com].

Ear, Nose, Throat, and Oral Cavity

The key dental findings in our study (Figs. 1e and 2c) included macrodontia¹ of upper incisors in most of our patients (23/27 with secondary dentition) and 82% of patients with KBG syndrome overall (Fig. 1i). Palatal abnormalities occurred in a quarter of patients (8/32); four had sub-mucous cleft palate, two had bifid uvulas, one had velopharyngeal incompetence and one had a short palate necessitating two pharyngoplasties. Two further patients had hyper-nasal speech, but neither was formally assessed by a palate team nor had a comprehensive examination under anaesthetic, which was the case for the majority of patients. We may therefore have underestimated the frequency of palatal abnormalities.

Permanent conductive hearing loss of varying degrees was reported in a quarter of patients (8/32), all of whom had a history of recurrent otitis media and/or tympanic membrane perforation treated with grommet insertion. A further seven reported recurrent middle ear infections without lasting hearing loss. One patient required tympanic membrane grafts at the age of 13 and another had sensorineural hearing loss in the context of a positive maternal family history.

Skeleton

The most consistent findings were in the distal limbs including brachydactyly especially of the 5th fingers with striking clinodactyly (Fig. 1f). Hand function was normal. The feet were also affected with brachydactyly; often with wide spaces between the 1st and 2nd toes. Interestingly seven (22%) of patients had a large anterior fontanelle with delayed closure; in one patient it still remains patent aged 4.5 years. Ockeloen et al. [2015] reported the same finding in patient 6 of their cohort. A further two patients were reported to have a very large fontanelle as a baby, but this did not persist. A skeletal survey was available in six patients and three of these revealed significant but non-specific radiological signs. The first showed a copper-beaten appearance of the skull with small anterior beaks on the vertebrae at the thoracolumbar junction. Another showed radial-head dislocation with a wide carrying angle at the elbow, short distal phalanges, and a thoracic kyphosis. The third showed shortening of the pedicles of the lumbar vertebrae, brachycephaly, and a relatively long fibula on the side examined. A fused metopic suture was reported in one patient.

Gastrointestinal System

Gastrointestinal problems were seen in several patients. Feeding difficulties were the most notable affecting eight patients. These started in the neonatal period and resolved in some cases (Fig. 1h) but continued in three cases. One girl, who failed to thrive required nasogastric feeding until the age of 4 years and her weight remained below the 0.4th centile at 13 years. One boy was unable to feed orally at the age of 4.5 years and is dependent on a gastrostomy tube

¹It is acceptable to assess macrodontia subjectively for apparently increased maximum width of the tooth; or objectively by measuring mesiodistal tooth diameter (width) more than 2 SD above mean for age. [Carey et al., 2009]. In our cohort macrodontia was assessed subjectively by the referring clinician and then again on review of the images.

(he also has a sensory processing disorder and autism). Gastro-oesophageal reflux and severe constipation were also observed. One adult has a life-long history of diarrhea and an abnormal lack of thirst which has resulted in two hospital admissions for dehydration. One requires a patch for continual drooling.

Other Aspects of Phenotype

Half the patients (17/32) had eye abnormalities, most commonly strabismus and refractive errors (Fig. 1g). Congenital heart abnormalities occurred in 4/32 (13%), including a patent ductus arteriosus, an AVSD and two patients had small VSDs. One further patient had a structurally normal heart but second degree heart block on ECG (aged 6).

Many patients displayed mild hypertrichosis with thick hair, full eyebrows with lateral interruptions and mild synophrys (Fig. 2a and b). Skin abnormalities were found in several patients. One had a speckled hyperpigmented patch over his inguinal region. He also had two hair whorls on left and right sides of the occiput and a low V-shaped posterior hairline. One patient had a large hyperpigmented patch on his left arm and another had an asymmetric hair line, a tendency to skin bruising, and delayed wound healing. One had keloid scarring with rapid-growing thick hair and nails. Another had generalised ichthyosis and predominantly lower limb lymphoedema. Cutis marmorata was described in one patient. Dystrophic nails, particularly on the toes, occurred in 6/32 patients. Recurrent infections (largely respiratory) were described in 6/32 patients although immune function studies did not detect any specific deficit. Five patients had surgery for cryptorchidism and one also had an inguinal hernia repair.

Adulthood

Eight of the cohort were aged 18 or over including a parent identified after his daughter was diagnosed. He reported slow educational progress at primary school and did not achieve qualifications but was living independently as a father of three with no external help. Similarly, a mother diagnosed after KBG syndrome was identified in her son through DDD, described mild learning difficulties but was living independently, working in a kitchen and looking after her family without support. Her son achieved a few GCSE-level qualifications and was living in his own accommodation with some help from social services. The other five young adults were all living with their parents who considered it was unlikely they would be able to live alone, although they had many skills such as being able to operate computers. Two were embarking on college courses at the time of the study.

Genotype–Phenotype Correlations

Within our cohort, we identified 20 different *ANKRD11* mutations in 32 patients with KBG syndrome (Supplementary Table SI). All mutations reside within exon 9, consistent with it containing >80% of the coding sequence and are predicted to cause truncation of the ANKRD11 protein concurring with the mechanism of pathogenicity previously described [Walz et al., 2015]. Of the 20 mutations we identified, 18 were previously unreported. Previously

TABLE I. Recurrent ANKRD11 Mutations

Recurrent ANKRD11 mutations c.1903_1907del, p.(Lys635Glnfs*26)^a

Patient	19 (9y 6m) M	26 (9y 9m) F	4 (13y 3m) M	5 (21y) F*	6 (19y) F*	7 (12y) F*	8 (47y) M*	33 (3y 3m) M	Ockeloen et al. [2015] pt 6 (38y) F	Ockeloen et al. [2015] pt 10 (14y) M	Walz et al. [2015] pt B (13y) M
Macrodontia/other dental abnormalities	+	+	+	+	+	+	+	+	+	+	+
Short stature	+	+	+	+	+	+	+	+	+	+	+
Learning difficulties	+	+	+	+	+	+	+	+	+	+	+
Skeletal abnormalities	+	+	+	+	+	+	+	+	+	+	+
Delayed bone age	+	+	+	+	+	+	+	+	+	+	+
Hand anomalies	+	+	+	+	+	+	+	+	+	+	+
Ocular	+	+	+	+	+	+	+	+	+	+	+
Hearing Loss	+	+	+	+	+	+	+	+	+	+	+
Neurological	+	+	+	+	+	+	+	+	+	+	+
Behavioural abnormalities	+	+	+	+	+	+	+	+	+	+	+
Palate	+	+	+	+	+	+	+	+	+	+	+
Heart	+	+	+	+	+	+	+	+	+	+	+
Other	+	+	+	+	+	+	+	+	+	+	+

Recurrent ANKRD11 mutations c.2408_2412 del, p.(Lys803Argfs*5)^b

Patient	1 (6y 4m) F	18 (6y 10m) M	20 (13y 3m) F	13 (17y) F	27 (23y 1m) M
Macrodontia/other dental abnormalities	+	+	+	+	+
Short stature	+	+	+	+	+
Learning difficulties	+	+	+	+	+
Skeletal abnormalities	+	+	+	+	+
Delayed bone age	+	+	+	+	+
Hand anomalies	+	+	+	+	+
Ocular	+	+	+	+	+
Hearing Loss	+	+	+	+	+
Neurological	+	+	+	+	+

TABLE 1. (Continued)
Recurrent ANKRD11 mutations c.2408_2412 del, p.(Lys603Argfs 5)^b

Patient	1 (6y 4m) F DDD Artistic like	18 (6y 10m) M DDD Sensory processing disorder, "challenging"	20 (13y 3m) F DDD Artistic like; obsessions	13 (17y) F DDD	27 (23y 1m) M Challenging; no social awareness, outbursts. Stress/anxiety
Behavioural abnormalities					
Palate	—	—	—	—	—
Heart	—	—	—	—	—
Other	DCF; ant. placed anus	Fused metopic suture, crytochidism	Small apical VSD Severe feeding issues; infections, lacrimal duct surgery; pain insensitivity.	Puffy feet as neonate	Hypermobile elbows. Keloid scarring.

SEN, statement of education need; PVL, periventricular leukomalacia on MRI; ant, anterior; AF, anterior fontanelle; dn, de novo; DCF, delayed closure of fontanelle age >3y; EH, enamel hypoplasia; AF, anterior fontanelle; TM, tympanic membrane; SNHL(mat), sensorineural hearing loss (maternal history); SPC, single palmar crease.
 Skeletal abnormalities: costovertebral abnormalities; postnatal short stature, <2nd centile. Hand abnormalities to include—brachydactyly, 5th finger clinodactyly, Ocular abnormalities = strabismus, hypermetropia, astigmatism; myopia. Palate to include: cleft lip and/or palate; submucous cleft, velopharyngeal insufficiency; bifid uvula.
 *Patient 5–8 members of one family. **Patient 29–31 members of one family.
^aComparison of phenotypic findings in seven patients with recurrent c.1903_1907del (5–7 are daughters of eight) with those reported with the same mutation [Ockeloen et al., 2015; Walz et al., 2015].
^bComparison of phenotypic findings in three patients with c.2408_2412del and in two patients with c.1801C>T.

described mutations were found in eight patients (four from the same family) with c.1903_1907del [Ockeloen et al., 2015; Parenti et al., 2016] (Table Ia) and one patient with c.2398_2401del [Walz et al., 2015]. Novel recurrent mutations identified in this study included a deletion in three patients, c.2408-2412del and a nonsense mutation, c.1801C>T, p.(Arg601*) in two patients (Table Ib). The additional novel mutations occurred in single patients. Comparison of the phenotypes of all reported cases of recurrent mutations did not highlight specific correlations, although we noted that none of the eight c.1903_1907del patients had hearing loss or severe otitis media which was common in the cohort overall.

In addition to the patients in this study, we have analyzed additional samples for patients who were referred for TS or for validation after DDD. In this group an additional six mutations have been identified, two of which are of interest because they are outside exon 9. The first, a de novo frameshift in exon 6, occurred in a patient (DDD266108) with developmental delay, speech impairment, clinodactyly, anisocoria, and unusual face shape. The other, a nonsense mutation in exon 4, occurred in a child and mother with typical features of KBG Syndrome.

DISCUSSION

This study of British patients with a molecular genetic diagnosis of KBG syndrome is the largest to date, bringing the total number of published cases to >100. Our findings show that the characteristic facial phenotype of KBG syndrome can be recognized as indicated in previous reports [Herrmann et al., 1975; Smithson et al., 2000; Brancati et al., 2006; Skjei et al., 2007; Sirmaci et al., 2011; Youngs et al., 2011; Ockeloen et al., 2015]. On review of a panel of patient photographs, (Fig. 2) we identified in some but not all patients a triangular face, almond-shaped palpebral fissures which may slant upwards, synophrys, prominent ears, a high nasal bridge and bulbous nasal tip, prominent cheekbones, a smooth philtrum with thin upper lip, prominent corners of the mouth, and facial hypertrichosis/low hairlines. Our impression from available data is that the features can evolve in childhood and become more obvious in time. We agree with other authors that while some KBG patients can be recognized by gestalt, others may resemble Cornelia de Lange syndrome (CDLS). Ansari et al. [2014] found that 3/163 CDLS-like patients had ANKRD11 mutations and Busa et al. [2015] reported on a French cohort of 20 mutation/deletion cases and commented that the facial features were reminiscent of CDLS. Parenti et al. [2016] also found an overlap in phenotyping and advised ANKRD11 analysis for CDLS mutation negative patients. In our patients, we observed some facial similarities with CDLS, particularly the appearance of the eyebrows and eyelashes.

When we compared the facial phenotypes of patients in the two diagnostic pathways, we observed that in the targeted group the gestalt was present in a higher proportion of cases than in the DDD group, where it was only clear in three. In some patients, we considered that the specific facial features were not present, consistent with the high ascertainment from the DDD study (19/32 cases).

In 22% of patients, we found a wide anterior fontanelle with delayed closure and speculate this may contribute to the triangular

face shape in KBG syndrome. In some cases this was the presenting feature and led to consideration of cleidocranial dysostosis (MIM # 119600). Furthermore, delayed closure of the anterior fontanelle is one of the most distinctive features according to the Decipher gene browser view for *ANKRD11*. Other conditions with cranial involvement and a similarly broad forehead include Robinow syndrome (MIM # 180700) which was investigated in one patient in this study. Five of the patients were referred for investigation of possible craniosynostosis on the basis of the broad forehead and brachycephaly, but no evidence of fused cranial sutures was found. Although, patients with KBG syndrome can be relatively short in stature, we did not find evidence of a generalised skeletal dysplasia. The most consistent finding was brachydactyly with short tubular bones in the hands and feet radiographically. One of our patients (13) has been treated with growth hormone and some success has been reported elsewhere [Reynaert et al., 2015].

Our study clearly shows a predisposition for recurrent otitis media in KBG syndrome. This is important to identify and treat to reduce hearing loss and optimise speech development. We also identified palatal abnormalities/dysfunction in a quarter of patients, which may be missed without specialist assessment. Based on the findings of this study and others, we recommend some key measures for all patients with a new diagnosis of KBG syndrome (summarized in Table II) to assist pediatric and other colleagues involved in the care of patients.

All patients in this study had a mutation in *ANKRD11*, a member of a family of Ankyrin repeat containing cofactors described a decade ago [Zhang et al., 2004] which is expressed in a wide variety of tissues including the brain. It is implicated in activation of transcription by interaction with p160 coactivator and nuclear receptor complex by recruiting histone deacetylases [Sirmaci

et al., 2011]. This may account for the pleiotropic effects of *ANKRD11* alterations. Subsequently, *Ankrd11* has been shown to be a crucial chromatin regulator that controls histone acetylation and gene expression during neural development [Gallagher et al., 2015]. This provides a likely explanation for the widespread finding of variable learning difficulties and other cognitive dysfunction in our cohort. Yoda mice, heterozygous for a missense mutation in *ankrd11* show craniofacial anomalies, wide skulls, kyphoscoliosis, and reduced bone mineral density, but no other costovertebral anomalies or macrodontia [Sirmaci et al., 2011]. The precise role of *ANKRD11* in bone formation and modelling is also currently unknown.

Like other authors, we did not establish a predictive value for specific mutations, but it is possible that c.1903_1907del may be associated with a relatively mild phenotype with respect to ear infections and hearing loss and further work will be required to clarify this. This lack of genotype–phenotype correlation suggests that the variable nature of the KBG syndrome cannot be attributed to the position of truncation of the protein. This supports a common mechanism of pathogenicity in which truncating mutations in the C terminal domain would be expected to affect protein degradation [Walz et al., 2015]. Comparison of the phenotypes in patients 5–8 and 29–31 demonstrate intra-familial variability and variable penetrance.

In addition to comparing the facial features, we revisited the whole phenotype in the two ascertainment groups. If sufficient characteristics are present, such as macrodontia in a child with developmental/speech delay, or learning difficulties, it is likely the diagnosis of KBG syndrome will be considered and TS initiated. It is clear from this study that the presentation is variable and the diagnosis can be missed [Ashraf et al., 2015], especially in infants

TABLE II. Recommendations for Clinical Management of Patients With KBG Syndrome

Recommended investigations/follow up

Diagnosis	Molecular testing <i>ANKRD11</i> Array CGH for 16q24 microdeletion if KBG phenotype and <i>ANKRD11</i> sequencing normal
Care coordinated by key clinician	Hospital or community paediatrician for children and GP for adults
Investigations to consider in all patients	Echocardiogram Palatal assessment with specialist team Hearing and vision assessment Specialist dental review
Referrals and on-going surveillance that may be required	Neurology for seizures and movement disorders Cardiology if heart lesion identified Dietician for feeding issues Endocrinology for investigation of short stature if present Surgery for cryptorchidism and hernia Respiratory/sleep studies for apnoea ENT/audiology for recurrent otitis media and/or hearing loss
Management of learning and behaviour	Paediatric MDT assessment for developmental delay, ASD/ADHD and complex behaviour patterns Educational support where required
Investigations that may be indicated in individual patients	Skeletal survey/assessment of bone age Renal ultrasound MRI brain scan

A diagnosis of KBG Syndrome should be considered in a patient with developmental delay/learning difficulties, speech delay or significant behavioural issues with at least TWO major criteria OR one major AND two minor criteria	
Major criteria	Proportion of patients in cohort with feature
Macrodonia-or Gestalt of KBG in child with primary dentition	85%
Height below 10 th centile	66%
Recurrent otitis media and/or hearing loss	44%
1 st degree relative with KBG Syndrome	22%
Minor criteria	
Brachydactyly or relevant hand anomaly	50%
Seizures	43%
Cryptorchidism	31%
Feeding problems	34%
Palate abnormality	25%
Formal diagnosis of Autism	24%
Large anterior fontanelle and/or delayed closure	22%

FIG. 3. Diagnostic aid for KBG syndrome.

and young children. For KBG, ES methodology revealed that a non-specific *ANKRD11*-related phenotype exists, as found for other conditions [Need et al., 2012]. To date (January 2016), in addition to those reported here, 14 further patients with a variant in *ANKRD11* have been identified by the DDD study (32 intragenic SNV/indels and one intragenic deletion in the first 4294 trios). This is consistent with our finding that the gestalt can be unclear and easily missed.

Skjei et al. [2007] suggested diagnostic criteria for KBG syndrome which required at least 4/8 features including delayed bone age and costovertebral anomalies to be present. Ockeloen et al. [2015] suggested that only 3/8 may be necessary based on their Dutch cohort. However, in our experience and in previous reports, many patients have not had an estimation of bone age or skeletal survey as they were not clinically indicated. Furthermore, despite consensus regarding the gestalt of KBG, our data demonstrate that the facial features can be very subtle or unrecognizable. We have therefore, devised a diagnostic aid for KBG syndrome (Fig. 3) which does not rely on the original features-this would have identified all but one (patient 22 who is notably atypical in his overgrowth parameters) in our cohort. DDD has detected some mild cases with few features and colleagues who are experienced in clinical dysmorphology did not consider KBG syndrome in their differential diagnosis (personal communications).

CONCLUSIONS

This study is the largest to date of patients with KBG syndrome. Detailed phenotyping has provided data on the range of physical features in this condition including wide anterior fontanelle with delayed closure, not previously described. We report on the significant range of developmental and behavioral problems that affect patients. The presentation of KBG may be non-specific and the diagnosis made for the first time through ES; therefore, we recommend that *ANKRD11* is included in gene panels used to

investigate developmental delay. KBG syndrome is emerging as a Mendelian cause of learning difficulties with important implications for genetic counselling given the autosomal dominant inheritance of this condition. We have devised a revised diagnostic aid for clinical use and suggest clinical investigations and follow up that are likely to improve patient care.

ACKNOWLEDGMENTS

We thank the patients and families for their helpful participation in this work and all colleagues who have contributed to the data presented here. The DDD study presents independent research commissioned by the Health Innovation Challenge Fund [grant number HICF-1009-003], a parallel funding partnership between the Wellcome Trust and the Department of Health, and the Wellcome Trust Sanger Institute [grant number WT098051]. The views expressed in this publication are those of the author(s) and not necessarily those of the Wellcome Trust or the Department of Health. The study has UK Research Ethics Committee approval (10/H0305/83, granted by the Cambridge South REC, and GEN/284/12 granted by the Republic of Ireland REC). The research team acknowledges the support of the National Institute for Health Research, through the Comprehensive Clinical Research Network.

REFERENCES

- Ansari M, Poke G, Ferry Q, Williamson K, Aldridge R, Meynert AM, Bengani H, Yee Chan C, Kayserili H, Avci S, Hennekam RCM, Lampe AK, Redeker E, Homfray T, Ross A, Falkenberg Smeland M, Mansour S, Parker MJ, Cook JA, Splitt M, Fisher RB, Fryer A, Magee AC, Wilkie A, Barnicoat A, Brady AF, Cooper NS, Mercer C, Deshpande C, Bennett CP, Pilz DT, Ruddy D, Cilliers D, Johnson DS, Josifova D, Rosser E, Thompson EM, Wakeling E, Kinning E, Stewart F, Flinter F, Girisha KM, Cox H, Firth HV, Kingstn H, Wee JS, Hurst JA, Clayton-Smith J,

- Tolmie J, Vogt J, Tatton-Brown K, Chandler K, Prescott K, Wilson L, Behnam M, McEntagart M, Davidson R, Lynch S, Sisodiya S, Mehta SG, McKee SA, Mohammed S, Holden S, Park S, Holder SE, Harrison V, McConnell V, Lam WK, Green AJ, Donnai D, Bitner-Glindzicz M, Donnelly DE, Nellaker C, Taylor MS, Fitzpatrick DR. 2014. Genetic heterogeneity in Cornelia de Lange syndrome (CdLS) and CdLS-like phenotypes with observed and predicted levels of mosaicism. *J Med Genet* 51:659–668.
- Ashraf T, Irving M, Canham N, Holder S, Foulds N, Magee A, McConnell V, Fisher R, McEntagart M, Tolmie J, Joss S, Clayton-Smith J, Study DDD. 2015. KBG syndrome: A DDD front-runner? *Eur Soc Hum Genet Abstract* PM08 38 159.
- Brancati F, Sarkozy A, Dallapiccola B. 2006. KBG syndrome. *Orphanet J Rare Dis* 12:1–50.
- Busa T, Tessier A, Riccardi F, Jacquette A, Genevieve D, Gatinois V, Lacombe D, Michaud V, Rossi M, Lebreton C, Leheup B, Vincent-Delorme C, Delahaye A, Van Maldergem L, Cacciagli P, Scheffer H, Saugier-Weber P, Goldenburg A, Philip N. 2015. KBG syndrome: A series of 20 French patients. *Eur Soc Hum Genet Abstract* PS11 79 223.
- Carey JC, Cohen MM, Jr, Curry CJR, Devriendt K, Holmes LB, Verloes A. 2009. Elements of morphology: Standard terminology for the lips, mouth, and oral region. *Am J Med Genet Part A* 149A:77–92.
- Firth HV, Richards SM, Bevan AP, Clayton S, Corpas M, Rajan D, Van Vooren S, Moreau Y, Pettett RM, Carter NP. 2009. DECIPHER: Database of chromosomal imbalance and phenotype in humans using ensembl resources. *Am J Hum Genet* 84:524–533.
- Gallagher D, Voronova A, Zander MA, Cancino GI, Bramall A, Krause MP, Abad C, Tekin M, Neilsen PM, Callen DF, Scherer SW, Keller GM, Kaplan DR, Walz K, Miller FD. 2015. *Ankrd11* is a chromatin regulator involved in autism that is essential for neural development. *Dev Cell* 32:31–42.
- Herrmann J, Pallister PD, Tiddy W, Opitz M. 1975. The KBG syndrome—a syndrome of short stature, characteristic facies, mental retardation, macrodontia and skeletal anomalies. *Birth Defects Orig Artic Ser* 11:7–18.
- Isrie M, Hendriks Y, Gielissen N, Sijm EA, Willemsen MH, Peeters H, Vermeesch JR, Kleefstra T, Van Esch H. 2012. Haploinsufficiency of *ANKRD11* causes mild cognitive impairment, short stature and minor dysmorphisms. *Eur J Hum Genet* 20:131–133.
- Kim HJ, Cho E, Park JB, Wy I, Kim HJ. 2015. A Korean family with KBG syndrome identified by *ANKRD11* mutation, and phenotypic comparison of *ANKRD11* mutation and 16q24.3 microdeletion. *Eur J Med Genet* 58:86–94.
- Koessaar T, Remm M. 2007. Enhancements and modifications of primer design program Primer 3. *Bioinformatics* 23:1289–1291.
- Lo-Castro A, Brancati F, Digilio MC, Garaci FG, Bollero P, Alfieri P, Curatolo P. 2013. Neurobehavioral phenotype observed in KBG syndrome caused by *ANKRD11* mutations. *Am J Neuropsychiatr Genet* 162B:17–23.
- Need AC, Shashi V, Hitomi Y, Schoch K, Shianna KV, McDonald MT, Meisler MH, Goldstein DB. 2012. Clinical application of exome sequencing in undiagnosed genetic conditions. *J Med Genet* 49:353–361.
- Ockeloen CW, Willemsen MH, de Munnik S, van Bon BW, de Leeuw N, Verrips A, Kant SG, Jones EA, Brunner HG, van Loon RL, Smeets EE, van Haelst MM, van Haaften G, Nordgren A, Malmgren H, Grigelioniene G, Vermeer S, Louro P, Ramos L, Maal TJ, van Heumen CC, Yntema HG, Carels CE, Kleefstra T. 2015. Further delineation of the KBG syndrome phenotype caused by *ANKRD11* aberrations. *Eur J Hum Genet* 23:1270.
- Parenti I, Gervasini C, Pozojevic J, Graul-Neumann L, Azzollini J, Braunholz D, Watrin E, Wendt KS, Cereda A, Cittaro D, Gillissen-Kaesbach G, Lazarevic D, Mariani M, Russo S, Werner R, Krawitz P, Larizza L, Selicorni A, Kaiser FJ. 2016. Broadening of cohesinopathies: Exome sequencing identifies mutations in *ANKRD11* in two patients with Cornelia de Lange-overlapping phenotype. *Clin Genet* 89:74–81.
- Reynaert N, Ockeloen CW, Säwendahl L, Beckers D, Devriendt K, Kleefstra T, Carels CE, Grigelioniene G, Nordgren A, Francois I, de Zegher F, Casteels K. 2015. Short stature in KBG syndrome: First responses to growth hormone treatment. *Horm Res Paediatr* 83:361–364.
- Sacharow S, Li D, Fan YS, Tekin M. 2012. Familial 16q24.3 microdeletion involving *ANKRD11* causes a KBG-like syndrome. *Am J Med Genet Part A* 185A:547–552.
- Sirmaci A, Spiliopoulos M, Brancati F, Powell E, Duman D, Abrams A, Bademci G, Agolini E, Guo S, Konuk B, Kavaz A, Blanton S, Digilio MC, Dallapiccola B, Young J, Zuchner S, Tekin M. 2011. Mutations in *ANKRD11* cause KBG syndrome, characterized by intellectual disability, skeletal malformations, and macrodontia. *Am J Hum Genet* 89:289–294.
- Skjei KL, Martin MM, Slavotinek AM. 2007. KBG syndrome: Report of twins, neurological characteristics, and delineation of diagnostic criteria. *Am J Med Genet Part A* 143A:292–300.
- Smithson SF, Thompson EM, McKinnon AG, Smith IS, Winter RM. 2000. The KBG syndrome. *Clin Dysmorphol* 9:87–91.
- Untergasser A, Cutcutache I, Koessaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012. Primer3—new capabilities and interfaces. *Nucleic Acids Res* 40:e115.
- Walz K, Cohen D, Neilsen PM, Foster J, 2nd, Brancati F, Demir K, Fisher R, Moffat M, Verbeek NE, Bjorgo K, Lo Castro A, Curatolo P, Novelli G, Abad C, Lei C, Zhang L, Diaz-Horta O, Young JJ, Callen DF, Tekin M. 2015. Characterization of *ANKRD11* mutations in humans and mice related to KBG syndrome. *Hum Genet* 134:181–190.
- Willemsen MH, Fernandez BA, Bacino CA, Gerkes E, de Brouwer AP, Pfundt R, Sikkema-Raddatz B, Scherer SW, Marshall CR, Potocki L, van Bokhoven H, Kleefstra T. 2010. Identification of *ANKRD11* and *ZNF778* as candidate genes for autism and variable cognitive impairment in the novel 16q24.3 microdeletion syndrome. *Eur J Hum Genet* 18:429–435.
- Wright CF, Fitzgerald TW, Jones WD, Clayton S, McRae JF, van Kogelenberg M, King DA, Ambridge K, Barrett DM, Bayzatinova T, Bevan AP, Bragin E, Chatzimichali EA, Gribble S, Jones P, Krishnappa N, Mason LE, Miller R, Morley KI, Parthiban V, Prigmore E, Rajan D, Sifrim A, Swaminathan GJ, Tivey AR, Middleton A, Parker M, Carter NP, Barrett JC, Hurles ME, FitzPatrick DR, Firth HV, study DDD. 2015. Genetic diagnosis of developmental disorders in the DDD study: A scalable analysis of genome-wide research data. *Lancet* 385:1305–1314.
- Youngs EL, Hellings JA, Butler MG. 2011. *ANKRD11* deletion in a 17-year-old male. *Clin Dysmorphol* 20:170–171.
- Zhang A, Yeung PL, Li C-W, Tsai SC, Dinh GK, Wu X, Li H, Chen JD. 2004. Identification of a novel family of ankyrin repeats containing cofactors for p160 nuclear receptor coactivators. *J Biol Chem* 279:33799–33805.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's web-site.