



Ferrata Storti Foundation

Haematologica 2019
Volume 104(1):128-137

Clinical and molecular characteristics of *MEF2D* fusion-positive B-cell precursor acute lymphoblastic leukemia in childhood, including a novel translocation resulting in *MEF2D-HNRNPH1* gene fusion

Kentaro Ohki,¹ Nobutaka Kiyokawa,¹ Yuya Saito,^{1,2} Shinsuke Hirabayashi,^{1,3} Kazuhiko Nakabayashi,⁴ Hitoshi Ichikawa,⁵ Yukihide Momozawa,⁶ Kohji Okamura,⁷ Ai Yoshimi,^{1,8} Hiroko Ogata-Kawata,⁴ Hiromi Sakamoto,⁵ Motohiro Kato,¹ Keitaro Fukushima,⁹ Daisuke Hasegawa,³ Hiroko Fukushima,¹⁰ Masako Imai,¹¹ Ryosuke Kajiwara,¹² Takashi Koike,¹³ Isao Komori,¹⁴ Atsushi Matsui,¹⁵ Makiko Mori,¹⁶ Koichi Moriwaki,¹⁷ Yasushi Noguchi,¹⁸ Myoung-ja Park,¹⁹ Takahiro Ueda,²⁰ Shohei Yamamoto,²¹ Koichi Matsuda,²² Teruhiko Yoshida,⁵ Kenji Matsumoto,²³ Kenichiro Hata,⁴ Michiaki Kubo,⁶ Yoichi Matsubara,²⁴ Hiroyuki Takahashi,²⁵ Takashi Fukushima,²⁶ Yasuhide Hayashi,²⁷ Katsuyoshi Koh,¹⁶ Atsushi Manabe³ and Akira Ohara²⁵ for the Tokyo Children's Cancer Study Group (TCCSG)

¹Department of Pediatric Hematology and Oncology Research, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²Department of Hematology/Oncology, Tokyo Metropolitan Children's Medical Center, Fuchu-shi;

³Department of Pediatrics, St. Luke's International Hospital, Chuo-ku, Tokyo; ⁴Department of Maternal-Fetal Biology, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ⁵Fundamental Innovative Oncology Core, National Cancer Center Research Institute, Chuo-ku, Tokyo; ⁶Laboratory for Genotyping Development, RIKEN Center for Integrative Medical Sciences (IMS), Yokohama-shi, Kanagawa; ⁷Department of Systems BioMedicine, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ⁸Division of Pediatric Hematology and Oncology, Ibaraki Children's Hospital, Mito-shi; ⁹Department of Pediatrics, Dokkyo Medical University, Tochigi; ¹⁰Department of Pediatrics, University of Tsukuba Hospital, Ibaraki; ¹¹Department of Pediatrics, Japanese Red Cross Musashino Hospital, Tokyo; ¹²Department of Pediatrics, Yokohama City University Hospital, Kanagawa; ¹³Department of Pediatrics, Tokai University School of Medicine, Kanagawa; ¹⁴Department of Pediatrics, Matsudo City Hospital, Chiba; ¹⁵Department of Pediatrics, Japanese Red Cross Maebashi Hospital, Gunma; ¹⁶Department of Hematology/Oncology, Saitama Children's Medical Center; ¹⁷Department of Pediatrics, Saitama Medical Center, Saitama Medical University; ¹⁸Department of Pediatrics, Japanese Red Cross Narita Hospital, Chiba; ¹⁹Department of Hematology/Oncology, Gunma Children's Medical Center, Shibukawa-shi; ²⁰Department of Pediatrics, Nippon Medical School, Bunkyo-ku, Tokyo; ²¹Department of Pediatrics, Showa University Fujigaoka Hospital, Yokohama-shi, Kanagawa; ²²Laboratory of Clinical Genome Sequencing Department of Computational Biology and Medical Sciences Graduate School of Frontier Sciences, The University of Tokyo, Minato-ku; ²³Department of Allergy and Clinical Immunology, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²⁴Director, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²⁵Department of Pediatrics, Toho University Omori Medical Center, Tokyo; ²⁶Department of Child Health, Faculty of Medicine, University of Tsukuba, Ibaraki and ²⁷Institute of Physiology and Medicine, Jobu University, Takasaki-shi, Gunma, Japan

³Department of Pediatrics, St. Luke's International Hospital, Chuo-ku, Tokyo;

⁴Department of Maternal-Fetal Biology, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ⁵Fundamental Innovative Oncology Core, National Cancer Center Research Institute, Chuo-ku, Tokyo; ⁶Laboratory for Genotyping Development, RIKEN Center for Integrative Medical Sciences (IMS), Yokohama-shi, Kanagawa; ⁷Department of Systems BioMedicine, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ⁸Division of Pediatric Hematology and Oncology, Ibaraki Children's Hospital, Mito-shi; ⁹Department of Pediatrics, Dokkyo Medical University, Tochigi; ¹⁰Department of Pediatrics, University of Tsukuba Hospital, Ibaraki; ¹¹Department of Pediatrics, Japanese Red Cross Musashino Hospital, Tokyo; ¹²Department of Pediatrics, Yokohama City University Hospital, Kanagawa; ¹³Department of Pediatrics, Tokai University School of Medicine, Kanagawa; ¹⁴Department of Pediatrics, Matsudo City Hospital, Chiba; ¹⁵Department of Pediatrics, Japanese Red Cross Maebashi Hospital, Gunma; ¹⁶Department of Hematology/Oncology, Saitama Children's Medical Center; ¹⁷Department of Pediatrics, Saitama Medical Center, Saitama Medical University; ¹⁸Department of Pediatrics, Japanese Red Cross Narita Hospital, Chiba; ¹⁹Department of Hematology/Oncology, Gunma Children's Medical Center, Shibukawa-shi; ²⁰Department of Pediatrics, Nippon Medical School, Bunkyo-ku, Tokyo; ²¹Department of Pediatrics, Showa University Fujigaoka Hospital, Yokohama-shi, Kanagawa; ²²Laboratory of Clinical Genome Sequencing Department of Computational Biology and Medical Sciences Graduate School of Frontier Sciences, The University of Tokyo, Minato-ku; ²³Department of Allergy and Clinical Immunology, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²⁴Director, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²⁵Department of Pediatrics, Toho University Omori Medical Center, Tokyo; ²⁶Department of Child Health, Faculty of Medicine, University of Tsukuba, Ibaraki and ²⁷Institute of Physiology and Medicine, Jobu University, Takasaki-shi, Gunma, Japan

¹²Department of Pediatrics, Yokohama City University Hospital, Kanagawa; ¹³Department of Pediatrics, Tokai University School of Medicine, Kanagawa; ¹⁴Department of Pediatrics, Matsudo City Hospital, Chiba; ¹⁵Department of Pediatrics, Japanese Red Cross Maebashi Hospital, Gunma; ¹⁶Department of Hematology/Oncology, Saitama Children's Medical Center; ¹⁷Department of Pediatrics, Saitama Medical Center, Saitama Medical University; ¹⁸Department of Pediatrics, Japanese Red Cross Narita Hospital, Chiba; ¹⁹Department of Hematology/Oncology, Gunma Children's Medical Center, Shibukawa-shi; ²⁰Department of Pediatrics, Nippon Medical School, Bunkyo-ku, Tokyo; ²¹Department of Pediatrics, Showa University Fujigaoka Hospital, Yokohama-shi, Kanagawa; ²²Laboratory of Clinical Genome Sequencing Department of Computational Biology and Medical Sciences Graduate School of Frontier Sciences, The University of Tokyo, Minato-ku; ²³Department of Allergy and Clinical Immunology, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²⁴Director, National Research Institute for Child Health and Development, Setagaya-ku, Tokyo; ²⁵Department of Pediatrics, Toho University Omori Medical Center, Tokyo; ²⁶Department of Child Health, Faculty of Medicine, University of Tsukuba, Ibaraki and ²⁷Institute of Physiology and Medicine, Jobu University, Takasaki-shi, Gunma, Japan

Correspondence:

oki-kn@ncchd.go.jp

Received: December 11, 2017.

Accepted: August 29, 2018.

Pre-published: August 31, 2018.

doi:10.3324/haematol.2017.186320

Check the online version for the most updated information on this article, online supplements, and information on authorship & disclosures: www.haematologica.org/content/104/1/128

©2019 Ferrata Storti Foundation

Material published in *Haematologica* is covered by copyright. All rights are reserved to the Ferrata Storti Foundation. Use of published material is allowed under the following terms and conditions:

<https://creativecommons.org/licenses/by-nc/4.0/legalcode>.

Copies of published material are allowed for personal or internal use. Sharing published material for non-commercial purposes is subject to the following conditions:

<https://creativecommons.org/licenses/by-nc/4.0/legalcode>,

sect. 3. Reproducing and sharing published material for commercial purposes is not allowed without permission in writing from the publisher.



ABSTRACT

Fusion genes involving *MEF2D* have recently been identified in precursor B-cell acute lymphoblastic leukemia, mutually exclusive of the common risk stratifying genetic abnormalities, although their true incidence and associated clinical characteristics remain unknown. We identified 16 cases of acute lymphoblastic leukemia and 1 of lymphoma harboring *MEF2D* fusions, including *MEF2D-BCL9* (n=10), *MEF2D-HNRNPU1* (n=6), and one novel *MEF2D-HNRNPH1* fusion. The incidence of *MEF2D* fusions overall was 2.4% among consecutive precursor B-cell acute lymphoblastic leukemia patients enrolled onto a single clinical trial. They frequently showed a cytoplasmic μ chain-positive pre-B immunophenotype, and often expressed an aberrant CD5 antigen. Besides up- and down-regulation of *HDAC9* and *MEF2C*, elevated *GATA3* expression was also a characteristic feature of *MEF2D*

fusion-positive patients. Mutations of *PHF6*, recurrent in T-cell acute lymphoblastic leukemia, also showed an unexpectedly high frequency (50%) in these patients. *MEF2D* fusion-positive patients were older (median age 9 years) with elevated WBC counts (median: 27,300/ μ l) at presentation and, as a result, were mostly classified as NCI high risk. Although they responded well to steroid treatment, *MEF2D* fusion-positive patients showed a significantly worse outcome, with 53.3% relapse and subsequent death. Stem cell transplantation was ineffective as salvage therapy. Interestingly, relapse was frequently associated with the presence of *CDKN2A/CDKN2B* gene deletions. Our observations indicate that *MEF2D* fusions comprise a distinct subgroup of precursor B-cell acute lymphoblastic leukemia with a characteristic immunophenotype and gene expression signature, associated with distinct clinical features.

Introduction

Precursor B-cell acute lymphoblastic leukemia (B-ALL) is a heterogeneous disease characterized by a variety of genetic abnormalities.¹ In approximately one-quarter of B-ALL patients, known as the B-other-subgroup, the known major risk-stratifying cytogenetic abnormalities are absent.² However, recent studies using advanced analytical approaches have described a range of novel genetic subgroups among B-other-ALL.³⁻¹¹

The *myocyte enhancer factor 2D* (*MEF2D*) gene, located at 1q22, is present among these newly identified rearrangements in B-other-ALL.^{6,7,10,12-14} Seven known fusion partners: *B-cell CLL/lymphoma 9* (*BCL9*, 1q21), *heterogeneous nuclear ribonucleoprotein U-like 1* (*HNRNPUL1*, 19q13.2), *deleted in azoospermia-associated protein 1* (*DAZAP1*, 19p13.3), *colony stimulating factor 1 receptor* (*CSF1R*, 5q32), *synovial sarcoma translocation, chromosome 18* (*SS18*, 18q11.2), *signal transducer and activator of transcription 6* (*STAT6*, 12q13.3), and *Forkhead Box J2* (*FOXJ2*, 12p13.31) have been described, mostly among childhood and young adult B-ALL. The *MEF2D* gene encodes a member of the transcription factor family involved in the control of muscle and neuronal cell differentiation and development, which is regulated by class II histone deacetylases.¹⁵⁻¹⁷ It has been reported that rearrangements result in enhanced *MEF2D* transcriptional activity and lymphoid transformation, thus contributing to the development of a distinct subtype of high-risk leukemia.^{7,10} However, the true incidence and clinical characteristics, including outcome, of patients with B-ALL harboring *MEF2D* fusion genes remains unknown.

In this study, we report the detailed analysis of a subgroup of B-ALL with *MEF2D* fusions within a Japanese pediatric ALL cohort. The *heterogeneous nuclear ribonucleoprotein H1* gene (*HNRNP1*), encoding another family member of heterogeneous nuclear ribonucleoprotein,¹⁸ was identified as a new fusion partner of the *MEF2D* gene. Novel immunophenotypic characteristics and accompanying genetic abnormalities as well as distinctive clinical features of B-ALL harboring *MEF2D* fusions are evaluated and discussed.

Methods

Patient selection and sample preparation

RNA and DNA samples, obtained from pediatric B-ALL patients and stored in the Tokyo Children's Cancer Study Group (TCCSG) biobank^{5,11} were used in this study (*Online Supplementary Figure S1, Table S1*). As indicated in *Online Supplementary Table S1*, the majority of cases originated from the

TCCSG L04-16 study¹⁹ while others, including 2 B-lymphoblastic lymphoma (LBL) patients, originated from different cohorts. All investigations were approved by the institutional review boards and informed consent or assent was obtained from parents or guardians based on their age and level of understanding, as described previously.^{5,11} *Online Supplementary Figure S1* shows the analysis carried out on each case.

Total RNA and genomic DNA were extracted from bone marrow or peripheral blood of patients using the miRNeasy Mini Kit and the QIAamp DNA Mini Kit (Qiagen, Inc., Valencia, CA, USA), respectively.

In this paper, B-other-ALL is defined as B-ALL lacking the major risk stratifying genetic abnormalities, including high hyperdiploidy (≥ 51 chromosomes or DNA index ≥ 1.16), low hypodiploidy/near haploidy (≤ 44 chromosomes), fusions of *ETV6-RUNX1*, *TCF3-PBX1*, *TCF3-HLF*, *BCR-ABL1*, and *MLL* rearrangements as well as more recently identified genetic abnormalities, including rearrangements of *CRLF2* and *ZNF384*, Ph-like ALL-related tyrosine kinase fusions as well as *MEF2D* fusions (*Online Supplementary Table S1*).

Whole transcriptome sequencing and RT-PCR

From previous whole transcriptome sequencing (WTS) of B-other-ALL, we identified cases with *ZNF384* fusions¹¹ as well as other abnormalities (*Online Supplementary Figure S1, Table S1*).¹⁹⁻²¹ We re-analyzed remaining 153 WTS data manipulated by "deFuse",²² an algorithm for gene fusion discovery, and investigated the presence of *MEF2D* fusions. Details of this data analysis have been described previously.¹¹ RT-PCR followed by Sanger sequencing was performed to confirm and screen for fusion transcripts, as described previously,^{5,11} using the primers listed in *Online Supplementary Table S2*.

Multiplex Ligation-dependent Probe Amplification (MLPA)

MLPA analyses were performed on genomic DNA using two types of SALSA Reference Kits, P335 and P383 (MRC Holland, Amsterdam, the Netherlands), according to the manufacturer's instructions. After separation of amplified products, using the ABI3130 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA), the results were analyzed using Gene Mapper Software (Applied Biosystems) and data, including informative headers, electropherograms, ratio plots, validation boxes, and report tables were obtained. In this study, we present only results of deletions of the exons targeted in these kits.

Whole exome sequencing (WES)

Exome libraries prepared from 100 ng of genomic DNA were sequenced using SBS v.4 reagents with the HiSeq2500 sequencing system. Details of whole exome data analyses have been described previously.¹¹

Microarray and data analyses

The gene expression signature for *MEF2D* fusion-positive B-ALL was investigated by DNA microarray-based expression profiling using Human Genome U133 Plus 2.0 Arrays (Affymetrix, Santa Clara, CA, USA). Data were normalized and filtered as described previously.¹¹ Further details are provided in *Online Supplementary Methods*.

Statistical analysis

Mutual univariate analyses of characteristics were conducted using Fisher's exact test or the χ^2 test for qualitative variables. Overall survival (OS) and event-free survival (EFS) were estimated by the Kaplan-Meier method and compared by the log-rank test. Analyses were performed using Prism software, version 6.0 (GraphPad Software, Inc., La Jolla, CA, USA).

Table 1. Characteristics of *MEF2D* fusion-positive cases.

Case	Fusion partner	Identified	Age (years)	Sex	Diagnosis	Initial WBC (/L)	Karyotype	Initial risk (NCI)	Day8 blasts (/L)	Relapse (site)	Relapse date (years)	Salvage therapy after relapse	FUP (years)	Current status	Samples obtained at	Fusion points
1	<i>BCL9</i>	RT-PCR	8	F	B-ALL	3,400	46,XX,-X,add(3)(q27), add(9)(q13),add(9)(q13), add(10)(q22),-12,del(12)(q13), del(13)(q12q14) -20,+22,+2mar[cp3]/46,XX[2]	IR	0	No	-	-	10.9+	1 st CR	Newly diagnosed	Ex4-4bp* ins-Ex9
2	<i>BCL9</i>	RT-PCR	9	F	B-ALL	29,400	46,XX[20]	IR	40	Yes (BM)	1.0	Chemo	1.2	Dead	Newly diagnosed	Ex6-Ex10
3	<i>BCL9</i>	WTS/RT-PCR	15	F	B-ALL	63,200	46,XX[20]	HR	0	Yes (CNS)	1.1	SCT	2.0	Dead	Newly diagnosed	Ex9-Ex9, Ex5-Ex9
4	<i>BCL9</i>	WTS/RT-PCR	10	M	B-ALL	124,100	NA	HR	63	No	-	-	9.1+	1 st CR	Newly diagnosed	Ex6-Ex10
5	<i>BCL9</i>	WTS	7	M	B-ALL	77,300	45,XY,t(1;2)(q21;p13),-9,i(9)(q10),add(16)(q13)[20]	IR	240	Yes (BM)	2.0	SCT	2.6	Dead	Newly diagnosed	Ex5-Ex9
6	<i>BCL9</i>	RT-PCR	9	F	B-ALL	41,400	NA	IR	30	Yes (CNS)	0.8	Chemo	1.9	Dead	Newly diagnosed	Ex6-Ex10
7	<i>BCL9</i>	RT-PCR	9	F	B-ALL	46,200	NA	IR	18	Yes (BM)	1.6	SCT	4.2	Dead	Newly diagnosed, 1st relapse	Ex5-Ex10, Ex6-Ex10
8	<i>BCL9</i>	WTS/RT-PCR	10	F	B-ALL	NA	NA	HR	NA	NA	NA	NA	NA	NA	Newly diagnosed	Ex5-Ex9, Ex6-Ex9*2
9	<i>BCL9</i>	RT-PCR	3	M	B-ALL	5,400	NA	SR	25	No	-	-	13.1+	1 st CR	Newly diagnosed	Ex5-Ex9
10	<i>BCL9</i>	WTS/RT-PCR	7	F	B-LBL	NA	46,XX[4]	Stage IV	NA	No	-	-	5.4+	1 st CR	Newly diagnosed	Ex6-Ex10
11	<i>HNRNPUL1</i>	WTS/RT-PCR	9	M	B-ALL	5,300	NA	IR	0	No	-	-	8.6+	1 st CR	Newly diagnosed	Ex9-Ex12
12	<i>HNRNPUL1</i>	RT-PCR	8	F	B-ALL	8,200	NA	IR	54	No	-	-	9.9+	1 st CR	Newly diagnosed	Ex9-Ex12
13	<i>HNRNPUL1</i>	WTS/RT-PCR	5	F	B-ALL	27,300	46,XX[18]	SR	736	Yes (BM)	1.3	SCT	1.9	Dead	Newly diagnosed	Ex9-Ex12
14	<i>HNRNPUL1</i>	WTS/RT-PCR	15	M	B-ALL	6,200 fail	NA	IR	0	No	-	-	4.5+	1 st CR	Newly diagnosed	Ex9-Ex12
15	<i>HNRNPUL1</i>	WTS/RT-PCR	10	F	B-ALL	21,900	47,XX,t(3;9)	HR	750	Yes (BM)	1.9	Chemo, SCT	2.4	Dead	Newly diagnosed	Ex9-Ex12
16	<i>HNRNPUL1</i>	RT-PCR	14	M	B-ALL	10,800	46,XY,?ins(9)(q13p22p24)[20]	HR	1,134	No	-	-	7.5+	1 st CR	Newly diagnosed	Ex9-Ex12
17	<i>HNRNPUL1</i>	WTS/RT-PCR	7	F	B-ALL	76,100	46,XX,t(1;5)(q21;q35)[8]/47,idem,+8[8]/46,XX[2]	IR	NA	Yes (BM)	1.2	Chemo	1.8	Dead	Newly diagnosed	Ex4-Ex5, Ex7-21bp*4 ins-Ex5

RT-PCR: Reverse Transcription Polymerase Chain Reaction; WTS: Whole Transcriptome Sequencing; WBC: white blood cells; CNS: central nervous system; FUP: follow up; B-ALL: precursor B-cell acute lymphoblastic leukemia; B-LBL: precursor B-cell lymphoblastic lymphoma; NCI: National Cancer Institute; SR: standard risk; IR: intermediate risk; HR: high risk; Ex: exon; BM: bone marrow; CR: complete remission; NA: data not available; SCT: stem-cell transplantation; Chemo: chemotherapy. *1: (TGTC). *2: Ex6-Ex9 fusion was detected only by WTS. *3: The *PAX5-FOXP1* translocation was not detected by whole transcriptome sequencing. *4: (CCCGACCAGCTTGTTGCCG).

Results

Detection of *MEF2D* fusions in pediatric B-ALL patients

Among the 328 selected RNA samples from B-ALL patients (Online Supplementary Figure S1) analyzed by WTS and/or RT-PCR followed by Sanger sequencing, we iden-

tified 9 and 6 patients with *MEF2D-BCL9* and *MEF2D-HNRNPUL1*, respectively (Table 1, Online Supplementary Table S1 and S3, Figure 1 and Online Supplementary Figure S1). Of note, one case of each abnormality was identified by gene expression profiling (details in Online Supplementary Information). We also identified an additional case with *MEF2D-BCL9* (Table 1, Case 10) in B-LBL. As




Figure 1. Structures of the *MEF2D* fusions. Structures of fusion proteins and nucleotide sequences of (a) - (e) *MEF2D-BCL9*, (f) *MEF2D-HNRNPUL1*, (g) and (h) *MEF2D-HNRNP1*. The red arrowhead shows the donor and acceptor site breakpoint. The number of patients in whom a particular fusion isoform was found is indicated on the right.

well as the known *MEF2D* fusions, we identified *MEF2D-HNRNP1* as a novel fusion in 1 patient (Figure 1, Table 1, Case 17). Among the L04-16/L06-16 cohort,^{11,19} comprising a consecutive series of 290 B-ALL patients, including 126 classified as B-other-ALL, 5 *MEF2D-BCL9* and 2 *MEF2D-HNRNP1* patients were identified (Online Supplementary Table S1 and Figure S1). The incidence of *MEF2D* fusions in childhood ALL, calculated from this

cohort, was 5.6% in B-other-ALL and 2.4% in B-ALL overall. *MEF2D-BCL9* was the most recurrent, at a frequency of 4.0% in B-other-ALL and 1.7% in B-ALL overall.

Structure of *MEF2D* fusions

The structure and sequences of *MEF2D-BCL9* as well as a schematic representation of the predicted fusion proteins, are depicted in Figure 1, Table 1 and Online




Figure 2. Immunophenotypic characteristics of B-ALL patients with *MEF2D* fusions. (A) The positivity (percentage) of CD10, cytoplasmic μ , CD5, CD38, and CD33 of *MEF2D* fusion-positive, *TCF3-PBX1*-positive, and B-other patients are plotted as a scattergram. A detailed list of positivity for each immunophenotypic marker of the patients is presented in Online Supplementary Table S4. (B) Typical histograms of CD10, CD19, cytoplasmic μ , aberrant CD5, CD33, and CD38 of *MEF2D-BCL9*, *MEF2D-HNRNP1*, *TCF3-PBX1*, and B-other patients are indicated with a positive rate (%). X-axis, fluorescence intensity; Y-axis, relative cell number.

Supplementary Table S3. Five isoforms of *MEF2D-BCL9* fusion were identified in 10 patients, including one case in which different isoforms were present together. Most frequently, exon 6 of *MEF2D* was fused in-frame to exon 10 of *BCL9* in 5 patients (Figure 2A, Type (a)), in accordance with previous reports.¹⁰ Exons 5 and 6 of *MEF2D* were also fused to exon 9 of *BCL9* in 4 and 1 patient, respectively (Types (b) and (c)). In addition, we identified two novel breakpoints (Types (d) and (e)). The predicted protein from all fusions retains the DNA-binding MADS domain of the *MEF2D* protein, while it lacks most of the c-terminal portion of *MEF2D* as well as most of the functional domains of *BCL9*.

The structure and sequences of *MEF2D-HNRNPUL1* are also presented in Figure 1. Distinct from previous reports, only one isoform joining exon 9 of *MEF2D* to exon 12 of *HNRNPUL1* (Type (f)) was observed among our 6 patients. In the case of *MEF2D-HNRNP1*, two isoforms, in-frame fusions joining exons 4 and 7 of *MEF2D* to exon 5 of *HNRNP1* (Type (g) and (h)), respectively, were identified in the same patient.

Immunophenotypic characteristics of B-ALL patients with *MEF2D* fusions

It has been reported that B-ALL patients with *MEF2D* fusions show dull or negative expression of CD10 and overexpression of CD38 antigens, based on immunophenotypic examination.¹⁰ Among our *MEF2D* fusion-positive cases, CD10 expression was lower than that seen in *TCF3-PBX1*-positive patients, but not B-others-ALL, as shown in Figure 2, *Online Supplementary Figure S3 and Table S4*. On the other hand, CD38 expression in *MEF2D* fusion-positive cases was higher than B-other-ALL patients, but not *TCF3-PBX1*-positive patients. We also observed that expression of the cytoplasmic μ chain was higher in *MEF2D* fusion-positive cases than that in B-

other-ALL patients, with more than 10% of blasts positive for cytoplasmic μ in 8/12 patients (ranging from 5.50 to 99.74%, mean: $54.14 \pm 41.38\%$). Moreover, B-ALL patients with *MEF2D* fusions frequently exhibited aberrant CD5 expression, with more than 10% of the blasts positive for CD5 in 7/12 patients (ranging from 0.60 to 73.30%, mean: $27.01 \pm 26.67\%$).

Additional genetic abnormalities in *MEF2D* fusion-positive patients

To elucidate the frequency of additional genetic abnormalities in B-ALL with *MEF2D* fusions, we initially performed MLPA on 16 DNA samples from patients with *MEF2D* fusions. As shown in *Online Supplementary Table S5*, MLPA analysis revealed that deleted or amplified regions of *IKZF1*, *CRLF2*, *EBF1*, *BTG1*, *PHF6*, *NF1*, *EZH2*, *SUZ12*, or *PTEN* were absent from *MEF2D* fusion-positive cases, although 11/16 exhibited *CDKN2A/CDKN2B* deletions at a frequency significantly higher than that among the B-other-ALL patients enrolled on L04-16/L06-16 study ($P < 0.001$, *data not shown*). Heterozygous deletions of *LEF1*, *PAX5*, and *ETV6* were detected in 1 case each.

To further investigate the presence of additional genetic abnormalities affecting coding sequences in B-ALL with *MEF2D* fusions, we carried out WES on 3 DNA samples. We identified 16 mutations within genes that had been previously recognized within cancer: *ALK*, *ARFGER3*, *BRCA1*, *BRMS1*, *C8orf4*, *ITIH1*, *MAPK13*, *NCOR2*, *NLE*, *NOTCH1*, *PHF3*, *PHF6*, *PHF10*, *PIK3R5*, *RB1*, and *TET1* (*Online Supplementary Table S6*).

As mutations involving *NOTCH1* and *PHF6* are known to be recurrent in T-ALL²³⁻²⁶ and *NLE* encodes the modulator of *NOTCH1*,²⁷ we were encouraged to investigate abnormalities in *NOTCH1* signaling pathway genes and other genes reported as targets of recurrent genetic abnormalities in T-ALL. Thus, we performed RT-PCR and

Table 2. Additional genetic abnormalities of *MEF2D* fusion-positive cases.

Case	Fusion partner	Gene Chromosome	<i>FBXW7</i> 4q31.3	<i>NLE</i> 17q12	<i>NOTCH1</i> 9q34.3	<i>PHF6</i> Xq26.2	<i>PTEN</i> 10q23.31	<i>WT1</i> 11p13
1	<i>BCL9</i>		WT	WT	6764 T C, M2255T	WT	WT	WT
2	<i>BCL9</i>		WT	WT	WT	WT	WT	WT
3	<i>BCL9</i>		1282 C G, Q428E	WT	WT	R15 fs	WT	WT
4	<i>BCL9</i>		WT	WT	6920 A G, Q2307R,	S66C	WT	WT
5	<i>BCL9</i>		WT	WT	WT	WT	WT	WT
6	<i>BCL9</i>		WT	WT	WT	K16I	692 C → T, P231L	WT
7	<i>BCL9</i>		WT	WT	WT	WT	WT	1192 T → A, C398S
8	<i>BCL9</i>		WT	WT	WT	WT	WT	WT
9	<i>BCL9</i>		WT	WT	WT	D309H	WT	WT
11	<i>HNRNPUL1</i>		WT	WT	WT	ex2 del (no start codon), C82Y	WT	WT
12	<i>HNRNPUL1</i>		WT	WT	WT	ex2 del (no start codon)	WT	WT
13	<i>HNRNPUL1</i>		WT	WT	WT	WT	WT	WT
14	<i>HNRNPUL1</i>		WT	WT	WT	WT	WT	WT
15	<i>HNRNPUL1</i>		WT	WT	WT	WT	WT	WT
16	<i>HNRNPUL1</i>		WT	WT	WT	D309H	WT	WT
17	<i>HNRNP1</i>		WT	904 C G, R302G	WT	955 C → T, R319X	WT	WT
		Frequency	1/16	1/16	2/16	8/16	1/16	1/16

ex: exon; WT: wild-type; fs: frame shift; del: deletion.

Sanger sequencing of selected genes, including *FBXW7*, *NLE*, *NOTCH1*, *PHF6*, and *PTEN*, and combined these data with the results of WES. As shown in Table 2, mutations in *NOTCH1* signaling pathway genes, including *FBXW7*, *NLE*, and *NOTCH1*, were detected in 4/16 patients. In addition, 8 and 1 of 16 patients exhibited mutations in *PHF6* and *PTEN*, respectively.

Gene expression signature of *MEF2D* fusion-positive patients

We have microarray data of B-ALL as indicated in *Online Supplementary Figure S4*,¹¹ and, in addition, we performed WTS to search for new fusion genes specifically among cases of B-other-ALL. To further assess the func-

tional aspects of *MEF2D* fusions, we performed DNA microarray-based expression profiling. Upon supervised hierarchical clustering analysis using selected gene probe sets (*Online Supplementary Table S7 and S8*), we observed distinct clustering of *MEF2D* fusion cases, with clear separation from cases with other genetic abnormalities, indicating that *MEF2D* fusion cases have a distinct gene expression signature (Figure 3A). Of note was that this cluster of *MEF2D* fusion cases was close to that of *TCF3-PBX1*-positive cases. Principal component analysis (PCA) also revealed clear clustering of *MEF2D* fusion cases separate from those cases with other genetic abnormalities, but close to the cluster of *TCF3-PBX1*-positive cases (Figure 3B).




Figure 3. Characteristics of gene expression profile in *MEF2D*-fusion-positive ALL. (A) Two-way hierarchical clustering and (B) principal component analysis (PCA) were performed on the microarray data, including B-ALLs with *MEF2D* fusion-positive and other types of genetic abnormalities, using the probe sets of differentially expressed genes between B-ALL with *MEF2D* fusions and other types of genetic abnormalities selected from filtered microarray probes (presented in *Online Supplementary Tables S7 and S8*). The results of clustering analysis are displayed using a heatmap as a dendrogram.

As *TCF3-PBX1*-positive cases also express the cytoplasmic μ chain-positive pre-B ALL immunophenotype,^{28,29} we subsequently compared the gene expression of *MEF2D* fusion cases with *TCF3-PBX1*-positive cases as well as B-other-ALL. As shown in *Online Supplementary Figure S4*, *MEF2D* fusion patients were clearly separate, within a distinct cluster from *TCF3-PBX1*-positive and B-other-ALL cases, based on unsupervised hierarchical clustering. Up-regulated genes common to both *MEF2D* fusion and *TCF3-PBX1*-positive cases compared to B-other-ALL included: *IKZF2*, *IRF4*, *TCL6*, *IGHG*, *IGHV5-78*, *IIGLL1*, *VPREB3*, and *BCL2L11*, while *RUNX2*, *IRF9*, *CRLF2*, *CD34*, and *CCND2* were common down-regulated genes (summarized in *Online Supplementary Figure S5*). *HDAC9* was also identified as a common up-regulated gene in both *MEF2D* fusion and *TCF3-PBX1*-positive cases, whereas its expression was significantly higher in *MEF2D* fusion than *TCF3-PBX1*-positive cases. On the other hand, *MEF2D*, *MME* (coding CD10), and *RAG1* were down-regulated in *MEF2D* fusion, but not in *TCF3-PBX1*-positive cases, thus the low-level expression of CD10 seen in *MEF2D* fusion cases was identified at the gene-expression level. Interestingly, *GATA3* was identified as a highly expressed gene in *MEF2D* fusion cases, yet it was significantly down-regulated in *TCF3-PBX1*-positive cases.

To explore the gene expression characteristics of *MEF2D* fusion ALL connected with B-cell differentiation, we performed GSEA using 18 curated gene sets of B lymphocytes at various differentiation stages, as well as 7 early hematopoietic stages including stem cells. Firstly, we compared the gene expression signatures of B-ALL cases with different types of genetic abnormalities. Using B-other-ALL as a reference control, we observed that the majority of gene expression signatures found in B lymphocytes at various differentiation stages were enriched in ALL cases with *TCF3-PBX1* (*Online Supplementary Table S10 and S11*). In contrast, only three gene sets were enriched in *MEF2D* fusion ALL. We further examined the gene expression characteristics of *MEF2D* fusion ALL by direct comparison with *TCF3-PBX1*, and observed that most of the signatures of differentiation stage-specific B lymphocytes as well as early hematopoietic populations were enriched in *TCF3-PBX1*, but not *MEF2D* fusion ALL (*Online Supplementary Table S10*).

Clinical characteristics and outcomes of *MEF2D* fusion-positive patients

The clinical findings and outcome of patients with *MEF2D* fusions are summarized in Table 1. *MEF2D* fusion B-ALL patients were aged between 3 and 15 years (median: 9 years) at presentation and comprised 6 males and 10 females. Their initial white blood cell (WBC) counts at presentation ranged from 3,400 to 124,100 (median: 27,300/ μ L). Analysis of fluids obtained by lumbar puncture revealed no indication of central nervous system (CNS) involvement (data not shown). Among 13 patients, 10 (62.5%) and 4 (25.0%) were classified with an intermediate risk (IR) and high risk (HR), respectively, based on advanced age, elevated WBC counts, or both, and thus standard risk (SR) was assigned to only 2 patients. The response to steroid monotherapy, using the cutoff of 1,000/ μ L for the blast count in peripheral blood on day 8 was not poor in *MEF2D* fusion patients; however, among 15 patients, 8 (53.3%) showed bone marrow or CNS relapse, and all of the relapsed patients died. As shown in Figure 4, both the 5-year EFS and OS rates for *MEF2D* fusion patients were significantly lower compared with a consecutive series of B-other-ALL patients enrolled onto the L04-16/L06-16 study ($P=0.0306$ and $P=0.0013$, respectively), indicating that outcomes for *MEF2D* fusion-positive patients were significantly less favorable than those with B-other-ALL.

Discussion

In this study, we identified *MEF2D* fusions, including a novel fusion gene, *MEF2D-HNRNP11*, at an incidence of approximately 2% in our B-ALL cohort. It was notable that we identified *MEF2D* fusions in B-LBL. B-ALL patients with *MEF2D* fusions showed unique clinical and biological characteristics. They exhibited an older age at presentation and elevated WBC counts, thus were mostly classified into the IR or HR groups. Although their response to steroid treatment was not poor, *MEF2D* fusion patients showed a significantly worse prognosis with more than half of them relapsing and dying within 1 year. It is noteworthy that stem cell transplantation was not effective in any of the five cases where it was administered as a salvage therapy for relapsed patients in our cohort. Therefore, the establish-




Figure 4. Outcomes of patients with *MEF2D* fusions. (A) Kaplan-Meier estimates of event-free survival (EFS) of patients with *MEF2D* fusions, and B-other, log-rank $P=0.0306$. (B) Overall survival (OS) for the same as above (log-rank $P=0.0013$).

— B-others
 - - - *MEF2D* fusions
 *, $P=0.0306$ ** , $P=0.0013$

ment of an early diagnostic method and a new therapeutic strategy are necessary for this type of B-ALL. As *ex vivo* sensitivity of xenografted leukemic cells harboring *MEF2D* fusions to HDAC inhibitors has been recently reported,¹⁰ it may offer a plausible therapeutic option for this type of B-ALL. In relation to diagnosis, we have already established a qPCR based detection assay for the frequent *MEF2D* fusions, which may be useful for rapid diagnosis in combination with FISH.

Another interesting observation is that B-ALL cases with *MEF2D* fusions have a characteristic immunophenotype, most typically presenting as CD5- and cytoplasmic μ chain-positive. Although it was previously reported that their immunophenotypes were characterized by weak or absent expression of CD10 and high expression of CD38,¹⁰ we found that CD10 expression was not significantly lower than B-other-ALL, and that CD38 expression level was similar to that of *TCF3-PBX1*-positive cases. Therefore, B-ALL patients with *MEF2D* fusions could be more effectively diagnosed by combining genetic testing with immunophenotyping.

As shown in Table 1, *MEF2D* fusion cases were mutually exclusive of the known risk stratifying chromosomal translocations, at least at the cytogenetic level. Among additional genetic alterations in *MEF2D* fusion cases, we identified a significantly higher frequency of *CDKN2A/CDKN2B* deletions. At initial diagnosis, 7/8 (87.5%) relapsed patients had *CDKN2A/CDKN2B* deletions, while they were seen in only 2/7 (28.6%) non-relapse patients. Thus, the prognostic impact of *CDKN2A/CDKN2B* deletions in *MEF2D* fusion B-ALL needs to be assessed in larger patient cohorts. Interestingly, we also identified that *MEF2D* fusion patients had mutations in genes known to be recurrent in T-ALL; in particular, there was an unexpectedly high frequency of mutations in *PHF6* (8/16). *PHF6* encodes a plant homeodomain (PHD) factor with a proposed role in gene expression control.³⁰ As *PHF6* has been proposed to play a role as a tumor suppressor gene,²⁴ it may participate in the pathogenesis of *MEF2D* fusion B-ALL. Thus, further investigation is needed to clarify this role. Additionally, mutations of genes in the RAS pathway (*NRAS*, *KRAS*, *NF1*, and *PTPN11*) or *IKZF1* alterations were not observed in our cases, in conflict with previous reports.

Our data also showed that *MEF2D* fusion B-ALL has a gene expression signature significantly distinct from other genetic subtypes of B-ALL. Supervised analysis led to separation of *MEF2D* fusion cases into isolated clusters both in PCA and hierarchical clustering, and these clusters were the closest neighbors to *TCF3-PBX1*-positive cases. This observation is consistent with the knowledge that both types of B-ALL tend to show a pre-B ALL immunophenotype, and high-level expression of pre-BCR components, including *IGHG/IGHV5-78* and *IGLL1*, were commonly observed in both subtypes.³¹ However, when we examined the gene expression signatures closely, there were significant differences between them.

It has recently been shown that MEF2 family proteins, including MEF2C and MEF2D, play a critical role in early B-cell differentiation. It was demonstrated that B-cell development was blocked at the pre-B-cell stage in *Mef2c/d*-deficient mice, indicating that they are essential for progression of B-cell precursors (large to small pre-B-cell transition).³² It was further shown that, upon activation *via* pre-B-cell receptor signaling, *Mef2c/d* induces target genes, including *interferon regulatory factor 4* (*Irf4*; stimulates the expression of

Irf4) and *histone deacetylase 5/9* (*Hdac5/9*; encoding the protein known to repress Mef2c activity) as we have summarized schematically in *Online Supplementary Figure S4*. On the other hand, *MEF2D* fusion B-ALL cases exhibited elevated expression of *IRF4* and *HDAC9* with down-regulation of *MEF2C*, although it has been suggested that deregulated expression of the N terminus of MEF2D should induce the up-regulation of the target genes of MEF2C/D, whereas subsequent excess of HDAC9 activity represses MEF2C and its downstream gene expression cascade.¹⁰

Our existing data further revealed that up-regulation of *GATA3* is another gene expression characteristic of *MEF2D* fusion ALL. *GATA3* is a critical transcription factor in early T-cell development³³⁻³⁵ and its transcriptional repression is essential for early B-cell commitment,³⁶⁻³⁷ while it has also been reported that *GATA3* exhibits myeloid-inducing activity in committed B-lymphocytes under the defect of *PAX5* function.³⁸⁻⁴¹ Furthermore, significantly increased *GATA3* mRNA levels were associated with *ZNF384* fusion-positive cases and a higher risk of relapse in childhood B-ALL with the Ph-like phenotype.⁴²⁻⁴³ As our data also indicated down-regulation of *GATA3* expression in *TCF3-PBX1*-positive cases, ectopic overexpression of *GATA3* appears to participate in establishment of a characteristic gene expression signature as well as the biological signature of *MEF2D* fusion ALL distinct from that of other genetic subtypes of B-ALL.

The roles of the *MEF2D* fusion partners in terms of their biological effects within the fusion molecules of *MEF2D* fusion ALL are largely unknown. The genes known to be fused to *MEF2D* in B-ALL have a variety of biological functions. For example, *BCL9* is known to be related to WNT/ β -catenin signaling, whereas the *MEF2D-BCL9* fusion retains only the last one or two exons of the *BCL9* gene, thus lacking the functional domains required for WNT/ β -catenin signaling, suggesting a role other than deregulation of WNT/ β -catenin signaling.¹⁰ *HNRNPUL1* encodes a nuclear RNA-binding protein of the heterogeneous nuclear ribonucleoprotein family that may play a role in nucleocytoplasmic RNA transport and DNA repair,¹⁸ whereas its role as a portion of the fusion molecule in the pathogenesis of B-ALL remains unclear. As *HNRNP1* is a member of the same protein family as *HNRNPUL1*, both *MEF2D-NRNPUL1* and *MEF2D-HNRNP1* likely share the same unknown function.

In conclusion, we have shown that ALL patients harboring *MEF2D* fusion genes possess a characteristic immunophenotype and gene expression signature as well as distinct clinical features, defining them as a distinct genetic subtype among B-other-ALL. Although additional studies are required to elucidate the biological function of the *MEF2D* fusion protein in leukemogenesis, our data has allowed improved characterization of this new B-ALL subtype.

Acknowledgments

The authors would like to thank K. Itagaki, H. Yagi, Y. Katayama, A. Tamura, K. Takeda, K. Hayashi, and the staff of the Laboratory for Genotyping Development, Riken Center for Integrative Medical Sciences for their excellent data management and experimental assistance. We thank all members of the Committees of ALL and of Research and Diagnosis of TCCSG.

Funding

This work was supported in part by a Health and Labour Sciences Research Grant (3rd-term comprehensive 10-year

strategy for cancer control H22-011), the Grant of the National Center for Child Health and Development (26-20), and the Advanced Research for Medical Products Mining Programme of the National Institute of Biomedical Innovation (NIBIO, 10-41, -42, -43, -44, -45), and Biobank Japan project funded by the Ministry of Education, Culture, Sports, Science and Technology

(MEXT) and the Japan Agency for Medical Research and Development (AMED), and the Practical Research for Innovative Cancer Control from AMED.

These funding sources played no role in the collection, analysis, or interpretation of the results, or in the writing of the manuscript and decision to submit it.

References

- Hunger SP, Mullighan CG. Acute lymphoblastic leukemia in children. *N Engl J Med*. 2015;373(16):1541-1552.
- Schwab C, Harrison CJ. Advances in B-cell precursor acute lymphoblastic leukemia genomics. *HemaSphere*. 2018;2(4):e53
- Roberts KG, Morin RD, Zhang J, et al. Genetic alterations activating kinase and cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell*. 2012;22(2):153-166.
- Roberts KG, Li Y, Payne-Turner D, et al. Targetable kinase-activating lesions in Ph-like acute lymphoblastic leukemia. *N Engl J Med*. 2014;371(11):1005-1015.
- Gocho Y, Kiyokawa N, Ichikawa H, et al. A novel recurrent EP300-ZNF384 gene fusion in B-cell precursor acute lymphoblastic leukemia. *Leukemia*. 2015;29(12):2445-2448.
- Liu YF, Wang BY, Zhang WN, et al. Genomic profiling of adult and pediatric B-cell acute lymphoblastic leukemia. *EBioMedicine*. 2016;8:173-183.
- Yasuda T, Tsuzuki S, Kawazu M, et al. Recurrent DUX4 fusions in B cell acute lymphoblastic leukemia of adolescents and young adults. *Nat Genet*. 2016;48(5):569-574.
- Lilljebjörn H, Henningson R, Hyrenius-Wittsten A, et al. Identification of ETV6-RUNX1-like and DUX4-rearranged subtypes in paediatric B-cell precursor acute lymphoblastic leukaemia. *Nat Commun*. 2016;7:11790.
- Zhang J, McCastlain K, Yoshihara H, et al. Deregulation of DUX4 and ERG in acute lymphoblastic leukemia. *Nat Genet*. 2016;48(12):1481-1489.
- Gu Z, Churchman M, Roberts K, et al. Genomic analyses identify recurrent MEF2D fusions in acute lymphoblastic leukaemia. *Nat Commun*. 2016;7:13331.
- Hirabayashi S, Ohki K, Nakabayashi K, et al. ZNF384-related fusion genes define a subgroup of childhood B-cell precursor acute lymphoblastic leukemia with a characteristic immunotype. *Haematologica*. 2017;102(1):118-129.
- Prima V, Gore L, Caires A, et al. Cloning and functional characterization of MEF2D/DAZAP1 and DAZAP1/MEF2D fusion proteins created by a variant t(1;19)(q23;p13.3) in acute lymphoblastic leukemia. *Leukemia*. 2005;19(5):806-813.
- Lilljebjörn H, Ågerstam H, Orsmark-Pietras C, et al. RNA-seq identifies clinically relevant fusion genes in leukemia including a novel MEF2D/CSF1R fusion responsive to imatinib. *Leukemia*. 2014;28(4):977-979.
- Suzuki K, Okuno Y, Kawashima N, et al. MEF2D-BCL9 fusion gene is associated with high-risk acute B-cell precursor lymphoblastic leukemia in adolescents. *J Clin Oncol*. 2016;34(28):3451-3459.
- Breitbart RE, Liang CS, Smoot LB, Laheru DA, Mahdavi V, Nadal-Ginard B. A fourth human MEF2 transcription factor, hMEF2D, is an early marker of the myogenic lineage. *Development*. 1993;118(4):1095-1106.
- Mao Z, Bonni A, Xia F, Nadal-Vicens M, Greenberg M E. (1999) Neuronal activity-dependent cell survival mediated by transcription factor MEF2. *Science*. 1999;286(5440):785-790.
- Miska EA, Karlsson C, Langley E, Nielsen SJ, Pines J, Kouzarides T. HDAC4 deacetylase associates with and represses the MEF2 transcription factor. *EMBO J*. 1999; 18(18): 5099-5107.
- Honoré B, Rasmussen HH, Vorum H, et al. Heterogeneous nuclear ribonucleoproteins H, H', and F are members of a ubiquitously expressed subfamily of related but distinct proteins encoded by genes mapping to different chromosomes. *J Biol Chem*. 1995; 270(48):28780-28789.
- Takahashi H, Kajiwara R, Kato M, et al. Treatment outcome of children with acute lymphoblastic leukemia: the Tokyo Children's Cancer Study Group (TCCSG) Study L04-16. *Int J Hematol*. 2018;108:98-108.
- Masuzawa A, Kiyotani C, Osumi T, et al. Poor responses to tyrosine kinase inhibitors in a child with precursor B-cell acute lymphoblastic leukemia with SNX2-ABL1 chimeric transcript. *Eur J Haematol*. 2014; 92(3):263-267.
- Kobayashi K, Mitsui K, Ichikawa H, et al. ATF7IP as a novel PDGFRB fusion partner in acute lymphoblastic leukaemia in children. *Br J Haematol*. 2014;165(6):836-841.
- McPherson A, Hormozdiari F, Zayed A, et al. deFuse: an algorithm for gene fusion discovery in tumor RNA-Seq data. *PLoS Comput Biol*. 2011;7(5):e1001138.
- Weng AP, Ferrando AA, Lee W, et al. Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia. *Science*. 2004;306(5694):269-271.
- Van Vlierberghe P, Palomero T, Khiabani H, et al. PHF6 mutations in T-cell acute lymphoblastic leukemia. *Nat Genet*. 2010; 42(4):338-342.
- Wang Q, Qiu H, Jiang H, et al. Mutations of PHF6 are associated with mutations of NOTCH1, JAK1 and rearrangement of SET-NUP214 in T-cell acute lymphoblastic leukemia. *Haematologica*. 2011;96(12): 1808-1814.
- Spinella JF, Cassart P, Richer C, et al. Genomic characterization of pediatric T-cell acute lymphoblastic leukemia reveals novel recurrent driver mutations. *Oncotarget*. 2016;7(40):65485-65503.
- Royet J, Bouwmeester T, Cohen SM. Notchless encodes a novel WD40-repeat-containing protein that modulates Notch signaling activity. *EMBO J*. 1998; 17(24):7351-7360.
- Carroll AJ, Crist WM, Parnley RT, Roper M, Cooper MD, Finley WH. Pre-B cell leukemia associated with chromosome translocation 1;19. *Blood*. 1984;63(3):721-724.
- Borowitz MJ, Hunger SP, Carroll AJ, et al. Predictability of the t(1;19)(q23;p13) from surface antigen phenotype: implications for screening cases of childhood acute lymphoblastic leukemia for molecular analysis: a Pediatric Oncology Group study. *Blood*. 1993;82(4):1086-1091.
- Lower KM, Turner G, Kerr BA, et al. Mutations in PHF6 are associated with Börjeson-Forsman-Lehmann syndrome. *Nat Genet*. 2002;32(4):661-665.
- Chen D, Zheng J, Gerasimcik N, et al. The expression pattern of the Pre-B cell receptor components correlates with cellular stage and clinical outcome in acute lymphoblastic leukemia. *PLoS One*. 2016; 11(9):e0162638.
- Herglotz J, Unrau L, Hauschildt F, et al. Essential control of early B-cell development by Mef2 transcription factors. *Blood*. 2016;127(5):572-581.
- Landry DB, Engel JD, Sen R. Functional GATA 3 binding sites within murine CD8 alpha upstream regulatory sequences. *J Exp Med*. 1993;178(3):941-949.
- Ting CN, Olson MC, Barton KP, Leiden JM. Transcription factor GATA-3 is required for development of the T-cell lineage. *Nature*. 1996;384(6608):474-478.
- Gao J, Chen YH, Peterson LC. GATA family transcriptional factors: emerging suspects in hematologic disorders. *Exp Hematol Oncol*. 2015;4:28.
- Banerjee A, Northrup D, Boukarabila H, Jacobsen SE, Allman D. Transcriptional repression of Gata3 is essential for early B cell commitment. *Immunity*. 2013; 38(5):930-42.
- Somasundaram R, Prasad MA, Ungerback J, Sigvardsson M. Transcription factor networks in B-cell differentiation link development to acute lymphoid leukemia. *Blood*. 2015;126(2):144-152.
- Chen D, Zhang G. Enforced expression of the GATA-3 transcription factor affects cell fate decisions in hematopoiesis. *Exp Hematol*. 2001;29(8):971-980.
- Ku CJ, Hosoya T, Maillard I, Engel JD. GATA 3 regulates hematopoietic stem cell maintenance and cell cycle entry. *Blood*. 2012;119(10):2242-2251.
- Frelin C, Herrington R, Janmohamed S, et al. GATA 3 regulates the self renewal of long term hematopoietic stem cells. *Nat Immunol*. 2013;14(10):1037-1044.
- Heavey B, Charalambous C, Cobaleda C, Busslinger M. Myeloid lineage switch of Pax5 mutant but not wild-type B cell progenitors by C/EBPalpha and GATA factors. *EMBO J*. 2003;22(15):3887-3897.
- Perez-Andreu V, Roberts KG, Harvey RC, et al. Inherited GATA3 variants are associated with Ph-like childhood acute lymphoblastic leukemia and risk of relapse. *Nat Genet*. 2013;45(12):1494-1498.
- Perez-Andreu V, Roberts KG, Xu H, et al. A genome-wide association study of susceptibility to acute lymphoblastic leukemia in adolescents and young adults. *Blood*. 2015; 125(4):680-686.