

## Clinical characteristics of patients with anti-TIF1- $\gamma$ antibodies

Anna Masiak<sup>1</sup>, Julia Kulczycka<sup>2</sup>, Zenobia Czuszyńska<sup>1</sup>, Zbigniew Zdrojewski<sup>1</sup>

<sup>1</sup>Department of Internal Diseases, Connective Tissue Diseases and Geriatrics, University of Gdansk, Gdansk, Poland

<sup>2</sup>Clinical Immunology i Transplantology Unit, UCK Gdańsk, University of Gdansk, Gdansk, Poland

### Abstract

**Objectives:** Inflammatory myopathies are a group of idiopathic, heterogeneous systemic diseases affecting predominantly skeletal muscles, though they can also involve the skin and internal organs. The association between cancer and idiopathic inflammatory myopathies, particularly dermatomyositis, which is termed cancer-associated myositis (CAM), has been reported in the medical literature. A newly described autoantibody to a 155-kDa nuclear protein, identified as transcription intermediary factor 1-gamma (TIF1- $\gamma$ ), has proven useful for cancer screening in patients with dermatomyositis.

**Material and methods:** Based on our database of laboratory results, between November 2014 and January 2016, we found 80 patients with a positive autoimmune inflammatory myopathy immunoblot profile.

**Results:** Eleven of 80 patients revealed the presence of anti-TIF1- $\gamma$  antibodies: 8 women and 3 men with average age 54.2 years. Dermatomyositis (DM) was diagnosed in 6 cases, polymyositis in 1 case, myositis limited to ocular muscles and rhabdomyolysis in 1 case each, and undifferentiated connective tissue disease in 2 cases. Neoplasm was found in 4 cases. All of those patients had DM. The average time between DM and diagnosis of neoplasm was 7.5 months (from 1 to 18 months).

**Conclusions:** The association between cancer and idiopathic inflammatory myopathies, particularly DM, is well known, and cancer screening should be obligatory in such patients. So far there is no consensus as to the method or frequency with which patients with an idiopathic inflammatory myopathy should be tested to rule out neoplasm. Detection of anti-TIF1- $\gamma$  antibodies in patients with DM gives the clinicians the very important suggestion of CAM. It seems reasonable that these patients should have more detailed and often repeated differential diagnostics.

**Key words:** neoplasm, anti-TIF1- $\gamma$ , clinical characteristic.

### Introduction

Inflammatory myopathies are a group of idiopathic, heterogeneous systemic diseases affecting predominantly skeletal muscles, though they can also involve the skin and internal organs. In adults they include mainly polymyositis (PM), dermatomyositis (DM) and inclusion body myositis (IBM) [1, 2]. The association between cancer and idiopathic inflammatory myopathies, particularly dermatomyositis, which is termed cancer-associated myositis (CAM), has been reported in the medical literature. A newly described autoantibody to a 155-kDa

nuclear protein, identified as transcription intermediary factor 1- $\gamma$  (TIF1- $\gamma$ ), has proven useful for cancer screening in patients with dermatomyositis.

The aim of this study was to evaluate the clinical picture of patients with the presence of TIF1- $\gamma$  antibodies in one center.

### Material and methods

We searched our database of laboratory results with the aim of finding patients with a positive autoimmune inflammatory myopathy profile (EUROLINE, EUROIM-

---

#### Address for correspondence:

Dr Anna Masiak, Department of Internal Diseases, Connective Tissue Diseases and Geriatrics, University of Gdansk, Dębinki 7, 80-001 Gdansk, Poland, e-mail: anna.masiak@wp.pl

Submitted: 10.02.2016; Accepted: 26.02.2016

MUN). Between November 2014 and January 2016, we found 80 such patients.

## Results

In a group of 80 Caucasian patients, 11 revealed the presence of anti-TIF1- $\gamma$  antibodies, but in 3 cases anti-TIF1- $\gamma$  were only low positive. There were 8 women and 3 men with average age 54.2 years (from 27 to 84 years). Dermatomyositis was diagnosed in 6 cases, polymyositis in 1 case, myositis limited to ocular muscles in 1 case and rhabdomyolysis in 1 case. In 2 patients with low positive antibodies undifferentiated connective tissue disease was recognized. In this group of patients with anti-TIF1- $\gamma$  antibodies neoplasm was found in 5 cases. All of those patients had DM. The average time between DM and diagnosis of neoplasm was 7.5 months (from 1 to 18 months). In 2 cases neoplasm was recognized concurrently with myositis. In 2 other cases, the neoplasm was detected 8 and 18 months after diagnosis of myositis. None of the patients had lung involvement, but 3 of them had severe skin changes and 2 of them dysphagia. Clinical characteristics of patients with TIF1- $\gamma$  positivity are presented in Table I.

In some patients the coexistence of anti-TIF1- $\gamma$  and other antibodies was found: in 2 cases anti-Ro52, anti-MDA5 in another 2 cases, and anti-OJ and anti-Ku in individual cases. In patients with low positive TIF1- $\gamma$ , anti-Mi2 and anti-DFS70 were found. It is worth pointing out that in patients with neoplasm, anti-TIF1- $\gamma$  were highly positive and were the only positive antibodies. The immune profile of patients with the presence of anti-TIF1- $\gamma$  antibodies is presented in Table II.

## Discussion

The association between cancer and idiopathic inflammatory myopathies is well described in the literature [3, 4]. The incidence of cancer in published series of patients with idiopathic inflammatory myopathy ranges from 9% to 42%. A great variety of cancer types may occur: the most frequent are ovarian, breast, lung, gastric, colorectal tumors and lymphomas in dermatomyositis, and lung, urinary bladder cancers and lymphomas in polymyositis [5]. Cancer can develop before, concurrently, or subsequent to the onset of idiopathic inflammatory myopathy, but is usually recognized within 3 years of myositis diagnosis, with most diagnoses within 12 months [5]. The mean period in our group of patients was 7.5 months (from 1 to 18 months). In 2 cases neoplasm was recognized concurrently with myositis.

Malignant disease is also one of the main causes of mortality in patients with inflammatory myopathies. Thus cancer screening is obligatory in such patients, but there is no consensus as to the method or frequency

with which patients with an idiopathic inflammatory myopathy should be tested to rule out neoplasm during the follow-up. Clinical research studies have shown that older age, male sex, dysphagia, and skin manifestations, such as skin necrosis, periungual erythema, and the 'V' or 'shawl' sign, are associated with occult malignancy in patients with myositis. Refractory or recurrent disease has also been related to cancer-associated myositis. In contrast, the presence of interstitial lung disease seems to be protective for the development of cancer. None of our patients with cancer had lung involvement. An absence of autoantibodies seems to be predictive of a high risk of occult malignancy, whereas the presence of anti-synthetase antibodies seems to have a protective value against cancer.

In 2006, Targoff et al. [6] described a novel autoantibody against a 155-kDa protein in a large series of patients with myositis. This antibody was then identified as human TIF1- $\gamma$  [7, 8]. It seemed to be a myositis-specific autoantibody, as it was not detected in other autoimmune diseases or in other noninflammatory myopathies [7]. Among the 85 patients with idiopathic inflammatory myopathy, Trallero-Araguás et al. [9] found anti-p155 (TIF1- $\gamma$ ) autoantibody in 10 of the 16 patients classified as having CAM. They found that in DM, the negative and positive predictive value of presence of the TIF1- $\gamma$  autoantibody for a diagnosis of CAM was 92% and 66.7%, respectively. No myositis-specific autoantibodies were detected in patients positive for anti-p155 (TIF1- $\gamma$ ) autoantibody. In contrast, myositis-associated autoantibodies were identified in some anti-TIF1- $\gamma$ -positive patients [9]. None of the patients with CAM in our group had myositis-specific and only one had myositis-associated autoantibodies. Chinoy et al. [10] found the coexistence of TIF1- $\gamma$  antibodies and other myositis-specific antibodies in only 3 patients, 2 of whom had anti-Mi-2 autoantibodies. Some authors have suggested an association between anti-TIF1- $\gamma$  autoantibody and a specific clinical phenotype in adult DM, characterized by severe skin disease [11, 12]. Three of our patients had such a clinical presentation (Fig. 1).

Vincent et al. [13] in their study proposed that TIF1- $\gamma$  is involved in the transforming growth factor  $\beta$  (TGF- $\beta$ ) signaling pathway, which is inactivated in some malignancies [13]. TIF1- $\gamma$  acts on TGF- $\beta$  and is essential in chromatin-mediated transcriptional regulation, functioning as a positive (agonist) or negative (antagonist) regulator of this pathway, which seems to be related to carcinogenesis.

Saleva-O'Callaghan et al. in their meta-analysis concluded that testing positive for TIF1- $\gamma$  has an 18-fold higher association with cancer than TIF1- $\gamma$ -negative status, that is, 93 of every 100 negative dermatomyositis patients will

Table 1. Clinical manifestations in our patients who exhibited anti-TIF1- $\gamma$  positivity

N	Gender	Age at disease onset	Diagnosis	Observation (months)	Cancer	Death	Smoking	Muscle symptoms	Skin lesions	Skeletal symptoms	Lung involvement	GI involvement	CK [U/l]	LDH [U/l]	AST [U/l]
1	F	33	NDCTD	9	no	no	no	mild	no	arthralgia	no	no	65	170	27
2	F	51	limited ocular myositis	5	no	no	no	localized	no	no	no	no	34	122	19
3	M	55	DM	1	stomach	no data	yes	no	swelling of the eyelids	no	no	no	2337	NP	149
4	F	27	no	4	no	no	no	mild	no	no	no	no	1414	263	87
5	F	59	PM	102	no	no	no	severe	no	severe	no	dysphagia	25630	365	720
6	F	57	DM	18	pulmonitis	no	yes	severe	heliotrope rash, V-sign, erythematous and oedematous rashes on face, rashes on the trunk	no	no	no	305	258	31
7	M	74	rhabdomyolysis	1	no	no	yes	severe	no	no	no	no	145924	4221	1767
8	M	84	DM	1	no	yes	no	severe	erythematous rashes Gottron's signs V-sign, scarf-sign periungual erythema	no	fibrotic changes	dysphagia	307	349	65
9	F	41	DM	9	no	no	no	no	erythematous and oedematous rashes on face, rashes on the trunk; Raynaud	arthralgia	no	no	101	328	46
10	F	53	DM	8	ovarian	yes	no	severe	erythematous rashes on face, heliotrope rash, V-sign, scarf-sign; arm and leg oedema	no	no	dysphagia	17288	871	504
11	F	62	DM	3	colon	yes	no	severe	swelling of the eyelids, heliotrope rash, V-sign, scarf-sign	no	no	dysphagia	1280	333	75

DM – dermatomyositis; NDCTD – undifferentiated connective tissue disease; NP – not performed

**Table II.** Immune profile of patients with the presence of anti-TIF1-γ antibodies

No.	Sex	Antinuclear antibodies	TIF1-γ	Mi-2	Ku	PM100	PM75	Jo1	SRP	PL7	PL12	EJ	OJ	MDA5	Other
1	F	1 : 2560	low positive	+++	+	-	-	-	-	-	-	-	-	-	-
2	F	NP	+	-	-	-	-	-	-	-	-	-	-	-	-
3	M	NP	+++	-	-	-	-	-	-	-	-	-	-	-	-
4	F	1 : 2560	low positive	-	-	-	-	-	-	-	-	-	-	+	DFS70
5	F	1 : 320	low positive	-	-	-	-	-	-	-	-	-	-	-	-
6	F	1 : 2560	+++	-	-	-	-	-	-	-	-	-	-	-	Ro52
7	M	1280	+	-	-	-	-	-	-	-	-	-	-	+	-
8	M	2560	+++	-	-	-	-	-	-	-	-	-	+++	-	Ro52
9	F	1280	+	-	-	-	-	-	-	-	-	-	-	-	-
10	F	2560	++	-	-	-	-	-	-	-	-	-	-	-	-
11	F	negative	+++	-	-	-	-	-	-	-	-	-	-	-	-

NP – not performed; Mi-2 – anti-helicase protein antibodies; PM100 – anti-human exosome complex PM100; PM75 – anti-human exosome complex PM75; SRP – anti-signal recognition particle antibodies; PL7 – anti-threonyl tRNA; PL12 – anti-alanyl tRNA; EJ – anti-glycyl tRNA; OJ – anti-isoleucyl tRNA; MDA5 – melanoma; differentiation-associated gene 5 antibodies; DFS70 – autoantibodies to dense fine speckles; Anti-Jo1 – anti-histidyl tRNA



**Fig. 1.** Clinical presentation of DM.

not develop cancer. They also proposed a reasonable approach for detecting occult malignancy in myositis [14]. Testing for the anti-TIF1-γ antibody alone had 50% sensitivity and 96% specificity for detection of CAM. The risk of developing cancer-associated DM in anti-TIF1-γ-positive patients has also been assessed by a systematic review and meta-analysis on six cohort studies including a total of 312 adult patients with DM [15]. It was concluded that when the test for anti-TIF1-γ determination is negative, it reasonably rules out the presence of associated cancer. It was suggested, however, that it would be desirable to perform a single PET/CT study.

**Conclusions**

The association between cancer and idiopathic inflammatory myopathies, particularly dermatomyositis, is well known, and cancer screening should be obligatory in such patients. As yet there is no consensus as to the method or frequency with which patients with an idiopathic inflammatory myopathy should be tested to rule out neoplasm. Detection of anti-TIF1-γ antibodies in patient with DM gives the clinicians the very important suggestion of CAM. It seems reasonable that these patients should have more detailed and often repeated differential diagnostics.

*The authors declare no conflict of interest.*

**References**

1. Dobloug C, Garen T, Bitter H, et al. Prevalence and clinical characteristics of adult polymyositis and dermatomyositis; data from a large and unselected Norwegian cohort. *Ann Rheum Dis* 2015; 74: 1551-1556.
2. Vánca A, Gergely L, Ponyi A, et al. Myositis-specific and myositis-associated antibodies in overlap myositis in comparison to primary dermatopolymyositis: Relevance for clinical classi-

- fication: Retrospective study of 169 patients. *Joint Bone Spine* 2010; 77: 125-130.
3. Huang YL, Chen YJ, Lin MW, et al. Malignancies associated with dermatomyositis and polymyositis in Taiwan: a nationwide population-based study. *Br J Dermatol* 2009; 161: 854-860.
  4. Antiochos BB, Brown LA, Li Z, et al. Malignancy is associated with dermatomyositis but not polymyositis in Northern New England, USA. *J Rheumatol* 2009; 36: 2704-2710.
  5. Dankó K, Ponyi A, Molnar AP, et al. Paraneoplastic myopathy. *Curr Opin Rheumatol* 2009; 21: 594-598.
  6. Targoff IN, Mamyrova G, Trieu EP, et al. A novel autoantibody to a 155-kd protein is associated with dermatomyositis. *Arthritis Rheum* 2006; 54: 3682-3689.
  7. Targoff IN, Trieu EP, Levy-Neto M, et al. Autoantibodies to transcriptional intermediary factor-1 gamma (TIF-1gamma) in dermatomyositis [abstract]. *Arthritis Rheum* 2006; 54: S518.
  8. Fujimoto M, Hamaguchi Y, Kaji K, et al. Myositis-specific anti-155/140 autoantibodies target transcription intermediary factor 1 family proteins. *Arthritis Rheum* 2012; 64: 513-522.
  9. Trallero-Araguás E, Labrador-Horrillo M, Selva-O'Callaghan A, et al. Cancer-associated myositis and anti-p155 autoantibody in a series of 85 patients with idiopathic inflammatory myopathy. *Medicine* 2010; 89: 47-52.
  10. Chinoy H, Fertig N, Oddis CV, et al. The diagnostic utility of myositis autoantibody testing for predicting the risk of cancer-associated myositis. *Ann Rheum Dis* 2007; 66: 1345-1349.
  11. Kaji K, Fujimoto M, Hasegawa M, et al. Identification of a novel autoantibody reactive with 155 and 140 kDa nuclear proteins in patients with dermatomyositis: an association with malignancy. *Rheumatology (Oxford)* 2007; 46: 25-28.
  12. Gunawardena H, Wedderburn LR, North J, et al. Clinical associations of autoantibodies to a p155/140 kDa doublet protein in juvenile dermatomyositis. *Rheumatology (Oxford)* 2008; 47: 324-328.
  13. Vincent DF, Yan KP, Treilleux I, et al. Inactivation of TIF1gamma cooperates with Kras to induce cystic tumors of the pancreas. *PLoS Genet* 2009; 5: e1000575.
  14. Selva-O'Callaghan A, Trallero-Araguás E, Grau-Junyent JM, Labrador-Horrillo M. Malignancy and myositis: novel autoantibodies and new insights. *Curr Opin Rheumatol* 2010; 22: 627-632.
  15. Trallero-Araguás E, Rodrigo-Pendás JÁ, Selva-O'Callaghan A, et al. Usefulness of anti-p155 autoantibody for diagnosing cancer-associated dermatomyositis. A systematic review and meta-analysis. *Arthritis Rheum* 2012; 64: 523-532.