

Clinical Effectiveness of School Screening for Adolescent Idiopathic Scoliosis: A

Large Population-Based Retrospective Cohort Study

Keith D. K. Luk*, C. F. Lee*, Kenneth M. C. Cheung*, Jack C. Y. Cheng, Bobby K.

W. Ng, T. P. Lam, K. H. Mak, Paul S. F. Yip, Daniel Y. T. Fong*

* These authors contributed equally to this paper.

Author affiliations

Department of Orthopaedics and Traumatology, The University of Hong Kong,

Pokfulam, Hong Kong SAR, China

Prof. Keith Dip Kei Luk, M.Ch. Orth. (e-mail: hrmoldk@hku.hk)

Prof. Kenneth Man Chee Cheung, M.D. (e-mail: cheungmc@hku.hk)

Department of Nursing Studies, The University of Hong Kong, Pokfulam, Hong Kong

SAR China

Mr. Chun Fan Lee, M.Phil. (e-mail: fanlee@hku.hk)

Dr. Daniel Yee Tak Fong, Ph.D. (e-mail: dtyfong@hku.hk)

Department of Orthopaedics and Traumatology, The Chinese University of Hong

Kong, Hong Kong SAR, China

Prof. Jack Chun Yiu Cheng, M.D. (e-mail: jackcheng@uab.cuhk.edu.hk)

Dr. Tsz Ping Lam, M.D. (e-mail: tplam@ort.cuhk.edu.hk)

Department of Orthopaedics and Traumatology, Prince of Wales Hospital, Hong Kong
SAR, China

Dr. Bobby Kin Wah Ng, M.D. (e-mail: bobng@ort.cuhk.edu.hk)

Department of Health, Hong Kong SAR, China

Dr. Kwok Hang Mak, M.Sc. PH (e-mail: drkhmak@dh.gov.hk)

Department of Social Work and Social Administration, The University of Hong Kong,
Hong Kong SAR, China

Prof. Paul Siu Fai Yip, Ph.D. (e-mail: sfpyip@hkucc.hku.hk)

Correspondence to:

Daniel Y. T. Fong

Postal address: Department of Nursing Studies, The University of Hong Kong, 4/F

William M.W. Mong Block, Faculty of Medicine Building, 21 Sassoon Road,

Pokfulam, Hong Kong (dytfong@hku.hk)

Telephone: 852-2819 2645

Fax number: 852-2872 6079

E-mail: dytfong@hku.hk

Acknowledgement

We thank Dr. LY Tse and Dr. SP Mak, for their valuable contribution in setting up the Hong Kong scoliosis screening program. We also thank Ms SY Lam for coordinating the data collection and the registered nurses and nursing students who collected the records of patients with scoliosis from the two specialist hospitals.

Financial support

The work was financially supported by the Central Policy Unit of the Government of the Hong Kong Special Administrative Region and the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No.: HKU 7006-PPR-20051).

ABSTRACT

Study Design. Retrospective cohort study

Objective. To evaluate the clinical effectiveness of school scoliosis screening using a large and long-term-followed cohort of students in Hong Kong.

Summary of Background Data. School screening for adolescent idiopathic scoliosis (AIS) has been criticized as resulting in over-referrals for radiography and having low predictive values. Indeed, all but one previous retrospective cohort studies had no follow-up assessments of students until their skeletal maturity, leaving any late-developed curves undetected. The one study that completed this follow-up was well conducted but had low precisions due to its small sample size.

Methods. A total of 157,444 students were eligible for a biennial scoliosis screening, and their screening results and medical records up to 19 years of age were available. Screening tests included a forward bending test (FBT), angle of trunk rotation (ATR), and moiré topography for those who showed signs of AIS. Students with an ATR $\geq 15^\circ$, ≥ 2 moiré lines, or significant clinical signs were referred for radiography and had their Cobb angle measured.

Results. Of the 115,190 screened students in the cohort, 3,228 (2.8%, 95% CI = 2.7% to 2.9%) were referred for radiography. At the final follow-up, the positive predictive values were 43.6% (41.8% to 45.3%) for a Cobb angle $\geq 20^\circ$ and 9.4%

(8.4% to 10.5%) for needing treatment, while the sensitivities were 88.1% (86.4% to 89.6%) and 80.0% (75.6% to 83.9%), respectively.

Conclusions. This is the largest study that has demonstrated that school scoliosis screening in Hong Kong is predictive and sensitive with a low referral rate.

Screening should thus be continued in order to facilitate early administration of conservative treatments.

Key words: adolescent idiopathic scoliosis, school screening program, clinical effectiveness

INTRODUCTION

Scoliosis is a lateral deviation of the spine.¹ Although the disorder has been recognized for decades, the etiology is unknown for over 80% of patients and is therefore labeled as idiopathic. Apart from the cosmetic concerns, patients with severe spinal deformity may also suffer from a higher risk of mortality or morbidity.² Aggressive spinal fusion is currently the only treatment option for severe curves, but if they are detected early, curve progression may be prevented with bracing. Hence, because the majority of spinal curves are detectable during adolescence, school screening for early detection of adolescent idiopathic scoliosis (AIS) was initiated. The first program was started in Delaware in the late 1950s,³ and other programs were subsequently started elsewhere, either by legislation or voluntarily.⁴⁻¹²

The use of school scoliosis screening remains controversial. Some programs were discontinued, perhaps due to the unfavorable recommendations by some professional bodies.¹³ In particular, the United States Preventive Services Task Force (USPSTF) stated in 1996 that there was insufficient evidence to either recommend or refute routine screening for AIS. In 2004, the USPSTF advised against screening on the grounds of reasonable evidence of unnecessary brace prescriptions and referrals for specialty care.^{14, 15} This policy change was criticized as being based not on new evidence but on a change of rating methodology on the

available evidence.¹⁶ In contrast, the American Academy of Orthopaedic Surgeons, the Scoliosis Research Society, the Pediatric Orthopaedic Society of North America, and the American Academy of Pediatrics have continually supported scoliosis screening for detecting reversible spinal curves before they progress.¹⁶

The controversy over school scoliosis screening was due mainly to over-referral of students who do not require follow-up or treatment for radiography,³ leading to a positive predictive value (PPV) as low as 0.5% for identifying curves $\geq 20^\circ$.⁶ Indeed, there was a large variation across studies, with a reported PPV of 64%.¹⁷ The variation is likely due to the diversity in study design, referral criteria, screening tests used, frequency of screening, and duration of follow-up.¹⁸ In particular, all studies except one did not have all screened students followed until skeletal maturity. Insufficient follow-up may lower the PPV, since referred students may at first show insignificant curves but later progress. Moreover, whether or not scoliosis develops during adolescence is not known in all screened students. Hence, the sensitivity and specificity of the tests (which are measures of screening accuracy robust to the disorder prevalence) cannot be obtained.¹⁹ The exception was a retrospective cohort study conducted in Rochester, US.¹¹ It followed 2,242 screened students until they were 19 years old or they left school. In the 68 students referred for radiography, the PPV and sensitivity for identifying curves $\geq 20^\circ$ were 17.4% (95% CI = 10.3% to

26.7%) and 64.0% (42.5% to 82.0%), respectively. These amounted to errors of 9.3% and 21.5%, which are large because of the small sample size. Therefore, we needed a large retrospective cohort study with a sufficient sample size to ensure proper and reliable evaluation of school scoliosis screening.

In Hong Kong, school screening for AIS was introduced in 1995 by the Department of Orthopaedics and Traumatology of the University of Hong Kong in collaboration with the Duchess of Kent Children's Hospital and the Department of Health (DH). By 2006, 886,906 students had been screened. We thus aimed to determine the clinical effectiveness of school screening for AIS.

METHODS

Study Design

This was a retrospective cohort study. Students in grade 5 (mostly 10 years old) in the academic years of 1995/96 or 1996/97 were included. For each eligible student, screening history and all medical records related to scoliosis were traced until they reached 19 years of age. The study was approved by the IRBs of the investigating universities and the relevant health authorities.

School Scoliosis Screening Program

The screening program for AIS in Hong Kong was administered by the

Department of Health using a standardized protocol designed by the Department of Orthopaedics and Traumatology of the University of Hong Kong. Participation in the screening was voluntary, and students in grade 5 or aged 10 years or above were eligible for screening until they either reached grade 13 or were 19 years old. The flowchart for the screening protocol is shown in Figure 1.

AIS screening was first performed at the Student Health Service Centres (SHSC) using the FBT and measurement of the angle of trunk rotation (ATR) using a scoliometer.^{20,21} Students in grades 5, 7 and 9 (mostly 10, 12 and 14 years old, respectively) were screened by trained doctors or trained registered nurses. Students with an ATR between 0° and 2° had the tests repeated biennially, and those with an ATR of 3° or 4° repeated them annually. When students had an ATR between 5° and 14° or obvious signs of trunk or shoulder asymmetry, or there was special concern from medical staff or parents, they were further evaluated in a Special Assessment Centre (SAC). Students with an ATR $\geq 15^\circ$ were referred directly to one of the two specialist hospitals that manage spinal deformities, in which they would have X-ray examinations of the whole spine.

In the SAC, the students were assessed by a specially trained doctor using ATR and moiré topography. Moiré topography is a biostereometric technique that projects contour lines, or moiré fringes, on a subject's back.²² A moiré photograph is

then taken, from which the number of fringes that deviate from symmetry between the left and right sides are counted; this reflects the severity of the back deformity. This method was reported to be more sensitive than conventional clinical screening.²³ In the current protocol, students who had <1 moiré line were referred back to the SHSC for yearly screening but were referred to the SAC again if their ATR deteriorated by 1° or more within one year. Students with 1 to <2 moiré line differences underwent a repeat assessment every 6 to 12 months. For students who had ≥2 moiré line differences or showed significant clinical signs, including uneven shoulder height, pelvic tilt, rib or loin hump, or a scapular prominence and/or truncal shift, a standing posteroanterior X-ray of the whole spine was taken, from which the Cobb angle was measured. Students with a Cobb angle under 20° had an ATR and moiré assessments repeated every 6 to 12 months and had X-rays repeatedly taken when they showed a deterioration of 1 or more moiré lines. Students who had a Cobb angle ≥20° were referred to a specialist hospital for follow-up and treatment.

In summary, students who resulted an ATR ≥15°, ≥2 moiré lines, or showed significant clinical signs of scoliosis would be referred for radiography, and those who were found a Cobb angle ≥20° would be followed-up in a specialist hospital until skeletal maturity.

Treatment and follow-up of AIS

Students referred to a specialist hospital received a standing posteroanterior X-ray and were assessed by orthopedists. All patients were followed up every 3 to 6 months and were observed or treated according to the severity of the spinal curvature, the rapidity of progression and their skeletal maturity. Immature patients with curve progression of at least 5° or with a Cobb angle between 30° and 45° were required to wear a brace for prevention of progression. Patients with a Cobb angle over 45° were offered surgical correction.

Data Collection

For each student in our cohort, the demographic information, school grade, date and results of tests performed at each visit and the Cobb angle measurements were obtained from SHSC and SAC. For those who visited the two specialist hospitals, the date, Cobb angle and body height were measured at each follow-up visit, and the type of treatment (brace or operation) was recorded. Students who did not take part in the screening program but were diagnosed with AIS and referred from other sources were also identified from records at the two specialist hospitals. They were included in the cohort if they were born in 1985 or 1986, since they should have been in grade 5 in 1995/96 or 1996/97.

Statistical Analysis

Based on the students referred for radiography, the prevalence and measures of

clinical effectiveness were estimated for different conditions, including different spinal curvatures and treatment. These measures included the PPV, the negative predictive value (NPV), and the sensitivity and specificity. For detecting a condition, the PPV was the proportion of students with the condition in those referred by screening and NPV was the proportion of students without the condition in those not referred. The sensitivity was the proportion of students who had the condition correctly detected by screening, and specificity was the proportion of students without the condition not referred by screening. There were students referred for radiography who did not show up for the assessment. They were considered to be non-scoliotic, which is conservative for the estimation of the PPV and sensitivity.

Per protocol, students were referred for radiography when they had an ATR $\geq 15^\circ$, ≥ 2 moiré lines or significant clinical signs. Unlike the ATR and moiré topography, clinical signs cannot be objectively measured, but were subjectively judged by the screeners. Moreover, some students who had borderline screening results had visited a specialist hospital by themselves and might have eventually met the referral criteria had they stayed in the screening program. To assess the impact of the use of clinical signs for referral and the borderline cases on the clinical effectiveness, we performed a sensitivity analysis on four groups of students. Group A comprised students who were referred for radiography with an ATR $\geq 15^\circ$ or ≥ 2 moiré lines (objective referral

criteria) only. Group B1 consisted of students who had 1 to <2 moiré lines and were referred because they had significant clinical signs of scoliosis. Group B2 consisted of those who had 1 to <2 moiré lines but had visited a specialist hospital without referral and did not have clinical signs. Students in these two groups might eventually have met the objective referral criteria had they continued being screened with the ATR and moiré topography. Thus, we optimistically considered these students as having met the objective referral criteria. Finally, we defined Group C as the students who were referred due to the presence of clinical signs. Again, they might have met the objective referral criteria had they continued being screened. Hence, we also accepted them as if they had been referred by the objective referral criteria.

All estimates were accompanied by exact 95% confidence intervals based on a binomial distribution, and a 5% level of significance was used in all significance tests. The data management and analysis were performed by the Statistical Analysis System (SAS) version 9.1.²⁴

Role of the funding source

The funding agencies financially supported a research postgraduate student who performed the study design, data collection and analysis, interpretation of results, and writing of the article, but they were not involved in the study. The corresponding

author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

RESULTS

According to the Hong Kong Education Bureau, there were 81,173 and 76,271 students enrolled at grade 5 (mostly 10 years old) in the academic years of 1995/96 and 1996/97, respectively. Consequently, 157,444 students eligible for screening were included in our cohort.

The numbers of students who participated in the screening program, who were referred for radiography, and in whom AIS was detected are summarized in Figure 2. Prior to the commencement of screening, 51 students had already detected scoliosis and were not considered to have been identified by screening. Among the 42,203 (26.8%) non-participants, 71 (0.17%) had AIS detected by the age of 19, 4 cases of which were severe enough to require surgery despite an unknown Cobb angle. For the 115,190 (73.2%) participants, 12 were diagnosed non-idiopathic scoliosis (4 congenital, 1 neuromuscular, 4 Marfan syndrome, and 3 neurofibromatosis). A total of 3,228 (2.8%, 95% CI = 2.7% to 2.9%) students were referred for radiography by 19 years of age, 2,425 objectively and 803 due to clinical signs, and 271 students had self-initiated radiography performed, probably as a result of parental concerns. A

detailed disposition of screened students by their ATR results obtained in the SHSC by the age of 19 can be found in Table 1.

The prevalence rates of AIS for different curvatures and treatment in Hong Kong are shown in Table 2. AIS was more common in girls than in boys ($p < 0.001$ by Fisher's exact test), and the girls-to-boys ratio increased with the severity. The prevalence of treatment was 0.33% (95% CI = 0.30% to 0.36%), and girls were 8.4 times more likely to have treatment than boys ($p < 0.001$ by Fisher's exact test).

Table 3 compares the treatment outcomes for the objective and protocol referral criteria. The additional use of clinical signs for referral identified an additional 107 students who eventually required treatment. A good agreement (unweighted kappa = 0.79, 95% CI = 0.75 to 0.82) was found between the treatment outcomes of the students referred with and without the use of clinical signs. Nevertheless, use of the protocol criteria identified 0.09% (95% CI = 0.08% to 0.11%, $p < 0.001$ by McNemar's test) more students requiring treatment than use of the objective criteria alone.

The clinical effectiveness of the Hong Kong school screening program for AIS is summarized in Table 4. In the cohort, 252 students who had 1 to < 2 moiré lines were also X-rayed in an SAC due to the presence of clinical signs (Group B1), and another 16 students who had 1 to < 2 moiré lines were X-rayed in a specialist hospital without meeting the protocol referral criteria (Group B2). These students might

have attained ≥ 2 moiré lines had they continued the screening. Taking them as if they met the objective criteria, the accuracy measures substantially improved. Further improvement was observed when referral by clinical signs was also considered (Groups A+C).

DISCUSSION

This was the largest retrospective cohort study in the area of scoliosis screening, and added to the only study in the literature that adequately followed screened students in a school scoliosis screening program in Rochester.¹¹ Contrary to the Rochester study, the Hong Kong screening program appears to be sensitive and predictive for screening AIS patients with only a low referral rate.

Using an ATR $\geq 15^\circ$ or ≥ 2 moiré lines as the referral criteria for radiography, the PPVs for curves with a Cobb angle $\geq 20^\circ$ and treatment were 36.5% and 8.1%, respectively, at a referral rate of 2.1%. In other screening programs that used moiré topography, the PPVs for curves $\geq 20^\circ$ and treatment may range from 3.0% to 10.8% and 0.4% to 4.8%, respectively, with referral rates in the range of 3.0% to 8.4%.^{5, 9, 12} A screening program using moiré topography and low-dose roentgenography referred only 0.3% of all screened students and resulted in a PPV of 64.0% for curves $\geq 20^\circ$.¹⁷ Another program using the same screening tools evaluated in a much larger sample

size had a referral rate of 1.0% and a PPV of 24.2%.⁸

In the Rochester study, which used only FBT/ATR for screening but followed students till skeletal maturity, the PPVs for curves $\geq 20^\circ$ and treatment were 17.4% and 5.4%, respectively, with 4.1% rate of referral.¹¹ The school scoliosis screening program in Hong Kong was thus more clinically effective than other comparable programs.

By following all screened students, the sensitivity of scoliosis screening in Hong Kong was estimated as 55.5% for curves $\geq 20^\circ$, and 51.7% for treatment. These values fall in the confidence intervals of those reported in the Rochester study, the only evaluation that reported sensitivity. However, the Rochester study size was small, which resulted in 21% and 18% error in the sensitivities for curves $\geq 20^\circ$ and treatment, respectively. In contrast, our estimates had at most 2.5% error.

In the screening protocol of the Hong Kong program, apart from the objective criteria, a student would also be referred for radiography when there are significant signs of scoliosis, including uneven shoulder height, pelvic tilt, rib or loin hump, or scapular prominence and truncal shift. Referral by clinical signs was also used in many other screening programs, especially in those that used FBT only.^{7, 25, 26} These programs were much less effective than the Hong Kong screening program, which had PPVs of 43.6% for curves $\geq 20^\circ$ and 9.4% for treatment, with corresponding

sensitivities of at least 80%. These were attained with only a slightly increased referral rate of 2.8%. This shows a clear clinical effectiveness of the Hong Kong scoliosis screening program whether or not it is compared with other existing programs.

The reasons for the high clinical effectiveness of school scoliosis screening in Hong Kong are manifold. First, moiré topography was utilized for screening as opposed to the use of FBT/ATR alone in many other programs. Indeed, moiré topography has been demonstrated to be more accurate than FBT/ATR alone.²⁷ Second, since treatment would be considered for curves exceeding 25°,¹ our program, aiming to detect curvature $\geq 20^\circ$, is less likely to unnecessarily refer students who require no treatment than most other programs targeting a common cut-off of 10°. Third, we had follow-up information for all screened students, which resulted in a higher PPV, as students may at first have insignificant curves that later progress. Fourth, students in this program were screened by trained doctors and registered nurses, who were more skilled and experienced than the non-professionals who carried out the screening in some previous studies.⁴ Finally, this screening program is highly centralized and coordinated by the DH, and the two specialist hospitals are within the only two medical school teaching hospitals in Hong Kong. This triad is the most appropriate combination to provide public health and medical services for

scoliosis patients.

Participation in the Hong Kong screening program was satisfactory, with 73.2% of eligible students screened at least once. This screening program was voluntary, but participation was comparable to or better than that in other regions such as Singapore, Minnesota and the Netherlands, where participation rates were 51%, 75% and 80% (estimated), respectively.^{10, 28, 29}

The estimated prevalence of AIS with curves $\geq 10^\circ$ by the age of 19 years in Hong Kong students is 2.5%. Although this is within the range reported in the literature,³ the true prevalence is probably higher, since the Hong Kong program aimed to detect curves $\geq 20^\circ$ and the number of undiscovered curves between 10° and 20° was unknown. On the other hand, the estimated prevalence of curves $\geq 20^\circ$ was 1.4%, which is higher than that reported elsewhere.³ However, most other studies estimated point prevalence, i.e. prevalence at the time of screening, but students who developed AIS later were not accounted for. In contrast, we estimated a period prevalence by the age of 19 years, and all screened students who were ever diagnosed with AIS with a Cobb $\geq 20^\circ$ during adolescence were included.

We have not examined if our scoliosis screening program attained the ultimate goal of minimizing the operation rate.³⁰ However, a reduction in the operation rate depends not only on the effectiveness of screening but also on the efficacy of

conservative bracing treatments.³¹ Its efficacy was supported by a well-known prospective cohort study performed by the Scoliosis Research Society.³² A more recent review also concluded its long term effect even after treatment.³³ However, proper randomized controlled trials (RCTs) are still lacking. Two multi-center RCTs were recently initiated and their results will be important.^{28, 34}

School scoliosis screening provides information essential to the understanding of the epidemiology and the etiology of AIS, which can be a life-long disorder if managed improperly or too late.³⁵ Moreover, it sets up a platform to facilitate research on improving conservative treatments. The scoliosis screening program in Hong Kong, which screened both boys and girls for the whole adolescent period, and referred them for radiography when an $ATR \geq 15^\circ$, ≥ 2 moiré lines were resulted or significant clinical signs were observed, was clearly predictive and sensitive for detecting curves requiring follow-up or treatment. It is better than what has been reported in the literature in terms of both clinical effectiveness and the way these programs were evaluated. The current evidence supports the continuation of school scoliosis screening. The screening protocol being used in Hong Kong could be regarded as a model for further evaluation or refinement in other places.

REFERENCES

1. Scoliosis Research Society. Available at: <http://www.srs.org>. Accessed 7 July 2008.
2. Pehrsson K, Larsson S, Oden A, et al. Long-term follow-up of patients with untreated scoliosis. A study of mortality, causes of death, and symptoms. *Spine*. 1992;17(9):1091-6.
3. Karachalios T, Roidis N, Papagelopoulos PJ, et al. The efficacy of school screening for scoliosis. *Orthopedics*. 2000;23(4):386-91.
4. Gore DR, Passehl R, Sepic S, et al. Scoliosis screening: Results of a community project. *Pediatrics*. 1981;67:196-200.
5. Hansen TB. [Adolescent idiopathic scoliosis among girls in the Herning region. A follow-up of girls with adolescent idiopathic scoliosis found in an earlier screening at school]. *Ugeskr Laeger*. 1994;156(35):4979-82.
6. Juma AHA, Mursal AM, Mangoud AM, et al. Adolescent Idiopathic Scoliosis in School-Children. *Saudi Medical Journal*. 1989 May;10(3):213-5.
7. Morais T, Bernier M, Turcotte F. Age- and sex-specific prevalence of scoliosis and the value of school screening programs. *Am J Public Health*. 1985;75:1377-80.
8. Ohtsuka Y, Yamagata M, Arai S, et al. School screening for scoliosis by the

- Chiba University Medical School Screening Program. Results of 1.24 million students over an 8-year period. *Spine*. 1988;13(11):1251-7.
9. Pruijs JEH, van der Meer R, Hageman MAPE, et al. The benefits of school screening for scoliosis in the central part of the Netherlands. *Eur Spine J*. 1996;5:374-9.
 10. Wong HK, Hui JHP, Rajan U, et al. Idiopathic scoliosis in Singapore schoolchildren. A prevalence study 15 years into the screening program. *Spine*. 2005;30(10):1188-96.
 11. Yawn BP, Yawn RA, Hodge D, et al. A population-based study of school scoliosis screening. *JAMA*. 1999;282:1427-32.
 12. Zhang GP, Li ZR, Wei XR, et al. Screening for scoliosis among school children in Beijing. *Chin Med J*. 1988;101(2):151-4.
 13. Burwell G. The British decision and subsequent events. *Spine*. 1988;13(10):1192-4.
 14. U.S. Preventive Services Task Force (USPSTF). Screening for adolescent idiopathic scoliosis. Policy statement. *JAMA*. 1993;269(20):2664-6.
 15. U.S. Preventive Services Task Force (USPSTF). Screening for idiopathic scoliosis in adolescents: recommendation statement, 2004. Available at: www.preventiveservices.ahrq.gov. Accessed 5 Feb 2008.

16. Grivas TB, Wade MH, Negrini S, et al. SOSORT consensus paper: school screening for scoliosis: Where are we today? *Scoliosis*. 2007;2(17).
17. Motohashi R, Mori I, Kurosawa Y, et al. [Study on an efficient school screening system for scoliosis]. *Yobo Igaku Janaru*. 1999(346):13-7.
18. Fong DYT, Lee CF, Cheung KMC, et al. A meta-analysis of the clinical effectiveness of school scoliosis screening (Unpublished manuscript). 2008.
19. Lonstein JE. Why school screening for scoliosis should be continued. *Spine*. 1988;13(10):1198-200.
20. Bunnell WP. An objective criterion for scoliosis screening. *J Bone Joint Surg Am*. 1984;66(9):1381-7.
21. Lonstein JE. Screening for spinal deformities in Minnesota schools. *Clin Orthop Relat Res*. 1977;126:33-42.
22. Adair IV, van Wijk MC, Armstrong GWD. Moiré topography in scoliosis screening. *Clin Orthop Relat Res*. 1977;129:165-71.
23. Laulund T, Søjbjerg JO, Hørlyck E. Moiré topography in school screening for structural scoliosis. *Acta Orthop Scand*. 1982;53(5):765-8.
24. SAS Institute Inc. SAS/STAT user's guide, version 8. Cary, N.C: SAS Institute Inc; 1999.
25. Rogala EJ, Drummond DS, Gurr J. Scoliosis: Incidence and natural history. A

- prospective epidemiological study. *J Bone Joint Surg Am.* 1978;60:173-6.
26. Soucacos PN, Soucacos PK, Zacharis KC, et al. School-screening for scoliosis. A prospective epidemiological study in Northwestern and Central Greece. *J Bone Joint Surg Am.* 1997;79(1498-1503).
 27. Willner S. A comparative study of the efficiency of different types of school screening for scoliosis. *Acta Orthop Scand.* 1982;53:769-74.
 28. Bunge EM, de Koning HJ, brace trial group. Bracing patients with idiopathic scoliosis: design of the Dutch randomized controlled treatment trial. *BMC Musculoskelet Disord.* 2008;9(57).
 29. Lonstein JE, Bjorklund S, Wanninger MH, et al. Voluntary school screening for scoliosis in Minnesota. *J Bone Joint Surg Am.* 1982;64(4):481-8.
 30. Auke WP, Hofman A, Zielhuis GA. The effect of school screening on surgery for adolescent idiopathic scoliosis. *Eur J Public Health.* 1998;8:237-40.
 31. Lonstein JE. Adolescent idiopathic scoliosis. *Lancet.* 1994;344(8934):1407-12.
 32. Nachemson AL, Peterson LE. Effectiveness of treatment with a brace in girls who have adolescent idiopathic scoliosis. A prospective, controlled study based on data from the Brace Study of the Scoliosis Research Society. *J Bone Joint Surg Am.* 1995;77(6):815-22.
 33. Maruyama T. Bracing adolescent idiopathic scoliosis: a systematic review of the

literature of effective conservative treatment looking for end results 5 years after weaning. *Disabil Rehabil.* 2008;30(10):786-91.

34. ClinicalTrials.gov. Bracing in Adolescent Idiopathic Scoliosis Trial (BrAIST).

Available at: <http://Clinicaltrials.gov>. Accessed 29 Feb 2008.

35. Grivas TB, Vasiliadis E, Savvidou OD, et al. What a school screening program

could contribute in clinical research of idiopathic scoliosis aetiology. *Disabil*

Rehabil. 2008;30(10):752-62.

Figure 1. Protocol of the school screening programme for AIS in Hong Kong





Figure 3. School screening for adolescent idiopathic scoliosis (AIS)



* Numbers of students referred due to (i) ATR $\geq 15^\circ$ or ≥ 2 moiré lines / (ii) clinical signs.

Table 1. Disposition of students by angle of trunk rotation (ATR) results at Student Health Centres (SHSC) by age of 19 years

| Subsequent screening | Screening result in SHSC | | |
|--|--------------------------|-----------------------|--------------------|
| | Only had ATR < 5° | Only had ATR 5° - 14° | Ever had ATR ≥ 15° |
| | (n = 105393) N (%) | (n = 9614) N (%) | (n = 171) N (%) |
| Visited a Special Assessment Centre (SAC) | | | |
| Moiré topography not performed | 73 (0.1) | 538 (5.6) | 1 (0.6) |
| Moiré topography performed, but result unknown | 2 (0.0) | 41 (0.4) | 0 (0.0) |
| Less than 1 moiré line | 22 (0.0) | 521 (5.4) | 3 (1.8) |
| 1 to less than 2 moiré lines | 57 (0.1) | 3078 (32.0) | 6 (3.5) |
| 2 or more moiré lines | 28 (0.0) | 2226 (23.2) | 10 (5.8) |
| Less than 2 moiré lines and X-rayed | 6 (0.0) | 797 (8.3) | 8 (4.7) |
| Visited a specialist hospital | | | |
| Directly from SHSC | 125 (0.1) | 102 (1.1) | 41 (24.0) |
| Through at SAC | 15 (0.0) | 1215 (12.6) | 65 (38.0) |
| Did not visited SAC nor specialist hospital | 105080 (99.7) | 2311 (24.0) | 47 (27.5) |

Table 2. Prevalence of AIS by age of 19 in Hong Kong

| | Total (Exact 95% CI) | Boys (Exact 95% CI) | Girls (Exact 95% CI) | Girls : Boys Ratio |
|------------------------|----------------------|----------------------|----------------------|-----------------------|
| Curves $\geq 10^\circ$ | 2.49% (2.40%, 2.58%) | 1.34% (1.25%, 1.44%) | 3.59% (3.45%, 3.75%) | 2.7 |
| Curves $\geq 20^\circ$ | 1.39% (1.32%, 1.45%) | 0.50% (0.44%, 0.56%) | 2.24% (2.12%, 2.36%) | 4.5 |
| Curves $\geq 40^\circ$ | 0.23% (0.20%, 0.26%) | 0.05% (0.03%, 0.07%) | 0.40% (0.35%, 0.45%) | 8.1 |
| Treatment | 0.33% (0.30%, 0.36%) | 0.07% (0.05%, 0.09%) | 0.58% (0.52%, 0.64%) | 8.4 |
| Brace only | 0.28% (0.25%, 0.31%) | 0.06% (0.04%, 0.08%) | 0.48% (0.43%, 0.54%) | 8.0 |
| Surgery only | 0.02% (0.01%, 0.03%) | 0.01% (0.00%, 0.02%) | 0.03% (0.02%, 0.05%) | 5.8 |
| Brace and Surgery | 0.04% (0.03%, 0.05%) | 0.00% (0.00%, 0.01%) | 0.07% (0.05%, 0.09%) | 18.8 |

Table 3. Treatment outcomes of students by both objective and protocol referral criteria

| | Total (n = 115178) N (%) | Boys (n = 56566) N (%) | Girls (n = 58612) N (%) | Girls : Boys Ratio |
|---|--------------------------------|------------------------------|-------------------------------|-----------------------|
| Referral criteria for radiography: ATR $\geq 15^\circ$ or ≥ 2 moiré lines | | | | |
| Referred | 2425 (2.11) | 771 (1.36) | 1654 (2.82) | 2.1 |
| Brace only | 175 (7.22) | 14 (1.82) | 161 (9.73) | 11.1 |
| Surgery only | 5 (0.21) | 2 (0.26) | 3 (0.18) | 1.4 |
| Brace and Surgery | 16 (0.66) | 0 (0.0) | 16 (0.97) | N.A. |
| Not referred | 112753 (97.89) | 55795 (98.64) | 56958 (97.18) | 1.0 |
| Brace only | 142 (0.13) | 20 (0.04) | 122 (0.21) | 5.9 |
| Surgery only | 16 (0.01) | 1 (0.002) | 15 (0.03) | 14.5 |
| Brace and Surgery | 25 (0.02) | 2 (0.004) | 23 (0.04) | 11.1 |
| Referral criteria for radiography: ATR $\geq 15^\circ$, ≥ 2 moiré lines or presence of clinical signs | | | | |
| Referred | 3228 (2.80) | 976 (1.73) | 2252 (3.84) | 2.2 |
| Brace only | 264 (8.04) | 27 (2.62) | 237 (10.52) | 8.5 |
| Surgery only | 10 (0.3) | 2 (0.19) | 8 (0.36) | 3.9 |
| Brace and Surgery | 29 (0.88) | 0 (0.0) | 29 (1.29) | N.A. |
| Not referred | 111950 (97.20) | 55590 (98.27) | 56360 (96.16) | 1.0 |
| Brace only | 53 (0.05) | 7 (0.01) | 46 (0.08) | 6.3 |
| Surgery only | 11 (0.01) | 1 (0.002) | 10 (0.02) | 9.7 |
| Brace and Surgery | 12 (0.01) | 2 (0.004) | 10 (0.02) | 4.8 |

Table 4. Clinical effectiveness of school screening for adolescent idiopathic scoliosis in Hong Kong

| | Students taken as those referred for radiography by screening | | | |
|----------------------------------|---|--|---|---|
| | Those who had ATR \geq 15° or \geq 2 moiré lines (A) | (A) + Those who had 1 to <2 moiré lines and referred by clinical signs (B1) | (A) + (B1) + Those non- referrals who had 1 to <2 moiré lines but X-rayed in a specialist hospital (B2) | (A) + Those referred by clinical signs (C) |
| Sensitivity | | | | |
| Curves \geq 10° | 64.7% (62.9%, 66.5%) | 72.8% (71.1%, 74.4%) | 73.2% (71.6%, 74.8%) | 91.4% (90.3%, 92.4%) |
| Curves \geq 20° | 55.5% (53.0%, 58.0%) | 63.0% (60.6%, 65.3%) | 63.5% (61.1%, 65.8%) | 88.1% (86.4%, 89.6%) |
| Curves \geq 40° | 40.1% (34.1%, 46.3%) | 45.8% (39.7%, 52.0%) | 46.2% (40.0%, 52.4%) | 75.6% (69.9%, 80.7%) |
| Treatment | 51.7% (46.6%, 56.9%) | 58.8% (53.7%, 63.8%) | 59.4% (54.2%, 64.4%) | 79.9% (75.6%, 83.9%) |
| Positive Predictive Value | | | | |
| Curves \geq 10° | 76.5% (74.7%, 78.1%) | 77.9% (76.3%, 79.5%) | 77.9% (76.3%, 79.5%) | 81.1% (79.7%, 82.5%) |
| Curves \geq 20° | 36.5% (34.6%, 38.5%) | 37.5% (35.7%, 39.4%) | 37.6% (35.8%, 39.5%) | 43.6% (41.8%, 45.3%) |
| Curves \geq 40° | 4.3% (3.6%, 5.2%) | 4.5% (3.7%, 5.3%) | 4.5% (3.7%, 5.4%) | 6.1% (5.3%, 7.0%) |
| Treatment | 8.1% (7.0%, 9.2%) | 8.3% (7.3%, 9.4%) | 8.4% (7.3%, 9.5%) | 9.4% (8.4%, 10.5%) |

The specificity and negative predictive values were larger than 95% for curves \geq 10°, \geq 20°, \geq 40° and for treatment in all scenarios, and thus were not presented here.