

5-9-2018

Cloning and Characterization of a Pyrethroid Pesticide Decomposing Esterase Gene, *Est3385*, from *Rhodopseudomonas palustris* PSB-S

Xiangwen Luo

Hunan Academy of Agricultural Science, China

Deyong Zhang

Hunan Academy of Agricultural Science, China

Xuguo Zhou

University of Kentucky, xuguozhou@uky.edu

Jiao Du


Hunan Academy of Agricultural Science, China

Songbai Zhang

Hunan Academy of Agricultural Science, China, zsongb@hotmail.com

Follow this and additional works at: https://uknowledge.uky.edu/entomology_facpub

See next page for additional authors

 Part of the [Agronomy and Crop Sciences Commons](#), [Entomology Commons](#), [Environmental Health Commons](#), [Environmental Microbiology and Microbial Ecology Commons](#), [Food Biotechnology Commons](#), [Molecular Biology Commons](#), and the [Other Ecology and Evolutionary Biology Commons](#).
Right-click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Luo, Xiangwen; Zhang, Deyong; Zhou, Xuguo; Du, Jiao; Zhang, Songbai; and Liu, Yong, "Cloning and Characterization of a Pyrethroid Pesticide Decomposing Esterase Gene, *Est3385*, from *Rhodopseudomonas palustris* PSB-S" (2018). *Entomology Faculty Publications*. 151.

https://uknowledge.uky.edu/entomology_facpub/151

Authors

Xiangwen Luo, Deyong Zhang, Xuguo Zhou, Jiao Du, Songbai Zhang, and Yong Liu

Cloning and Characterization of a Pyrethroid Pesticide Decomposing Esterase Gene, *Est3385*, from *Rhodopseudomonas palustris* PSB-S

Notes/Citation Information

Published in *Scientific Reports*, v. 8, 7384, p. 1-8.


© The Author(s) 2018

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Digital Object Identifier (DOI)

<https://doi.org/10.1038/s41598-018-25734-9>

SCIENTIFIC REPORTS



OPEN

Cloning and characterization of a pyrethroid pesticide decomposing esterase gene, *Est3385*, from *Rhodopseudomonas palustris* PSB-S

Xiangwen Luo¹, Deyong Zhang¹, Xuguo Zhou², Jiao Du¹, Songbai Zhang¹ & Yong Liu¹

Full length open reading frame of pyrethroid detoxification gene, *Est3385*, contains 963 nucleotides. This gene was identified and cloned based on the genome sequence of *Rhodopseudomonas palustris* PSB-S available at the GneBank. The predicted amino acid sequence of *Est3385* shared moderate identities (30–46%) with the known homologous esterases. Phylogenetic analysis revealed that *Est3385* was a member in the esterase family I. Recombinant *Est3385* was heterologous expressed in *E. coli*, purified and characterized for its substrate specificity, kinetics and stability under various conditions. The optimal temperature and pH for *Est3385* were 35 °C and 6.0, respectively. This enzyme could detoxify various pyrethroid pesticides and degrade the optimal substrate fenpropathrin with a *K_m* and *V_{max}* value of $0.734 \pm 0.013 \text{ mmol}\cdot\text{l}^{-1}$ and $0.918 \pm 0.025 \text{ U}\cdot\mu\text{g}^{-1}$, respectively. No cofactor was found to affect *Est3385* activity but substantial reduction of enzymatic activity was observed when metal ions were applied. Taken together, a new pyrethroid degradation esterase was identified and characterized. Modification of *Est3385* with protein engineering toolsets should enhance its potential for field application to reduce the pesticide residue from agroecosystems.

Synthetic pyrethroids have been used extensively worldwide to control insect pests damaging cereal, potato, cotton, and fruit crops¹. To date, synthetic pyrethroids have been used in field for more than 30 years due mainly to their high toxicities to insect pests, allowing each field application with a relatively low dosage². Furthermore, these pesticides were reported to be less toxic to mammals and thus became the top choices of agricultural industries to replace other more toxic pesticides, including organochlorine and organophosphorus pesticides.

However, the ever increasing use of synthetic pyrethroids in the agro-ecosystem has raised significant concerns in the recent years. Synthetic pyrethroids could be highly toxic to non-targeted organisms such as ladybeetles (*Cycloneda sanguinea* L. and *Harmonia axyridis* Pallas)³, bees (*Apis mellifera*)⁴, and arthropods and vertebrates in the aquatic subsystem⁵. Moreover, some synthetic pyrethroids were reported to have neurotoxicity⁶, reproductive toxicity⁷, cytotoxicity⁸, or teratogenicity and carcinogenicity⁹. These public concerns promoted us to investigate the potential routes that can be used to reduce the amount of synthetic pyrethroid residues in the agro-ecosystem to provide a more safe and healthier environment for agricultural industry.

Residues of synthetic pyrethroids can be broken down by both biotic and abiotic approaches. For biotic approaches, synthetic pyrethroid residues can be decomposed through degradation by microbes. To date, multiple soil resident bacteria have been identified to have the abilities to degrade synthetic pyrethroid residues and several bacterial genes controlling this function are cloned and characterized¹⁰. These enzymes would be play a potential application in subsequent detoxification of the synthetic pyrethroid residues.

Several photosynthetic bacteria in genus *Rhodopseudomonas* have been demonstrated to have the ability to degrade various synthetic pyrethroids¹¹. In 2009, Zhang and his colleagues cloned a gene encoding 2OG-Fe(II) oxygenase from *Rhodopseudomonas* sp. PSB07-21 and showed that this enzyme was capable of degrading synthetic pyrethroids¹². So far, no esterase gene from genus *Rhodopseudomonas* has been reported to have the ability to degrade synthetic pyrethroids.

¹Key Laboratory of Pest Management of Horticultural Crop of Hunan Province, Hunan Plant Protection Institute, Hunan Academy of Agricultural Science, Changsha, 410125, China. ²Department of Entomology, University of Kentucky, Lexington, KY, 40546, USA. Correspondence and requests for materials should be addressed to S.Z. (email: zsongb@hotmail.com) or Y.L. (email: haoasliu@163.com)

In this study, we identified and cloned an esterase gene from *R. palustris* PSB-S after searching the existing genome sequences of this strain deposited at the NCBI database. This identified gene was cloned and expressed in *E. coli* Rosetta cells. Analysis of this recombinant esterase (i.e., Est3385) indicated that this enzyme was indeed capable of decomposing various synthetic pyrethroids, and fenpropathrin as its optimal substrate. We consider that this enzyme can play a significant role in detoxification of pyrethroid residues in field.

Methods

Sources of bacteria strains, and cloning and expression vectors. *Rhodopseudomonas palustris* strain PSB-S was isolated, characterized and stored in our lab. *Escherichiacoli* strain Rosetta and the cloning vector pMD-18T were purchased from Takara Biotechnology (Dalian, China). The expression vector, pGEX-3x, was from Dr. Jian Yan (Virology and Biotechnology Institute, Zhejiang Academy of Agricultural Sciences, Hangzhou, China). For expression of the recombinant protein, *E. coli* Rosetta was cultured in Luria-Bertani medium containing proper antibiotics. Culture medium for *R. palustris* PSB-S contained 0.2 g K_2HPO_4 , 0.8 g KH_2PO_4 , 0.2 g $MgSO_4$, 0.1 g $CaSO_4 \cdot 2H_2O$, 0.0033 g $NaMoO_4 \cdot 2H_2O$, 0.005 g $FeSO_4 \cdot 7H_2O$, and 1.5 g yeast extract in one liter H_2O , pH 7.2.

Chemical reagents and restriction enzymes. Technical-grade synthetic pyrethroids were all purchased from the Hainan Zhengye Pesticide Chemical Co., Ltd (Hainan, China). Other grades of synthetic pyrethroids were from the Tianjin Orient Green Technology and Development Co., Ltd (Tianjin, China). All endonucleases and reagents used in the study were from the Transgen Biotech (Beijing, China). Chemicals used for the analytical analyses were analytical grades and were purchased from the Shanghai Sango Biological Engineering Technology & Services Co., Ltd (Shanghai, China).

Cloning, expression, and analysis of recombinant *Est3385* gene. Sequences of three published synthetic pyrethroids degrading esterases (e.g., pyrethroid-hydrolyzing carboxylesterase pytH)¹³, pyrethroid-hydrolyzing enzyme Sys410 (sys410)¹⁴, and esterase (pytZ)¹⁵ were aligned and the conserved sequence regions were used to blast search for new candidate synthetic pyrethroids degrading esterase genes in the genome of *R. palustris* strain PSB-S deposited at the GenBank (www.ncbi.nlm.nih.gov). Nucleotide sequences and amino acid sequences obtained initially were aligned using the MEGA 6.0 software¹⁶ and then compared with other known esterase and lipase sequences found in different pyrethroid degrading bacterial strains to identify the sequence of esterase *Est3385* in *R. palustris* strain PSB-S. The deduced molecular weight of *Est3385* was estimated using the ExPASy software (http://web.expasy.org/compute_pi/). The protein structure of *EST3385* was predicted and constructed with homology-modelling (<https://www.swissmodel.expasy.org/>).

The resulting full length open reading frame of *Est3385* was PCR amplified using primers *est3385-F* and *est3385-R* (*est3385-F* 5'-CGCGGATCCGCATGAAGGACGTGGCG-3' and *est3385-R* 5'-CCGGAATTCGssCATCGCCGGTAGGT-3', the underlined sequences are the *BamH* I and *EcoR* I restriction sites). The amplified PCR fragment was digested with *BamH* I and *EcoR* I restriction enzymes, and inserted into the pre-digested *BamH* I/*EcoR* I site in the pGEX-3x vector to produce plasmid pGEX-3x-3385. The plasmid DNA was transformed into *E. coli* Rosetta cells, and the recombinant protein was produced and then purified using the Glutathione Sepharose 4B Kit as instructed (GE Healthcare, Beijing, China). Quality of the purified *Est3385* recombinant protein was analyzed in SDS-PAGE gel. Concentration of the purified recombinant protein was estimated by the BCA method using bovine albumin as a standard control¹⁷. To confirm the recombinant protein, protein bands in the SDS-PAGE gels were transferred to PVDF membrane followed probing using an anti-GST antibody diluted 1:1000 (v/v) in a PBS buffer. The detection signal was visualized using nitro blue tetrazolium and 5-bromo-4-chloro-3-indolylphosphate (BCIP).

Substrate specificity and kinetics assays. Synthetic pyrethroids, ρ -nitrophenyl esters and short-chain fatty acids were used as substrates for substrate specificity assays. *Est3385* was tested for its hydrolytic capacity on different synthetic pyrethroids, ρ -nitrophenyl esters and short-chain fatty acids according to a previously published method^{18,19}. Treatments with glutathione S-transferase (GST) were used as a control.

Enzyme kinetics of *Est3385* was determined using five different concentrations ($10 \mu\text{g} \cdot \text{ml}^{-1}$ to $50 \mu\text{g} \cdot \text{ml}^{-1}$) of fenpropathrin as the substrate, and the kinetic constant of *Est3385* was calculated using the Michaelis-Menten equation published previously²⁰. For each treatment, the final non-degraded fenpropathrin was less than 10% of the total fenpropathrin added to the reactions. Each experiment was repeated three times.

Effects of temperature and pH value on enzyme activity. To investigate the effect of temperature on *Est3385* activity, we incubated purified recombinant *Est3385* protein with fenpropathrin in $0.1 \mu\text{mol} \cdot \text{ml}^{-1}$ PBS buffer, pH 7.2, at different temperatures (e.g. 15, 25, 35, 45, 55 and 65 °C). For thermostability assays, the mixed *Est3385* and fenpropathrin samples were incubated first at 35 °C for 1 h followed by 5 min in a 100 °C water bath to end the reaction. The treated samples were then tested again for the residual, and the results were compared with the activity of non-pretreated *Est3385*. The residual activity of the enzyme was presented as the percentage of the activity from the pretreated versus the non-pretreated *Est3385* enzyme. The effect of different pH value on the *Est3385* activity was determined by conducting the activity assays in PBS buffers adjusted to various pH values (e.g., pH 4.0, 5.0, 6.0, 7.0, 8.0 and 9.0) for 1 h. For all the assays, reactions in the absence of *Est3385* enzyme were used as controls. Each experiment was repeated three times.

Effect of metal ions and chemical reagents on *Est3385* activity. Metal ions were added to a PBS buffer, pH 7.2, to produce PBS solutions containing Na^+ , K^+ , Mg^{2+} , Zn^{2+} , Cu^{2+} , Mn^{2+} or Fe^{3+} ($1 \mu\text{mol} \cdot \text{ml}^{-1}$ each). Several other reagents were also added to the PBS solution to produce PBS solutions with 1.0% Tween-80, 1.0% Tween-20, $1 \mu\text{mol} \cdot \text{ml}^{-1}$ ethylenediaminetetraacetic acid (EDTA), $1 \mu\text{mol} \cdot \text{ml}^{-1}$ β -mercaptoethanol, $1 \mu\text{mol} \cdot \text{ml}^{-1}$ phenylmethane sulfonyl fluoride (PMSF) or $1 \mu\text{mol} \cdot \text{ml}^{-1}$ diethyl pyrocarbonate (DEPC). Purified recombinant




Figure 1. Phylogenetic relationship between Est3385 and other proteins belonging to different esterase families. The number in the brackets are the accession numbers of individual protein sequences, while the boxed sequence is Est3385.

Est3385 ($1 \mu\text{g}\cdot\text{ml}^{-1}$) was then added to each reaction and the reactions were incubated at 35°C for 10 min. During these assays, reactions without addition of a metal ion or a chemical reagent were used as controls. Activity of the enzyme in each reaction was calculated as the percentage between the enzyme activity found in a specific reaction versus that from its control. Each experiment was repeated three times.

Gas chromatography (GC) analysis. The residual pyrethroid pesticides, ρ -nitrophenyl esters and short-chain fatty acids left in individual reactions were quantified by GC as described previously^{19,21}. Briefly, samples from above assays were extracted individually three times with n-hexane (10 ml each time). The organic phase from a sample was filtrated using the SampliQ Florisil PR 1GM 6ML as instructed (Agilent Technologies, CA). The resulting residue was dissolved in n-hexane and then quantified by an Agilent 6890N system equipped with a HP-5 chromatographic column ($30 \text{ m} \times 0.32 \text{ mm} \times 0.25 \mu\text{m}$) as instructed (Agilent, USA). Temperatures set for the injection port and the electron capture detector (μECD) were at 250°C and 320°C , respectively. The pipe was kept at 160°C for 5 min and then ramped at $10^\circ\text{C}\cdot\text{min}^{-1}$ to 200°C for 1 min. Subsequently, the pipe was ramped again at $10^\circ\text{C}\cdot\text{min}^{-1}$ to 280°C for another 8 min. High purity nitrogen gas ($>99.999\%$) was introduced at $1 \text{ ml}\cdot\text{min}^{-1}$ and the injection volume was $1 \mu\text{l}$. For the concentrations of pesticide substituents, ρ -nitrophenyl esters and short chain fatty acids, the peak areas mantra was used to analyze gas chromatography using the known corresponding standards as controls.

Results

Identification of Est3385 through Sequence Analysis. Conserved sequences from three known pyrethroid pesticide degrading esterase were identified through sequence alignment, and were used to search new candidate esterase genes in a pyrethroid pesticide decomposing bacteria, *R. palustris* strain PSB-S. A total of six candidate genes (gene ID: PSB-SGL000612, PSB-SGL000613, PSB-SGL000878, PSB-SGL003706, PSB-SGL003752, and PSB-SGL0003385) were found and gene PSB-SGL0003385 (referred to Est3385 thereafter) was selected for further analysis. Result of phylogenetic analysis of Est3385 is shown in Fig. 1. The full length open reading frame of Est3385 contains 963 nucleotides and is now deposited in the GenBank (accession number: KU377526). Although the conserved region in Est3385 sheared 53.2–76.2% nucleotide sequence similarities with the sequences of the three known pyrethroid pesticide degrading esterase genes [e.g., *pytH*, *pytZ* and *est843-sys410*, Fig. S1], the protein sequence of Est3385 sheared only 24.1% sequence identity with *pytH*, 13.9% identity with *pytZ*, and 20% identity with *est843-sys410* (Fig. S2). Phylogenetic analysis indicated that Est3385 belonged to the esterase family cluster I, and was together with an esterase from *Geobacillus thermocatenulatus* (CAA64621.1) and an esterase from *Staphylococcus epidermidis* (AAC67547.1). The 3D structure of protein Est3385 contains six α -helices and nine β -sheets (Fig. S6).

Expression and purification of the recombinant Est3385. Coding sequence of Est3385 was cloned into the pGEX-3x expression vector and the recombinant Est3385 protein was expressed in *E. coli* Rosetta. The




Figure 2. Expression, purification and SDS-PAGE of the recombinant protein Est3385. M: protein marker; lane 1: purified Est3385 protein; lane 2: total protein from *E. coli* Rosetta harboring pGEX-3x-3385.

expressed recombinant protein was then purified by affinity chromatography and examined in SDS-PAGE gels (Fig. 2). A single protein band of approximately 59 kDa was found in lanes loaded with the soluble protein fraction or purified recombinant Est3385 protein. Expression of the recombinant protein was then confirmed by probing the membrane with an anti-GST antibody. Based on the size of protein markers, the molecular mass of this recombinant protein was similar to the estimated molecular mass for Est3385 (33.94 kDa plus ~26 kDa GST tag, Fig. S3).

Substrate specificity and enzyme kinetics of Est3385. Substrate specificity was determined using various pyrethroid pesticides, ρ -nitrophenyl esters, and short-chain fatty acids in the assays. The results shown in Fig. 3 indicated that although Est3385 could degrade all the four tested pyrethroid substrates, fenpropathrin was the most sensitive substrate for Est3385. In this study, control treatments using only GST protein or no enzyme was added did not yield a clear degradation of any of the assayed substrates (Fig. S4). In this study we also determined that Est3385 was able to degrade ρ -nitrophenyl acetate and formic acid effectively. This enzyme had, however, very little effect on ρ -nitrophenyl butyrate and ρ -nitrophenyl caproate, and isobutyric acid (Fig. S5). Our enzyme kinetic studies using fenpropathrin as a substrate showed that the kinetic constants of K_m and V_{max} were at $0.734 \pm 0.013 \text{ mmol}\cdot\text{l}^{-1}$ and $0.918 \pm 0.025 \text{ U}\cdot\mu\text{g}^{-1}$, respectively (Fig. S7).

Effect of temperature and pH value on Est3385 activity. When enzyme activity was tested under various temperature conditions, the highest degradation efficiency of Est3385 was observed at 35 °C (Fig. 4). When the reaction temperature was changed to 25, 45, 55 or 65 °C, the degradation efficiency was found to be moderately reduced. Even when the reaction temperature was set at 15 °C, the relative enzyme activity was still at about 60% compared with that observed at 35 °C. Results shown in Fig. 5 indicated that the optimal pH value for Est3385 activity was at 6.0. The results also showed that under the elevated pH conditions (pH 8.0 to 9.0), Est3385 still showed about 50% enzyme activity compared that observed under the optimal pH condition. These observations indicated that Est3385 had a good tolerance to temperature changes and alkaline conditions.




Figure 3. Substrate specificity analysis for purified Est3385 protein. The substrate activity was set as 100% for reactions with fenpropathrin as the substrate. Each result is the mean of three independent experiments \pm standard deviation.




Figure 4. Temperature effect on the activity of purified Est3385 protein. The highest enzyme activity was observed at 35°C and was set as 100%. Each result is the mean of three independent experiments \pm standard deviation.




Figure 5. Effect of pH value on the activity of purified Est3385 protein. The highest enzyme activity was observed at pH 6.0 and was set as 100%. Each result is the mean of three independent experiments \pm standard deviation.

Effects of metal ions and chemical reagents on Est3385 activity. To determine if metal ions could affect Est3385 activity, we incubated purified Est3385 with fenpropathrin in a PBS buffer containing various metal ions. We then assayed Est3385 activity in these reactions as described above. Results shown in Fig. 6 indicated clearly that all the metal ions tested could significantly suppress but not completely eliminate the activity of Est3385. Among these six metal ions, Fe^{3+} showed the strongest inhibition effect (~63%) followed by Cu^{2+} , Mn^{2+} and Mg^{2+} (~50%). These results demonstrated that the higher the metal ion valence had stronger effect on




Figure 6. Effect of metal ions and chemical reagents on the activity of purified Est3385 enzyme. Each metal ion was used at a concentration of $1.0 \mu\text{mol}\cdot\text{ml}^{-1}$, and the concentration of each chemical reagent was used at 1.0%. Relative enzyme activity in a reaction without the addition of metal ions and chemical reagents was set at 100%. Each result is the mean of three independent experiments \pm standard deviation.

Inhibitors/modifiers	Concentration ($\mu\text{mol}\cdot\text{ml}^{-1}$)	Relative enzyme activity (\pm SD)*
Mercaptoethanol	1.0	100 \pm 5.36
PMSF	1.0	103.97 \pm 7.64
DEPC	1.0	91.83 \pm 3.25

Table 1. Effects of amino acid, protease inhibitor or protease modifier on the activity of purified Est3385 enzyme. *Relative enzyme activity in a reaction without the addition of amino acid, protease inhibitor or protease modifier was set at 100%. SD represents standard deviation.

Est3385 activity. In this study we also tested five chemical reagents for their effects on the Est3385 activity. Results showed that tween-80 could indeed reduce the activity of Est3385 while Tween-20 and EDTA showed no clear effect on the activity of Est3385 (Fig. 6). The assays using protease inhibitor (PMSF) or protease modifier (DEPC or β -mercaptoethanol, Table 1) showed that Est3385 activity was only slightly affected by DEPC, a histidine modifier, but was not affected by PMSF, a serine protease inhibitor, or β -mercaptoethanol, a cysteine modifier (Table 1).

Discussion

It was reported that the key step involved in degradation of pyrethroid pesticides is the cleavage of the ester-bond in the fenpropathrin compound^{13,14,21}. To target this degradation step, pyrethroid hydrolyzing esterase 3 (pye3) was previously identified through direct screening of esterase activities on ester, an indicator commonly used for the esterase activity assays. This esterase screening assay was recommended by the authors as a time-saving approach for identification of pyrethroid hydrolyzing esterases. In addition, this method did not require expensive quantification assays for pyrethroids residues²². In the present study, we searched the available genome sequence of *R. palustris* PSB-S using the conserved sequences from three published pyrethroid degradation esterase genes from other bacteria species in the GenBank^{13–15} and then cloned Est3385 from *R. palustris* PSB-S. Through enzyme kinetics studies, the ability of Est3385 to decompose pyrethroid pesticides was confirmed. We consider that this gene cloning and characterization approach should allow us to screen more new candidate genes important for pesticide residue detoxification in a more straight forward and time-saving manner.

Phylogenetic study (Fig. 1) indicated that Est3385 is a member belonging to the esterase family I. Several other characterized pyrethroid pesticide degrading enzymes were also found in the esterase family VI cluster (i.e., pyt Y¹⁸ and pyt Z¹⁵) or in the family V cluster (i.e., est843-Sys410¹⁴). The conserved pentapeptide motif, Gly-X-Ser-X-Gly, was found in many pyrethroid pesticide degrading enzymes^{13,14,18}, and residue Ser in this motif was shown as a nucleophile catalyzing the hydrolysis of ester bonds²². Because EstP, a pyrethroid pesticide decomposition protein, does not have this conserved motif²³, we consider that pyrethroid pesticide degradation enzymes contain other motif(s) may also serve as nucleophile(s). Interestingly, analysis of Est3385 sequence did not identify any conserved pentapeptide motif. Although the amino acid residues Ser117-119 and Ser125-128 exist in surface of protein Est3385 (Fig. S6), the fact that neither PMSF nor DEPC abolished Est3385 activity during the assays (Table 1), indicating that no histidine or serine presented in the active site of Est3385. Thus, the amino acid residue(s) in Est3385 needed for the breakdown of ester bonds in the pyrethroid pesticide compound remained unknown.

Results shown in this paper indicated that Est3385 is a moderate pyrethroid detoxifying enzyme with a deduced molecular mass of 33.94 kDa (Fig. 2). This molecular mass is smaller than that reported for

permethrinase (61 kDa) from *Bacillus cereus* SM3²⁴, pyrethroid hydrolase (56 kDa) from *Aspergillus niger* ZD11²⁵, carboxylesterase (60 kDa) from mouse liver microsome²⁶, carboxylesterase E3 (58.6 kDa) from *Nephotettix cincticeps* Uhler²⁷, EstP (73 kDa) from *Klebsiella* sp. ZD112²³, and PytY (41.7 kDa) from *Ochrobactrum anthropi* YZ-1¹⁸. This molecular mass is, however, bigger than that reported for carboxylesterase (31 kDa) from *Sphingobium* sp. JZ-1¹³, esterase (31.15 kDa) from the metagenome²², and carboxylesterase PytZ (24.2 kDa) from *Ochrobactrum anthropi* YZ-1¹⁵. Similar to other recombinant pyrethroid degrading enzymes, Est3385 was quickly deployed by IPTG at 35 °C. The operational performance contributed to an identification of characteristics and potential utilization for large-scale manufacturing of the recombinant protein.

This enzyme Est3385 showed a remarkable activity and stability at a broad range of temperature (25 to 65 °C) and pH conditions (pH 5 to 8). Our results have indicated that Est3385 could decompose pyrethroids in the absence of other cofactors, based on the fact that the activity of Est3385 was not inhibited by chelating reagents (Fig. 6), which is similar to the degrading enzyme EstSt7²⁸.

Taken together, a new pyrethroid decomposing enzyme has been identified and cloned in this study. The novelty of this newly identified protein is its lack of the conserved pentapeptide motif, Gly-X-Ser-X-Gly, and AT domain²⁹, found in other pyrethroids degrading enzymes (Table S1). Utilizing fenpropathrin as a substrate, we have demonstrated that this novel enzyme is thermostable and can tolerate a wide range of pH conditions. We propose that Est3385 is a potential pyrethroid pesticides decomposing enzyme for agricultural industry. It is noteworthy that Est3385 can be moderately inhibited by metal ions (Fig. 6), a potential limiting factor for commercial application of this enzyme in field. It is, however, reasonable to speculate that modification of this enzyme through current biotechnologies should eliminate or at least reduce this barrier to minimal as previously discussed³⁰.

References

- Li, H. Z. *et al.* Global occurrence of pyrethroid insecticides in sediment and the associated toxicological effects on benthic invertebrates: an overview. *J Hazard Mat* **324**, 258–271 (2016).
- Steven, A. B. & David, E. R. Structure-activity and interaction effects of 14 different pyrethroids on voltage-gated chloride ion channels. *Toxicol Sci* **77**, 341–346 (2004).
- Michaud, J. P. Relative toxicity of six insecticides to *Cycloneda sanguinea* and *Harmonia axyridis* (Coleoptera: Coccinellidae). *J Entomol Sci* **37**, 83–93 (2002).
- Collet, C. *et al.* Differential action of pyrethroids on honey bee and bumble bee voltage-gated sodium channels. *Biophys J* **110**, 112a (2016).
- Wang, J. Z. *et al.* Occurrence, compositional distribution, and toxicity assessment of pyrethroid insecticides in sediments from the fluvial systems of Chaohu Lake, Eastern China. *Environ Sci Pollut Res* **23**, 10406–10414 (2016).
- Halstead, N. T., Civitello, D. J. & Rohr, J. R. Comparative toxicities of organophosphate and pyrethroid insecticides to aquatic macroarthropods. *Chemosphere* **135**, 265–271 (2015).
- Tu, W. Q. *et al.* Acute exposure to synthetic pyrethroids causes bioconcentration and disruption of the hypothalamus–pituitary–thyroid axis in zebrafish embryos. *Sci Total Environ* **42**, 876–885 (2016).
- Costa, C. *et al.* Immunomodulatory effect of occupational exposure to synthetic pyrethroids. *Toxicol Lett* **238**, S222–S223 (2015).
- Radwan, M. *et al.* The association between environment exposure to pyrethroids and sperm aneuploidy. *Chemosphere* **128**, 42–48 (2015).
- Yang, C. R. & Jian, J. Z. Recent advance in biodegradation in China: New microorganism and pathway, biodegradation engineering, and bioenergy from pollutant biodegradation. *Process Biochem* **45**, 1937–1943 (2010).
- Zhang, S. B. *et al.* Degradation characteristics and pathway of fenpropathrin by *Rhodospseudomonas* sp. strain PSB07-6. *Fresenius Environ Bull* **11**, 2060–2065 (2009).
- Zhang, S. B. *et al.* Isolation, identification and degrading gene cloning of a pyrethroid-degrading bacterium. *Acta Microbi Sinica* **49**, 1520–1526 (2009).
- Wang, B. Z. *et al.* Cloning of a novel pyrethroid-hydrolyzing carboxylesterase gene from *Sphingobium* sp. strain JZ-1 and characterization of the gene product. *Appl Environ Microbiol* **75**, 5496–5500 (2009).
- Fan, X. J. *et al.* Identification and characterization of a novel thermostable pyrethroid-hydrolyzing enzyme isolated through metagenomic approach. *Microbial Cell Factories* **11**, 33 (2012).
- Zhai, Y. *et al.* Molecular cloning, purification and biochemical characterization of a novel pyrethroid-hydrolyzing carboxylesterase gene from *Ochrobactrum anthropi* YZ-1. *J Hazard Mat* **221–222**, 206–212 (2012).
- Tamura, K. *et al.* MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* **30**, 2725–2729 (2013).
- Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248–252 (1976).
- Ruan, Z. *et al.* Molecular cloning and characterization of a newly isolated pyrethroid-degrading esterase gene from a genomic library of *Ochrobactrum anthropi* YZ-1. *PLoS ONE* **8**, e77329 (2013).
- Chen, S. H. *et al.* Pathway and kinetics of cyhalothrin biodegradation by *Bacillus thuringiensis* strain ZS-19. *Sci Rep* **5**, 8784 (2015).
- Bezerra, R. M. F. *et al.* Enzyme inhibition studies by integrated Michaelis-Menten equation considering simultaneous presence of two inhibitors when one of them is a reaction product. *Comput Meth Prog Bio* **125**, 2–7 (2016).
- Zhang, S. B. *et al.* Cometabolic biotransformation of fenpropathrin by *Clostridium* species strain ZP3. *Biodegradation* **22**, 869–875 (2011).
- Li, G., Wang, K. & Liu, Y. H. Molecular cloning and characterization of a novel pyrethroid-hydrolyzing esterase originating from the metagenome. *Microbial Cell Factories* **7**, 38 (2008).
- Wu, P. C. *et al.* Molecular cloning, purification, and biochemical characterization of a novel pyrethroid-hydrolyzing esterase from *Klebsiella* sp. strain ZD112. *J Agric Food Chem* **54**, 836–842 (2006).
- Maloeny, S. E., Maule, A. & Smith, A. R. Purification and preliminary characterization of permethrinase from a pyrethroid-transforming strain of *Bacillus cereus*. *Appl Environ Microbiol* **59**, 2007–2013 (1993).
- Liang, W. Q. *et al.* Purification and characterization of a novel pyrethroid hydrolase from *Aspergillus niger* ZD11. *J Agric Food Chem* **53**, 7415–7420 (2005).
- Stock, J. E. *et al.* Identification, expression, and purification of a pyrethroid-hydrolyzing carboxylesterase from mouse liver microsomes. *J Biol Chem* **279**, 29863–29869 (2004).
- Chiang, S. W. & Sun, C. N. Purification and characterization of carboxylesterase of a rice green leafhopper *Nephotettix cincticeps* uhler. *Pestic Biochem Physiol* **54**, 181–189 (1996).
- Wei, T. *et al.* Characterization of a novel thermophilic pyrethroid-hydrolyzing carboxylesterase from *Sulfolobus tokodaii* into a new family. *J Mol Catal B: Enzy* **97**, 225–232 (2013).

29. Cai, X. H. *et al.* Autotransporter domain-dependent enzymatic analysis of a novel extremely thermostable carboxylesterase with high biodegradability towards pyrethroid pesticides. *Sci Rep* **7**, 3461 (2017).
30. Coppina, C. W. *et al.* Testing the evolvability of an insect carboxylesterase for the detoxification of synthetic pyrethroid insecticides. *Insect Biochem Mol Biol* **42**, 343–352 (2012).

Acknowledgements

This work was supported by the National Key R&D Program of China (2017YFD0800702), the National Natural Science Foundation of China (31272067, 3140110978), the Agriculture Research System of China (CARS-25-B-05), Hunan Science and technology project (2017JC44) and the Hunan Talent Project (2016RS2019).

Author Contributions

S.Z., X.G. and Y.L. conceived and designed the experiments. X.W., D.J. and D.Y. performed the experiments, analyzed the data, contributed reagents and materials. S.Z., X.G. and Y.L. wrote the paper.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-25734-9>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018