

2003

Cluster of type IV secretion genes in *Helicobacter pylori*'s plasticity zone

Dangeruta Kersulyte

Washington University School of Medicine in St. Louis

Billie Velapatiño

Washington University School of Medicine in St. Louis

Asish K. Mukhopadhyay

Washington University School of Medicine in St. Louis

Lizbeth Cahuayme

Washington University School of Medicine in St. Louis

Alejandro Bussalleu

Universidad Peruana Cayetano Heredia

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Kersulyte, Dangeruta; Velapatiño, Billie; Mukhopadhyay, Asish K.; Cahuayme, Lizbeth; Bussalleu, Alejandro; Combe, Juan; Gilman, Robert H.; and Berg, Douglas E., "Cluster of type IV secretion genes in *Helicobacter pylori*'s plasticity zone." *Journal of Bacteriology*. 185,13. 3764-3772. (2003).
https://digitalcommons.wustl.edu/open_access_pubs/2805

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Dangeruta Kersulyte, Billie Velapatiño, Asish K. Mukhopadhyay, Lizbeth Cahuayme, Alejandro Bussalleu, Juan Combe, Robert H. Gilman, and Douglas E. Berg

Cluster of Type IV Secretion Genes in *Helicobacter pylori*'s Plasticity Zone

Dangeruta Kersulyte, Billie Velapatiño, Asish K. Mukhopadhyay, Lizbeth Cahuayme, Alejandro Bussalleu, Juan Combe, Robert H. Gilman and Douglas E. Berg
J. Bacteriol. 2003, 185(13):3764. DOI:
10.1128/JB.185.13.3764-3772.2003.

Updated information and services can be found at:
<http://jb.asm.org/content/185/13/3764>

	<i>These include:</i>
REFERENCES	This article cites 51 articles, 26 of which can be accessed free at: http://jb.asm.org/content/185/13/3764#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Cluster of Type IV Secretion Genes in *Helicobacter pylori*'s Plasticity Zone

Dangeruta Kersulyte,¹ Billie Velapatiño,^{1,2} Asish K. Mukhopadhyay,^{1†} Lizbeth Cahuayme,²
Alejandro Bussalleu,³ Juan Combe,³ Robert H. Gilman,² and Douglas E. Berg^{1*}

Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri 63110,¹
and Department of Microbiology² and Department of Gastroenterology, Facultad de Medicina,³
Universidad Peruana Cayetano Heredia, Lima 31, Peru

Received 21 February 2003/Accepted 21 April 2003

Some genes present in only certain strains of the genetically diverse gastric pathogen *Helicobacter pylori* may affect its phenotype and/or evolutionary potential. Here we describe a new 16.3-kb segment, 7 of whose 16 open reading frames are homologs of type IV secretion genes (*virB4*, *virB7* to *virB11*, and *virD4*), the third such putative secretion gene cluster found in *H. pylori*. This segment, to be called *tfs3*, was discovered by subtractive hybridization and chromosome walking. Full-length and truncated *tfs3* elements were found in 20 and 19%, respectively, of 94 strains tested, which were from Spain, Peru, India, and Japan. A *tfs3* remnant (6 kb) was found in an archived stock of reference strain J99, although it was not included in this strain's published genome sequence. PCR and DNA sequence analyses indicated the following. (i) *tfs3*'s ends are conserved. (ii) Right-end insertion occurred at one specific site in a chromosomal region that is varied in gene content and arrangement, the "plasticity zone." (iii) Left-end insertion occurred at different sites in each of nine strains studied. (iv) Sequences next to the right-end target in *tfs3*-free strains were absent from most strains carrying full-length *tfs3* elements. These patterns suggested insertion by a transposition-like event, but one in which targets are chosen with little or no specificity at the left end and high specificity at the right end, thereby deleting the intervening DNA.

Helicobacter pylori is a genetically diverse gastric pathogen that chronically infects more than half of all people worldwide, typically for decades. It is a major cause of gastritis and peptic ulcer disease and is an early risk factor for gastric cancer (for reviews, see references 19, 25, and 40). Comparison of full genome sequences of two strains (26695 and J99) showed that some 6 to 7% of genes of one strain were absent from the other and vice versa (3, 48). Similarly, subtractive hybridization and microarray experiments indicated that 5% or more of genes known from strains 26695 and J99 were absent from the genomes of many other strains (2, 45) and that they, in turn, contain genes not found in the fully sequenced genomes. Some two-thirds of *H. pylori*'s strain-specific genes have no predicted function, and most have been found to date only in this species (3, 48).

Genome sequence comparisons indicated that nearly half of *H. pylori*'s strain-specific genes are located in its plasticity zone (3). This zone is about 45 kb long in strain J99 and 68 kb long in strain 26695, where it is split in two by a chromosomal rearrangement (3, 48). Many plasticity zone genes are transcribed, suggesting that they are functional (43). The zone contains several insertion sequences, as well as genes with protein-level homology to various recombinases, integrases, and topoisomerases. Their presence implicated specialized recombination events, including transposition, as a cause of differences among strains in gene content and arrangement. Two

other genes found in the plasticity zone of one strain, 26695, potentially encode VirB4- and VirD4-type ATPases, which, in other systems interact in multiprotein assemblies to help translocate specific proteins and/or DNAs through the cell envelope (type IV secretion). Type IV secretion systems are involved in various ways in pilus formation and nucleoprotein transfer from donor to recipient bacterial or plant cells (in case of tumor induction by *Agrobacterium tumefaciens*), DNA transformation in *H. pylori*, toxin secretion, cell signaling, and intracellular survival of various pathogens (14, 18, 27, 38, 46). Each type IV secretion system is quite specific for a particular set of macromolecule (protein or nucleic acid) substrates and probably has its own distinct role.

Illustration that natural differences in gene content can affect *H. pylori* phenotypes came from studies of the *cag* pathogenicity island (PAI), a 37-kb (27 genes) DNA segment that encodes a type IV secretion system. Although the *cag* PAI is present in nearly all East Asian strains, it is epidemiologically associated with virulence in the West: many strains that cause only benign infections lack this gene cluster, whereas nearly all of those implicated in overt disease contain it (19). Mutational tests indicated that 17 of the 27 *cag* PAI-encoded proteins, including its six VirB homologs, participate in translocation of CagA protein (also encoded in this PAI) to target epithelial cells, where it is tyrosine phosphorylated and disrupts cell signaling. Fourteen of these 17 proteins help promote gastric epithelial synthesis of the proinflammatory cytokine interleukin-8 and thereby potentially damaging inflammatory responses (28). A second putative type IV secretion system, which seems to be present in essentially all *H. pylori* strains, is needed for competence in DNA transformation ("*comB*" cluster) (30).

Here we describe (i) a new gene cluster in *H. pylori* that,

* Corresponding author. Mailing address: Department of Molecular Microbiology, Campus Box 8230, Washington University School of Medicine, St. Louis, MO 63110. Phone: (314) 362-2772. Fax: (314) 362-1232. E-mail: berg@borcim.wustl.edu.

† Present address: National Institute of Cholera and Enteric Diseases, Scheme-XM, Beliaghata, Calcutta-700010, India.

based on homology, probably encodes a type IV secretion system, the third such system found in this species; (ii) its location in the plasticity zone; (iii) its distribution among *H. pylori* strains; and (iv) evidence suggesting it is an unusual transposon that generates large nested deletions during insertion.

MATERIALS AND METHODS

General methods. Standard procedures were used for *H. pylori* growth on brain heart infusion agar (Difco) containing 10% horse blood in a microaerobic atmosphere (1, 34). High-molecular-weight genomic DNA was isolated by a hexadecyltrimethylammonium method (9). A subtractive DNA library was made with the PCR-Select bacterial genome subtraction kit (Clontech) (2) and the strains described below.

Specific PCR was carried out in 20- μ l volumes containing 5 to 10 ng of DNA, 0.25 to 0.5 U of *Taq* polymerase ("Biolase" Midwest Scientific, St. Louis, Mo.), 2.5 pM each primer, and 0.25 mM each deoxynucleoside triphosphate (dNTP) in a standard buffer for 30 cycles with the following cycling parameters: denaturation at 94°C for 30 s, annealing as appropriate for the primer sequence (generally 52°C) for 30 s, and DNA synthesis at 72°C for the appropriate time (1 min per kilobase). PCR products for sequencing were purified with a PCR purification kit (Qiagen, Chatsworth, Calif.) or extracted from agarose by centrifugation with Ultrafree-DA (Amicon, Millipore). DNA sequencing was carried out with a Big Dye Terminator DNA sequencing kit (Perkin-Elmer) and ABI automated sequencers (gel models 373 and 377; capillary model 3100). Direct sequencing of PCR products was done with 5 μ g of PCR fragment (about 100 ng of DNA), 1 μ l of primer (1.6 pM), and 4 μ l of Big Dye (version 2) under the following conditions: 25 cycles of denaturation at 96°C for 10 s, annealing at 50°C for 5 s, and extension at 60°C for 4 min under oil-free conditions (Perkin-Elmer 2400). Direct sequencing on chromosomal DNA was done with 5 μ l of chromosomal DNA (0.5 to 1 μ g), 1 μ l of primer (10 pM), and 6 μ l of Big Dye (version 2) under the following conditions: 96°C for 5 min and then 90 cycles with denaturation at 96°C for 10 s, annealing at 50°C for 5 s, and extension at 60°C for 4 min under oil-free conditions (Perkin-Elmer 2400). For capillary sequencers, the reaction mixture was cleaned through a column (Performa DTR gel filtration cartridges; Edge BioSystems) instead of ethanol precipitation. DNA sequence editing and analysis were performed with programs in the GCG package (Genetics Computer Group, Madison, Wis.) and Vector NTI (Informax, Bethesda, Md.) programs and data in *H. pylori* genome sequence databases (3, 48) and BLAST and Pfam (version 5.3) homology search programs (<http://www.ncbi.nlm.nih.gov/blast/blast.cgi>; <http://pfam.wustl.edu/hmmsearch.shtml>). Primer sequences used for PCR and sequencing are available from the authors on request.

Gene knockout mutations in *tfs3* were made by PCR (without cloning) using a kanamycin resistance cassette (51), essentially as described previously (16). A *comB* deletion allele, in which the *comB7*-to-*comB10* segment was replaced with a chloramphenicol resistance cassette, was kindly provided by K. Ogura (unpublished observations). These mutant alleles were introduced into *H. pylori* by electroporation and selection for resistance to kanamycin or chloramphenicol, as appropriate.

Natural transformation to test for competence was carried out with 16S ribosomal DNA (rDNA) containing the Aus108 tetracycline resistance (*tet*) allele (AGA to TTC in tetracycline binding site) (20, 50). Bacteria from about 1/4 of a confluent plate culture were transferred to fresh agar medium and incubated for 24 h. Exponentially growing cells from this plate were collected with a loop (generally from half or all of the plate), spread in a small circle (1 cm in diameter) on fresh medium, and incubated for 2 h. A 5- μ l aliquot containing about 0.5 to 1 μ g of a 16S rDNA PCR fragment containing the *tet* allele (20) was mixed with this patch of growth and incubated for 4 h, and then the cells were distributed over the entire plate and incubated for 20 h. Cells were then harvested and transferred to fresh medium containing 2 μ g of tetracycline per ml and incubated to select Tet^r transformants (3 to 6 days).

Bacterial strains. The *H. pylori* strains used for subtractive library construction were PeCan18B from a Peruvian gastric cancer patient (tester) and a pool of strains SJM180A, SJM184A, and SJM189A from Peruvian gastritis patients (driver). An earlier analysis starting with another subtractive clone from this set of strains led to the finding of insertion sequence ISHp608 (35). All Peruvian strains were cultured from biopsies from persons with gastroduodenal complaints at the Servicio Universitario de Apoyo, Universidad Peruana Cayetano Heredia (UPCH), in Lima. Biopsies were obtained with written informed consent under protocols approved by the UPCH Human Studies Committee. Most other *H. pylori* strains used here were from the Berg laboratory collection and

have been described in detail elsewhere (11, 15, 35, 41). The ethnicities of the patients and clinical disease associations are as follows. The 12 Peruvian gastric cancer strains and the 12 Peruvian gastritis strains were from persons of primarily Amerindian ancestry living in the shanty town of San Juan de Miraflores, in Lima, Peru. Eighteen of 22 Indian strains were from middle or lower middle class urban (12 strains) or rural (6 strains) residents of West Bengal; all were from patients with peptic ulcer disease. The other four strains were from Chennai (South India): three from patients with peptic ulcer and one from a patient with dyspepsia. Twelve of the 24 Japanese strains were from gastric cancer patients, and the other 12 were from gastritis patients.

Nucleotide sequence accession number. The 20,457-nucleotide (nt) sequence we determined from strain PeCan18B was deposited in GenBank under accession no. AF487344. *tfs3* extends from nt 253 to 18397. A total of 1,833 nt of this sequence (from nt 1752 to 3584) consists of ISHp608 (35), which was not found in *tfs3* elements in other strains. A 6,983-nt sequence found in reference strain J99, 5,795 nt of which (from nt 905 to 6699) was not included in its published genome sequence, was deposited in GenBank under accession no. AY128679. (The *tfs3* component extends from nt 460 to 6699.) A 1,968-nt sequence from strain CPY6081, which includes the gene designated "*jhp926like*" and the putative empty site for *tfs3* insertion, was deposited in GenBank under accession no. AY128680.

RESULTS

Discovery of *tfs3*. The studies presented here define a 16-kb putative transposable element (Fig. 1A; detailed in Table 1), seven of the open reading frames (ORFs) of which encode homologs of core type IV secretion proteins. This element will be called "*tfs3*" to connote that it is the third putative type IV secretion system found in *H. pylori*. These studies began with a subtractive hybridization, carried out to search for sequences in an *H. pylori* strain from a Peruvian patient with gastric cancer (PeCan18B) that were not present in strains from three Peruvians with more benign infections. One subtractive clone exhibited 28% protein identity to a *virB11* homolog in the *H. pylori* *cag* PAI (*hp525* in strain 26695 sequence), which is implicated in CagA protein translocation, induction of proinflammatory cytokine synthesis, and virulence (28, 48). It was more closely related (64% DNA identity, 55% protein identity, and 74% protein similarity) to another *virB11* homolog (*hp1421*) whose role in *H. pylori* is not known. We decided to examine the genomic context of this third *virB11* homolog and its distribution among *H. pylori* strains, based on our interest in genome evolution and VirB11-mediated secretion processes.

***tfs3* sequence.** The set of genes containing this new *virB11* homolog in strain PeCan18B was sequenced by primer walking from the original subtractive clone, with the intent to continue until genes known from published sequences were encountered. The first such genes were *jhp927* and then *jhp928* (known from strain J99) on the right, and *hp460* and then *hp459* (known from strain 26695) on the left (Fig. 1A). PCR tests with DNAs from 94 representative *H. pylori* strains (see "Geographic distribution," below) identified *jhp927* in 22 of 37 strains that contained any *tfs3* genes and in 28 of 57 strains that seemed to be *tfs3* free. This pattern was taken as initial evidence that *jhp927* is to the right of *tfs3*, not within it. In contrast, PCR tests for *hp459* and *hp460* indicated that they were present in each of 37 strains that contained *tfs3* sequences, but not in any of 57 strains that lacked them. These data suggested that *hp459* and *hp460* were part of *tfs3* and that *tfs3* was truncated in strain 26695. We therefore continued sequencing leftward and checked sequences obtained by PCR using DNAs from a panel of other representative *H. pylori* strains at each step. The results suggested that *tfs3* extended




FIG. 1. Organization of *tfs3* in *H. pylori*. The homologies diagrammed here are detailed in Table 1. (A) *tfs3* in strain PeCan18B. The *tfs3* segment extends from nt 253 to 18397 of the 20,457 nt deposited under GenBank accession no. AF487344 (Details of PCR and genomic sequencing used to determine this sequence are available from the authors upon request.) ISHp608 (35) is 60 nt from the 3' end of *orfC* in PeCan18B but was not found in any of 37 other *tfs3* elements tested. LJ, left junction; RJ, right junction. (B) Organization of the *tfs3* remnant in archived strain J99. This remnant extends from nt 460 to 6699 of the 6,983 nt deposited under GenBank accession no. AY128679. Two copies of *jhp927* were found in J99: one connected to *jhp926* (as published in the full genome sequence in reference 3) and another connected to the right end of *tfs3* (present data) (GenBank accession no. AY128679 [absent in the J99 full genome sequence data]). (C) Map of truncated *tfs3* element of strain 26695. This remnant corresponds to nt 253 to 7622 of *tfs3* in strain PeCan18B (GenBank accession no. AF487344). *orfD* (94 codons) and *orfG2* (47 codons) were not annotated as ORFs in the 26695 genome sequence, but are present in its DNA sequence.

leftward through *orfB* (*hp455*) and the 3' end of *hp454*. Of 20,457 nt sequenced (GenBank accession no. AF487344), 16,312 nt were assigned to the *tfs3* element itself; another 1,833 nt consisted of ISHp608 (35), which was inserted near the 3' end of *orfC*. The leftmost 252 nt and rightmost 2,060 nt were considered flanking sequences, just outside the left and right ends, respectively, of *tfs3*. Seven of *tfs3*'s 16 ORFs encoded homologs of VirB proteins that act in type IV secretion: four that form a membrane-traversing transporter channel (VirB7, VirB8, VirB9, and VirB10) and three that are cytoplasmic membrane-associated ATPases and that efficiently move their cognate macromolecular substrates to and through the channel (VirB4, VirB11, and VirD4) (Fig. 1A and Table 1). We also note that *orfB* (*hp455*), at the left end of *tfs3*, was nearly identical in the first third of its length to part of *H. pylori* gene *jhp941*, another segment of which exhibits high protein-level homology to members of the XerC/XerD-type integrase-recombinase-transposase family.

Geographic distribution. A set of 94 *H. pylori* strains from four geographic regions (Japan, Peru, Spain, and India) were tested by PCR for 15 *tfs3* ORFs. One-fifth of the strains carried an apparently full-length *tfs3* cluster, as defined in relation to the sequence in strain PeCan18B (Fig. 1A), and another one-fifth of the strains contained just part of this segment (most often the left half) (Table 2). No correlation of *tfs3* carriage with disease status was observed: *tfs3* was equally present in

strains from Peruvian and Japanese patients with gastric cancer and with gastritis. Similarly, no correlation with *cag* PAI status was seen with Spanish strains (half of which carried and half of which lacked the *cag* PAI).

***tfs3* in strain J99.** DNA from an archived stock of reference strain J99 had been chosen initially as a negative control for tests of the geographic distribution of *tfs3* elements, because no *tfs3* genes had been included in this strain's published genome sequence (GenBank accession no. AE001439). Our PCR tests indicated, however, that parts of *tfs3* were present in the strain J99 genome. Our DNA sequencing then identified a 6.2-kb segment consisting of 741-, 402-, and 5,097-nt segments from the left end, interior, and right end, respectively, of full-length *tfs3* (GenBank accession no. AY128679). This structure can be ascribed to deletion of 1.3 kb between sites in *orfB* and *orfE* and deletion of 8.7 kb between sites in *orfG* and *orfK* (Fig. 1B).

We found *orfP* of this *tfs3* remnant in strain J99 next to *jhp927* (as in strain PeCan18B), although another gene, *jhp926*, had been placed next to *jhp927* in the published J99 sequence. Direct PCR tests identified both connections (both *jhp926*- and *tfs3*-*jhp927*), and DNA sequencing showed that a 750-nt segment containing the *jhp926*-*jhp927* junction was identical to that in the published genome sequence. The implication that J99 contained two copies of *jhp927* was confirmed by Southern blot hybridization (data not shown). Further PCR tests of DNAs from two single-colony isolates from this J99 stock iden-

TABLE 1. Description of *tfs3* from strain PeCan18B

ORF (aa)	Region of homology (aa)	Homolog	Length (aa)	Protein comparison (%identity/%similarity)	%DNA	Description (reference)	ORFs in <i>tfs3</i> ^a
A (78>)							<i>jhp949</i> (98% DNA), left of <i>tfs3</i> , 2 transmembrane helices
B (262)	49–134	<i>jhp941</i>	331	96/96	99	Integrase-resolvase	<i>hp455</i> (97% DNA)
C (164)							<i>hp456</i> (87% DNA), ISHp608 at 145/146 aa
D (94)	6–94	<i>hp15</i>	93	57/74	64	Hypothetical	<i>hp456-57</i> intergenic region (98% DNA), 3 transmembrane helices
E (87)	1–87	<i>hp16</i>	87	60/81	67	Hypothetical	<i>hp457</i> (99% DNA), 1 transmembrane helix
	1–87	<i>hp442</i>	88	38/56		Hypothetical, plasticity zone	
	6–87	<i>Cjp52</i>	80	27/49		<i>C. jejuni</i> , pVir (10)	
F (78)							<i>hp458</i> (100% DNA)
G (858)	1–837	<i>hp17</i>	787	46/64		<i>comB4</i>	<i>hp459</i> (99% DNA), <i>virB4</i> homolog, 2 transmembrane helices
	11–843	<i>hp441</i>	807	37/58		<i>virB4</i> , plasticity zone	
	22–848	<i>Cjp53</i>	822	31/53		<i>virB4</i> in <i>C. jejuni</i> , pVir	
	211–697	<i>jhp917</i>	475	36/57		<i>virB4-2</i> , plasticity zone	
	708–833	<i>jhp918</i>	140	34/55		<i>virB4-3</i> , plasticity zone	
	38–717	<i>hp544</i>	983	21/39		<i>cag23</i> (<i>virB4</i>), <i>cag</i> PAI	
G2 (47)	5–33	<i>orf2</i>	37	41/61		<i>comB7</i> in P1 (30)	<i>hp459-60</i> intergenic region (98% DNA), <i>virB7</i> homolog
	7–32	<i>Cjp54</i>	42	48/52		<i>virB7</i> in <i>C. jejuni</i> , pVir	
H (352)	116–336	<i>hp38</i>	245	43/66		<i>comB8</i>	<i>hp460</i> (5' half) (98% DNA), <i>virB8</i> homolog (3' half), 1 transmembrane helix
	39–345	<i>jhp921</i>	328	36/56		Putative, plasticity zone	
	122–343	<i>Cjp01</i>	225	34/57		<i>virB8</i> in <i>C. jejuni</i> , pVir	
	4–337	<i>hp439</i>	366	32/52		Hypothetical, plasticity zone	
I (551)	153–551	<i>hp39m</i>	326	53/67		<i>comB9</i>	<i>virB9</i> homolog, 1 transmembrane helix
	142–545	<i>jhp922m</i>	511	37/52		Conjugation, plasticity zone	
	87–546	<i>Cjp02</i>	356	30/50		<i>virB9</i> in <i>C. jejuni</i> , pVir	
J (402)	13–143	<i>jhp36</i>	376	26/32	69	<i>comB10</i> +	<i>virB10</i> homolog, 2 transmembrane helices
	156–384			77/83			
	27–385	<i>Cjp03</i>	378	36/54		<i>virB10</i> in <i>C. jejuni</i> , pVir	
K (735)	7–508	<i>jhp945</i>	668	34/54		Putative, plasticity zone	1 transmembrane helix
L (320)							No homology, 1 transmembrane helix
M (92)							No homology
N (314)	6–310	<i>hp1421</i>	304	53/72		<i>virB11-12</i>	<i>virB11</i> homolog
	12–309	<i>Cjp05</i>	317	40/59		<i>virB11</i> in <i>C. jejuni</i> , pVir	
	27–302	<i>hp525</i>	330	28/50		<i>virB11</i> , <i>cag</i> PAI	
O (172)							No homology
P (747)	449–612	<i>hp1006</i>	177	62/79		<i>traG</i> (<i>virD4-2</i>), plasticity zone	<i>virD4</i> homolog, 4 transmembrane helices
	14–636	<i>Cjp06</i>	628	37/56		<i>virD4</i> in <i>C. jejuni</i> , pVir	
	150–604	<i>hp524</i>	748	23/46		<i>virD4</i> , <i>cag</i> PAI	
Q (701>)							<i>jhp927</i> (90% DNA) + <i>jhp928</i> (93% DNA), <i>tfs3</i> right junction is in 9th codon

^a See Fig. 1 for the overall structure of *tfs3*. *orfA* and *orfQ* are adjacent to *tfs3* in PeCan18B (*orfQ* in PeCan18B is a fusion of *jhp927* and *jhp928*) *orfB* through *orfP* are within *tfs3*.

tified the same two connections to *jhp927*. Thus, archived J99 had a duplication of the region containing *jhp927*; it was not a mixture of two strains with different *jhp927* connections.

The structure of *tfs3* and its context were examined further

by using an aliquot of the genomic DNA that had been provided years earlier to Genome Therapeutics Corporation for genome sequencing and that was kindly given to us by T. L. Cover. PCR tests of this DNA revealed the internally deleted

TABLE 2. Geographic distribution of *tfs3* elements^a

Country	No. of strains tested	No. (%) containing <i>tfs3</i>	
		Full	Partial
Japan	24	4 (17)	7 (29)
Peru	24	6 (25)	4 (17)
Spain	24	6 (25)	1 (4)
India	22	3 (14)	6 (27)
Total	94	19 (20)	18 (19)

^a This distribution was determined by PCR with primers specific for each of 15 ORFs in *tfs3* (see Fig. 1). Primer sequences are available from the authors upon request.

copy of *tfs3* and the *tfs3-hp927* and *jhp926-jhp927* connections noted above. However, this DNA also seemed to contain full-length *tfs3*, based on PCR tests for each *tfs3* ORF. These results imply that the DNA used for J99 genome sequencing and the original J99 culture may have contained two closely related strains: one with full-length *tfs3* and one with the *tfs3* remnant. Presumably, only the strain with the remnant was archived or survived long-term frozen storage.

Limited divergence among *H. pylori* isolates recovered from individual patients, reminiscent of that found in DNA from the original J99 culture, had been reported sporadically in the past (26, 34, 53) and emerged again with further studies of *H. pylori* that were recovered from patient J99 by re-endoscopy and culture 6 years after his initial endoscopy (31). This yielded a complex population of strains that were closely related to, but not identical to, the original J99 strain (31). Our PCR analysis of 14 of these new isolates (DNAs kindly provided by D. Israel) identified three main variants with respect to *tfs3* region markers (diagrammed as variants A, B, and C in Fig. 2), none of which was perfectly matched to the *tfs3* remnant in archived strain J99 (line D in Fig. 2).




FIG. 2. Structures of *tfs3* regions in J99 variants from patient J99, recovered in the second endoscopy (31) 6 years after isolation of archived strain J99. Gene designations here (B, C, D, etc.) are as in Fig. 1, and are annotated in Table 1. RJ and LJ, right and left junctions, respectively, of *tfs3* with flanking sequences. (A) Variant A. Shown is the full-length *tfs3* region, as in strain PeCan18B, containing two copies of *jhp927* (clones A1, A7, A9, C5, C6, C8, and D1) (31). (B) Variant B. Variant B is similar to variant A, except it lacks the *jhp926-jhp927* segment and has deletion or rearrangement at *tfs3*'s left end (clones A3, A5, C10, C12, and D3). (C) Variant C. Variant C is a truncated *tfs3*, lacking *tfs3* sequences that extend rightward from the 5' end of *orfG* (*hp459*) and also *jhp927* (clones Ca1 and C2). (D) Variant D. *tfs3* remnant in archived stock of strain J99 found here (Fig. 1B).

Strain	left flanking sequence	right flanking sequence
	J99	tacacaTgcacaaaaaagaacccaagaacc
SpainB43
SpainB49
PeCan9A
India-Chennai2g.T.a.....	a.....g.....
SpainB50	aca.a.....g.T.a.....	a.....c.t.....g.
JapCPY2363g.T.a.....	a.....
JapHU131g.T.a.....	a.....
JapCPY6081g.T.a.....	a.....c.t.....g.
India39Ag.T.a.....	a.....g.....
SpainB70g.T.a.....	a.....g.....

Strain	<i>tfs3</i> right end	right flanking sequence
	PeCan18B	CTAACAAAAAGAGTAAAGAAATTAAGAGA
SpainB84c.c.....c.t.....g.
PeruSJM148c.c.....c.t.....g.
PeruSJM9	.C.G.....AC.TC..ACT.....	ag.....c.....a.
PeruSJM91	.C.G.....AC.TC..ACT.....	ag.....c.....a.
India75	.C.G.....AC.TC..ACT.....	ag.....c.....t.....g.
J99archived

Strain	left flanking sequence	<i>tfs3</i> left end
	PeCan18B	gtggagcggccatgttatttaactcttaaca
PeruSJM148	taggggttattcttattcttattcttattctta
SpainB84	cttcttcttcttctataactttaaaggcttg
PeCan10B	gactactcttctataactttagacgcctca
SpainB84	atcgaaacttagagaacacttaaaacttaca
J99arch	tttattcttggggagaatattagccta
26695	tcgttctagtctgctgttactaaacca
JapHU87	ctttatgaaataagaatgataataaaccag.....
JapCPY1672	taaggttagtggaacgccttaagtcttca	a.....
India77A

FIG. 3. Empty site in the 5' end of *jhp927* and left and right *tfs3* junctions. *tfs3* sequences are in uppercase, and flanking sequences are in lowercase. (A) The empty site is the site of *tfs3* right-end insertion in codon 9 of *jhp927* (probable translation start is underlined). This start is 33 codons downstream from the start assigned in strain J99 due to the presence of one extra nucleotide in the annotated *jhp927* sequence (3). The underline designates the first in-frame ATG in these 10 strains. PCR and sequence analysis indicated that *jhp927* is adjacent to *jhp926* in strains J99, SpainB43, SpainB49, PeCan9A and India-Chennai2 and is adjacent to *jhp926like* (GenBank accession no. AY128680) in strains SpainB50, JapCPY2362, JapHU131, JapCPY6081, India39A, and SpainB70. (B) Right junction sequences of the predominant type (Table 3). (C) Left junctions of *tfs3*. The sequences are adjacent to the following: in PeCan18B, to *jhp949*; in archived J99, to the intergenic region of *jhp941-jhp942*; in SJM148(Peru), to a site 38 nt upstream of a 5S ribosomal DNA gene; in PeCan10B, to an unknown sequence (no homology in database); in SpainB84, to a weak homolog of *hp1383* and *jhp1422* (restriction modification system S subunit, 55 to 66% protein identity and similarity); in 26695, to the 3' end of *hp454*; in JapHU87, to *hp513*; in JapCPY1672, to the *jhp644-jhp645* intergenic region; and in India77A, to a *jhp1297* homolog.

***tfs3* ends, flanking sequences, and site of insertion.** PCR and sequencing further identified the left and right *tfs3* ends and empty sites in strains lacking *tfs3* and gave insights into insertion specificity. First, we tested whether the putative *tfs3* right-end-*jhp927* junction of strain PeCan18B was typical or unique. PCR using *orfP*- and *jhp927*-specific primers resulted in a product of the expected size from 25 of 41 strains with full-length *tfs3* elements that were tested. DNA sequencing of PCR products from five such strains and comparison with the published strain J99 genome sequence identified *tfs3*'s right end at the ninth codon of *jhp927* (Fig. 3, right junction). Eight of the remaining 16 *tfs3*-containing strains yielded PCR products with primers specific for the *tfs3* right end and *jhp931* instead of *jhp927*. Products from six of these eight strains were of one size class (variant 1a in Table 3), and DNA sequences from three of them suggested that these variants had arisen by deletion involving a direct repeat of 22 nt, ATTTTAAAAGGAGCAACAAATG, present near the *tfs3* right end and again near the *jhp931* start; a second variant end (variant 1b) was ascribed to a shorter direct repeat, GCAACC(G/C)AA, at the *tfs3* right

TABLE 3. *tfs3* right junction types

Country	Total no. of strains	No. of strains with complete <i>tfs3</i>	No. (%) of strains with right junction type ^a				
			1	Variant			Unidentified
				1a	1b	1c	
Japan	24	4 (17)	3 (75)	0	0	0	1 (25)
Peru	24	6 (25)	5 (83)	0	0	1 (17)	0
Spain	24	6 (25)	3 (50)	3 (50)	0	0	0
India	121	25 (21)	14 (56)	3 (12)	1 (4)	0	7 (28)
Total	193	41 (21)	25 (61)	6 (15)	1 (2)	1 (2)	8 (20) ^b

^a Predominant type 1 is described in the legend to Fig. 3. Variant types 1a and 1b contain short deletions that can be ascribed to slippage between direct repeats (see text). Variant type 1c contains a 23-nt deletion from the right end of *tfs3* to a site 110 nt from the 3' end of *jhp930*.

^b About 20% of strains did not amplify with available primers, suggesting that they might contain other rearrangements. However, chromosomal sequencing of three such strains identified a *tfs3-jhp927* connection (as in Fig. 3) in each case.

end and upstream of *jhp931*; and a third variant (1c) did not seem to be associated with repeat sequences.

The lack of amplification with *orfP*- and *jhp927*- or *jhp931*-specific primers in cases of eight *tfs3*-containing strains ("unidentified" column in Table 3) was attributable to chromosome rearrangement or to sequence divergence affecting primer binding sites. Direct sequencing of genomic DNAs from three of these eight strains identified a typical *tfs3-jhp927* connection in each case. This outcome further supports the inference that the *tfs3* right end inserts preferentially at the ninth codon of *jhp927*.

We next tested the possibility that *jhp926* and *jhp927* might be adjacent to one another unless disrupted by *tfs3* insertion. PCR tests of strains that contained *jhp927* but not *tfs3* identified this connection in only 11 of 41 strains tested. Primer walking was carried out to identify a sequence adjacent to the 5' end of *jhp927* by using genomic DNA from 1 of the 30 other strains. The sequence obtained was distantly related to *jhp926* and will be called "*jhp926like*" (34% protein identity, 43% protein similarity, no significant DNA identity; GenBank accession no. AY128680). Further PCR tests identified a *jhp926like-jhp927* connection in 28 of the remaining 29 strains that carried *jhp927* but not canonical *jhp926*. The *jhp926like* sequence can be considered a divergent allele of *jhp926* based on (i) its connection to *jhp927*, (ii) the inability to find any strain among 94 tested that contained both *jhp926like* and *jhp926*, and (iii) the protein-level homology noted above. Both *jhp926* and *jhp926like* alleles were found in Peruvian, Spanish, and Indian populations, although *jhp926like* was twice as common as canonical *jhp926* itself; only *jhp926like* (9 strains) was found among the 24 Japanese strains tested. Neither *jhp926* nor *jhp926like* was found next to the *tfs3* left end in any of 37 strains containing partial or complete *tfs3* elements, nor were these genes found at all in the genomes of 16 of 19 strains with apparently full-length *tfs3* elements; the three exceptions contained duplication of *jhp927*, equivalent to that in strain J99. Eight of 18 strains with only partial *tfs3* elements (mainly the left end) also lacked *jhp926* sequences, and the other 10 contained either *jhp926* (1 strain) or *jhp926like* (9 strains) connected to *jhp927*.

The left end of *tfs3* had been identified provisionally at 79 nt upstream of the 3' end of *hp454*, based on divergence between

sequences from strains PeCan18B and 26695 (Fig. 3). Consistent with this, primer walking along *tfs3* in archived J99 showed that 76 nt matching the 3' end of *hp454* were joined to a segment that seemed to correspond to the intergenic region between *jhp941* and *jhp942*. However, none of these three connections (*jhp949-orfB* in PeCan18B, *jhp941/2* intergenic region-*orfB* in J99, *hp454* [5' end]-*orfB* in 26695) was found in any of 37 other *tfs3*-containing strains tested by PCR. Left-end junction sequences were determined by primer walking from *hp455* by using three additional strains containing full-length *tfs3* elements and three others containing left-end remnants of *tfs3*. In each case, the 79-nt segment from *hp454*'s 3' end was joined to a different sequence (Fig. 3). These results established again that the left end of *tfs3* was about 79 nt from *hp454*'s 3' end, that this end was well conserved, and that it could be joined to many target sites.

Test for possible *tfs3* function. The closest homologs of four *tfs3* genes were *comB* genes of *H. pylori* (*comB7* to *comB10*, Table 1), which are needed for competence in DNA transformation (30). Accordingly, three mutant derivatives of strain PeCan18B were constructed to test if these *tfs3* genes might participate in natural transformation, such that the corresponding *comB* genes would be redundant. The three derivatives contained (i) a deletion of the entire *comB7*-to-*comB10* region and replacement with a *cat* (chloramphenicol resistance) gene, (ii) a deletion of the entire *tfs3* segment and replacement with an *aphA* (kanamycin resistance) gene, and (iii) replacement of both gene clusters by *cat* and *aphA* genes. These three derivatives were then used as recipients for natural transformation using a 16S rDNA PCR fragment that confers tetracycline resistance (due to mutation in the tetracycline binding site) (20). Tet^r transformants were obtained at frequencies of about 10⁻⁶ by using the PeCan18B wild type and its *tfs3* deletion derivative as recipients, but at ≤10⁻⁹ when using PeCan18B derivatives with the *comB* gene deletion alone or in combination with the *tfs3* deletion. Thus, these data did not support the idea that *tfs3 virB* homologs contribute to transformability.

Tests for *tfs3* in several additional strains gave no support for models in which *tfs3* might have a critical role in general bacterial conjugation, in intracellular entry or survival, or in mouse colonization. First, no *tfs3* sequences were found in HPK5, a strain that seemed to serve as a donor in conjugation (36). Second, no *tfs3* sequences were found in strain G27, which is quite invasive (4), and, as noted above, only remnant *tfs3* elements were found in strains J99 and 26695, two other strains that also enter epithelial cells quite well (37). Third, neither of the two special and widely used mouse-colonizing strains, SS1 or X47-2AL, was found to contain *tfs3* sequences.

DISCUSSION

We have described a new gene cluster in *H. pylori*, designated *tfs3*, 7 of whose 16 genes are protein-level homologs of genes involved in type IV secretion. Four of these genes are transmembrane pore genes (*virB7*, *virB8*, *virB9*, and *virB10*), and three code for ATPases that move cognate macromolecule substrates to and through the pore with specificity and efficiency (*virB4*, *virB11*, and *virD4*). None of the nine other *tfs3* ORFs had homologs in current databases, but transmembrane

motifs encoded in four of them also suggest possible type IV secretion involvement. Full-length and truncated *tfs3* elements were each found in about one-fifth of *H. pylori* strains—obtained from Peru, Spain, India, and Japan. In all cases analyzed, they were inserted in the plasticity zone, a genomic region rich in insertions and deletions. This is the third type IV secretion gene complex found in *H. pylori*, but the first found to be in a putative transposable element (discussed below) or in *H. pylori*'s special plasticity zone.

tfs3's role in *H. pylori* is not known. The closest known homologs of some *tfs3* genes are needed in *H. pylori* for competence in natural transformation; more distant homologs form part of the *cag* PAI and contribute to virulence. There are also striking homologies with genes in a virulence plasmid of *Campylobacter jejuni* that are needed for this pathogen's intracellular growth and competence for transformation (10). Our mutational test did not support a possible *tfs3* involvement in transformation, nor was an association of *tfs3* carriage and overt disease detected: *tfs3* was either entirely absent or was present only in truncated form in several strains that can persist in cultured epithelial cells and was absent from a strain that acts as a donor in conjugation. In terms of other possible roles, *H. pylori* strains are diverse phenotypically, especially in their interactions with host cells and tissues during chronic infection (23, 39, 44), and it is thus tempting to imagine that *tfs3*-encoded proteins may help secrete macromolecules involved in some of these interactions.

An unusual *tfs3* derivative with two internal deletions was found in an archived copy of reference strain J99, and both this vestigial element and full-length *tfs3* were found in an aliquot of J99 DNA that had been provided for genome sequencing, but no *tfs3* genes were reported in J99's full genome sequence. This discrepancy might be explained by supposing that a J99 variant, putatively free of *tfs3*, was used to complete the genome sequence and that inconsistent sequences were omitted from the final contig. (This project was started by one company and finished by another several years later [3, 6].) Another explanation invokes difficulty in *tfs3* segment cloning, possibly related to its genes for membrane proteins, and then in silico assembly of a circular contig without *tfs3* and one *jhp927* region, due to the duplicated *jhp927* region.

The principle that *H. pylori* infections can be mixed and can involve clusters of related strains (26, 34, 53) was well illustrated by the study by Israel et al. (31) of *H. pylori* isolates recovered from "patient J99" at the second endoscopy, 6 years after the one that had yielded the original reference strain. We found three *tfs3* variants in 14 new J99-related isolates, none of which completely matched the *tfs3* element found in the archived strain. This was in accord with the diversity Israel et al. found at other loci. Possible explanations for such variant strains include (i) mutant or recombinant formation in the years between the first and second endoscopies (as postulated in reference 31); (ii) the strain's being present at the time of the first endoscopy, but not recovered by culture; and (iii) the strain being transmitted to the patient from other family members during the intervening 6 years.

The diversity in gene content among *H. pylori* strains stimulates interest in how DNA is gained and lost during *H. pylori* evolution. *tfs3*'s lower G+C content (35% versus 39% for the *H. pylori* genome overall) suggests transfer from another bac-




FIG. 4. Motorized transposition model for *tfs3* acquisition. This model envisions that two at least subtly different sets of transposition proteins (hatched ovals) act on *tfs3*'s two ends and then bind to a specific recognition site in target DNA (solid oval). DNAs adjacent to this recognition site are pulled progressively through the protein complex until they encounter sequences or DNA-bound proteins that *tfs3* uses as signals for left-end joining to one of many potential target sites and right-end joining to its specific *jhp927* site. jcn, junction.

terial species, as does its presence in only a subset of *H. pylori* strains. *tfs3* entry into *H. pylori* gene pools might have occurred long ago, because *tfs3* is common among strains from both East Asia and the West. Alternatively, it might have been transferred into *H. pylori* recently and could be increasing in abundance by lateral DNA transfer and insertion into new genomes ("molecular drive") (21, 22) at rates that also reflect any effects of this element on bacterial fitness.

In terms of possible acquisition mechanisms, our analyses showed (i) that left and right ends of full-length *tfs3* elements were conserved; (ii) that *tfs3*'s right end was joined to the same target site in *jhp927* in most *H. pylori* strains; whereas (iii) its left end could be joined to many different target sequences; and (iv) that the *jhp926* or *jhp926like* genes to the left of *tfs3*'s fixed insertion site were missing from most strains with full-length *tfs3*. These findings fit with a formal model of clean insertion into a fixed site in *jhp927*, followed by formation of leftward deletions, equivalent to those generated by elements such as IS1, Tn5, and $\gamma\delta$ (32, 49, 52, 54). Because such deletions are rare in other systems, however, we prefer a model in which the deletions observed here result directly from *tfs3* insertion and the joining of its right and left ends to different chromosomal sites. Although none of the proteins or specific DNA sites needed for *tfs3* transposition are known, we are attracted to a model that entails "motorized" tracking of recipient DNA through the transposition complex (Fig. 4), prompted in part by knowledge of how type I restriction endonucleases act (13, 29, 42, 47). Insertion would begin with binding of a *tfs3*-transposition protein complex to a specific

target site. Adjacent recipient DNA would then be pulled by the complex from both sides, perhaps coupled to ATP hydrolysis, until signals for DNA cleavage and *tfs3* end joining are encountered. Signals for cleavage might be inherent in the DNA sequence itself, as is seen with RecBC and related recombination enzymes (5, 7, 17), or it might consist of other proteins that the transposition complex encounters along the recipient DNA, as is seen with type I restriction endonucleases (13, 29, 42, 47). Activation of latent nucleases, as in MutSLH-mediated mismatch repair (12), also merits consideration. Finally, differences between right and left junctions suggest that different signals may be used to trigger the processing of sites for left and right end joining.

The truncated (left-end containing) *tfs3* elements can be explained by this tracking model, based on a finding that half of *tfs3*-free *H. pylori* strains lacked the *jhp927* site for right-end insertion. We propose that transposition in these strains also entails transposition protein binding to the preferred target site, tracking bidirectionally along target DNA, and left-end joining to one of its many possible target sites, essentially as in Fig. 4. Right-end joining might fail, however, either because of deletion or point mutation of its specific target sequence or because of the lack of a putative protein signal. Joining of just one end would linearize the chromosome, but recircularization would be needed for viability, and this might be achieved by illegitimate recombination, which can occur by various mechanisms (8, 24, 33).

Because most deletions are either potentially deleterious or at best only neutral, transposable elements that regularly generate them would usually be lost during evolution. *tfs3* may be an exception, however, able to escape such contraselection in two ways. First, if the initial target is between sites of *tfs3* joining to recipient DNAs, as postulated (Fig. 4), *tfs3* insertion would remove this site. This might be selected because it would lower the chance of displacing resident *tfs3*s by "superinfecting" *tfs3* elements. Second, many deletions formed by *tfs3* insertion might not be deleterious because they occur in the plasticity zone, where DNA loss and rearrangement are the norm, and flexibility and diversity in gene content may contribute to bacterial fitness in different members of the diverse human host population.

ACKNOWLEDGMENTS

We thank Timothy Cover and Dawn Israel for providing DNAs from J99 and related *H. pylori* strains and stimulating discussion, Daiva Dailidienė and Keiji Ogura for mutant DNAs, Patricia Guerry for access to her *C. jejuni* pVir plasmid data before publication, and anonymous reviewers for valuable suggestions.

This research was supported by grants from the Public Health Service (AI38166, AI49161, DK53727, P30 DK52574, and 3 D43 TW 00910).

REFERENCES

- Akopyants, N. S., S. W. Clifton, D. Kersulyte, J. E. Crabtree, B. E. Youree, C. A. Reece, N. O. Bukanov, E. S. Drazek, B. A. Roe, and D. E. Berg. 1998. Analyses of the *cag* pathogenicity island of *Helicobacter pylori*. *Mol. Microbiol.* **28**:37–53.
- Akopyants, N. S., A. Fradkov, L. Diatchenko, P. D. Siebert, S. Lukyanov, E. D. Sverdlov, and D. E. Berg. 1998. PCR-based subtractive hybridization and differences in gene content among strains of *Helicobacter pylori*. *Proc. Natl. Acad. Sci. USA* **95**:13108–13113.
- Alm, R. A., L. S. Ling, D. T. Moir, B. L. King, E. D. Brown, P. C. Doig, D. R. Smith, B. Noonan, B. C. Guild, B. L. deJonge, G. Carmel, P. J. Tummino, A. Caruso, M. Uria-Nickelsen, D. M. Mills, C. Ives, R. Gibson, D. Merberg,
- S. D. Mills, Q. Jiang, D. E. Taylor, G. F. Vovis, and T. J. Trust. 1999. Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* **397**:176–180.
- Amieva, M. R., N. R. Salama, L. S. Tompkins, and S. Falkow. 2002. *Helicobacter pylori* enter and survive within multivesicular vacuoles of epithelial cells. *Cell Microbiol.* **4**:677–690.
- Amundsen, S. K., and G. R. Smith. 2003. Interchangeable parts of the *Escherichia coli* recombination machinery. *Cell* **112**:741–744.
- Anonymous. 1995. The Gold Bug: *Helicobacter pylori*. *Science* **267**:173.
- Arnold, D. A., N. Handa, I. Kobayashi, and S. C. Kowalczykowski. 2000. A novel, 11 nucleotide variant of chi, chi*: one of a class of sequences defining the *Escherichia coli* recombination hotspot chi. *J. Mol. Biol.* **300**:469–479.
- Ashizawa, Y., T. Yokochi, Y. Ogata, Y. Shobuike, J. Kato, and H. Ikeda. 1999. Mechanism of DNA gyrase-mediated illegitimate recombination: characterization of *Escherichia coli* *gyrA* mutations that confer hyper-recombination phenotype. *J. Mol. Biol.* **289**:447–458.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A., Smith, and K. A. Struhl. 1994. Current protocols in molecular biology; supplement 27 CPMB, p. 2.4.1. Greene Publishing and Wiley Interscience, New York, N.Y.
- Bacon, D. J., R. A. Alm, L. Hu, T. E. Hickey, C. P. Ewing, R. A. Batchelor, T. J. Trust, and P. Guerry. 2002. DNA sequence and mutational analyses of the pVir plasmid of *Campylobacter jejuni* 81–176. *Infect. Immun.* **70**:6242–6250.
- Berg, D. E., R. H. Gilman, J. Lelwala-Guruge, K. Srivastava, Y. Valdez, J. Watanabe, J. Miyagi, N. S. Akopyants, A. Ramirez-Ramos, T. H. Yoshiwara, S. Recavarren, and R. Leon-Barua. 1997. *Helicobacter pylori* populations in individual Peruvian patients. *Clin. Infect. Dis.* **25**:996–1002.
- Blackwell, L. J., K. P. Bjornson, D. J. Allen, and P. Modrich. 2001. Distinct MutS DNA-binding modes that are differentially modulated by ATP binding and hydrolysis. *J. Biol. Chem.* **276**:34339–34347.
- Bourniquel, A. A., and T. A. Bickle. 2002. Complex restriction enzymes: NTP-driven molecular motors. *Biochimie* **84**:1047–1059.
- Cao, T. B., and M. H. Saier, Jr. 2001. Conjugal type IV macromolecular transfer systems of gram-negative bacteria: organismal distribution, structural constraints and evolutionary conclusions. *Microbiology* **147**:3201–3214.
- Chalkauskas, H., D. Kersulyte, I. Cepulienė, V. Urbonas, D. Ruzeviciene, A. Barakauskiene, A. Raudonikiene, and D. E. Berg. 1998. Genotypes of *Helicobacter pylori* in Lithuanian families. *Helicobacter* **3**:296–302.
- Chalker, A. F., H. W. Minehart, N. J. Hughes, K. K. Koretke, M. A. Lonetto, K. K. Brinkman, P. V. Warren, A. Lupas, M. J. Stanhope, J. R. Brown, and P. S. Hoffman. 2001. Systematic identification of selective essential genes in *Helicobacter pylori* by genome prioritization and allelic replacement mutagenesis. *J. Bacteriol.* **183**:1259–1268.
- Chedin, F., and S. C. Kowalczykowski. 2002. A novel family of regulated helicases/nucleases from gram-positive bacteria: insights into the initiation of DNA recombination. *Mol. Microbiol.* **43**:823–834.
- Christie, P. J. 2001. Type IV secretion: intercellular transfer of macromolecules by systems ancestrally related to conjugation machines. *Mol. Microbiol.* **40**:294–305.
- Cover, T. L., D. E. Berg, M. J. Blaser, and H. L. T. Mobley. 2001. *H. pylori* pathogenesis, p. 509–558. In E. A. Groisman (ed.), Principles of bacterial pathogenesis. Academic Press, New York, N.Y.
- Dailidienė, D., M. T. Bertoli, J. Miciuleviciene, A. K. Mukhopadhyay, G. Dailide, M. A. Pascasio, L. Kupcinskias, and D. E. Berg. 2002. Emergence of tetracycline resistance in *Helicobacter pylori*: multiple mutational changes in 16S ribosomal DNA and other genetic loci. *Antimicrob. Agents Chemother.* **46**:3940–3946.
- Doolittle, W. F., and C. Sapienza. 1980. Selfish genes, the phenotype paradigm and genome evolution. *Nature* **284**:601–603.
- Dover, G., and W. F. Doolittle. 1980. Modes of genome evolution. *Nature* **88**:646–647.
- Dubreuil, J. D., G. Del Giudice, and R. Rappuoli. 2002. *Helicobacter pylori* interactions with host serum and extracellular matrix proteins: potential role in the infectious process. *Microbiol. Mol. Biol. Rev.* **66**:617–629.
- Ehrlich, S. D., H. Biernie, E. d'Alencon, D. Vilette, M. Petranovic, P. Noirot, and M. Michel. 1993. Mechanisms of illegitimate recombination. *Gene* **135**:161–166.
- Ernst, P. B., and B. D. Gold. 2000. The disease spectrum of *Helicobacter pylori*: the immunopathogenesis of gastroduodenal ulcer and gastric cancer. *Annu. Rev. Microbiol.* **54**:615–640.
- Falush, D., C. Kraft, N. S. Taylor, P. Correa, J. G. Fox, M. Achtman, and S. Suerbaum. 2001. Recombination and mutation during long-term gastric colonization by *Helicobacter pylori*: estimates of clock rates, recombination size, and minimal age. *Proc. Natl. Acad. Sci. USA* **98**:15056–15061.
- Fischer, W., R. Haas, and S. Odenbreit. 2002. Type IV secretion systems in pathogenic bacteria. *Int. J. Med. Microbiol.* **292**:159–168.
- Fischer, W., J. Puls, R. Buhrdorf, B. Gebert, S. Odenbreit, and R. Haas. 2001. Systematic mutagenesis of the *Helicobacter pylori* *cag* pathogenicity island: essential genes for *CagA* translocation in host cells and induction of interleukin-8. *Mol. Microbiol.* **42**:1337–1348.
- Garcia, L. R., and I. J. Molineux. 1999. Translocation and specific cleavage

- of bacteriophage T7 DNA in vivo by EcoKI. *Proc. Natl. Acad. Sci. USA* **96**:12430–12435.
30. Hofreuter, D., S. Odenbreit, and R. Haas. 2001. Natural transformation competence in *Helicobacter pylori* is mediated by the basic components of a type IV secretion system. *Mol. Microbiol.* **41**:379–391.
 31. Israel, D. A., N. Salama, U. Krishna, U. M. Rieger, J. C. Atherton, S. Falkow, and R. M. Peek, Jr. 2001. *Helicobacter pylori* genetic diversity within the gastric niche of a single human host. *Proc. Natl. Acad. Sci. USA* **98**:14625–14630.
 32. Jilk, R. A., J. C. Makris, L. Borchardt, and W. S. Reznikoff. 1993. Implications of Tn5-associated adjacent deletions. *J. Bacteriol.* **175**:1264–1271.
 33. Juhala, R. J., M. E. Ford, R. L. Duda, A. Youton, G. F. Hatfull, and R. W. Hendrix. 2000. Genomic sequences of bacteriophages HK97 and HK022: pervasive genetic mosaicism in the lambdoid bacteriophages. *J. Mol. Biol.* **299**:27–51.
 34. Kersulyte, D., H. Chalkauskas, and D. E. Berg. 1999. Emergence of recombinant strains of *Helicobacter pylori* during human infection. *Mol. Microbiol.* **31**:31–43.
 35. Kersulyte, D., B. Velapatiño, G. Dailide, A. K. Mukhopadhyay, Y. Ito, L. Cahuayme, A. J. Parkinson, R. H. Gilman, and D. E. Berg. 2002. Transposable element ISHp608 of *Helicobacter pylori*: nonrandom geographic distribution, functional organization, and insertion specificity. *J. Bacteriol.* **184**:992–1002.
 36. Kuipers, E. J., D. A. Israel, J. G. Kusters, and M. J. Blaser. 1998. Evidence for a conjugation-like mechanism of DNA transfer in *Helicobacter pylori*. *J. Bacteriol.* **180**:2901–2905.
 37. Kwok, T., S. Backert, H. Schwarz, J. Berger, and T. F. Meyer. 2002. Specific entry of *Helicobacter pylori* into cultured gastric epithelial cells via a zipper-like mechanism. *Infect. Immun.* **70**:2108–2120.
 38. Lee, V. T., and O. Schneewind. 2001. Protein secretion and the pathogenesis of bacterial infections. *Genes Dev.* **15**:1725–1752.
 39. Mahdavi, J., B. Sonden, M. Hurtig, F. O. Olfat, L. Forsberg, N. Roche, J. Angstrom, T. Larsson, S. Teneberg, K. A. Karlsson, S. Altraja, T. Wadstrom, D. Kersulyte, D. E. Berg, A. Dubois, C. Petersson, K. E. Magnusson, T. Norberg, F. Lindh, B. B. Lundskog, A. Arnqvist, L. Hammarstrom, and T. Boren. 2002. *Helicobacter pylori* SabA adhesin in persistent infection and chronic inflammation. *Science* **297**:573–578.
 40. Mobley, H. L. T., G. L. Mendez, and S. L. Hazell (ed.). 2001. *Helicobacter pylori*: physiology and genetics. ASM Press, Washington, D.C.
 41. Mukhopadhyay, A. K., D. Kersulyte, J.-Y. Jeong, S. Datta, Y. Ito, A. Chowdhury, S. Chowdhury, A. Santra, S. K. Bhattacharya, T. Azuma, G. B. Nair, and D. E. Berg. 2000. Distinctiveness of genotypes of *Helicobacter pylori* in Calcutta, India. *J. Bacteriol.* **182**:3219–3227.
 42. Murray, N. E. 2000. Type I restriction systems: sophisticated molecular machines (a legacy of Bertani and Weigle). *Microbiol. Mol. Biol. Rev.* **64**:412–434.
 43. Occhialini, A., A. Marais, R. Alm, F. Garcia, R. Sierra, and F. Mégraud. 2000. Distribution of open reading frames of plasticity region of strain J99 in *Helicobacter pylori* strains isolated from gastric carcinoma and gastritis patients in Costa Rica. *Infect. Immun.* **68**:6240–6249.
 44. Rhen, M., S. Eriksson, M. Clements, S. Bergström, and S. J. Normark. 2003. The basis of persistent bacterial infections. *Trends Microbiol.* **11**:80–86.
 45. Salama, N., K. Guillemin, T. K. McDaniel, G. Sherlock, L. Tompkins, and S. Falkow. 2000. A whole-genome microarray reveals genetic diversity among *Helicobacter pylori* strains. *Proc. Natl. Acad. Sci. USA* **97**:14668–14673.
 46. Sexton, J. A., and J. P. Vogel. 2002. Type IVB secretion by intracellular pathogens. *Traffic* **3**:178–185.
 47. Studier, F. W., and P. K. Bandyopadhyay. 1988. Model for how type I restriction enzymes select cleavage sites in DNA. *Proc. Natl. Acad. Sci. USA* **85**:4677–4681.
 48. Tomb, J. F., O. White, A. R. Kerlavage, R. A. Clayton, G. G. Sutton, R. D. Fleischmann, K. A. Ketchum, H. P. Klenk, S. Gill, B. A. Dougherty, K. Nelson, J. Quackenbush, L. Zhou, E. F. Kirkness, S. Peterson, B. Loftus, D. Richardson, R. Dodson, H. G. Khalak, A. Glodek, K. McKenney, L. M. Fitzgerald, N. Lee, M. D. Adams, E. K. Hickey, D. E. Berg, J. D. Gocayne, T. R. Utterback, J. D. Peterson, J. M. Kelley, M. D. Cotton, J. M. Weidman, C. Fujii, C. Bowman, L. Watthey, E. Wallin, W. S. Hayes, M. Borodovsky, P. D. Karp, H. O. Smith, C. M. Fraser, and J. C. Venter. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **388**:539–547.
 49. Tomcsanyi, T., C. M. Berg, S. H. Phadnis, and D. E. Berg. 1990. Intramolecular transposition by a synthetic IS50 (Tn5) derivative. *J. Bacteriol.* **172**:6348–6354.
 50. Trieber, C. A., and D. E. Taylor. 2002. Mutations in the 16S rRNA genes of *Helicobacter pylori* mediate resistance to tetracycline. *J. Bacteriol.* **184**:2131–2140.
 51. Trieu-Cuot, P., G. Gerbaud, T. Lambert, and P. Courvalin. 1985. In vivo transfer of genetic information between gram-positive and gram-negative bacteria. *EMBO J.* **4**:3583–3587.
 52. Turlan, C., and M. Chandler. 1995. IS1-mediated intramolecular rearrangements: formation of excised transposon circles and replicative deletions. *EMBO J.* **14**:5410–5421.
 53. van der Ende, A., E. A. Rauws, M. Feller, C. J. Mulder, G. N. Tytgat, and J. Dankert. 1996. Heterogeneous *Helicobacter pylori* isolates from members of a family with a history of peptic ulcer disease. *Gastroenterology* **111**:638–647.
 54. Wang, G., R. W. Blakesley, D. E. Berg, and C. M. Berg. 1993. pDUAL: a transposon-based cosmid cloning vector for generating nested deletions and DNA sequencing templates in vivo. *Proc. Natl. Acad. Sci. USA* **90**:7874–7878.