

# Clustering Induces a Lateral Redistribution of $\alpha 2\beta 1$ Integrin from Membrane Rafts to Caveolae and Subsequent Protein Kinase C–dependent Internalization

Paula Upla,\* Varpu Marjomäki,\* Pasi Kankaanpää,\* Johanna Ivaska,<sup>†</sup>  
Timo Hyypiä,<sup>‡</sup> F. Gisou van der Goot,<sup>§</sup> and Jyrki Heino\*<sup>¶</sup>

\*Cell Biology, University of Jyväskylä, FIN-40351 Jyväskylä, Finland; <sup>†</sup>VTT Medical Biotechnology, FIN-20520 Turku, Finland; <sup>‡</sup>Department of Microbiology, University of Oulu, FIN-90220 Oulu, Finland; and <sup>§</sup>Department of Genetics and Microbiology, University of Geneva, 1211 Geneva 4, Switzerland

Submitted August 13, 2003; Revised October 29, 2003; Accepted November 13, 2003  
Monitoring Editor: Carl-Henrik Heldin

**Integrin  $\alpha 2\beta 1$  mediates the binding of several epithelial and mesenchymal cell types to collagen. The composition of the surrounding plasma membrane, especially caveolin-1- and cholesterol-containing membrane structures called caveolae, may be important to integrin signaling. On cell surface  $\alpha 2\beta 1$  integrin was located in the raft like membrane domain, rich in GPI-anchored proteins, rather than in caveolae. However, when antibodies were used to generate clusters of  $\alpha 2\beta 1$  integrin, they started to move laterally on cell surface along actin filaments. During the lateral movement small clusters fused together. Finally  $\alpha 2\beta 1$  integrin was found inside caveolae and subsequently internalized into caveosome-like perinuclear structures. The internalization process, unlike cluster formation or lateral redistribution, was dependent on protein kinase C $\alpha$  activity. Caveolae are known to be highly immobile structures and  $\alpha 2\beta 1$  integrin clusters represent a previously unknown mechanism to activate endocytic trafficking via caveolae. The process was specific to  $\alpha 2\beta 1$  integrin, because the antibody-mediated formation of  $\alpha V$  integrin clusters activated their internalization in coated vesicles and early endosomes. In addition to natural ligands human echovirus-1 (EV1) gains entry into the cell by binding to  $\alpha 2\beta 1$  and taking advantage of  $\alpha 2\beta 1$  internalization via caveolae.**

## INTRODUCTION

Caveolae are cave-like invaginations of the cell surface (for reviews see Kurzchalia and Parton, 1999; Pelkmans and Helenius, 2002; Parton 2003). They have been described as highly immobile and not involved in constitutive endocytic trafficking (Thomsen *et al.*, 2002). Their internalization can be activated, for example, by the phosphatase inhibitor okadaic acid (Parton *et al.*, 1994). In endothelial cells the interaction of albumin docking protein gp60 and caveolin-1 initiates the endocytosis of caveolae (Minshall *et al.*, 2000). Caveolins are membrane proteins that are molecular markers of caveolae and shown to be crucial for the formation of these structures (Fra *et al.*, 1995). Caveolins can be associated with cell surface growth factor receptors (Couet *et al.*, 1997) and cell adhesion receptors (Wary *et al.*, 1996, 1998; Wei *et al.*, 1999) and caveolae might therefore play a role in cellular signaling. In addition, caveolae have been suggested to participate in the uptake of folate by pinocytosis, but this is still under

discussion (Parton, 2003). Caveolin-deficient mice have given novel insight into the function of caveolae. Caveolin-1 gene knockout leads to pulmonary defects, vasoconstriction, or dilatation abnormalities and resistance to diet-induced obesity (Drab *et al.*, 2001; Razani *et al.*, 2001; Schubert *et al.*, 2001; Sotgia *et al.*, 2002; Razani *et al.*, 2002). Simian virus 40 (SV40) has been shown to be internalized through caveolae and detailed studies focused on the virus entry mechanism have produced a lot of new information about the biology of caveolae (Pelkmans *et al.*, 2001, 2002). We have recently shown that human echovirus 1 (EV1) is another virus using caveolae in its entry (Marjomäki *et al.*, 2002). Subsequent to internalization via caveolae, EV1 was shown to be localized in perinuclear structures, which contained caveolin-1 (Marjomäki *et al.*, 2002). Such structures include caveosomes, recently identified endocytic organelles, which participate in the entry of SV40. However, the entry routes of SV40 and EV1 seem not to be identical (Marjomäki *et al.*, 2002) and EV1 was found to be a unique tool to study the cell biology of caveolae. Importantly, EV1 uses  $\alpha 2\beta 1$  integrin, a collagen receptor, to bind to cell surface (Bergelson *et al.*, 1992) and therefore, the entry of EV1 can be used to study the role of integrins in caveolae function. Previous studies have shown that integrins can be coprecipitated with caveolin-1 (Wary *et al.*, 1996, 1998), suggesting that caveolae form a platform for integrin-mediated signaling.

Here, we show that  $\alpha 2\beta 1$  integrin is located in raft like membrane domains rather than in caveolae. However, the formation of  $\alpha 2\beta 1$  integrin clusters triggers their lateral redistribution along cell surface to caveolae and consequently

Article published online ahead of print. Mol. Biol. Cell 10.1091/mbc.E03-08-0588. Article and publication date are available at [www.molbiolcell.org/cgi/doi/10.1091/mbc.E03-08-0588](http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E03-08-0588).

<sup>¶</sup> Corresponding author. E-mail address: [jyrki.heino@utu.fi](mailto:jyrki.heino@utu.fi).

Abbreviations used: CA, constitutively active; DN, dominant negative; DRM, detergent resistant membranes; EEA1, early endosome associated protein; ERK, extracellular signal regulated kinase; EV1, echovirus-1; GPI-APs, glycosyl phosphatidyl inositol anchored proteins; MAPK, mitogen activated protein kinase; MOI, multiplicity of infection; PKC, protein kinase C; SV40, simian virus 40.

activates the internalization of caveolae in a protein kinase C $\alpha$  (PKC $\alpha$ )-dependent manner. Integrin  $\alpha 2\beta 1$  represents a novel mechanism to activate caveolae-mediated endocytosis. The molecular mechanisms of endocytosis and recycling of  $\beta 1$ ,  $\beta 2$ , and  $\alpha V$  integrins have been studied in detail (Bretscher 1992; Fabbri *et al.*, 1999; Ng *et al.*, 1999a; Laukaitis *et al.*, 2001), but integrin trafficking has been described to take place in endosomes and in an endocytic recycling pathway. Thus  $\alpha 2\beta 1$  seems to have unique activities compared with other integrins. In agreement with this our results show that the clusters of  $\alpha V$  integrin are internalized by a different mechanism, namely in clathrin-coated entry vesicles and early endosomes.

## MATERIALS AND METHODS

### Cells, Viruses, and Antibodies

SAOS cells (ATCC, Manassas, VA) were transfected with an expression construct encoding  $\alpha 2$  integrin (SAOS- $\alpha 2\beta 1$  cells; Ivaska *et al.*, 1999). Cells transfected with an empty expression vector (SAOS-pAW) were used as a control. EV1 (Farouk strain, ATCC) was propagated in GMK cells and purified in sucrose gradients as described previously (Marjomäki *et al.*, 2002). The following antibodies were used: rabbit antisera against purified EV1 (Marjomäki *et al.*, 2002), EEA1 (Mu *et al.*, 1995), transferrin (Zymed, South San Francisco, CA), ERK (Santa Cruz Biotechnology, Santa Cruz, CA), caveolin-1 (Transduction Laboratories, Lexington, KY; N20 antibody, Santa Cruz), and phosphorylated protein kinase C $\alpha$  (Upstate Biotechnology, Lake Placid, NY) as well as monoclonal antibodies (mAb) against total protein kinase C $\alpha$  (Upstate), caveolin-1 (Transduction Laboratories and Zymed), tubulin (Sigma, St. Louis, MO), myc (9E10, ATCC), integrin  $\alpha 2$  subunit (MCA2025, Serotec, Raleigh, NC), integrin  $\alpha V$  subunit (L230, ATCC), and phosphorylated ERK (Transduction Laboratories). Conjugation of anti- $\alpha 2$  and anti- $\alpha V$  antibodies to Alexa 488 was done using a mAb labeling kit (Molecular Probes, Inc., Eugene, OR).

### Transfections

SAOS- $\alpha 2\beta 1$  cells were transfected with FuGENE reagent (Boehringer Mannheim, Indianapolis, IN) and the cells were used for experiments after an expression time of 25–40 h. The GFP constructs of actin (CLONTECH, Palo Alto, CA), caveolin-1 (caveolin-GFP; from Dr. Ari Helenius, Institute of Biochemistry, ETH-Hoenggerberg, Switzerland; Pelkmans *et al.*, 2001) wild-type Eps15 (DIIId2; from Dr. Alice Dautry-Varsat, Pasteur Institute, Paris; Benmerah *et al.*, 1998), dominant negative (DN) Eps15 (d95/925; from Dr. Alice Dautry-Varsat, Pasteur Institute, Paris; Benmerah *et al.*, 1999), wild-type PKC $\alpha$  (Dr. Peter J. Parker, Cancer Research, UK; Ng *et al.*, 1999b), DN PKC $\alpha$  (T497A kinase-dead, substrate-binding mutant; Dr. Peter J. Parker, Cancer Research, UK; Mostafavi-Pour *et al.*, 2003), and DN PKC $\epsilon$  (kinase dead form; Ivaska *et al.*, 2002b) were used. DN MEK (1E8; from Dr. Natalie Ahn, University of Colorado; Holmström *et al.*, 1999), constitutively active MEK (1R4F; from Dr. Natalie Ahn, University of Colorado; Holmström *et al.*, 1999), and DN Ras (asn-17-ras; from Dr. Larry Feig, Tufts University; Feig and Cooper, 1988) were cotransfected to the cells with pEGFP-C2 (CLONTECH). A myc tagged AP180C (from Dr. Dieter Blaas, University of Vienna; Ford *et al.*, 2001) was revealed from transfected cells by myc labeling.

### Flotation Gradient Centrifugation

SAOS- $\alpha 2\beta 1$  cells were lysed with 1% Triton X-100 in PBS supplemented with protease inhibitors for 30 min on ice. The homogenate was adjusted to a sucrose density of 40.6%, overlaid with 35% sucrose and then filled with 5% sucrose. The gradient was centrifuged for 35,000 rpm at 4°C in a Beckman Sw 41Ti rotor for 18 h.

### Integrin Clusters

Antibody against  $\alpha 2$  integrin was added in DMEM (supplemented with 0.1% serum, low-DMEM), incubated for 1 h on ice, and washed. Cells were subsequently incubated with Alexa-conjugated goat anti-mouse IgG antibody on ice and washed. Formation of integrin clusters was allowed to occur at 37°C for 30–90 min. Clusters of  $\alpha V$  integrins were allowed to form using an anti- $\alpha V$  subunit mAb and Alexa-conjugated anti-mouse IgG antibody. As a control, anti- $\alpha 2$  and anti- $\alpha V$  mAbs, directly labeled with Alexa 488, were used alone without the clustering secondary antibodies.

### Measurement of Internalization

The internalization of EV1 or integrin clusters was estimated by confocal microscopy using the three-dimensional (3D) LSM program, version 1.4.2 (Carl Zeiss, Jena, Germany). Cells were incubated with EV1 for 1 h and fixed with 3% paraformaldehyde for 20 min. After fixation, EV1 was labeled

without permeabilization using a primary antibody and then the anti-rabbit Alexa 546 conjugate. After a 5-min permeabilization with 0.2% Triton X-100 EV1 was again labeled but now using anti-rabbit Alexa 488 conjugate. Thus, plasma membrane associated EV1 was stained with both Alexa 546 and 488 conjugates and was seen as yellow when the red and green channels were merged. Green signal representing internalized EV1 was measured. First the double-labeled cells were imaged as z-stacks with the confocal microscope. Then, in the 3D for LSM program the volume of internalized vesicles and the total amount of fluorescence was measured. Altogether 20 cells were measured in the experiment. In parallel experiments the clusters of  $\alpha 2$  or  $\alpha V$  integrins were allowed to form in the presence of Alexa-conjugated secondary antibody and to be internalized. Samples were fixed as above. The internalized integrins had green color, and the plasma membrane associated integrins were labeled with Alexa 546 conjugate. Again yellow color in merge images represented integrin clusters on the plasma membrane and green signal represented internalized clusters.

### Protein Kinase C Activation

SAOS- $\alpha 2\beta 1$  cells were starved overnight in low-DMEM. Cells were then treated in different ways: 1) stimulated with 1  $\mu$ M PMA 30 min at 37°C; 2) incubated with EV1 for 1 h on ice, washed, and incubated for 30 min at 37°C; 3) pretreated with 10  $\mu$ M safinolol at 37°C for 30 min, followed by EV1 treatment together with 10  $\mu$ M safinolol as described above; 4) incubated with anti- $\alpha 2$  mAb Fab fragment (2  $\mu$ g) for 1 h on ice, washed, and incubated for 30 min at 37°C; 5)  $\alpha 2\beta 1$  integrin clustering as explained above and incubated for 30 min at 37°C; 6) control SAOS- $\alpha 2\beta 1$  cells were treated with low-DMEM for 1 h on ice and then incubated for 30 min at 37°C. The treated cells were lysed with 30 mM octylglucopyranoside (Sigma) with protease inhibitors for 30 min on ice.

### SDS-PAGE and Immunoblotting

Samples were separated in 12% SDS-polyacrylamide gels and electroblotted onto PVDF membrane (Millipore, Bedford, MA). Aerolysin overlay assay was performed as described (Fivaz *et al.*, 2002). Primary antibodies and horseradish peroxidase-conjugated secondary antibodies (Bio-Rad, Richmond, CA) were used. Bands were detected by chemiluminescence (Pierce, Rockford, IL).

### Immunofluorescence and Confocal Microscopy

Subconfluent SAOS- $\alpha 2\beta 1$  cultures were incubated with EV1 for various time periods and then fixed with methanol at  $-20^{\circ}\text{C}$  for 6 min or with 4% paraformaldehyde for 20 min at room temperature. Cross-adsorbed goat secondary antibodies against rabbit (Alexa 488, 546, or 633 nm; Molecular Probes, Inc.) and mouse (Alexa 488, 546, or 633 nm; Molecular Probes, Inc.) antibodies were used for labeling. The cells were examined with an Axiovert 100 M SP epifluorescence microscope (Carl Zeiss) equipped with a confocal setup (Zeiss LSM510). Images were acquired using a Plan Neofluar objective (63 $\times$ , 1.25 oil) and a digital resolution of 512  $\times$  512. False colocalization signals were avoided by scanning fluorescence from different excitation wavelengths separately.

### Live Cell Microscopy

SAOS- $\alpha 2\beta 1$  cells were stained with Alexa 546-conjugated mutant aerolysin (ASSP; Fivaz *et al.*, 2002) together with an anti- $\alpha 2$  mAb on ice for 1 h. Alternatively, cells transfected with cDNA encoding caveolin-1-GFP (green fluorescent protein) or actin-GFP were treated with anti- $\alpha 2$  mAb. After washes the cells were transferred to a preheated sample stage (27/37°C) at a confocal microscope (Carl Zeiss Axiovert 100M with LSM510). Anti-mouse Alexa 488 (aerolysin-containing cells) or anti-mouse Alexa 546 (GFP-containing cells) was added and z-stacks through the cells were imaged at 5- or 10-min intervals. CO $_2$  independent medium (Sigma) was used in all steps. 3D projections of the selected images were created for each time point using LSM 510 3.0 and AxioVision Inside4D 3.0 (Carl Zeiss). These image series were processed and edited using Corel Photo-Paint 8, Ulead Media Studio ProVideo Editor 5.02a VE, and Quick Time Pro.0.2.

### Electron Microscopy

SAOS- $\alpha 2\beta 1$  cells were fixed in 2.5% glutaraldehyde in 0.1 M phosphate buffer, pH 7.4, for 1 h, and then postfixed in 1% osmium tetroxide, dehydrated, stained with uranyl acetate, and embedded in LX-112. For visualizing clusters of  $\alpha 2$  or  $\alpha V$  integrin, cells were treated with anti- $\alpha 2$  or  $\alpha V$  mAb for 1 h on ice, then with rabbit antibodies against mouse IgG (Sigma), and finally with protein A gold (10-nm particles, G. Posthuma and J. Slot, Utrecht, The Netherlands), both for 1 h on ice. Cells were then either fixed immediately or incubated for 1 min to 2 h at 37°C in complete culture medium. Further preparation for electron microscopy was performed as described above.

### Statistical Analysis

For analyzing the transfected samples in Figures 5C and 6B the binomial *t* test (comparison of two experimental percentual figures) was used. For analyzing the differences in binding of EV1 to cells (Figure 7B) multivariate analysis of

variance (MANOVA) was used with the least significant difference (LSD) as a post hoc test.

## RESULTS


### *On Cell Surface $\alpha 2\beta 1$ Integrin Is in Membrane Rafts*

Flotation gradient centrifugation indicated that  $\alpha 2\beta 1$  integrin in SAOS- $\alpha 2\beta 1$  cells (human osteosarcoma cells transfected with  $\alpha 2$  integrin cDNA) is found in fractions containing detergent resistant membranes (DRM; Figure 1A). The same fractions contained glycosyl phosphatidyl inositol anchored proteins (GPI-APs), well-known components of raft type cholesterol-sphingolipid-rich microdomains and caveolin-1, a hallmark protein for caveolae and caveosomes (Figure 1A). GPI-APs were recognized using Alexa 546-conjugated mutant aerolysin (ASSP; Fivaz *et al.*, 2002). Because flotation gradient centrifugation cannot be used to separate different DRM domains, confocal microscopy was used. When cells were plated on plastic in the presence of serum, patches of ASSP were found at the cell peripheries, suggesting the presence of raft domains. In addition, some intracellular staining was seen (Figure 1B). In SAOS- $\alpha 2\beta 1$  cells  $\alpha 2\beta 1$  integrin colocalized with ASSP positive membrane domains (Figure 1B), indicating that ligand free  $\alpha 2\beta 1$  is present in raft domains. Caveolin-1 was mainly seen inside the cell, most clearly around the nucleus. Some caveolin-1 was also seen on the cell surface in accordance with the fact that the SAOS cell surface is rich in caveolae (Marjomäki *et al.*, 2002). No clear colocalization of caveolin-1 and  $\alpha 2\beta 1$  integrin was detected (Figure 1B).


### *Clustering of $\alpha 2\beta 1$ Integrin Leads to Lateral Redistribution following Actin Filaments and Internalization in Caveolae-like Structures*

Cluster formation was first induced using primary anti- $\alpha 2$  integrin antibodies together with secondary antibodies. The fate of the clustered integrins was followed by live cell confocal microscopy, and the data were processed to obtain projections of the 3D images of the cells at different time points. Concomitant staining of cells with Alexa 546-conjugated ASSP was used to mark the GPI-APs containing domains. During the follow-up period integrins formed clusters (Figure 2A) that started to move along the cell surface. Concomitantly, small clusters fused together (Figure 2B, Video 1). Moving integrin clusters followed cortical actin microfilaments, i.e., filaments close to the cell surface (Figure 2, C-E, Video 2). Destruction of microfilaments with cytochalasin D (5  $\mu\text{g}/\text{ml}$ ) inhibited the movement of  $\alpha 2\beta 1$  integrins out of lipid rafts (our unpublished results). The presence of  $\alpha 2\beta 1$  integrins on the cell surface was also confirmed by measuring the relative distribution of internalized vs. cell surface integrins using differential staining before and after permeabilization (our unpublished results).

Integrin clusters were subsequently internalized (Figure 3A, Movie 3). Importantly, under the same conditions no integrin clustering or internalization was detected if cells were treated with fluorophore-labeled primary antibody without the secondary antibody (Figure 3B). Furthermore, the conditions used in live cell microscopy had no effects on cell viability or morphology (our unpublished results). Integrin clusters were internalized in caveolin-1-positive vesicles (Figure 3C, Video 4). Integrin  $\alpha 2\beta 1$  was immunostained before and after permeabilization of the cells to distinguish between internalized and cell surface integrin. Two hours after antibody-mediated clustering  $71 \pm 3\%$  of  $\alpha 2\beta 1$  integrin was intracellular. Destruction of microfilaments with cytochalasin D (5  $\mu\text{g}/\text{ml}$ ), which inhibited the movement of



**Figure 1.** Integrin  $\alpha 2\beta 1$  is located in lipid rafts. (A) In flotation gradient centrifugation of 1% Triton X-100 lysed cell homogenate  $\alpha 2\beta 1$  integrin, caveolin-1, and GPI-APs (ASSP) localize at the 5–35% sucrose interphase. The localization of detergent-resistant membranes (DRM) have been indicated. Western blot of gradient fractions 1–9 (from top to bottom). (B) Lipid rafts containing GPI-APs (ASSP) are enriched in  $\alpha 2\beta 1$  integrin in SAOS- $\alpha 2\beta 1$  cells. SAOS-pAW cells are  $\alpha 2\beta 1$  negative control cells. Indirect immunofluorescence labeling was used to reveal  $\alpha 2\beta 1$  integrin and caveolin-1, whereas GPI-APs were labeled directly with an ASSP-Alexa 546 conjugate. Confocal images are 3D projections of scanned cells. Scale bar, 10  $\mu\text{m}$ . In merge images yellow color indicates colocalization of GPI-APs and  $\alpha 2\beta 1$  integrin.



**Figure 2.** Formation of  $\alpha 2\beta 1$  clusters initiates redistribution of  $\alpha 2\beta 1$  out of lipid rafts. (A) In SAOS- $\alpha 2\beta 1$  cells  $\alpha 2\beta 1$  integrin (green) is colocalized with GPI-APs (ASSP, red; control), but after incubation with anti- $\alpha 2$  mAb and secondary antibody (+a- $\alpha 2$ )  $\alpha 2\beta 1$  integrin is located outside the lipid rafts. Scale bar, 10  $\mu\text{m}$ . Blow-ups show lipid raft areas on cell edges (merge images). (B) A close-up about the formation of  $\alpha 2\beta 1$  integrin clusters (red) and integrin redistribution (see Video 1). Scale bar, 0.5  $\mu\text{m}$ . (C) A whole-cell view shows that integrin clusters (red) move along actin filaments (GFP-actin; see Video 2). Scale bar, 10  $\mu\text{m}$ . (D) A close-up of integrin clusters fusing together and moving along actin. Scale bar, 2  $\mu\text{m}$ . B, C, and D are all live images of the same cell, B and D being digitally isolated close ups of C (actin is not shown in B). The live imaging (B–D) was performed at low temperature (27°C) in order to increase temporal resolution. (E) A rectangular slice of the cell from the 70-min time point was taken to verify that the actin filaments were indeed cortical, i.e., close to the plasma membrane. The white rectangle shows the location of this slice.


$\alpha 2\beta 1$  integrins out of lipid rafts, did not completely prevent the  $\alpha 2\beta 1$ -mediated EV1 entry and infection (our unpublished results). GPI-APs were also internalized to some extent. However, this is not evident in the images because of the “shadow” rendering method used. In accordance with our previous observations (Marjomäki *et al.*, 2002), EV1 was rapidly internalized into similar vesicles that contained caveolin-1 and  $\alpha 2\beta 1$  integrin, and the vesicles had the tendency to accumulate close to the nucleus (Figure 3, C and D). The internalization of clustered integrins was further studied by EM using secondary antibodies and protein A gold. The cell surface invaginations containing clustered integrins appeared to be similar to caveolae and, accordingly, the integrin containing intracellular vesicles resembled caveosomes (Pelkmans *et al.*, 2001; Figure 3E). EV1 labeled with gold particles was shown to accumulate into identical vesicles (Figure 3F). Our data indicate that the formation of  $\alpha 2\beta 1$  clusters alone may trigger accumulation of the receptors into caveosome like organelles and that EV1 may behave as a

multivalent ligand that activates the same internalization process as the antibodies.

Importantly, in EM we could not see EV1 or  $\alpha 2\beta 1$  clusters inside coated vesicles, suggesting that clathrin-dependent processes are not involved in their entry. This was further studied by transient transfection with cDNA coding for dominant negative (DN) Eps15, known to inhibit the internalization of transferrin (Benmerah *et al.*, 1999). As shown in Table 1, DN Eps15 had no effect on  $\alpha 2\beta 1$  internalization, although it could inhibit the entry of transferrin. In SAOS- $\alpha 2\beta 1$  cells, transfected AP180C (Ford *et al.*, 2001) proved to be an even more effective inhibitor of transferrin internalization, but without an effect on  $\alpha 2\beta 1$  entry (Table 1).

#### *The Clusters of $\alpha V$ Integrins Are Internalized in Coated Vesicles and Endosomes*

We repeated the same experiment with antibodies against  $\alpha V$  integrin subunit and secondary antibodies. In 2 h  $93 \pm 1\%$  of  $\alpha V$  clusters had been internalized. If primary anti- $\alpha V$



**Figure 3.** Clusters of  $\alpha 2\beta 1$  integrin are internalized in caveolae. (A) Live cell imaging of cellular internalization of clustered  $\alpha 2\beta 1$  integrin. Integrin  $\alpha 2\beta 1$  is green and GPI-APs are red (ASSP). A rectangular slice was taken from the center area of one cell, and projections were viewed from above and the side (see Video 3). (B) Live cell imaging with fluorophore (green)-labeled anti- $\alpha 2$  integrin mAb without secondary antibody. A rectangular slice was taken from the center area of one cell, and the projection was viewed from the side. (C) In SAOS- $\alpha 2\beta 1$  cells incubated with anti- $\alpha 2$  mAb and secondary antibody or EV1,  $\alpha 2\beta 1$  integrin (green) is colocalized with caveolin-1 (red; see Video 4). Scale bar, 10  $\mu$ M. (D) Caveolin-1 colocalizes with EV1 2 h p.i. Scale bar, 10  $\mu$ M. (E) Electron microscopy of gold particles linked to secondary antibodies against anti- $\alpha 2$  mAb indicates that clustered  $\alpha 2\beta 1$  integrin is internalized in caveolae (left) and later is found in caveosomes (right). Protein A gold particles (5 nm) label the secondary antibodies that were used to cluster the primary anti- $\alpha 2$  antibodies. Scale bar, 100 nm. (F) A caveosome containing internalized EV1 that is labeled with gold particles (5 nm) is shown for comparison. Scale bar, 100 nm. For EV1 infections MOI 100 was used.

antibodies were used alone, no clustering or internalization of  $\alpha V$  integrins was detected (our unpublished results). The clusters of  $\alpha V$  integrins were internalized in caveolin-1-negative vesicles (Figure 4A). However, some vesicles containing  $\alpha V$  clusters were positive for early endosome associated protein (EEA1; Figure 4B) and for endocytosed transferrin (Figure 4C). Importantly, internalized  $\alpha 2\beta 1$  clusters were in EEA1 and transferrin negative vesicles (Figure 4, B and C). Finally, electron microscope images revealed that  $\alpha V$  clusters labeled with gold particles were found in structures that were morphologically identical to clathrin-coated vesicles (Figure 4D). Picornaviruses that bind to  $\alpha V$  integrins, such as parechovirus-1, are known to enter their host cells via the clathrin-dependant pathway (Joki-Korpela *et al.*, 2001).


#### **Protein Kinase C Activity Is Essential for the Internalization of $\alpha 2\beta 1$ Integrin via Caveolae**

The fact that integrins can mediate cellular signals (Schwartz and Ginsberg, 2002) may influence their internalization. We

**Table 1.** Inhibition of transferrin internalization has no effect on the internalization of  $\alpha 2\beta 1$  integrin clusters

cDNA	No. of cells containing internalized transferrin (% of all transfected cells)	No. of cells containing internalized $\alpha 2\beta 1$ integrin (% of all transfected cells)
Control	78 (39/50)	82 (41/50)
EPS15 WT	84 (42/50)	92 (46/50)
EPS15 DN	40 (20/50)	92 (46/50)
AP180C	24 (12/50)	90 (45/50)


The number of cells containing internalized transferrin or  $\alpha 2\beta 1$  integrin was counted 2 h after  $\alpha 2\beta 1$  integrin clustering was induced by antibodies. Transferrin was added 15 min before fixation. Only the cells that had successfully been transfected were analyzed. The figures in parentheses show the number of positive cells/number of cells analyzed.



**Figure 4.** Clusters of  $\alpha V$  and  $\alpha 2\beta 1$  integrins are internalized through distinct pathways. (A) Clusters of  $\alpha V$  integrins (green) are not targeted to caveolin-1-containing vesicles (red). Scale bar, 10  $\mu M$ . (B) Clusters of  $\alpha V$  integrins, unlike clusters of  $\alpha 2\beta 1$  integrins, are found in vesicles positive for EEA1, a molecular marker for early endosomes. Scale bar, 10  $\mu M$ . (C) Clusters of  $\alpha V$  integrins, unlike clusters of  $\alpha 2\beta 1$  integrins, are internalized *via* the same pathway as transferrin. Scale bar, 10  $\mu M$ . In A–C, white rectangles show the locations of the blow-ups. (D)  $\alpha V$  integrin clusters labeled with gold particles (10 nm) 1 min after internalization can be seen inside coated vesicles by electron microscopy. Clathrin coat is indicated by arrows. Scale bar, 100 nm.

have previously shown that mitogen activated protein kinases (MAPKs) are regulated by the collagen receptor integrins (Ravanti *et al.*, 1999). Here, extracellular signal regulated kinase (ERK) was activated at very early time points after  $\alpha 2\beta 1$  binding to EV1 (Figure 5A). Because a selective inhibitor of ERK activation (20  $\mu M$  PD98059, Calbiochem, La Jolla, CA; Figure 5B) or transient transfections with cDNAs coding for dominant negative or constitutively active MEK (DN MEK and CA MEK, respectively) had no effect on

EV1 infection (Figure 5C), and DN MEK inhibited the internalization of  $\alpha 2\beta 1$  integrin only 10% (Figure 5C), we tried to specify other components of the same pathway, upstream of ERK and MEK, using selective inhibitors of Ras and protein kinase C (PKC). A Ras inhibitor (500  $\mu M$  perillyl alcohol, POH, Aldrich, Milwaukee, WI) had little effect on ERK activation and its consequent transport into nucleus (Figure 5D). In addition, transfections with DN Ras had no effect on  $\alpha 2\beta 1$  internalization or EV1 infection (Figure 5C). In con-




**Figure 5.** Binding of  $\alpha 2\beta 1$  integrin to EV1 induces the phosphorylation of ERK MAP kinase in a PKC $\alpha$  activity-dependent manner. (A) ERK is transiently phosphorylated <15 min after  $\alpha 2\beta 1$  integrin binding to EV1 and the phosphorylation lasts ~1 h (Western blot). (B) Inhibitors of ERK activation (PD) and p38 kinase (SB) do not inhibit infection in contrast to methyl  $\beta$ -cyclodextrin (CYCLO), an inhibitor of cholesterol metabolism and caveolae. (C) Transient transfections with cDNAs encoding the dominant negative MEK (DN MEK) or dominant negative Ras (DN Ras) do not inhibit  $\alpha 2\beta 1$  integrin internalization or infection. Constitutively active MEK (CA MEK) does not increase the internalization of  $\alpha 2\beta 1$  integrin either. (D) ERK is translocated to the nucleus 4 h p.i. The general PKC inhibitor bisindolylmaleimide (5  $\mu$ M, BIS) inhibits the nuclear translocation of ERK, unlike a Ras inhibitor, perillyl alcohol (500  $\mu$ M, POH). Quantification of ERK positive nuclei after treatments with bisindolylmaleimide (BIS), another PKC inhibitor safinol (10  $\mu$ M, SAF) or Ras inhibitor (POH). Error bars show SE values (B and C: the graph on the left, D) or 95% confidence limits (C: the graph on the right).

trast, a PKC inhibitor (5  $\mu$ M bisindolylmaleimide, Calbiochem) almost completely prevented accumulation of ERK in nucleus (Figure 5D). The experiment was repeated with another PKC inhibitor (10–25  $\mu$ M safinol, Calbiochem) with the same results (Figure 5D).


On the basis of these findings, we deduced that PKC activity is needed for the activation of signaling pathways by

$\alpha 2\beta 1$  integrin. Furthermore, as shown in Figure 6A, the presence of PKC inhibitors reduced the number of EV1-infected cells to background levels. Safinol has been considered to be a selective inhibitor of the  $\alpha$  isoform of PKC (Masur *et al.*, 2001). Here, we also performed transfections with DN PKC $\alpha$  cDNA to confirm the role of this isoform in the process (Figure 6B). DN PKC $\alpha$  inhibited EV1 entry and



**A**




**B**



**C**



**D**



**Figure 6.** PKC $\alpha$  activity is crucial for EV1 replication and the clustering of  $\alpha$ 2 $\beta$ 1 increases PKC $\alpha$  phosphorylation. (A) EV1 infection is prevented after inhibition of PKC activity with either bisindolylmaleimide (BIS) or safinol (SAF). (B) Dominant negative PKC $\alpha$  inhibits infection ( $p < 0.001$ , binomial  $t$  test), whereas overexpression of wild-type PKC $\alpha$  increases infection to some extent ( $p < 0.05$ , binomial  $t$  test). Dominant negative PKC $\epsilon$  does not have any effect on infection. (C) PKC $\alpha$  is downregulated after a chronic treatment (6 h) with phorbol ester (PMA), which is shown by Western blotting. This treatment also inhibits EV1 infection. (D) Phosphorylation of PKC $\alpha$  detected by Western blotting during EV1 entry (EV1), a short treatment (30 min) with PMA and clustering of  $\alpha$ 2 $\beta$ 1 integrin by anti- $\alpha$ 2 mAb (a- $\alpha$ 2 + Sec) and Fab of anti- $\alpha$ 2 mAb as its control. Negative control is marked as C. In C and D antibodies against phosphorylated PKC $\alpha$ , total PKC $\alpha$ , tubulin and caveolin-1 were used. Error bars show SE values (A and C) or 95% confidence limits (B).

infection, whereas overexpression of wild-type PKC $\alpha$  promoted it (Figure 6B). DN PKC $\epsilon$  was used as a control and it had no effect (Figure 6B). The critical role of PKC in EV1 entry and infection was also confirmed in a third set of experiments (Figure 6C), in which chronic treatment of cells with a phorbol ester (PMA, 6 h) was used to downregulate

PKC expression. Chronic PMA treatment is known to induce the degradation of both classical and novel PKCs (Srivastava *et al.*, 2002), and it is not specific for PKC $\alpha$ . The effectiveness of the treatment was confirmed by Western blotting (Figure 6C). Concomitantly with a decrease in PKC $\alpha$  level, EV1 infection was prevented (Figure 6C). The amount of active



(phosphorylated) PKC $\alpha$  in SAOS- $\alpha$ 2 $\beta$ 1 cells was analyzed by Western blotting with phosphospecific antibodies. Control SAOS- $\alpha$ 2 $\beta$ 1 cells contained phosphorylated PKC $\alpha$  and the fact that short PMA treatment (30 min) could only slightly increase the phosphorylation (in five experiments on average 1.5-fold) suggested that most of it might be in the active form (Figure 6D). Some increase (maximally 1.5 fold) in the phosphorylation of PKC $\alpha$  was seen after the cells were exposed to EV1 (Figure 6D). A similar small increase in phosphorylation was seen in the presence of primary anti- $\alpha$ 2 and secondary antibodies when compared with the effect of Fab fragments of the same anti- $\alpha$ 2 antibody (Figure 6D). We suggest that the increase in PKC $\alpha$  phosphorylation may be a part of the signaling events after integrin clustering. In any case, its activity seems to be critical for integrin signaling.

We next tested the mechanism used by PKC $\alpha$  inhibitors to affect the  $\alpha$ 2 $\beta$ 1 integrin-mediated entry of EV1. If safingol was added to cells 5 min after EV1 almost complete inhibition was still obtained, whereas if added 30 min after EV1 it could no longer prevent infection (Figure 7A). This result limited the action of PKC $\alpha$  to the first minutes of the process. Experiments measuring the number of radioactive EV1 bound to the cell surface indicated that safingol does not inhibit ligand binding to  $\alpha$ 2 $\beta$ 1 integrin (Figure 7B). Actually, in repeated experiments safingol seemed to slightly increase the number of cell bound EV1 viruses (Figure 7B), making it possible to speculate that safingol may actually affect receptor recycling. However, the increase was not statistically significant (multivariate analysis of variance), and therefore it was not studied further. In the presence of safingol EV1 binding to  $\alpha$ 2 $\beta$ 1 integrin could still initiate the lateral redistribution of the integrin out of lipid rafts (our unpublished results). Furthermore, in the presence of safingol clustering antibodies could still initiate the process leading to the colocalization of  $\alpha$ 2 $\beta$ 1 integrin and caveolin-1 (Figure 7C). Thus, safingol had no effect on the lateral movement of  $\alpha$ 2 $\beta$ 1 integrin after cluster formation. Finally, we used confocal microscopy to quantify the number of EV1 particles inside cells and safingol could remarkably reduce the entry of EV1 (Figure 7D). Furthermore, safingol had no effect on the entry of clustered  $\alpha$ V integrins (Figure 7E). Thus, the results indicate that the crucial role of PKC $\alpha$  in the process is limited to the activation of caveola entry by  $\alpha$ 2 $\beta$ 1 integrins that have moved from lipid rafts to caveolae.

## DISCUSSION

Integrins anchor cells to the extracellular matrix. In addition, they signal and participate in the regulation of cellular functions, such as migration, differentiation, survival, and growth. Like most cell surface receptors also integrins can be internalized after ligand binding. Fibrinogen endocytosis is mediated by  $\alpha$ M $\beta$ 2 integrin in monocytoic cells (Simon *et al.*, 1993) and by  $\alpha$ IIb $\beta$ 3 integrin in platelets (Wencel-Drake *et al.*, 1996). Integrins  $\alpha$ V $\beta$ 3 and  $\alpha$ V $\beta$ 5 can mediate the internalization of vitronectin (Pijuan-Thompson and Gladson, 1997). Some integrins can also recycle (Bretscher 1992; Fabbri *et al.*, 1999). This process seems to be important for migrating epithelial cells (Ng *et al.*, 1999a; Laukaitis *et al.*, 2001). During cell migration  $\beta$ 1 integrin trafficking has been described to take place in endosomes and the endocytic recycling pathway (Ng *et al.*, 1999a; Laukaitis *et al.*, 2001) in a PKC $\alpha$ -dependent manner (Ng *et al.*, 1999a). We have recently reported that  $\alpha$ 2 $\beta$ 1 integrin is internalized via caveolae in the process of EV1 entry (Marjomäki *et al.*, 2002). This represents a novel internalization mechanism for integrins. Integrin  $\alpha$ 2 $\beta$ 1 is a collagen receptor. It is expressed on platelets, epithelial cells, and also on many mesenchymal cell types, including fibroblasts, chondrocytes, osteoblasts, and endo-


thelial cells. Integrin  $\alpha$ 2 $\beta$ 1 mediates internalization of collagen and therefore it may participate in the regulation of matrix turnover (Segal *et al.*, 2001). In endothelial cells  $\alpha$ 2 $\beta$ 1-mediated collagen endocytosis may play a special role in the formation of capillary tube and lumen (Davis and Camarillo, 1996). It is not known whether integrins that are internalized in caveolae can recycle.

Many members of the integrin family are used by different viruses for attachment to the cell surface and for subsequent internalization into the host cell (Wang, 2002; Bonsel and Alfsen, 2003). It has been suggested that viruses have evolved to recognize cell adhesion receptors because of their relatively low affinity to natural ligands (Wang, 2002). The integrins may, however, play an active role in the entry process. For example, Kaposi's sarcoma associated herpes virus (human herpes virus 8) has been shown to use  $\alpha$ 3 $\beta$ 1 integrin as its receptor and to activate  $\alpha$ 3 $\beta$ 1-dependent signaling pathways, including focal adhesion kinase (Akula *et al.*, 2002). Thus, virus binding may activate integrin associated cellular signaling, which could promote or modulate virus entry. Integrin  $\alpha$ 2 $\beta$ 1 differs from other integrin type virus receptors by being the only integrin known to mediate virus entry through caveolae instead of clathrin-coated vesicles (Marjomäki *et al.*, 2002). Caveolae are also involved in the entry of SV40 (Pelkmans *et al.*, 2001; Pelkmans *et al.*, 2002), mouse polyoma virus (Richterova *et al.*, 2001), filoviruses (Empig and Goldsmith, 2002), and some bacteria (Shin *et al.*, 2000).


Despite the fact that  $\alpha$ 2 $\beta$ 1 integrin is internalized in caveolae our results indicate that after synthesis it is targeted to lipid rafts together with GPI-APs. The binding of both natural ligands and EV1 to  $\alpha$ 2 $\beta$ 1 integrin may cause the formation of a receptor cluster. Here, using antibodies to induce the cluster formation we were able to show that  $\alpha$ 2 $\beta$ 1 clustering alone can initiate the lateral redistribution of  $\alpha$ 2 $\beta$ 1 out of raft domain. The integrins seem to follow actin microfilaments and during the process smaller clusters fuse together forming larger ones. Lateral redistribution of  $\alpha$ 5 $\beta$ 1 integrin on the cell surface has been detected during electric field directed fibroblast locomotion (Brown and Loew, 1994). When integrins moved toward the cathode they also seemed to form large clusters (Brown and Loew, 1994). However, typically  $\alpha$ 5 $\beta$ 1 is redistributed in endocytic vesicles in which the integrin colocalizes with transferrin receptor (Laukaitis *et al.*, 2001). Thus, the lateral movement of  $\alpha$ 2 $\beta$ 1 on cell surface from rafts to caveolae indicates that  $\alpha$ 2 $\beta$ 1 acts in a different way than fibronectin receptors or other integrins studied so far. In migrating cells it is not known, whether  $\alpha$ 2 $\beta$ 1 is among the  $\beta$ 1 integrins that are guided to the endocytic recycling pathway (Ng *et al.*, 1999a). However, integrin clustering after the binding of multivalent ligands seems to initiate a very different process.

When  $\alpha$ 2 $\beta$ 1 integrin clusters were found in caveolae like invaginations on the cell surface their colocalization with caveolin-1 was observed. Previous studies have shown that integrins can be coprecipitated with caveolin-1 (Wary *et al.*, 1996, 1998; Wei *et al.*, 1999). Our results indicate, that in the case of  $\alpha$ 2 $\beta$ 1 this association is obvious only after integrin clustering and redistribution. Caveolae are immobile and their internalization requires an activating stimulus (Thomsen *et al.*, 2002). For example, the interaction of albumin-docking protein gp60 and caveolin-1 in endothelial cells (Minshall *et al.*, 2000) and the phosphatase inhibitor okadaic acid (Parton *et al.*, 1994) can trigger the internalization of caveolae. Clustering of  $\alpha$ 2 $\beta$ 1 integrin represents a novel mechanism to activate caveolae.

The formation of integrin clusters is known to be important for signaling events (Miyamoto *et al.*, 1995). Furthermore, caveolin-1 may participate in integrin-mediated signaling (Wary *et al.*, 1996, 1998). The interaction of  $\alpha$ 2 $\beta$ 1 with collagen can activate several signaling pathways, including MAPKs



**Figure 7.** Integrin  $\alpha 2\beta 1$ -mediated internalization of EV1 *via* caveolae is dependent on PKC $\alpha$  activity. (A) Safingol can be added to cells 30 min before EV1 ( $-30'$ ), at the same time as EV1 ( $0'$ ), or 5 min after EV1 ( $+5'$ ) to prevent the infection. (B) Safingol does not affect the ability of  $\alpha 2\beta 1$  integrin to bind radioactive EV1. (C) Live cell imaging showing that safingol does not prevent the clustering-related redistribution leading to colocalization of  $\alpha 2\beta 1$  integrin (red) and caveolin-1 (green) on the plasma membrane. To show that safingol inhibits integrin (red) internalization a rectangular slice was taken from the center area of one safingol treated and one nontreated cell (both in the presence of clustering antibodies) and projections were viewed from the side. Caveolin-1 has been excluded from these images. Note that safingol is causing some rounding of cells. (D) Safingol prevents the internalization of EV1 (100 MOI). Internalization was quantified by labeling EV1 before and after permeabilization with different colors and then taking Z-scans through the cells with a confocal microscope and measuring the internalized vesicles using 3D for LSM software. (E) Safingol has no effect on the internalization of antibody induced  $\alpha V$  clusters. Error bars show SE values.



**Figure 8.** The proposed model of  $\alpha 2\beta 1$  integrin-mediated activation of caveolae. Antibody or virus-mediated formation of  $\alpha 2\beta 1$  integrin clusters leads to lateral movement on cell surface in which integrin clusters seem to follow microfilaments. Often smaller integrin clusters seem to fuse to each other. Clustering and lateral movement are followed by PKC $\alpha$  dependent internalization in caveolae.

(Ivaska *et al.*, 1999; Ravanti *et al.*, 1999), the  $\zeta$  isoform of PKC (PKC $\zeta$ ; Xu and Clark, 1997), and protein phosphatase 2A (PP2A; Ivaska *et al.*, 2002a). Here, PKC $\alpha$  activity was needed for the internalization of  $\alpha 2\beta 1$  integrin in caveolae. Inhibition of PKC $\alpha$  could not prevent the cluster formation or the lateral movement of  $\alpha 2\beta 1$  on cell surface. Its inhibitory effect was seen only during a limited time period, suggesting that PKC $\alpha$  activity is obligatory only for the entry of caveolae. Interestingly, PKC $\alpha$  may directly bind to the cytoplasmic domain of integrin  $\beta 1$  subunit (Ng *et al.*, 1999a). Furthermore, PKC $\alpha$  seems to be a resident protein of caveolae, which are major cell surface locations for this enzyme (Smart *et al.*, 1994; Mineo *et al.*, 1998). In caveolae PKC $\alpha$  might be constantly active (Smart *et al.*, 1994; Mineo *et al.*, 1998). Accordingly, control SAOS- $\alpha 2\beta 1$  cells con-

tained phosphorylated PKC $\alpha$ , and PMA could only slightly increase its phosphorylation. A similar small increase in the phosphorylation of PKC $\alpha$  was seen after the cells were exposed to EV1 or antibodies forming  $\alpha 2\beta 1$  clusters. Previous studies support the idea that PKC $\alpha$  is needed for caveola-related signaling functions (Smart *et al.*, 1994; Mineo *et al.*, 1998).

We have identified the formation of  $\alpha 2\beta 1$  integrin clusters as a key event leading to the relocation of  $\alpha 2\beta 1$  to caveolae. Furthermore,  $\alpha 2\beta 1$  clusters trigger the internalization of caveolae (the process has been summarized in Figure 8). Endocytosis via caveolae may be utilized by the natural ligands of  $\alpha 2\beta 1$  integrin. In addition, EV1 has evolved to take advantage of this phenomenon by binding to  $\alpha 2\beta 1$  on the cell surface. PKC $\alpha$  appears to be an important regulator of the entry process. The data indicate that the function of  $\alpha 2\beta 1$  is unique when compared with any other integrin.

## ACKNOWLEDGMENTS

We thank Arja Mansikkaviita and Raimo Pesonen for technical assistance and Dr. Daniel J. White for critical reading of the manuscript. We thank Dr. Dieter Blaas, Dr. Alice Dautry-Varsat, Dr. Natalie Ahn, Dr. Larry Feig, Dr. Ari Helenius, and Dr. Harald Stenmark for cDNAs and other reagents. The work was supported by grants from the Academy of Finland, the Sigrid Jusélius Foundation, and the Finnish Cancer Association. P.U. has a fellowship from the Finnish Cancer Institute.

## REFERENCES

- Akula, S.M., Pramod, N.P., Wang, F.Z., and Chandran, B. (2002). Integrin  $\alpha 3\beta 1$  (CD 49c/29) is a cellular receptor for Kaposi's sarcoma-associated herpesvirus (KSHV/HHV-8) entry into the target cells. *Cell* 108, 407–419.
- Benmerah, A., Lamaze, C., Begue, B., Schmid, S.L., Dautry-Varsat, A., and Cerf-Bensussan, N. (1998). AP-2/Eps15 interaction is required for receptor-mediated endocytosis. *J. Cell Biol.* 140, 1055–1062.
- Benmerah, A., Bayrou, M., Cerf-Bensussan, N., and Dautry-Varsat, A. (1999). Inhibition of clathrin-coated pit assembly by an Eps15 mutant. *J. Cell Sci.* 112, 1303–1311.
- Bomsel, M., and Alfsen, A. (2003). Entry of viruses through the epithelial barrier: pathogenic trickery. *Nat. Rev. Mol. Cell. Biol.* 4, 57–68.
- Bergelson, J.M., Shepley, M.P., Chan, B.M., Hemler, M.E., and Finberg, R.W. (1992). Identification of the integrin VLA-2 as a receptor for echovirus 1. *Science* 255, 1718–1720.
- Bretscher, M.S. (1992). Circulating integrins:  $\alpha 5\beta 1$ ,  $\alpha 6\beta 4$  and Mac-1, but not  $\alpha 3\beta 1$ ,  $\alpha 4\beta 1$  or LFA-1. *EMBO J.* 11, 405–410.
- Brown, M.J., and Loew, L.M. (1994). Electric field-directed fibroblast locomotion involves cell surface molecular reorganization and is calcium independent. *J. Cell Biol.* 127, 117–128.
- Couet, J., Sargiacomo, M., and Lisanti, M.P. (1997). Interaction of a receptor tyrosine kinase, EGF-R, with caveolins. Caveolin binding negatively regulates tyrosine and serine/threonine kinase activities. *J. Biol. Chem.* 272, 30429–30438.
- Davis, G.E., and Camarillo, C.W. (1996). An  $\alpha 2\beta 1$  integrin-dependent pinocytic mechanism involving intracellular vacuole formation and coalescence regulates capillary lumen and tube formation in three-dimensional collagen matrix. *Exp. Cell Res.* 224, 39–51.
- Drab, M. *et al.* (2001). Loss of caveolae, vascular dysfunction, and pulmonary defects in caveolin-1 gene-disrupted mice. *Science* 293, 2449–2452.
- Empig, C.J., and Goldsmith, M.A. (2002). Association of the caveola vesicular system with cellular entry by filoviruses. *J. Virol.* 76, 5266–5270.
- Fabbri, M., Fumagalli, L., Bossi, G., Bianchi, E., Bender, J.R., and Pardi, R. (1999). A tyrosine-based sorting signal in the  $\beta 2$  integrin cytoplasmic domain mediates its recycling to the plasma membrane and is required for ligand-supported migration. *EMBO J.* 18, 4915–4925.
- Feig, L.A., and Cooper, G.M. (1988). Inhibition of NIH 3T3 cell proliferation by a mutant ras protein with preferential affinity for GDP. *Mol. Cell. Biol.* 8, 3235–3243.
- Fivaz, M., Vilbois, F., Thumheer, S., Pasquali, C., Abrami, L., Bickel, P.E., Parton, R.G., and van der Goot, F.G. (2002). Differential sorting and fate of endocytosed GPI-anchored proteins. *EMBO J.* 21, 3989–4000.
- Ford, M.G., Pearse, B.M., Higgins, M.K., Vallis, Y., Owen, D.J., Gibson, A., Hopkins, C.R., Evans, P.R., and McMahon, H.T. (2001). Simultaneous binding of

- PtdIns(4, 5)P<sub>2</sub> and clathrin by AP180 in the nucleation of clathrin lattices on membranes. *Science* 291, 1051–1055.
- Fra, A.M., Williamson, E., Simons, K., and Parton, R.G. (1995). De novo formation of caveolae in lymphocytes by expression of VIP21-caveolin. *Proc. Natl. Acad. Sci. USA* 92, 8655–8659.
- Holmström, T.H., Tran, S.E., Johnson, V.L., Ahn, N.G., Chow, S.C., and Eriksson, J.E. (1999). Inhibition of mitogen-activated kinase signaling sensitizes HeLa cells to Fas receptor-mediated apoptosis. *Mol. Cell. Biol.* 19, 5991–6002.
- Ivaska, J., Reunanen, H., Westermarck, J., Koivisto, L., Kähäri, V.-M., and Heino, J. (1999). Integrin  $\alpha 2\beta 1$  mediates isoform-specific activation of p38 and upregulation of collagen gene transcription by a mechanism involving the  $\alpha 2$  cytoplasmic tail. *J. Cell Biol.* 147, 401–416.
- Ivaska, J., Nissinen, L., Immonen, N., Eriksson, J.E., Kähäri, V.-M., and Heino, J. (2002a). Integrin  $\alpha 2\beta 1$  promotes activation of protein phosphatase 2A and dephosphorylation of Akt and glycogen synthase kinase  $3\beta$ . *Mol. Cell. Biol.* 22, 1352–1359.
- Ivaska, J., Whelan, R.D., Watson, R., and Parker, P.J. (2002b). PKC epsilon controls the traffic of beta1 integrins in motile cells. *EMBO J.* 21, 3608–3619.
- Joki-Korpela, P., Marjomäki, V., Krogerus, C., Heino, J., and Hyypä, T. (2001). Entry of human parechovirus 1. *J. Virol.* 75, 1958–1967.
- Kurzchalia, T., and Parton, R. (1999). Membrane microdomains and caveolae. *Curr. Opin. Cell Biol.* 11, 424–431.
- Laukaitis, C.M., Webb, D.J., Donais, K., and Horwitz, A.F. (2001). Differential dynamics of  $\alpha 5$  integrin, paxillin, and  $\alpha$ -actinin during formation and disassembly of adhesions in migrating cells. *J. Cell Biol.* 153, 1427–1440.
- Marjomäki, V. *et al.* (2002). Internalization of echovirus 1 in caveolae. *J. Virol.* 76, 1856–1865.
- Masur, K., Lang, K., Niggemann, B., Zanker, K.S., and Entschladen, F. (2001). High PKC alpha and low E-cadherin expression contribute to high migratory activity of colon carcinoma cells. *Mol. Biol. Cell* 12, 1973–1982.
- Mineo, C., Ying, Y., Chapline, C., Jaken, S., and Anderson, R. (1998). Targeting of protein kinase C $\alpha$  to caveolae. *J. Cell Biol.* 141, 601–610.
- Minshall, R.D., Tirupathi, C., Vogel, S. M., Niles, W.D., Gilchrist, A., Hamm, H.E., and Malik, A.B. (2000). Endothelial cell-surface gp60 activates vesicle formation and trafficking via G(i)-coupled Src kinase signaling pathway. *J. Cell Biol.* 150, 1057–1070.
- Miyamoto, S., Akiyama, S., and Yamada, K. (1995). Synergistic roles for receptor occupancy and aggregation in integrin transmembrane function. *Science* 267, 883–885.
- Mostafavi-Pour, Z., Askari, J.A., Parkinson, S.J., Parker, P.J., Ng, T.T.C., and Humphries, M.J. (2003). Integrin-specific signaling pathways controlling focal adhesion formation and cell migration. *J. Cell Biol.* 161, 155–167.
- Mu, F.T. *et al.* (1995). EEA1, an early endosome-associated protein. EEA1 is a conserved alpha-helical peripheral membrane protein flanked by cysteine “fingers” and contains a calmodulin-binding IQ motif. *J. Biol. Chem.* 270, 13503–13511.
- Ng, T., Shima, D., Squire, A., Bastiaens, P.I., Gschmeissner, S., Humphries, M.J., and Parker, P.J. (1999a). PKC $\alpha$  upregulates  $\beta 1$  integrin-dependent cell motility through association and control of integrin traffic. *EMBO J.* 18, 3909–3923.
- Ng, T. *et al.* (1999b). Imaging protein kinase C $\alpha$  activation in cells. *Science* 283, 2085–2089.
- Parton, R.G., Joggerst, B., and Simons, K. (1994). Regulated internalization of caveolae. *J. Cell Biol.* 127, 1199–1215.
- Parton, R.G. (2003). Caveolae—from ultrastructure to molecular mechanisms. *Nat. Rev. Mol. Cell Biol.* 4, 162–167.
- Pijuan-Thompson, V., and Gladson, C.L. (1997). Ligand of integrin  $\alpha 5\beta 1$  is required for internalization of vitronectin by integrin  $\alpha v\beta 3$ . *J. Biol. Chem.* 272, 2736–2743.
- Pelkmans, L., Kartenbeck, J., and Helenius, A. (2001). Caveolar endocytosis of simian virus 40 reveals a new two-step vesicular-transport pathway to the ER. *Nat. Cell Biol.* 3, 473–483.
- Pelkmans, L., and Helenius, A. (2002). Endocytosis via caveolae. *Traffic* 3, 311–320.
- Pelkmans, L., Puntener, D., and Helenius, A. (2002). Local actin polymerization and dynamin recruitment in SV40-induced internalization of caveolae. *Science* 296, 535–539.
- Ravanti, L., Heino, J., Lopez-Otin, C., and Kähäri, V.-M. (1999). Induction of collagenase-3 (MMP-13) expression in human skin fibroblasts by three-dimensional collagen is mediated by p38 mitogen-activated protein kinase. *J. Biol. Chem.* 274, 2446–2455.
- Razani, B. *et al.* (2001). Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities. *J. Biol. Chem.* 276, 38121–38138.
- Razani, B. *et al.* (2002). Caveolin-1-deficient mice are lean, resistant to diet-induced obesity, and show hypertriglyceridemia with adipocyte abnormalities. *J. Biol. Chem.* 277, 8635–8647.
- Richterova, Z., Liebl, D., Horak, M., Palkova, Z., Stokrova, J., Hozak, P., Korb, J., and Forstova, J. (2001). Caveolae are involved in the trafficking of mouse polyomavirus virions and artificial VP1 pseudocapsids toward cell nuclei. *J. Virol.* 75, 10880–10891.
- Schubert, W., Frank, P.G., Razani, B., Park, D.S., Chow, C.W., and Lisanti, M.P. (2001). Caveolae-deficient endothelial cells show defects in the uptake and transport of albumin in vivo. *J. Biol. Chem.* 276, 48619–48622.
- Schwartz, M., and Ginsberg, M. (2002). Networks and crosstalk: integrin signaling spreads. *Nat. Cell Biol.* 4, E65–E68.
- Segal, G., Lee, W., Arora, P.D., McKee, M., Downey, G., and McCulloch, C.A. (2001). Involvement of actin filaments and integrins in the binding step in collagen phagocytosis by human fibroblasts. *J. Cell Sci.* 114, 119–129.
- Shin, J., Gao, Z., and Abraham, S. (2000). Involvement of cellular caveolae in bacterial entry into mast cells. *Science* 289, 785–788.
- Simon, D.I., Ezratty, A.M., Francis, S.A., Rennke, H., and Loscalzo, J. (1993). Fibrin(ogen) is internalized and degraded by activated human monocytoic cells via Mac-1 (CD11b/CD18): a nonplasmin fibrinolytic pathway. *Blood* 82, 2414–2422.
- Smart, E., Foster, D., Ying, Y.-S., Kamen, B., and Anderson, R. (1994). Protein kinase C activators inhibit receptor-mediated potocytosis by preventing internalization of caveolae. *J. Cell Biol.* 124, 307–313.
- Sotgia, F. *et al.* (2002). Intracellular retention of glycosylphosphatidyl inositol-linked proteins in caveolin-deficient cells. *Mol. Cell. Biol.* 22, 3905–3926.
- Srivastava, J., Procyk, K.J., Iturrioz, X., and Parker, P.J. (2002). Phosphorylation is required for PMA- and cell-cycle-induced degradation of protein kinase Cdelta. *Biochem. J.* 368, 349–355.
- Thomsen, P., Roepstorff, K., Stahlhut, M., and van Deurs, B. (2002). Caveolae are highly immobile plasma membrane microdomains, which are not involved in constitutive endocytic trafficking. *Mol. Biol. Cell* 13, 238–250.
- Wang, J. (2002). Protein recognition by cell surface receptors: physiological receptors versus virus interactions. *Trends Biochem. Sci.* 27, 122–126.
- Wary, K.K., Mainiero, F., Isakoff, S., Marcantonio, E., and Giancotti, F.G. (1996). The adaptor protein Shc couples a class of integrins to the control of cell cycle progression. *Cell* 87, 733–743.
- Wary, K.K., Mariotti, A., Zurzolo, C., and Giancotti, F.G. (1998). A requirement for caveolin-1 and associated kinase Fyn in integrin signaling and anchorage-dependent cell growth. *Cell* 94, 625–634.
- Wei, Y., Yang, X., Liu, Q., Wilkins, J.A., and Chapman, H.A. (1999). A role for caveolin and the urokinase receptor in integrin-mediated adhesion and signaling. *J. Cell Biol.* 144, 1285–1294.
- Wencel-Drake, J.D., Boudignon-Proudhon, C., Dieter, M.G., Criss, A.B., and Parise, L.V. (1996). Internalization of bound fibrinogen modulates platelet aggregation. *Blood* 87, 602–612.
- Xu, J., and Clark, R. (1997). A three-dimensional collagen lattice induces protein kinase C- $\zeta$  activity: role in  $\alpha 2$  integrin and collagenase mRNA expression. *J. Cell Biol.* 136, 473–483.