



Co-occurrence of Variants of *mcr-3* and *mcr-8* Genes in a *Klebsiella pneumoniae* Isolate From Laos

Linda Hadjadj^{1,2}, Sophie Alexandra Baron^{1,2,3}, Abiola Olumuyiwa Olaitan^{1,2}, Serge Morand⁴ and Jean-Marc Rolain^{1,2,3*}

¹ Aix Marseille Univ, IRD, MEPHI, Faculté de Médecine et de Pharmacie, Marseille, France, ² IHU Méditerranée Infection, Marseille, France, ³ Assistance Publique des Hôpitaux de Marseille, Marseille, France, ⁴ Institut des Sciences de l'Évolution, CNRS-IRD-UM2, CC065, Université Montpellier 2, Montpellier, France

Colistin is considered as a last resort antibiotic. The re-use of this antibiotic highlighted the emergence of colistin resistance mediated by chromosomal and plasmidic resistance mechanisms. Five colistin-resistant *Klebsiella pneumoniae* strains from Laos and Thailand were analyzed by Next Generation Sequencing (NGS) approaches to determine their colistin resistance mechanisms. Antimicrobial susceptibility testing, conjugation and transformation were performed on these strains. Moreover, whole genome sequencing (WGS) combining Illumina (MiSeq) and Oxford Nanopore technologies (MinION) was realized to obtain closed genomes and plasmids. Resistome analyses as well as location of *mcr* genes and its genetic environments were done *in silico*. All five strains had colistin MIC of 32 mg/L and were positive for *mcr-3* variants including additionally positive for a *mcr-8* variant gene. The novel variants were named *mcr-3.21*, *mcr-3.26*, *mcr-3.28*, and *mcr-8.3* genes. The *mcr-3* variants genes were located on plasmids IncP1, IncFII, and IncI1 type, while *mcr-8.3* gene was found on an IncFII type plasmid. The genetic environment of *mcr-3.21* and *mcr-3.26* genes were composed of a composite transposon ISKpn40- *mcr-3*-*dgkA*- ISKpn40. Concerning *mcr-8.3* gene, a similar genetic environment of *mcr-8.1* gene surrounded by ISIX2 and IS903B was observed. To the best of our knowledge, this is the first description of the novel variants *mcr-3.21*, *mcr-3.26*, *mcr-3.28* and *mcr-8.3* genes as well as the first study on co-occurrence of *mcr-3* and *mcr-8* genes. Spread and evolution of *mcr* genes should be monitored.

Keywords: colistin, *mcr*, whole genome sequencing, resistance, epidemiology – descriptive

INTRODUCTION

Antibiotic resistance is a major problem worldwide. This phenomenon has required adjustments in therapeutic approaches, including the use of powerful and broad-spectrum antibiotics, such as carbapenems against multi-drug resistant Gram-negative bacteria. The emergence of carbapenem resistant *Enterobacteriaceae* has led to the re-use of an old antibiotic: colistin (Biswas et al., 2012). This new therapeutic strategy promoted the apparition of colistin resistance mediated by chromosomal and plasmidic resistance mechanisms (Baron et al., 2016). In *Klebsiella pneumoniae*, *mgrB* inactivation appears to be the predominant mechanism of chromosomal resistance to colistin.

OPEN ACCESS

Edited by:

Laura Villa,
National Institutes of Health (NIH), Italy

Reviewed by:

Jian-Hua Liu,
South China Agricultural University,
China
Hong-Yu OU,
Shanghai Jiao Tong University, China

*Correspondence:

Jean-Marc Rolain
jean-marc.rolain@univ-amu.fr

Specialty section:

This article was submitted to
Antimicrobials, Resistance
and Chemotherapy,
a section of the journal
Frontiers in Microbiology

Received: 18 July 2019

Accepted: 08 November 2019

Published: 26 November 2019

Citation:

Hadjadj L, Baron SA, Olaitan AO,
Morand S and Rolain J-M (2019)
Co-occurrence of Variants of *mcr-3*
and *mcr-8* Genes in a *Klebsiella*
pneumoniae Isolate From Laos.
Front. Microbiol. 10:2720.
doi: 10.3389/fmicb.2019.02720

Changes in amino acids in the two-component control systems PmrAB, PhoPQ, and CrrAB are also known to decrease colistin sensitivity by adding 4-amino-4-deoxy-L-arabinose and/or phosphoethanolamine to lipopolysaccharide (Baron et al., 2016). The *mcr-1* gene encoding a phosphoethanolamine transferase is the most widespread mechanism of colistin resistance mediated by a plasmid reported in the literature. To date, nine different plasmid-mediated colistin resistance genes have been described, noted from *mcr-1* to *mcr-9*, which are isolated from bacteria in human, animal and environmental samples (Liu et al., 2016; Yin et al., 2017; Wang et al., 2018; Carroll et al., 2019). Although *mcr-1* has been mainly reported in *Escherichia coli* from animal origins, *K. pneumoniae* can also harbor *mcr* genes (Yin et al., 2017). In a previous study, we isolated 32 colistin-resistant *K. pneumoniae* strains from human and animal samples collected in Laos, Thailand and France (Olaitan et al., 2014). 20 of these isolates had an inactivation of the *mgrB* gene but 12 strains had no colistin resistance mechanisms identified. In this work, we re-analyzed five of these strains by a Next-Generation Sequencing (NGS) approach to finally identify the mechanism of colistin resistance of these isolates.

MATERIALS AND METHODS

Bacterial Strains and Antimicrobial Susceptibility Testing

Five colistin-resistant *K. pneumoniae* strains isolated in 2012 from stool samples of healthy humans from Laksip, Laos (LH102-A, LH94, LH375) and from Thailand (TH114, TH164) were reanalyzed by NGS approach to determine their colistin resistance mechanisms. Approval was given by the Ministry of Health Council of Medical Sciences, National Ethics Committee for Health Research (NHCHR), Lao PDR [no.51/NECHR] and by the Ethical Committee of Mahidol University, Bangkok, Thailand [no.0517.1116/661]. Each participant was informed and signed a consent form. Consents are kept at the Rodolphe Mérieux Laboratory in Laos, Vientiane, for Laos and at the Department of Helminthology, Faculty of Tropical Medicine, Mahidol University, Bangkok, for Thailand.

All strains had intact *mgrB* gene (Olaitan et al., 2014). Antimicrobial susceptibility testing (AST) was determined by minimum inhibitory concentrations (MICs) using microdilution with UMIC test (Biocentric, Bandol, France) for colistin and using the *E*-test method (bioMérieux, Marcy l'Etoile, France) for others antibiotics in accordance with EUCAST recommendations.

Plasmid Transferability

To confirm *mcr* genes location biologically, conjugation was performed with an azide-resistant *E. coli* J53 strain. Selection of transconjugants was done on MacConkey agar (Beckton Dickinson, Le Pont de Claix, France) supplemented with 120 mg/L sodium azide and 2 mg/L colistin. In case of unsuccessful conjugation, transformation of pure plasmid by electroporation method with Top10 electrocompetents *E. coli*

(Thermo Fisher Scientific, Waltham, MA, United States) was performed and transformants were selected on LB agar (Beckton Dickinson, France) with 2 mg/L colistin. Plasmid curing was performed at high temperature on *K. pneumoniae* LH94 strain (Letchumanan et al., 2015). Then, the presence of *mcr* genes and plasmid types were controlled by standard PCR (Carattoli et al., 2005; García-Fernández et al., 2009; Rebelo et al., 2018; Wang et al., 2018).

Genomic Analysis

All strains were sequenced using both MiSeq (Illumina Inc., San Diego, CA, United States) and MinION (Oxford Nanopore Technologies Inc., United Kingdom) technologies. Spades genome assembler was used to assemble long read sequencing data generated by Nanopore and short read data generated by Illumina (Bankevich et al., 2012). Genomes were annotated with RAST (Aziz et al., 2008), resistance gene using both ARG-ANNOT (Gupta et al., 2014) and blastN alignment, and the presence of plasmid was screened using PlasmidFinder software (Carattoli et al., 2014). oriFinder was used to detect the presence of transfer-related modules of the sequenced plasmids: oriT region, relaxase, T4SS and T4CP (Li et al., 2018). The sequence type (ST) of *K. pneumoniae* strains was performed *in silico* by multilocus sequence typing (MLST) analysis using the MLST database (Larsen et al., 2012). The presence of known mutations conferring colistin resistance on the *mgrB*, *pmrA*, *pmrB*, *phoP*, *phoQ*, *crrA* and *crrB* genes has been also investigated *in silico* using both nucleic and amino acids alignments. Mutations of *mcr* genes have been verified by Sanger sequencing. The genetic environment of *mcr* genes has been reconstituted by comparing the sequence of genes surrounding the *mcr*-like genes against the NCBI database, using blastX parameter. These environments were also compared to those of *mcr-3.1* (KY924928), *mcr-3.11* (MG552133) and *mcr-8.1* (MG736312) genes. The plasmid pKP91 (MG736312) carrying the first *mcr-8.1* gene described in the literature was compared with the plasmid pLH94-8 (CP035204) of the strain LH94 that harbor a *mcr-8.3* gene using a pairwise comparison of their Average Nucleotide Identity based on Blast (ANIb) on Jspecies software (Richter et al., 2016). CGViewServer was used to represent the complete sequences of plasmids carrying *mcr* genes (Stothard et al., 2017).

Genomes of LH102-A, LH94, LH375, TH114, and TH164 strains have been submitted to GenBank under accession numbers CP035194-CP035195, CP035202-CP035206, CP035196-CP035201, CP035207-CP035209, and CP035210-CP035213, respectively (Table 1). Novel variants of *mcr-3* and *mcr-8* genes were detected and assigned by NCBI (Partridge et al., 2018) as *mcr-3.21* (AWY10763), *mcr-3.26* (AWY10762), *mcr-3.28* (AXI82466) and *mcr-8.3* (AXI82467) (Figures 1A,B).

RESULTS

All *K. pneumoniae* strains had a colistin MIC at 32 mg/L (Table 1). They remain susceptible to ceftriaxone, ertapenem, imipenem, fosfomicin and amikacin but were resistant

TABLE 1 | Characteristics of *Klebsiella* spp. strains analyzed in this study.

Strains	Genome size (bp)	GC%	ST	Sensitive to (MIC mg/L)	Resistant to (MIC mg/L)	ARGs	Plasmid type	Conjugal modules predicted	Accession number
<i>K. pneumoniae</i> LH102-A	5,213,480	57.0	87	CRO (0.023), ERT (0.006), IMP (0.25), FF (32), AK (2), GN (0.38), SXT (0.064), CIP (0.047)	AMX (>256), AMC (4), CT (32)	<i>bla</i> _{SHV-27}	/	/	CP035194
						<i>bla</i> _{TEM-1B} , <i>mcr-3.28</i>	IncP1 (pLH102-A)	oriT, Relaxase, T4SS, T4CP	CP035195
<i>K. pneumoniae</i> LH94	5,709,714	57.0	39	CRO (0.032), ERT (0.006), IMP (0.19), FF (32), AK (2), GN (0.38)	AMX (>256), AMC (4), SXT (>32), CIP (1), CT (32)	<i>bla</i> _{SHV-11}	/	/	CP035202
						<i>bla</i> _{TEM-1B} , <i>qnrS1</i> , <i>aadA1</i> , <i>aadA2</i> , <i>floR</i> , <i>cmlA1</i> , <i>sul3</i> , <i>sul2</i> , <i>dfrA12</i> , <i>mcr-8.3</i>	IncFII (pLH94-8)	oriT, Relaxase, T4SS*, T4CP	CP035204
						<i>mcr-3.21</i>	IncI1 (pLH94-3)	oriT, Relaxase, T4SS, T4CP	CP035205
<i>K. pneumoniae</i> LH375	5,372,573	57.3	1315	CRO (0.032), ERT (0.008), IMP (0.25), FF (24), AK (2), GN (0.38), SXT (0.064)	AMX (>256), AMC (4), CIP (1), CT (32)	<i>bla</i> _{SHV-1}	/	/	CP035196
						<i>mcr-3.26</i>	IncFII (pLH375-3)	oriT, T4SS	CP035200
						<i>bla</i> _{TEM-1C} , <i>qnrS1</i> , <i>aadA2</i> , <i>lnu(F)</i> , <i>floR</i> , <i>tetA</i>	IncR (pLH375-2)	/	CP035198
<i>K. quasipneumoniae</i> TH114	5,242,749	57.7	1321	CRO (0.032), ERT (0.008), IMP (0.25), FF (32), AK (2), GN (0.38), CIP (0.032)	AMX (>256), AMC (4), SXT (>32), CT (32)	<i>bla</i> _{O_{KP}-B-7}	/	/	CP035207
						<i>mcr-3.21</i>	IncI1 (pTH114-3)	oriT, Relaxase, T4SS, T4CP	CP035208
						<i>bla</i> _{TEM-1B} , <i>aadA2</i> , <i>floR</i> , <i>sul2</i> , <i>tetA</i> , <i>dfrA12</i>	IncR (pTH114-1)	oriT	CP035209
<i>K. pneumoniae</i> TH164	5,656,506	57.3	873	CRO (0.023), ERT (0.006), IMP (0.25), FF (16), AK (2)	AMX (>256), AMC (4), SXT (>32), CIP (1), GN (48), CT (32)	<i>bla</i> _{SHV-27}	/	/	CP035210
						<i>mcr-3.21</i>	IncI1 (pTH164-3)	oriT, Relaxase, T4SS, T4CP	CP035213
						<i>bla</i> _{TEM-1B} , <i>qnrS1</i> , <i>aadA1</i> , <i>aadA2</i> , <i>aac(3)-IId</i> , <i>floR</i> , <i>cmlA1</i> , <i>sul3</i> , <i>sul2</i> , <i>tetA</i> , <i>dfrA12</i>	IncR (pTH164-1)	oriT	CP035212

MIC, minimum inhibitory concentration; ARGs, antibiotic resistance genes; ST, sequence type; AMX, amoxicillin; AMC, amoxicillin-clavulanic; CT, colistin; CRO, ceftriaxone; ERT, ertapenem; IMP, imipenem; FF, fosfomicin; AK, amikacin; GN, gentamicin; SXT, trimethoprim-sulfamethoxazole; CIP, Ciprofloxacin; *, incomplete.





FIGURE 1 | (A) Amino acid sequence alignments of the new MCR-3 and MCR-8 variants using Clustal W on NPS@ website. **(B)** Phylogenetic tree of MCR variants, including MCR-3 and MCR-8 variants. Sequences were aligned using MUSCLE and phylogenetic inferences were obtained using the neighbor-joining method within the MEGA 7 software. Numbers at the nodes are percentages of bootstrap values obtained by repeating the analysis 500 times to generate a majority consensus tree. Evolutionary distances were computed using the Poisson correction method and are expressed in units of the number of amino acid substitutions per site.




to amoxicillin and amoxicillin-clavulanic acid. Resistance to trimethoprim-sulfamethoxazole, ciprofloxacin and gentamicin was inconstantly observed (**Table 1**). The resistome analysis of these strains indicated the presence of genes encoding resistance for beta-lactams (*bla_{SHV}*, *bla_{OKP-B}*, *bla_{TEM}*), aminoglycosides (*aadA*) sulphonamides (*sul*), fluoroquinolones (*qnrS1*), fosfomycin, phenicol (*floR*), tetracycline (*tetA*), trimethoprim (*dfrA*), and colistin (*mcr*) (**Table 1**).

Genome size of the five *Klebsiella* spp. strains ranged from 5,213,480 to 5,709,714 bp with chromosomes and plasmids included for each strain. Blast analysis of RpoB protein and the specific presence of the *bla_{OKP-B-7}* gene instead of *bla_{SHV}* gene allow to reclassify TH114 isolate as a *Klebsiella quasipneumoniae* strain (**Table 1**). According to the MLST analysis, each strain belonged to different sequence types (ST39, ST97, ST873, ST1315, and ST1321).

We observed no disruption or known mutations in the *mgrB* gene or in the different two-component systems *pmrAB*, *phoPQ*, and *csrAB* of these five genomes. However, *K. pneumoniae* LH102-A and LH375 carried a *mcr-3.28* and a *mcr-3.26* gene, respectively. The strains LH94, TH164 and TH114 harbored a *mcr-3.21* gene, but LH94 isolate also carried *mcr-8.3* gene (**Table 1**). Compared to *mcr-3.1* gene, *mcr-3.21* displayed two amino acids substitution (N296D and Q468K), while *mcr-3.26* had only one amino acid change (N296D). For the *mcr-3.28* gene, two mutations (M23V and A457E) were identified. The *mcr-8.3* gene is distinguished from *mcr-8.1* by an amino acid substitution T391K (**Figures 1A,B**). In each strain, the *mcr-3* gene was present on one plasmid but lastly only conjugate in LH102-A, LH375 and TH114 strains. The transconjugants and transformants isolates had a colistin MIC of 2 mg/L and were susceptible to antibiotics previously tested except for amoxicillin in a transconjugant LH102-A strain which was

consistent with the presence of a *bla_{TEM-1B}* one the plasmid harboring the *mcr* gene (**Table 1**). The *mcr-8* and *mcr-3* genes of the LH94 strain were not found in the original strain after curing plasmid. In addition, the *mcr-8* gene was not detected in transformants of the LH94 strain carrying the *mcr-3* gene, which confirms its presence on another plasmid. The presence of potential transfer-related modules, including the origin of transfer (*oriT*) region, the relaxase gene, T4CP gene and the type IV secretion system gene cluster (T4SS) were detected in some strains (**Table 1**).

The reconstitution of plasmids was achieved using reads from both Miseq and Minion sequencer. The three *mcr-3.21* genes were carried by an IncI1 plasmid type (LH94, TH114 and TH164) whereas the *mcr-3.28* gene was harbored by an IncP1 (LH102-A) and the *mcr-3.26* by an IncFII (LH375). The *mcr-8.3* gene was also found on an IncFII plasmid type (LH94) (**Table 1**). Plasmid structure of the five plasmids carrying *mcr-3* genes and the plasmid containing the *mcr-8* gene was represented in **Figure 2**. Plasmids LH94, TH114, and TH164 belonging to IncI1 plasmid type were very similar, except for TH164 that carried more transposases genes around *mcr* gene. Although the close genetic environment of *mcr* genes was very similar between the five plasmids, the LH375 and LH102A ones belong to other plasmid type and so were different from the IncI1 plasmids. The genetic environments of *mcr-3.21* and *mcr-3.26* genes in strains TH114, LH94, TH164, and LH375 were similar and composed of a composite transposon ISKpn40-*mcr-3.21-dgkA*-ISKpn40 such as *mcr-3.11* gene (**Figure 3A**; Xiang et al., 2018). Concerning the *mcr-3.28* gene, it was framed by two mobile elements as follow: TnAs2-*mcr-3.28-dgkA*-IS26. For *mcr-8.3* gene, a similar genetic environment of *mcr-8.1* gene was observed (**Figure 3B**). It was interesting to note the presence of two genes annotated as a response regulator transcription factor CopR for the



first one (EQH49_28300) and as a HAMP domain-containing histidine kinase BaeS for the second one (EQH49_28305). The combination of these two protein families generally constitutes a two-component system. These systems are often involved in colistin resistance in *Enterobacteriaceae* (*pmrAB*, *crrAB* or *phoPQ*). The presence of a two-component system on a plasmid carrying the *mcr-8* gene requires further analysis to understand its role in colistin resistance. The plasmids pLH94-8 (CP035204) and pKP91 (MG736312) harboring *mcr-8* gene shared 95.42% of identity.

DISCUSSION

Following plasmid-mediated colistin resistance gene *mcr-1*, the *mcr-3* gene is the most widely described variants found in the literature, while *mcr-2*, and *mcr-4* to *mcr-8* remain uncommon. The MCR-3 protein shares only 32.5% amino acid with MCR-1 which makes impossible its detection with previously described *mcr-1* primers (Yin et al., 2017). Since its first description in 2017, thirty variants of *mcr-3* gene have been deposit in

NCBI and this gene has been observed in different countries and in different strains such as *E. coli*, *Aeromonas* spp., and *K. pneumoniae* (Figure 1B; Yin et al., 2017; Fukuda et al., 2018; Shen et al., 2018).

The presence of *K. pneumoniae* strains carrying the *mcr-3* gene in samples from humans in Thailand has already been described (Yin et al., 2017) but never in Laos. Similarly to the transposon Tn6330 (*ISAp11-mcr-1-ORF-ISAp11*), a composite transposon ISKpn40-*mcr-3-dgkA*-ISKpn40 has been reported in several type of plasmids (Xiang et al., 2018). It is known the integration of this type of transposon into a variable region is mediated by insertion sequences (IS), that are then lost to stabilize the integration of the transposon (Snesrud et al., 2018). The genetic environment of our *mcr-8.3* gene was similar to that of the *mcr-8.1* gene in the reference plasmid pK91 which was isolated from a swine fecal sample in China (Wang et al., 2018). However, unlike plasmid pK91, the pLH94-8 plasmid also harbors other antibiotic resistance genes. The co-occurrence of different antibiotic resistance genes could explain the presence of two *mcr* genes in a same isolate (Bi et al., 2015). Indeed,

this *mcr-8* gene could have been selected because it was harbored by a plasmid that also conferred resistance to beta-lactam, aminoglycoside, sulphonamide, fluoroquinolone, phenicol and trimethoprim.

The first description of the *mcr-8* gene was done in *K. pneumoniae* strains isolated from animals and human in 2016 in China. This gene was found on a conjugative IncFII type plasmid (Wang et al., 2018). To date, the *mcr-8* gene and its variants have only been described in China in *Stenotrophomonas* spp., *Raoultella ornithinolytica* and *K. quasipneumoniae* from animal and environmental origins (Li et al., 2019; Wang et al., 2019; Yang et al., 2019). Since MCR-8 had only 30.26 to 39.96% identity with other MCR, it is possible that the prevalence of *mcr-8* gene is underestimated. Reanalysis of previous works will probably respond to this question. In our study, the detection of a *mcr-8.3* gene in a *K. pneumoniae* strain isolated from a 2012 sample from Laos raises the question of the spread of this gene. Moreover, the emergence of novel variants for *mcr-3* and *mcr-8* gene indicates that this gene has evolved over time. Furthermore, co-location in the same strain on different plasmids encoding different *mcr* genes and others antibiotic resistance genes increases the risk of spread of colistin resistance (Long et al., 2019). Antibiotic resistance genes and their transferability are a major risk for human health. The “One health” approach promoting systemic coordination of three sectors: environment, animals and humans could be beneficial for the effective control of global antibiotic resistance problems (Ryu et al., 2017).

To the best of our knowledge, this is the first study describing the novel variants *mcr-3.21*, *mcr-3.26*, *mcr-3.28*, *mcr-8.3* genes and demonstrating the co-existence of MCR-3 and MCR-8 in a *K. pneumoniae* isolate from Laos.

REFERENCES

- Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., et al. (2008). The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. doi: 10.1186/1471-2164-9-75
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., et al. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* 19, 455–477. doi: 10.1089/cmb.2012.0021
- Baron, S., Hadjadj, L., Rolain, J. M., and Olaitan, A. O. (2016). Molecular mechanisms of polymyxin resistance: knowns and unknowns. *Int. J. Antimicrob. Agents* 48, 583–591. doi: 10.1016/j.ijantimicag.2016.06.023
- Bi, D., Jiang, X., Sheng, Z.-K., Ngmenterebo, D., Tai, C., Wang, M., et al. (2015). Mapping the resistance-associated mobilome of a carbapenem-resistant *Klebsiella pneumoniae* strain reveals insights into factors shaping these regions and facilitates generation of a ‘resistance-disarmed’ model organism. *J. Antimicrob. Chemother.* 70, 2770–2774. doi: 10.1093/jac/dkv204
- Biswas, S., Brunel, J.-M., Dubus, J.-C., Reynaud-Gaubert, M., and Rolain, J.-M. (2012). Colistin: an update on the antibiotic of the 21st century. *Expert Rev. Anti. Infect. Ther.* 10, 917–934. doi: 10.1586/eri.12.78
- Carattoli, A., Bertini, A., Villa, L., Falbo, V., Hopkins, K. L., and Threlfall, E. J. (2005). Identification of plasmids by PCR-based replicon typing. *J. Microbiol. Methods* 63, 219–228. doi: 10.1016/j.mimet.2005.03.018
- Carattoli, A., Zankari, E., Garcia-Fernandez, A., Voldby Larsen, M., Lund, O., Villa, L., et al. (2014). In silico detection and typing of plasmids using plasmidfinder

DATA AVAILABILITY STATEMENT

The datasets analyzed in this manuscript are publicly available on GenBank. Requests to access the datasets should be directed to linda.hadjadj@uni-amu.fr.

AUTHOR CONTRIBUTIONS

LH wrote the manuscript, performed the experiments, and analyzed the data. SB helped for bioinformatics analysis and to draft the manuscript. AO isolated the strains and helped to draft the manuscript. SM was responsible for the collection of samples and helped to draft the manuscript. J-MR conceived the study, participated in designing and coordination, and helped to draft the manuscript. All authors read and approved the final draft of the manuscript.

FUNDING

This work was supported by the ANR SEA-EU-NET FarmResist (IZJFZ3-177614) and the French ANR FutureHealthSEA (ANR-17-CE35-0003-02). It was also supported by the French Government under the “Investissements d’avenir” (Investments for the Future) program managed by the Agence Nationale de la Recherche (ANR, fr: National Agency for Research) (reference: Méditerranée Infection 10-IAHU-03), Région Provence-Alpes-Côte d’Azur, and European funding FEDER PRIMI.

ACKNOWLEDGMENTS

The authors thank CookieTrad for proofreading the text.

- and plasmid multilocus sequence typing. *Antimicrob. Agents Chemother.* 58, 3895–3903. doi: 10.1128/AAC.02412-14
- Carroll, L. M., Gaballa, A., Guldemann, C., Sullivan, G., Henderson, L. O., and Wiedmann, M. (2019). Identification of novel mobilized colistin resistance gene *mcr-9* in a multidrug-resistant, colistin-susceptible *Salmonella enterica* serotype *Typhimurium* isolate. *mBio* 10:e852-19. doi: 10.1128/mBio.00853-19
- Fukuda, A., Usui, M., Okubo, T., Tagaki, C., Sukpanyatham, N., and Tamura, Y. (2018). Co-harboring of cephalosporin (bla)/colistin (*mcr*) resistance genes among *Enterobacteriaceae* from flies in Thailand. *FEMS Microbiol. Lett.* 365:fny178. doi: 10.1093/femsle/fny178
- García-Fernández, A., Fortini, D., Veldman, K., Mevius, D., and Carattoli, A. (2009). Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. *J. Antimicrob. Chemother.* 63, 274–281. doi: 10.1093/jac/dkn470
- Gupta, S. K., Padmanabhan, B. R., Diene, S. M., Lopez-Rojas, R., Kempf, M., Landraud, L., et al. (2014). ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrob. Agents Chemother.* 58, 212–220. doi: 10.1128/AAC.01310-13
- Larsen, M. V., Cosentino, S., Rasmussen, S., Friis, C., Hasman, H., Marvig, R. L., et al. (2012). Multilocus sequence typing of total-genome-sequenced bacteria. *J. Clin. Microbiol.* 50, 1355–1361. doi: 10.1128/JCM.06094-11
- Letchumanan, V., Chan, K.-G., and Lee, L.-H. (2015). An insight of traditional plasmid curing in *Vibrio* species. *Front. Microbiol.* 6:735. doi: 10.3389/fmicb.2015.00735

- Li, J., Liu, S., Fu, J., Yin, J., Zhao, J., Zhong, C., et al. (2019). Co-occurrence of colistin and meropenem resistance determinants in a *Stenotrophomonas* strain isolated from sewage water. *Microb. Drug Resist.* 25, 317–325. doi: 10.1089/mdr.2018.0418
- Li, X., Xie, Y., Liu, M., Tai, C., Sun, J., Deng, Z., et al. (2018). oriTfinder: a web-based tool for the identification of origin of transfers in DNA sequences of bacterial mobile genetic elements. *Nucleic Acids Res.* 46, W229–W234. doi: 10.1093/nar/gky352
- Liu, Y.-Y., Wang, Y., Walsh, T. R., Yi, L.-X., Zhang, R., Spencer, J., et al. (2016). Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect. Dis.* 16, 161–168. doi: 10.1016/S1473-3099(15)00424-7
- Long, H., Feng, Y., Ma, K., Liu, L., McNally, A., and Zong, Z. (2019). The co-transfer of plasmid-borne colistin-resistant genes *mcr-1* and *mcr-3.5*, the carbapenemase gene *bla*NDM-5 and the 16S methylase gene *rmtB* from *Escherichia coli*. *Sci. Rep.* 9:696. doi: 10.1038/s41598-018-37125-1
- Olaitan, A. O., Diene, S. M., Kempf, M., Berrazeg, M., Bakour, S., Gupta, S. K., et al. (2014). Worldwide emergence of colistin resistance in *Klebsiella pneumoniae* from healthy humans and patients in Lao PDR, Thailand, Israel, Nigeria and France owing to inactivation of the PhoP/PhoQ regulator *mgrB*: an epidemiological and molecular study. *Int. J. Antimicrob. Agents* 44, 500–507. doi: 10.1016/j.ijantimicag.2014.07.020
- Partridge, S. R., Di Pilato, V., Doi, Y., Feldgarden, M., Haft, D. H., Klimke, W., et al. (2018). Proposal for assignment of allele numbers for mobile colistin resistance (*mcr*) genes. *J. Antimicrob. Chemother.* 73, 2625–2630. doi: 10.1093/jac/dky262
- Rebelo, A. R., Bortolaia, V., Kjeldgaard, J. S., Pedersen, S. K., Leekitcharoenphon, P., Hansen, I. M., et al. (2018). Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Euro. Surveill.* 23:17-00672. doi: 10.2807/1560-7917.ES.2018.23.6.17-00672
- Richter, M., Rosselló-Móra, R., Oliver Glöckner, F., and Peplies, J. (2016). JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32, 929–931. doi: 10.1093/bioinformatics/btv681
- Ryu, S., Kim, B. I., Lim, J.-S., Tan, C. S., and Chun, B. C. (2017). One health perspectives on emerging public health threats. *J. Prev. Med. Public Heal.* 50, 411–414. doi: 10.3961/jpmph.17.097
- Shen, Y., Xu, C., Sun, Q., Schwarz, S., Ou, Y., Yang, L., et al. (2018). Prevalence and genetic analysis of *mcr-3*-positive aeromonas species from humans, retail meat, and environmental water samples. *Antimicrob. Agents Chemother.* 62:e404-18. doi: 10.1128/AAC.00404-18
- Snesrud, E., McGann, P., and Chandler, M. (2018). The birth and demise of the ISApl1-*mcr-1*-ISApl1 composite transposon: the vehicle for transferable colistin resistance. *mBio* 9:e02381-17. doi: 10.1128/mBio.02381-17
- Stothard, P., Grant, J. R., and Van Domselaar, G. (2017). Visualizing and comparing circular genomes using the CGView family of tools. *Brief. Bioinform.* 20, 1576–1582. doi: 10.1093/bib/bbx081
- Wang, X., Wang, Y., Zhou, Y., Li, J., Yin, W., Wang, S., et al. (2018). Emergence of a novel mobile colistin resistance gene, *mcr-8*, in NDM-producing *Klebsiella pneumoniae*. *Emerg. Microbes Infect.* 7:122. doi: 10.1038/s41426-018-0124-z
- Wang, X., Wang, Y., Zhou, Y., Wang, Z., Wang, Y., Zhang, S., et al. (2019). Emergence of colistin resistance gene *mcr-8* and its variant in *Raoultella ornithinolytica*. *Front. Microbiol.* 10:228. doi: 10.3389/fmicb.2019.00228
- Xiang, R., Liu, B.-H., Zhang, A.-Y., Lei, C.-W., Ye, X.-L., Yang, Y.-X., et al. (2018). Colocation of the polymyxin resistance gene *mcr-1* and a variant of *mcr-3* on a plasmid in an *Escherichia coli* isolate from a chicken farm. *Antimicrob. Agents Chemother.* 62:e00501-18. doi: 10.1128/AAC.00501-18
- Yang, X., Liu, L., Wang, Z., Bai, L., and Li, R. (2019). Emergence of *mcr-8.2*-bearing *Klebsiella quasipneumoniae* of animal origin. *J. Antimicrob. Chemother.* 74, 2814–2817. doi: 10.1093/jac/dkz213
- Yin, W., Li, H., Shen, Y., Liu, Z., Wang, S., Shen, Z., et al. (2017). Novel plasmid-mediated colistin resistance gene *mcr-3* in *Escherichia coli*. *mBio* 8:e543-17. doi: 10.1128/mBio.00543-17

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Hadjadj, Baron, Olaitan, Morand and Rolain. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.