

Cocliques in the Kneser graph on the point-hyperplane flags of a projective space

A. Blokhuis, A. E. Brouwer, Ç. Güven

October 14, 2010

Abstract

We prove an Erdős-Ko-Rado-type theorem for the Kneser graph on the point-hyperplane flags in a finite projective space.

1 Introduction

Let V be a vector space of finite dimension $n > 0$ over \mathbf{F}_q , and consider in the associated projective space PV the flags (P, H) consisting of an incident point-hyperplane pair (that is, with $P \subseteq H$), and call two such flags (P, H) and (P', H') adjacent when $P \not\subseteq H'$ and $P' \not\subseteq H$. We study maximal cocliques in this graph $\Gamma(V)$.

Let $F = (X_1, \dots, X_{n-1})$ be a chamber (maximal flag) in PV , that is, X_i is a subspace of V of vector space dimension i for all i , and $X_i \subseteq X_j$ for $i \leq j$. From F we construct the coclique $C_F = \{(P, H) \mid \exists i : P \subseteq X_i \subseteq H\}$. This coclique is maximal, and $|C_F| = 1 + 2q + 3q^2 + \dots + (n-1)q^{n-2}$.

For a coclique C in $\Gamma(V)$, define $Z(C) = \{P \mid (P, H) \in C \text{ for some } H\}$ and $Z'(C) = \{H \mid (P, H) \in C \text{ for some } P\}$.

Theorem 1 *Let C be a coclique in $\Gamma(V)$ and let $Z = Z(C)$. Let $f(n) := 1 + 2q + 3q^2 + \dots + (n-1)q^{n-2}$ and $g(n) := 1 + q + q^2 + \dots + q^{n-2}$. Then*

- (i) $|C| \leq f(n)$, and equality holds iff $C = C_F$ for some chamber F , and
- (ii) $|Z| \leq g(n)$, and for $q > 2$ equality holds iff Z is a hyperplane.

Note that $g(n)$ is the number of points in a hyperplane, and $f(n) = qf(n-1) + g(n)$. Part (i) was conjectured, and partial results were obtained, in Mussche [2]. Both inequalities are sharp, and the theorem characterises equality in (i). What about equality in (ii)? In examples of type C_F the set Z is a hyperplane, and equality holds. When $q = 2$ there are further examples of equality, described below.

Example 1. ($n = 3$) In the plane, take three points P_i ($i = 1, 2, 3$) in general position. Let C consist of the flags $(P_i, P_i + P_{i+1})$ (indices mod 3). Then C is a coclique, and $|Z| = 3 = 2^{n-1} - 1$.

(This example also arises from the trivial one point coclique for $n = 2$ by the construction in Example 3.)

Example 2. ($n = 4$) Consider a plane π and a point ∞ outside π . For each point P in π , let P' be the third point of the line $P + \infty$. Label the points of π with the integers mod 7, so that the lines become $\{1, 2, 4\}$ (mod 7). The seven flags $(i', \langle i', i+1, i+2, i+4 \rangle)$ form a coclique C , and $|Z| = 7 = 2^{n-1} - 1$.

(This is an honest sporadic example.)

Example 3. Let H be a hyperplane in V , and let D be a coclique in $\Gamma(H)$ with $|Z(D)| = 2^{n-2} - 1$. Let G be a hyperplane of H , such that G is disjoint from $Z(D)$. Let Q be a point outside H . Let C consist of the flags (P, H) with $P \subseteq G$, and $(P, N + Q)$ with $(P, N) \in D$, and $(Q, G + Q)$. Then C is a coclique, and $Z(C) = Z(D) \cup G \cup \{Q\}$, so that $|Z| = 2^{n-1} - 1$.

(We'll see later that this only gives something for $n \leq 5$.)

Example 4. Let T be a t -space in V , $0 < t < n$. Let D be a coclique in $\Gamma(T)$. Let E be a coclique in $\Gamma(V/T)$ such that $Z(E)$ has maximal size $2^{n-1-t} - 1$. Construct a coclique C in $\Gamma(V)$ by taking (i) all flags (P, H) with $P \subseteq T \subseteq H$, (ii) the flags (P, K) where $(P, K \cap T) \in D$, and (iii) the flags (Q, H) where $(Q + T, H) \in E$. Then $|Z(C)| = (2^t - 1) + 2^t(2^{n-1-t} - 1) = 2^{n-1} - 1$.

Note that the cocliques C_F are of this form (for every T in F).

The above describes all cases of equality:

Proposition 2 *Let V be an n -dimensional vector space over \mathbf{F}_2 , and let C be a coclique in the graph $\Gamma(V)$. Put $Z := \{P \mid (P, H) \in C \text{ for some } H\}$. Then $|Z| \leq 2^{n-1} - 1$ with equality if and only if Z is a hyperplane, or C arises by the construction of Example 2, 3, or 4.*

1.1 Rank 1 matrices

The graph $\Gamma(V)$ can be described in terms of rank 1 matrices. Represent points by column vectors p and hyperplanes by row vectors h , then a point-hyperplane pair can be represented by the rank 1 matrix ph , and an incident point-hyperplane pair by the rank 1 matrix ph with $hp = 0$, that is, with $\text{tr } ph = 0$. Rank 1 matrices that differ by a constant only represent the same point-hyperplane pair.

Given two incident point-hyperplane pairs $x = ph$ and $x' = p'h'$, they are nonadjacent when $(hp')(h'p) = 0$, i.e., when $\text{tr } xx' = 0$.

Thus, finding the maximal cocliques in $\Gamma(V)$ is equivalent to finding the intersections of the maximal totally isotropic subspaces for the symmetric bilinear form $(x, y) = \text{tr } xy$ on the space $M_n(\mathbf{F}_q)$ (of matrices of order n over \mathbf{F}_q) with the space of trace 0 rank-1 matrices.

The extremal example C_F is conjugate to the example of all rank-1 strictly upper-triangular matrices.

1.2 The thin case

There is a thin analog ($q = 1$ version) of our problem. Consider for an n -set V the pairs (P, H) , where $|P| = 1$, $|H| = n - 1$, and $P \subseteq H \subset V$. Call (P, H) and (P', H') adjacent when $P \not\subseteq H'$ and $P' \not\subseteq H$, that is, when $(P', H') = (V \setminus H, V \setminus P)$. Here the graph is the union of $\binom{n}{2}$ components K_2 . A maximal coclique is obtained by taking a single vertex from each K_2 , so that $|C| \leq \binom{n}{2} = f(n)$ and $|Z| \leq n$. Here $g(n) = n - 1$, and $|Z| \leq g(n)$ holds only for $n = 1, 2$.

2 Maximum-size cocliques

Proof of Theorem 1. The problem is self-dual, so all that is proved for points and hyperplanes, also holds for hyperplanes and points. In particular, $g(n)$ will also be an upper bound for the number $|Z'(C)|$ of hyperplanes involved in C .

We may assume that C is maximal. It follows that C has a certain linear structure:

Lemma 3 *Let C be a maximal coclique in $\Gamma(V)$, and let $H \in Z'(C)$. Then*

H has a subspace $S(H)$ such that $(P, H) \in C$ if and only if $P \subseteq S(H)$. If $H, K \in Z'(C)$ then $S(H) \subseteq K$ or $S(K) \subseteq H$ (or both).

Proof. If $(P, H), (Q, H) \in C$ and R is a point on the line $P + Q$, then also $(R, H) \in C$ since C is maximal. If P is in $S(H)$ but not in K , and Q is in $S(K)$ but not in H , then (P, H) and (Q, K) are adjacent in $\Gamma(V)$, impossible. \square

We use induction on n . If W is a subspace of V , then $\{(P, H \cap W) \mid (P, H) \in C, P \subseteq W, W \not\subseteq H\}$ is a coclique in the graph $\Gamma(W)$. And if S is a subspace of V , then $\{(P + S, H) \mid (P, H) \in C, P \not\subseteq S, S \subseteq H\}$ is a coclique in the graph $\Gamma(V/S)$.

Let the maximal dimension of $S(H)$ (for varying H) be s , and let H be a hyperplane with $\dim S(H) = s$. We have $1 \leq s \leq n - 1$.

Lemma 4 *If $s = n - 1$, then $|C| \leq qf(n - 1) + g(n) = f(n)$ and $|Z| = g(n)$, and $Z = H$.*

Proof. If $s = n - 1$ then $S(H) = H$, so that $S(H) \not\subseteq K$ for $K \neq H$ and therefore $S(K) \subseteq H$ for all K , i.e., $Z = H$ and $|Z| = g(n)$. The coclique C consists of the $g(n)$ flags (P, H) together with at most $qf(n - 1)$ flags (P, K) with $P \subseteq H, K \neq H$, so that $|C| \leq qf(n - 1) + g(n) = f(n)$, as desired. \square

Lemma 5 *If $s < n - 1$, then $|C| < qf(n - 1) + g(n) = f(n)$.*

Proof. Count the three types of flags: a: those involving H , b: those involving M with $S(M)$ contained in H , c: those involving K with $S(K)$ not contained in H (and hence containing $S(H)$).

$$\text{a: } \frac{q^s - 1}{q - 1}; \quad \text{b: at most } qf(n - 1); \quad \text{c: at most } g(n - s) \frac{q^s - 1}{q - 1}.$$

It follows that $|C| \leq qf(n - 1) + 2 \cdot \frac{q^{n-2} - 1}{q - 1} < f(n)$, as desired. \square

Since there is strict inequality here, equality $|C| = f(n)$ only occurs for $s = n - 1$, where there is a hyperplane H such that $(P, H) \in C$ for all points $P \subseteq H$, and all other elements of C restrict to a coclique with equality in $\Gamma(H)$. By induction it follows that equality implies that $C = C_F$ for some flag F .

Lemma 6 *Let $q > 2$ and $s < n - 1$. Then $|Z| < g(n)$.*

Proof. We estimate $|Z|$. For the flags $(P, M) \in C$ with $P \subseteq H$ and $M \neq H$, the system $(P, M \cap H)$ forms a coclique in $\Gamma(H)$, so the number of points P involved is at most $g(n-1)$. For the flags $(Q, K) \in C$ with $Q \not\subseteq H$, the hyperplanes K contain $S(H)$, and the flags $(Q + S(H), K)$ form a coclique in $\Gamma(V/S(H))$, so there are at most $g(n-s)$ such hyperplanes K . For each K there are at most q^{s-1} points Q outside H with $(Q, K) \in C$. Finally, Z contains the points in $S(H)$. Altogether,

$$|Z| \leq g(n-1) + g(n-s) \cdot q^{s-1} + \frac{q^s - 1}{q - 1} < g(n).$$

□

For $q = 2$ and $s > 1$ more work is required, because now the above estimate of $|Z|$ is $2^{s-1} - 1$ larger than the desired upper bound $g(n)$.

As observed, there are at most $g(n-s)$ hyperplanes K involved in flags $(P, K) \in C$ with $P \not\subseteq S(H)$ and $S(H) \subseteq K$. If the number of such K is strictly smaller than this, or $S(K) \subseteq H$ for one of them, then the upper bound on $|Z|$ improves by 2^{s-1} , and $|Z| < g(n)$. Finally, if for at least two of them $\dim S(K) < s$, then again the same holds. So, we may assume that there are precisely $g(n-s)$ such hyperplanes K , for none of them $S(K) \subseteq H$, and for all of them with at most one exception $\dim S(K) = s$.

Let $E(H)$ be the set of points P in $S(H)$ that occur in a single flag only, namely in (P, H) . Let $e(H) = |E(H)|$. The (last) term $2^s - 1$ in the bound was an upper bound for $e(H)$ —the points of $S(H) \setminus E(H)$ are already covered by the term $g(n-1)$. If $e(H) \leq 2^{s-1}$ then $|Z| \leq g(n)$ as desired. So, we may assume that $e(H) \geq 2^{s-1} + 1$.

First consider the case $s = n - 2$. We have $g(n-s) = g(2) = 1$, and $S(H)$ is a hyperplane in the unique K . Now if $S(K)$ has dimension $t (\leq s)$, then $S(H)$ and $S(K)$ have $2^{t-1} - 1$ points in common, $S(K)$ contributes at most 2^{t-1} points outside H to $|Z|$ and $e(H) \leq 2^s - 2^{t-1}$, proving the bound. So, we may assume $2 \leq s \leq n - 3$.

Let \mathcal{H} be the collection of all hyperplanes H such that $\dim S(H) = s$. All we have said about H above holds for all $H \in \mathcal{H}$. Make a directed graph with vertex set \mathcal{H} and arrows $H \rightarrow K$ when $S(H) \subseteq K$. (As we have seen, now $K \not\rightarrow H$.)

The outdegree of the graph on \mathcal{H} is $g(n-s)$ or $g(n-s) - 1$ at each vertex. It follows that the indegree at some vertex is at least $g(n-s) - 1$.

First suppose that the indegree of H is at least $g(n-s)$. The number of points P involved in flags (P, M) where $M \rightarrow H$ is bounded above by $g(n-1)$.

On the other hand it is bounded from below by $(g(n-s)-1)(2^{s-1}+1)+(2^s-1) > g(n-1)$ since $e(M) \geq 2^{s-1}+1$ for each such M . This is a contradiction. It follows that all outdegrees and all indegrees of \mathcal{H} are precisely $g(n-s)-1$.

Our estimate now becomes $|Z| \leq g(n-1)+(g(n-s)-1)2^{s-1}+2^{s-2}+e(H)$ and $|Z| \leq g(n)$ will follow if $e(H) \leq 3 \cdot 2^{s-2}$. So, we may assume that $e(H) \geq 3 \cdot 2^{s-2} + 1$ for all $H \in \mathcal{H}$. Again we bound the number of points P from below. We find the contradiction $(g(n-s)-2)(3 \cdot 2^{s-2}+1)+(2^s-1) > g(n-1)$ if $s \leq n-4$. So, we may assume that $s = n-3$.

Now $S(H)$ has codimension 2 in each K , so codimension at most 2 in each $S(K)$, so that $\dim(S(H) \cap S(K)) \geq s-2$. It follows that $e(H) \leq 3 \cdot 2^{s-2}$, as desired.

This finally proves part (ii) of the theorem. \square

3 Maximum number of points

Proof of Proposition 2. The inequality was shown already. So assume we have equality, i.e., $|Z| = 2^{n-1}-1$. Recall the above proof. Given a hyperplane H , consider the three types of flags: (i) flags (Q, K) with $Q \not\subseteq H$, (ii) flags (P, M) with $P \subseteq H$ and $M \neq H$, (iii) flags (P, H) . Correspondingly we get three contributions to $|Z|$: the points Q from flags of type (i), the points P from flags of type (ii), and the points P from flags (P, H) that were not counted yet, i.e., that occur in flags (P, H) only. (The set of such P is called $E(H)$, and has size $e(H)$.)

Let H be one of the hyperplanes with $\dim S(H) = s$ maximal. We find $|Z| \leq g(n-s)2^{s-1} + g(n-1) + e(H) \leq 2^{n-1} + 2^{s-1} - 2$. This estimate is precisely $2^{s-1} - 1$ too large, so it cannot be improved by 2^{s-1} . It follows that there are precisely $g(n-s)$ hyperplanes K on $S(H)$, and all except at most one have $\dim S(K) = s$. None of these K satisfies $S(K) \subseteq H$. It also follows that $e(H) \geq 2^{s-1}$. By Lemma 4 we may suppose that $1 \leq s \leq n-2$.

Lemma 7 *If $s = 1$, then $n \leq 4$.*

Proof. For $s = 1$ the counting $|Z| \leq g(n-s)2^{s-1} + g(n-1) + 2^s - 1$ holds with equality. This means that if $(P, H) \in C$ then P is the only point in H with this property, because $s = 1$, but also H is the only hyperplane on P with this property, because the counting is exact. Hence C involves equally many points as hyperplanes. Above we saw that if $S(H) \subseteq K$, then $S(K) \not\subseteq$

H , which in this case simply means that for two flags (P, H) and (Q, K) exactly one of $P \subseteq K$ and $Q \subseteq H$ holds. Now consider the point/hyperplane nonincidence matrix of $PG(n-1, 2)$. The 2-rank of this matrix is n , but the submatrix M corresponding to the points and hyperplanes in C has the property that $M+M^T = J-I$ of 2-rank $2^{n-1}-2$. It follows that $2n \geq 2^{n-1}-2$, so $n \leq 4$. \square

Make a directed graph Δ on the hyperplanes H occurring in flags $(P, H) \in C$ with $\dim S(H) = s$, writing an arrow $H \rightarrow K$ if $S(H) \subseteq K$.

Lemma 8 *The graph Δ is a tournament, and we have one of three cases, where $k = g(n-s) = 2^{n-s-1} - 1$ and v is the number of vertices:*

- a) *All indegrees and all outdegrees equal k and $v = 2k + 1$,*
- b) *All indegrees and all outdegrees equal $k - 1$ and $v = 2k - 1$,*
- c) *All indegrees are $k - 1$ or k (and both occur) and all outdegrees are $k - 1$ or k (and both occur) and $v = 2k$.*

Proof. We saw that at each vertex H the outdegree equals either k or $k-1$, where $k = g(n-s)$. If Δ has v vertices, then at each vertex the indegree equals $v-1-k$ or $v-k$. Since the average indegree equals the average outdegree, we have one of the stated cases. \square

Lemma 9 *If H has outdegree $k-1$, so that one hyperplane K on $S(H)$ has $\dim S(K) = t < s$, then $e(H) \geq 2^s - 2^{t-1} \geq 3 \cdot 2^{s-2}$.*

Proof. The contribution to $|Z|$ from hyperplanes K is now at most $a := (g(n-s)-1)2^{s-1} + 2^{t-1}$, so $e(H) \geq |Z| - g(n-1) - a = 2^s - 2^{t-1}$. \square

Lemma 10 *If $k = 1$, and H is a vertex of Δ with indegree 1, then $e(H) = 2^{s-1}$, and H also has outdegree 1.*

Proof. If H has indegree 1, say $M \rightarrow H$, then $S(M)$ is a hyperplane in H (since $s = n-2$) and covers at least $2^{s-1} - 1$ of the points of $S(H)$. It follows that $e(H) = 2^{s-1}$, and H also has outdegree 1 by Lemma 9. \square

Lemma 11 *If H is a vertex of Δ with indegree k , then each of its inneighbours has outdegree k . If $k > 1$ then $e(M) = 2^{s-1}$ for each inneighbour M , and all inneighbours have the same $2^{s-1} - 1$ nonunique points.*

Proof. Look at the coclique D in $\Gamma(H)$ consisting of the flags $(P, M \cap H)$ for $M \rightarrow H$. We have $|Z(D)| \leq g(n-1)$. Each M has $e(M) \geq 2^{s-1}$ unique points, and contributes $2^s - 1$ points altogether to $Z(D)$, so we find at least $k \cdot 2^{s-1} + (2^{s-1} - 1) = 2^{n-2} - 1 = g(n-1)$ points in $Z(D)$. Equality must hold, so if $k > 1$ all inneighbours M have $e(M) = 2^{s-1}$. By Lemma 9 they all have outdegree k . Obviously, if $k = 1$, an inneighbour cannot have outdegree $k - 1$. \square

Lemma 12 *If all indegrees and all outdegrees are k , then there is a subspace T of V of dimension $s - 1$ such that for each H with $\dim S(H) = s$ we have $S(H) \setminus E(H) = T$.*

Proof. The counting of the previous lemma shows for $k > 1$ that each inneighbour M of H has the same set $S(M) \setminus E(M)$ of size $2^{s-1} - 1$. Since this covers $S(H) \setminus E(H)$ which has the same size, we conclude that $S(M) \setminus E(M) = S(H) \setminus E(H)$ when there is an arrow $M \rightarrow H$. If $k = 1$, the same conclusion follows from Lemma 10.

This set of size $2^{s-1} - 1$ is the intersection of all $S(H)$, so is a subspace. \square

Lemma 13 *If all indegrees and all outdegrees are k , and $s > 1$, then C arises by the construction of Example 4.*

Proof. We have $v = 2k + 1 = 2^{n-s} - 1$ vertices H . Each vertex H contributes $e(H) = 2^{s-1}$ unique points to Z , and all vertices contribute the same set of $|T| = 2^{s-1} - 1$ common points, $2^{n-1} - 1 = |Z|$ points altogether. Now consider another flag $(P, H) \in C$ (with $\dim S(H) < s$). The point P cannot be one of the unique points, so $P \subseteq T$. This shows that C arises as in Example 4. \square

Lemma 14 *If $s = n - 2$ then C arises as in Example 1, 3 or 4.*

Proof. If $s = n - 2$, then $k = 1$. Now Δ has at most 3 vertices. If $v = 3$, then it is a directed 3-cycle. By Lemma 13, either C arises by the construction of Example 4, or $s = 1$, $n = 3$, in which case we have Example 1.

By Lemma 10, if $v \neq 3$, then $v = 1$. Let K be the unique hyperplane in $Z'(C)$ with $S(H) \subseteq K$, and suppose $\dim S(K) = t$. Then $1 \leq t < s$. If $t = 1$, then we have Example 3 (with $G = S(H)$ and $Q = S(K)$). Let $t > 1$.

The subspace $S(H)$ is a hyperplane in both H and K , so $S(H) = H \cap K$, and the $2^{t-1} - 1$ points in $H \cap S(K)$ are not unique in H . By Lemma 9 we have $e(H) = 2^{n-2} - 2^{t-1}$. The situation is tight again, so the flags (P, M) contribute precisely $2^{n-2} - 1$ points. Put $T := S(K) \cap S(H)$, so that $\dim T = t - 1$. Since $t > 1$, $T \neq 0$. Compare (P, M) and (Q, K) . If $S(K) \not\subseteq M$, then $S(M) \subseteq K \cap H = S(H)$. Since $E(H) = S(H) \setminus T$, this means that $T \subseteq M$ or $S(M) \subseteq T$. This is Example 4 (applied to a coclique from Example 3). \square

Now assume $1 \leq s \leq n - 3$, so that $k \geq 3$.

Lemma 15 *Case c) of Lemma 8 does not occur.*

Proof. In case c), suppose that we have $v = 2k$ vertices, b of which have outdegree $k - 1$ (and indegree k) and $v - b$ of which have outdegree k (and indegree $k - 1$). The total number of arrows is $vk - b = vk - (v - b)$, so that $b = v/2 = k$. Let A be the set of vertices with outdegree k , B that with outdegree $k - 1$. By Lemma 11 we have all arrows from A to B , and this fills up all outarrows in A . Then there are no arrows inside A , contradiction. So case c) does not occur. \square

Lemma 16 *Case b) of Lemma 8 does not occur.*

Proof. In case b) we have $e(H) \geq 3 \cdot 2^{s-2}$ for each vertex H of Δ , by Lemma 9. The $k - 1$ inneighbours of any vertex H contribute at least $3 \cdot 2^{s-2}(k - 1) + 2^{s-2} - 1 = 3 \cdot 2^{n-3} - 5 \cdot 2^{s-2} - 1$ points in H , so $3 \cdot 2^{n-3} - 5 \cdot 2^{s-2} - 1 \leq 2^{n-2} - 1$, that is, $2^{n-3} \leq 5 \cdot 2^{s-2}$ and hence $s \geq n - 3$. Since we are assuming $s < n - 2$ this means $s = n - 3$. Now $k = 3$ and all vertices have indegree and outdegree 2. The graph Δ has 5 vertices. They can be labeled H_i , $i = 0, 1, 2, 3, 4$, such that $H_i \rightarrow H_j$ iff $j = i + 1$ or $j = i + 2 \pmod{5}$. Now H_2 contains S_{H_1} and S_{H_0} , both of dimension $s = n - 3$. Since H_1 contributes at least 2^{s-1} points Q , we have $S_{H_1} \neq S_{H_0}$. So $S_{H_0} + S_{H_1}$ is an $(n - 2)$ -space contained in the $(n - 2)$ -space $H_1 \cap H_2$. Hence $S_{H_0} + S_{H_1} = H_1 \cap H_2$. Since $S_{H_1} \not\subseteq H_0$, also $S_{H_0} = H_0 \cap H_1 \cap H_2$. Now S_{H_0} and S_{H_1} meet in an $(n - 4)$ -space, so at least $2^{n-4} - 1$ points are not unique in H_0 . The standard counting gives $|Z| \leq (2^{n-2} - 1) + (2 \cdot 2^{s-1} + 2^{s-2}) + (2^s - 2^{s-1}) = 2^{n-1} - 1 - 2^{n-5}$, contradiction. So case b) does not occur. \square

So far we saw: either we have Example 1, 3 or 4, or $s = 1$, $n = 4$.

Lemma 17 *If $s = 1$, $n = 4$, then we have Example 2.*

Proof. Let $Z' = Z'(C)$. Here $|Z| = |Z'| = 7$, and Δ is a tournament on seven vertices with all in and outdegrees equal to 3. We first show that Z is a cap, that is, no three points in Z are collinear. Indeed, if $(P, H), (Q, K), (R, L) \in C$, and P, Q, R are collinear, and $K \rightarrow H$, then $Q \subset H$ but also $P \subset H$, hence $R \subset H$, that is $L \rightarrow H$. But now there can be no consistent arrow between K and L . So the points form a cap of size 7. The only maximal caps of $PG(3, 2)$ are the elliptic quadric (of size 5) and the complement of a plane (of size 8), cf. [1, Theorem 18.2.1], so there is a (hyper)plane π disjoint from Z and a point A not in Z and not in π . If $(P, H), (Q, K) \in C$ and $H \cap K$ is a line in π , then $(P, H), (Q, K)$ are adjacent, impossible. So the seven planes in Z' meet π in seven different lines. The situation is self-dual, so there is a point not on any plane in Z' , necessarily the point A . So, the seven points of Z are the points different from A outside π , and the seven planes of Z' are the planes different from π not on A . One now quickly establishes that we have Example 2. \square

This finishes the proof of Proposition 2. \square

Remarks. The construction of Example 4 does not change $n - s$. The hyperplane has $n - s = 1$, in Example 3 we have $n - s = 2$ and in Example 2 we have $n - s = 3$. So, all examples of $|Z| = g(n)$ have $1 \leq n - s \leq 3$.

Example 3 needs as ingredient a hyperplane disjoint from $Z(D)$. It follows that $Z(D)$ cannot contain linear subspaces of dimension larger than 1. Hence the construction of Example 3 applies only for (i) the hyperplane example for $n = 2$, (ii) Example 1, (iii) Example 2. The resulting examples are Example 1, and two further examples with $n = 4$, $s = 2$ and $n = 5$, $s = 3$.

4 Acknowledgement

The authors thank Jan Draisma for the rank-1 matrix point of view.

References

- [1] J. W. P. Hirschfeld, *Finite projective spaces of three dimensions*, Oxford Univ. Press, 1985.

- [2] T. Mussche, *Extremal combinatorics in generalized Kneser graphs*, Ph.D. Thesis, Eindhoven University of Technology, 2009.

Address of the authors:

Department of Mathematics,
Eindhoven University of Technology,
P.O. Box 513, 5600 MB Eindhoven,
The Netherlands.

e-mail: aartb@win.tue.nl, aeb@cwil.nl, cicoguv@gmail.com