

CODING OF GEODESICS AND LORENZ-LIKE TEMPLATES FOR SOME GEODESIC FLOWS

PIERRE DEHORNOY AND TALI PINSKY

ABSTRACT. We construct a template with two ribbons that describes the topology of all periodic orbits of the geodesic flow on the unit tangent bundle to any sphere with three cone points with hyperbolic metric. The construction relies on the existence of a particular coding with two letters for the geodesics on these orbifolds.

1. INTRODUCTION

For p, q, r three positive integers— r being possibly infinite—satisfying $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$, we consider the associated hyperbolic triangle and the associated orientation preserving Fuchsian group $G_{p,q,r}$. The quotient $\mathbb{H}^2/G_{p,q,r}$ is a sphere with three cone points of angles $\frac{2\pi}{p}, \frac{2\pi}{q}, \frac{2\pi}{r}$ obtained by gluing two triangles. The unit tangent bundle $T^1\mathbb{H}^2/G_{p,q,r}$ is a 3-manifold that is a Seifert fibered space. It naturally supports a flow whose orbits are lifts of geodesics on $\mathbb{H}^2/G_{p,q,r}$. It is called the *geodesic flow on $T^1\mathbb{H}^2/G_{p,q,r}$* and is denoted by $\varphi_{p,q,r}$. These flows are of Anosov type [Ano67] and, as such, are important for at least two reasons: they are among the simplest chaotic systems [Had1898] and they are fundamental objects in 3-dimensional topology [Thu88]. Each of these flows has infinitely many periodic orbits, which are all pairwise non-isotopic. The study of the topology of these periodic orbits began with David Fried who showed that many collections of such periodic orbits form fibered links [Fri83]. More recently Étienne Ghys gave a complete description in the particular case of the modular surface—which corresponds to $p = 2, q = 3, r = \infty$ —by showing that the periodic orbits are in one-to-one correspondence with periodic orbits of the Lorenz template [Ghy07].

The goal of this paper is to extend this result by giving an explicit description of the isotopy classes of all periodic orbits of $\varphi_{p,q,r}$ for every p, q, r (with p, q finite). A *template* [BW83] is an embedded branched surface made of several ribbons and equipped with a semi-flow. A template is characterized by its embedding in the ambient manifold and by the way its ribbons are glued, namely by the *kneading sequences* that describe the left- and rightmost orbit of every ribbon [HS90, dMvS93]. Call $\mathcal{T}_{p,q,r}$ the template with two ribbons whose embedding in $T^1\mathbb{H}^2/G_{p,q,r}$ is depicted on Figure 1 and whose kneading sequences are the words u_L, u_R, v_L, v_R given by Table 1, where the letter a corresponds to travelling along the left ribbon and b to travelling along the right ribbon. If r is infinite, the template is open of both sides of every ribbon, so that the kneading sequences are not

Date: November 2014.

P. D. thanks Étienne Ghys, Jérôme Los, and Vincent Pit for interesting discussions related to templates and geodesic flows, and the Technion for inviting him to Haifa in 2011 where this project was initiated. We thank the anonymous referee for several suggestions that improved the readability of the paper.

realized by any orbit of the template. If r is finite, both ribbons are closed on the left and open on the right, so that u_L and v_L are realized, but not u_R nor v_R .

Theorem A. *Up to one exception when r is finite, there is a one-to-one correspondance between periodic orbits of the geodesic flow $\varphi_{p,q,r}$ on $T^1\mathbb{H}^2/G_{p,q,r}$ and periodic orbits of the template $\mathcal{T}_{p,q,r}$ such that its restriction to every finite collection is an isotopy. The exception consists of the two orbits of $\mathcal{T}_{p,q,r}$ whose codes w_L and w_R are given by Table 2 and which actually correspond to the same orbit of $\varphi_{p,q,r}$.*




FIGURE 1. The template $\mathcal{T}_{p,q,r}$ in $T^1\mathbb{H}^2/G_{p,q,r}$. The 3-manifold $T^1\mathbb{H}^2/G_{p,q,r}$ is obtained from \mathbb{S}^3 by surgeries on a three component Hopf link (dotted) with the given indices. The template $\mathcal{T}_{p,q,r}$ is characterized by its embedding in $T^1\mathbb{H}^2/G_{p,q,r}$ and by the kneading sequences that describe the orbits of the extremities of the ribbons. When r is infinite, $T^1\mathbb{H}^2/G_{p,q,r}$ is the open manifold obtained by removing the component labelled $r - 1$ of the 3-component Hopf link. When $p = 2$, the exit side of the left ribbon is strictly included into the entrance side of the right ribbon.

In subsequent works, we intend to use the template $\mathcal{T}_{p,q,r}$ for deriving important properties of the geodesic flow $\varphi_{p,q,r}$, in particular the fact that all periodic orbits form prime knots, or the left-handedness (a notion introduced in [Ghy09]) of $\varphi_{p,q,r}$ (see [Deh15-2]).

The kneading sequences (Table 1) may look complicated. In a sense, it is the cost for having a template with two ribbons only. The proof suggests that, in the case of r finite, there are infinitely many other possible words, thus leading to other templates (the difference lying in the kneading sequences, not in the embedding). But these words are not simpler than the ones we propose here.

Theorem A is to be compared with previous works by Ghys [Ghy07], Pinsky [Pin11], and Dehornoy [Deh15]. The results of [Ghy07, Pin11] deal with orbifolds of type $\mathbb{H}^2/G_{2,3,\infty}$ and $\mathbb{H}^2/G_{2,q,\infty}$ for all $q \geq 3$ respectively. The existence of a cusp in these cases allows the obtained templates to have trivial kneading sequences of the form a^∞ and b^∞ . These

	u_L	v_R
r infinite	$(a^{p-1}b)^\infty$	$(b^{q-1}a)^\infty$
$p, q, r > 2$		
r odd	$((a^{p-1}b)^{\frac{r-3}{2}} a^{p-1} b^2)^\infty$	$((b^{q-1}a)^{\frac{r-3}{2}} b^{q-1} a^2)^\infty$
r even	$((a^{p-1}b)^{\frac{r-2}{2}} a^{p-2} (ba^{p-1})^{\frac{r-2}{2}} b^2)^\infty$	$((b^{q-1}a)^{\frac{r-2}{2}} b^{q-2} (ab^{q-1})^{\frac{r-2}{2}} a^2)^\infty$
$p = 2, q > 2, r > 4$		
r odd	$((ab)^{\frac{r-3}{2}} ab^2)^\infty$	$b^{q-1} ((ab^{q-1})^{\frac{r-5}{2}} ab^{q-2})^\infty$
r even	$((ab)^{\frac{r-4}{2}} ab^2)^\infty$	$b^{q-1} ((ab^{q-1})^{\frac{r-4}{2}} ab^{q-2})^\infty$

$$v_L := bu_L,$$

$$u_R := av_R$$

TABLE 1. The kneading sequences of the template $\mathcal{T}_{p,q,r}$.

$p, q, r > 2$	w_L	w_R
r odd	$((a^{p-1}b)^{\frac{r-3}{2}} a^{p-2} b)^\infty$	$((b^{q-1}a)^{\frac{r-3}{2}} b^{q-2} a)^\infty$
r even	$((a^{p-1}b)^{\frac{r-2}{2}} a^{p-2} b (a^{p-1}b)^{\frac{r-4}{2}} a^{p-2} b)^\infty$	$((b^{q-1}a)^{\frac{r-2}{2}} b^{q-2} a (b^{q-1}a)^{\frac{r-4}{2}} b^{q-2} a)^\infty$
$p = 2, q > 2, r > 4$		
r odd	$((ab)^{\frac{r-3}{2}} ab^2)^\infty$	$((ab^{q-1})^{\frac{r-5}{2}} ab^{q-2})^\infty$
r even	$((ab)^{\frac{r-4}{2}} ab^2)^\infty$	$((ab^{q-1})^{\frac{r-4}{2}} ab^{q-2})^\infty$

TABLE 2. The codes of the two periodic orbits of $\mathcal{T}_{p,q,r}$ that correspond to the same periodic orbit of the geodesic flow $\varphi_{p,q,r}$.

constructions can be recovered from Theorem A. The construction in [Deh15] deals with compact orbifolds, including the case of $\mathbb{H}^2/G_{p,q,r}$, but the constructed templates have numerous ribbons instead of two here, and the kneading sequences are not determined. Thus, Theorem A is a generalization of the results of [Ghy07, Pin11] to compact orbifold of type $\mathbb{H}^2/G_{p,q,r}$, as well as a simplification of the construction of [Deh15] with more precise information.

The main idea for the proof of Theorem A is to distort all geodesics in $\mathbb{H}^2/G_{p,q,r}$ onto an embedded graph that is just a bouquet of two oriented circles. We perform the distortion in an equivariant way in \mathbb{H}^2 (or in fact in $T^1\mathbb{H}^2$), and this allows one to deform through the cone points. We use this freedom to fix a direction in which the deformed geodesic is allowed to wind around a cone point, and this allows us to choose such a simple graph and a template with only two ribbons. In order to have a one-to-one correspondence between orbits of the flow and orbits of the template, the distortion has to obey some constraints. We introduce the notion of *spectacles*, which allows one to verify that we always get at most one coding for any semi-infinite geodesic and that the coding is nice. Ultimately we introduce what we call an *accurate pairs of spectacles*, that ensures we do get some coding for any bi-infinite geodesic. The main intermediate step is the following

result which might be of independent interest (see the introduction of Section 3 for the definitions of the graph $\widehat{\mathcal{G}}_{p,q,r}$, of the code, and of the supershift operator $\hat{\sigma}$)

Theorem B. *Assume that η, ξ in $\partial\mathbb{H}^2$ are the two extremities of a lift in \mathbb{H}^2 of a periodic geodesic in $\mathbb{H}^2/G_{p,q,r}$, then there exists a bi-infinite path γ in $\widehat{\mathcal{G}}_{p,q,r}$ that connects η to ξ and whose code $w(\gamma)$ satisfies $u_L \leq \hat{\sigma}^k(w(\gamma)) < u_R$ (resp. $v_L \leq \hat{\sigma}^k(w(\gamma)) < v_R$) for every shift $\hat{\sigma}^k(w(\gamma))$ that begins with a power of the letter a (resp. b). This path is unique, except for the paths encoded by the words w_L and w_R that have the same extremities.*

Theorem B is reminiscent of results by Caroline Series [Ser81, Thm 2.7] and Jérôme Los [Los13, Thm 4.4]. The novelty here is that we deal with monoids instead of groups, and that we cover some cases that were not addressed by Series (for small values of p, q , and r).

The article is organized as follows. First we recall in Section 2 the topology of the space $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$ by giving a surgery presentation on a link in \mathbb{S}^3 . We then describe in Section 3 a coding with two letters only for geodesics on $\mathbb{H}^2/G_{p,q,r}$. In Section 4, we use the coding for constructing a template with two ribbons in $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$ and describe its kneading sequences in terms of the chosen coding. We conclude the article with a few questions in Section 5.

In the whole text, p, q, r denote three natural numbers (with r possibly infinite) satisfying $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$. We denote by $A_0B_0C_0$ a fixed hyperbolic triangle in \mathbb{H}^2 with respective angles $\pi/p, \pi/q$, and π/r . We denote by $G_{p,q,r}$ the orientation-preserving Fuchsian group generated by the rotations of angles $2\pi/p, 2\pi/q$, and $2\pi/r$ around A_0, B_0 , and C_0 . The quotient $\mathbb{H}^2/G_{p,q,r}$ is then a 2-dimensional orbifold, namely a sphere with three cone points (or a cusp when r is infinite), that we call $A_\downarrow, B_\downarrow, C_\downarrow$. The letters A and B with sub/superscripts will always denote points of \mathbb{H}^2 in the $G_{p,q,r}$ -orbit of A_0 and B_0 respectively.

2. THE TOPOLOGY OF $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$

The unit tangent bundle $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$ is a Seifert fibered space whose regular fibers correspond to fibers of regular points of $\mathbb{H}^2/G_{p,q,r}$ and exceptional fibers to the fibers of the cone points $A_\downarrow, B_\downarrow, C_\downarrow$. The goal of this section (Proposition 2.4) is to give a surgery presentation of the 3-manifold $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$ with explicit coordinates. This has already been done, for example in [Mon87, p. 183]. The presentation we give here is slightly different, but more suited to the description of the template $\mathcal{T}_{p,q,r}$. Using Kirby calculus (see [Rol76, p. 267]), one can check that our presentation yields the same manifold as Montesinos'. In this section, the numbers p, q, r can be finite or infinite.

We begin with some elementary lemmas that help us fixing some notation. Assume that γ is an oriented closed curve immersed in a surface. Then the restriction $\mathbb{T}^1\gamma$ to the fibers of the points of γ of the unit tangent bundle to the surface is a 2-torus which supports four particular homology classes, represented by four vector fields:

1. the fiber f of a given point of γ , oriented trigonometrically,
2. an index 0 vector field z_γ ; for example, a vector field that is everywhere tangent to γ ,
3. an index +1 vector field u_γ^+ , that is, a vector field that rotates once to the left when traveling along γ ,

4. an index -1 vector field u_γ^- , that is, a vector field that rotates once to the right when traveling along γ .

Lemma 2.1. *For γ an embedded oriented curve, the classes $[f]$ and $[z_\gamma]$ form a basis of $T^1\gamma$. In this basis, one has $[u_\gamma^-] = -[f] + [z_\gamma]$ and $[u_\gamma^+] = [f] + [z_\gamma]$.*

Proof. The vector fields f and z_γ have exactly one point in common where they intersect transversally, so they form a basis of $H_1(T^1\gamma)$. Rotating along a fiber amounts to turning to the left, so one get $[u_\gamma^+] = [f] + [z_\gamma]$. Similarly one obtains $[u_\gamma^-] = -[f] + [z_\gamma]$. \square

Let P be an oriented pair of pants whose three oriented boundary components are denoted by $\gamma_A, \gamma_B, \gamma_C$. Consider the vector field v on P given by Figure 2 and denote by v_A, v_B, v_C its respective restrictions to $\gamma_A, \gamma_B, \gamma_C$. Since v_A is tangent to γ_A , one has $[v_A] = [z_A]$ on $T^1\gamma_A$. Similarly one has $[v_B] = [z_B]$ on $T^1\gamma_B$. Now v_C is not tangent to γ_C . Since it rotates plus one times when traveling along γ_C , one has $[v_C] = [u_C^+]$ on $T^1\gamma_C$.




FIGURE 2. A pair of pants P and the vector field v . The orientations of the boundary components induced by the orientation of P are with larger red arrows.

We denote by $\mathcal{C}^3 = U^A \cup U^B \cup U^C$ the 3-component link in \mathbb{S}^3 that is a chain of 3 unknots, with U^C in the middle (see Figure 3).

Lemma 2.2. *The unit tangent bundle to the pair of pants P is homeomorphic to $\mathbb{S}^3 \setminus \mathcal{C}^3$, where the respective meridians of the components U^A, U^B, U^C of \mathcal{C}^3 are represented by the vector fields $-z_A, -z_B, f_C$ and the respective longitudes by f_A, f_B, u_C^+ .*

Proof. Since P is an open surface, the unit tangent bundle T^1P is homeomorphic to the product $P \times \mathbb{S}^1$, that is, to $\mathbb{S}^3 \setminus \mathcal{C}^3$ (see Figure 3). The vector field v given by Figure 2 yields a section for this product whose boundary consists of two meridians of U^A and U^B respectively and one longitude of U^C . The respective longitudes of U^A, U^B and the meridian of U^C correspond to three fibers (as can be checked on Figure 3). \square

The above presentation of T^1P is not symmetric in the three boundary components. This symmetry can be recovered using a twist. Denote by $\mathcal{H}_+^3 = H^A \cup H^B \cup H^C$ the positive 3-component Hopf link in \mathbb{S}^3 (the green link in Figure 1).




FIGURE 3. The unit tangent bundle to a pair of pants T^1P seen as a product $P \times \mathbb{S}^1$, where the section corresponding to the vector field v on P of Figure 2 is shaded. The vector fields f (double arrow, green), z (orange) and u^+ (purple) for the three boundary tori $T^1\gamma_A, T^1\gamma_B$ and $T^1\gamma_C$ are represented. Next to every torus is represented the homology classes of these vector fields and of u^- in the {meridian, longitude}-basis.

Lemma 2.3. *The unit tangent bundle to the pair of pants P is homeomorphic to $\mathbb{S}^3 \setminus \mathcal{H}_+^3$, where the respective meridians of the components H^A, H^B, H^C of \mathcal{C}^3 are represented by the vector fields $-z_A, -z_B, -z_C$ and the respective longitudes by u_A^+, u_B^+, u_C^+ .*

Proof. Starting from the presentation $T^1P \simeq \mathbb{S}^3 \setminus \mathcal{C}^3$ of Lemma 2.2, we perform a positive twist τ^C on the component U^C , that is, we cut \mathbb{S}^3 along a disc bounded by U^C and we glue back after performing one full positive turn along this disc (see Figure 4). We denote by H^A, H^B, H^C the images of U^A, U^B, U^C under τ^C . The tori H^A, H^B as well as the fibers of the points of P all get linked plus one times. On these two tori, the twist performs a positive transvection, adding a meridian to the longitude. Since the meridian of U^A is represented by the vector field $-z^A$, the meridian of H^A is also represented by $-z^A$ and the longitude by $f^A - (-z^A) = u_+^A$. The same holds on H^B . On the torus U^C , the twist performs another transvection, adding a longitude to the meridian. Therefore the longitude of H^C is represented by u_+^C , as is the longitude of U^C . Its meridian is represented by $f^C - u_+^C = -z^C$. \square

In the above presentation of T^1P as $\mathbb{S}^3 \setminus \mathcal{H}_+^3$, the fibers of all points of P are fibers of the positive Hopf fibration (see [AGL08] for explanations and movies).

We can now give the desired presentation of $T^1\mathbb{H}^2/G_{p,q,r}$. Recall that a Dehn filling with slope a/b on a knot amounts to gluing in a solid torus so that a meridian of the solid torus intersects a longitudes and b meridians of the knot. With this convention, a Dehn filling with slope ∞ amounts to filling the removed knot back in. Here we add the extra-convention that in that case of infinite slope, we keep the knot removed.




FIGURE 4. The unit tangent bundle to a pair of pants T^1P as the complement of a Hopf link. It is obtained from the previous one by a twist along the shaded disc. The vector fields f (double arrow, green), z (orange) and u^+ (purple) for the three boundary tori $T^1\gamma_A, T^1\gamma_B$ and $T^1\gamma_C$. On all three tori are represented the homology classes of these vector fields and u^- , in the {meridian, longitude}-basis. All meridians are of type $-z$ and all longitudes of type u^+ .

Proposition 2.4. *The 3-manifold $T^1\mathbb{H}^2/G_{p,q,r}$ can be obtained from $S^3 \setminus \mathcal{H}_+^3$ by Dehn filling the three components of \mathcal{H}_+^3 with respective slopes $p-1, q-1$, and $r-1$. Moreover, the fibers of $T^1\mathbb{H}^2/G_{p,q,r}$ correspond to the fibers of the Hopf fibration of $S^3 \setminus \mathcal{H}_+^3$ and the exceptional fibers to the cores of the Dehn fillings.*

Proof. Since $\mathbb{H}^2/G_{\infty,\infty,\infty}$ is a pair of pants, Lemma 2.3 gives the result in this case.

We now suppose p finite (and q, r arbitrary). Let D_p be the quotient of a disc D , by a rotation of order p . The orbifold $\mathbb{H}^2/G_{p,q,r}$ is obtained from $\mathbb{H}^2/G_{\infty,q,r}$ by gluing T^1D_p into the cusp γ_A (corresponding to the fiber of A_\downarrow), formally by first removing a neighbourhood of γ_A .

The unit tangent bundle T^1D_p is a solid torus whose meridian disc corresponds to the image on D_p of a non-singular vector field on D (see Figure 5). Such a vector field intersects each non singular fiber p times and intersects a vector field tangent to ∂D_p once (Figure 5 (d, e)). Therefore after gluing, the meridian of T^1D_p is glued to a curve with coordinates $(p-1, \pm 1)$ in the {meridian, longitude}-basis of H_A . The sign can be fixed either by working out explicitly the action of S^1 as in [Mon87], or simply by checking that in the $p=1$ case the meridian is glued to the curve u^+ as depicted in Figure 5 (a), hence has coordinates $(0, 1)$. Therefore the gluing corresponds to a Dehn filling of slope $p-1$.

The cases of q and r are treated similarly. \square

Note that, if we had not perform the negative Dehn twist in Lemma 2.3, we would obtain the complement of a chain of three unknots instead of the Hopf link, as depicted on Figure 3. In this case, a similar statement to Proposition 2.4 holds by replacing the




FIGURE 5. (a) The unit tangent bundle $T^1 D$ to a disc D is a solid torus $D \times \mathbb{S}^1$. The fiber $T^1\{*\} = f_*$ of a point $*$ of ∂D is shown in green, a vector field $z_{\partial D}$ tangent to ∂D in orange and a meridian disc given by a constant vector field in blue. When traveling along ∂D with the orientation given by the exterior (hence in the clockwise direction), the blue vector field has index $+1$, that is, it rotates to the left. It is then isotopic to $u_{\partial D}^+$. (b) the action of an order 3 rotation on $T^1 D$: the vector field $z_{\partial D}$ is invariant, while the meridian disc (in blue) is rotated in the fiber direction, and the fiber f_* (in green) is rotated around D . (c) the quotient of $T^1 D$ by an order 3 rotation: a fundamental domain is given by a horizontal slice of the full torus made of those tangent vectors that point between 9 o'clock and 1 o'clock, say. (d) the same picture expanded in the fiber direction. The top and bottom faces are identified using an order 3 rotation. (e) after a twist of $-1/3$ of a turn, $T^1 D_3$ is a solid torus with the standard identification of the opposite faces of a cylinder. The meridian disc is still given by the blue vector field. It intersects the fiber of a point (in green) 3 times, and the vector field tangent to ∂D_3 (in orange) minus one times.

surgery coefficients by p, q , and $1-1/r$ respectively (again, the change of coefficients is coherent with Rolfsen's move [Rol76, p.267]).

3. CODING OF THE GEODESIC FLOW

The geodesic flow ϕ on $T^1 \mathbb{H}^2 \simeq \mathbb{H}^2 \times \mathbb{S}^1$ is defined in the following way: every unit tangent vector to \mathbb{H}^2 is of the form $(\gamma(0), \dot{\gamma}(0))$ where γ is a geodesic traveled at speed 1, we then set $\phi^t(\gamma(0), \dot{\gamma}(0)) = (\gamma(t), \dot{\gamma}(t))$. For G a Fuchsian group, this definition is G -equivariant, and the geodesic flow then projects on $T^1 \mathbb{H}^2 / G$.

These flows are the oldest known example of Anosov flows [Ano67]. Their hyperbolic character implies the existence of a Markov partition [Rat73], that is, of a decomposition of the flow into flow boxes whose entry and exit faces glue nicely. However, describing such

a coding for explicit groups (for example for surface or triangular groups) is not so easy. This problem has a long history, see among others [MH38, BS79, Ser81, AF91, CN08, Kat08, Pit11] and the references therein.

Here we adapt to our needs the coding given by Caroline Series [Ser81] and completed by Jérôme Los [Los13] (who covers some cases that were left out by Series). The main idea in these codings is to distort geodesics in \mathbb{H}^2 on a planar Cayley graph of G . Here we do not even need a Cayley graph, but only a planar graph that is G -invariant. In this context, all properties of Series-Los' coding still hold. Our present generalisation works for general Fuchsian groups, but since stating the construction in full generality would make the notations heavier, we only state it in the case we are interested in.

We start with the triangle $A_0B_0C_0$ in \mathbb{H}^2 and the group $G_{p,q,r}$ as before. We denote by $\widehat{\mathcal{G}}_{p,q,r}$ the embedded graph in \mathbb{H}^2 whose vertices are the images of A_0 and B_0 by $G_{p,q,r}$, and whose edges are the images of the segment (A_0B_0) by $G_{p,q,r}$ (see Figure 6). All vertices of $\widehat{\mathcal{G}}_{p,q,r}$ have degree p or q . All components of the complement $\mathbb{H}^2 \setminus \widehat{\mathcal{G}}_{p,q,r}$ are polygons with $2r$ vertices, half of which are in the $G_{p,q,r}$ -orbit of A_0 and half of which in the orbit of B_0 .

Definition 3.1 (see Figure 6). Assume that $\gamma := (B^0, A^1, B^2, \dots)$ is a semi-infinite path in $\widehat{\mathcal{G}}_{p,q,r}$. The *code* of γ is the semi-infinite word $w(\gamma)$ defined as $a^{k_1}b^{k_2}a^{k_3}b^{k_4} \dots$, where $+2\pi k_{2i+1}/p$ (*resp.* $-2\pi k_{2i}/q$) is the value of the angle $B^{2i} \widehat{A^{2i+1}} B^{2i+2}$ (*resp.* $A^{2i-1} \widehat{B^{2i}} A^{2i+1}$). The code of a path starting in the orbit of A_0 is defined similarly, but starts with a power of the letter b . The code of a bi-infinite path $(\dots, A^{-1}, B^0, A^1, B^2, \dots)$ is the bi-infinite word $\dots a^{k_{-1}}b^{k_0}a^{k_1}b^{k_2} \dots$ defined in the same way.

Beware the choice of the opposite orientations for a and b . We denote by \leq^{lex} and $<^{\text{lex}}$ the lexicographic ordering on words on the alphabet $\{a, b\}$. This choice ensures that the lexicographic ordering on codes and the clockwise ordering on $\partial\mathbb{H}^2$ coincide (Lemma 3.6). Note that two paths obtained one from the other by an element of $G_{p,q,r}$ have the same code. Following an edge corresponds to suppressing the first block of a or b in the code of a path. We then define the *super-shift* operator $\hat{\sigma}$ on infinite words that suppresses, not only the first letter of a word but, the first block of similar letters of a word.

It is easy to see that every geodesic in \mathbb{H}^2 is quasi-isometric to a path in $\widehat{\mathcal{G}}_{p,q,r}$, but the latter is not unique. The words $u_L, u_R, v_L, v_R, w_L, w_R$ being given by Table 1 and 2, the goal of this section is to prove Theorem B of the introduction which states that for (almost) every closed geodesic there is a way to choose a unique code.

3.1. Pairs of spectacles and admissible paths.

Definition 3.2. A *base pair of spectacles* is a pair of disjoint semi-open intervals $\{I_{A_0 \rightarrow B_0}, I_{B_0 \rightarrow A_0}\}$ in $\partial\mathbb{H}^2$, such that all geodesic rays starting at A_0 and reaching $I_{A_0 \rightarrow B_0}$ span an angle $2\pi/p$ from A_0 , and all geodesic rays starting at B_0 and reaching $I_{B_0 \rightarrow A_0}$ span an angle $2\pi/q$ from B_0 . For every oriented edge (A, B) of $\widehat{\mathcal{G}}_{p,q,r}$, the *associated spectacle* $I_{A \rightarrow B}$ is the image of $I_{A_0 \rightarrow B_0}$ by the unique element of $G_{p,q,r}$ that maps (A_0, B_0) onto (A, B) . The spectacles associated to an oriented edge (B, A) is defined similarly.

Remark 3.3. It will usually make sense to choose the base pair of spectacles so that the geodesic rays from A_0 to $I_{A_0 \rightarrow B_0}$ include the ray passing through B_0 , and this will be the




FIGURE 6. The graph $\widehat{\mathcal{G}}_{p,q,r}$ in the case $p = 3, q = 4, r = 5$. In bold is a path through the graph whose code starts with aba^2b^3 . A choice of a base pair of spectacles $P = \{I_{A_0 \rightarrow B_0}, I_{B_0 \rightarrow A_0}\} \subset \partial\mathbb{H}^2$ is also depicted.

case for our choice of spectacles in the next section. However, we do not require this in the definition.

Deciding whether the intervals are open on the left or on the right is not important, but in order to avoid heavy case specifications, we arbitrarily decide that all intervals on $\partial\mathbb{H}^2$ are open on the right and closed on the left.

Since $I_{A_0 \rightarrow B_0}$ spans an angle $2\pi/p$ from A_0 , the p spectacles associated to the p edges starting at a point in the orbit of A_0 tessellate $\partial\mathbb{H}^2$ (and so do the q spectacles associated to the edges starting at a point in the orbit of B_0), for any choice of base pair of spectacles.

Definition 3.4. Given a base pair of spectacles S , a semi-infinite or bi-infinite path γ in $\widehat{\mathcal{G}}_{p,q,r}$ is S -admissible if it has a limit in $\partial\mathbb{H}^2$, say ξ , and if ξ belongs to all spectacles associated to all oriented edges of γ .

Given a pair of spectacles S and a point ξ on $\partial\mathbb{H}^2$, it is not clear whether there exists an S -admissible semi-infinite path connecting A_0 (say) to ξ . What is easy to check is that if there is such an admissible path, it is unique.

Lemma 3.5. *Given a pair of spectacles S , for every A in $\widehat{\mathcal{G}}_{p,q,r}$ and every ξ in $\partial\mathbb{H}^2$, there exists at most one S -admissible path in $\widehat{\mathcal{G}}_{p,q,r}$ joining A to ξ .*

Proof. The path can be constructed inductively: if the path begins by $(A^0, B^1, \dots, A^{2i})$ (resp. $(A^0, B^1, \dots, B^{2i+1})$), there is a unique oriented edge starting at A^{2i} (resp. B^{2i+1}) whose associated spectacle contains ξ . This gives a unique choice for B^{2i+1} (resp. A^{2i+2}). Note that it is not clear that the path so defined converges to ξ . \square

A nice feature of the coding with two letters given by Definition 3.1 is that the natural cyclic ordering on admissible paths starting at a given point is reflected in the lexicographic order of the codes.

Lemma 3.6. *Let S be a base pair of spectacles. Assume that γ, γ' are two S -admissible paths in $\widehat{\mathcal{G}}_{p,q,r}$ that begin with the same edge. Then γ is to the left of γ' after they diverge if and only if $w(\gamma) <^{\text{lex}} w(\gamma')$ holds.*

Proof. The main observation is that, after taking any edge of $\widehat{\mathcal{G}}_{p,q,r}$, the cyclic ordering on the $p-1$ or $q-1$ possible next edges coincide with the lexicographic ordering. Indeed, since a corresponds to a rotation of angle $+\frac{2\pi}{p}$, a path whose code begins with ab is on the right of a path whose code begins with a^2b , and so on, and the leftmost paths have codes beginning with $a^{p-1}b$. Similarly, since b corresponds to a rotation of angle $-\frac{2\pi}{q}$, a path whose code begins with ba is on the left of a path whose code begins with b^2a , and so on. Therefore, two paths coincide as long as their codes coincide. As soon as they diverge, the leftmost has the smallest code. \square

3.2. Accurate pairs of spectacles and existence of admissible paths. We have seen how a choice of a base pair of spectacles yields a coding for every path in $\widehat{\mathcal{G}}_{p,q,r}$, and leads to the notion of admissible infinite path. Now we want to find spectacles that ensure that every bi-infinite geodesic in \mathbb{H}^2 can be shadowed by an admissible bi-infinite path. This is the notion of *accurate spectacles*.

Definition 3.7. A base pair of spectacles S is said to be *accurate* if for every A in \mathbb{H}^2 and for every η, ξ in $\partial\mathbb{H}^2$ there exist an S -admissible semi-infinite path connecting A to ξ and an S -admissible bi-infinite path connecting η to ξ . These paths are denoted by $\gamma_{A \rightarrow \xi}^S$ and $\gamma_{\eta \rightarrow \xi}^S$ respectively.

Let us see which codes are yielded by accurate pairs of spectacles.

Definition 3.8. Assume that $S = \{I_{A_0 \rightarrow B_0}, I_{B_0 \rightarrow A_0}\}$ is an accurate pair of spectacles. Denote by u_L^S and u_R^S the respective codes of the admissible paths connecting A_0 to the left and right extremities of $I_{A_0 \rightarrow B_0}$, and by v_L^S and v_R^S the codes of admissible the paths connecting B_0 to the two extremities of $I_{B_0 \rightarrow A_0}$. The sequences $u_L^S, u_R^S, v_L^S, v_R^S$ are called the *kneading sequences* associated to the pair of spectacles S .

Lemma 3.9. *Given an accurate pair of spectacles S , a semi-infinite word $w = b^{k_1} a^{k_2} b^{k_3} \dots$ is the code of an S -admissible path if and only if it satisfies $u_L^S \leq^{\text{lex}} \hat{\sigma}^{2i+1}(w) <^{\text{lex}} u_R^S$ and $v_L^S \leq^{\text{lex}} \hat{\sigma}^{2i}(w) <^{\text{lex}} v_R^S$ for every i .*

Proof. Let ξ be a point $\partial\mathbb{H}^2$. By definition of accurate spectacles, there exists an S -admissible path $\gamma := (A^0, B^1, A^2, B^3, \dots)$ that connects A_0 to ξ and by Lemma 3.5 this S -admissible path is unique. Denote its code by w . For every i the code of the path $(A^{2i}, B^{2i+1}, \dots)$ is the word $\hat{\sigma}^{2i}(w)$. Since ξ belongs to the interval $I_{A^{2i} \rightarrow B^{2i+1}}$, by Lemma 3.6, we have $v_L^S \leq^{\text{lex}} \hat{\sigma}^{2i}(w) <^{\text{lex}} v_R^S$. Similarly, since ξ belongs to the interval $I_{B^{2i+1} \rightarrow A^{2i+2}}$, we have $u_L^S \leq^{\text{lex}} \hat{\sigma}^{2i+1}(w) <^{\text{lex}} u_R^S$.

Conversely, if w satisfies the above constraints, denote the associated path by γ and the limit of γ in $\partial\mathbb{H}^2$ by ξ . Lemma 3.5 implies that ξ belongs to all spectacles $I_{A^{2i} \rightarrow B^{2i+1}}$ and $I_{B^{2i+1} \rightarrow A^{2i+2}}$. This means exactly that w is S -admissible. \square

By the same argument, we immediately get

Lemma 3.10. *Given an accurate pair of spectacles S , a bi-infinite word $w = \dots b^{k_{-1}} a^{k_0} b^{k_1} \dots$ is the code of an S -admissible path if and only if it satisfies $u_L^S \leq^{\text{lex}} \hat{\sigma}^{2i+1}(w) <^{\text{lex}} u_R^S$ and $v_L^S \leq^{\text{lex}} \hat{\sigma}^{2i}(w) <^{\text{lex}} v_R^S$ for every i .*

3.3. An explicit choice of spectacles. Accurate pairs of spectacles yield codes that are easy to describe. However, it is not obvious that there exist accurate pairs of spectacles. For example, if the spectacles $I_{A_0 \rightarrow B_0}$ do not include the endpoint of the geodesic ray from A_0 through B_0 , then an admissible path $\gamma_{A \rightarrow \xi}^S$ would veer away from ξ instead of converging toward ξ . As another example, note that when r is infinite, there exists a unique pair of spectacles. Indeed in this case the graph $\widehat{\mathcal{G}}_{p,q,r}$ is a tree, and the only possibility for the intervals $I_{A_0 \rightarrow B_0}$ and $I_{B_0 \rightarrow A_0}$ is to be the two intervals that connect the extremities of the two horoballs meeting on the edge $(A_0 B_0)$.

The goal of this section is to construct explicitly an accurate pair of spectacles in the case where r is finite.

For the rest of this section, assume $p > 2$. The case $p = 2$ will be treated in Section 3.5. Assume F is a region of the complement of $\widehat{\mathcal{G}}_{p,q,r}$ in \mathbb{H}^2 . Since F has $2r$ vertices, it makes sense to say that two edges are *opposite* in F : if there are $r - 1$ edges between them in both directions.

Definition 3.11 (see Figures 7 and 8). Assume that A, B are two adjacent vertices of $\widehat{\mathcal{G}}_{p,q,r}$. Then the associated *bigon* β_{AB} is defined as the infinite sequence (F^1, F^2, \dots) of faces of the complement of $\widehat{\mathcal{G}}_{p,q,r}$, where F^1 is the face on the left of the edge (AB) , the edge $F^1 \cap F^2$ is opposite to (AB) in F^1 , and $F^i \cap F^{i+1}$ is opposite to $F^{i-1} \cap F^i$ in F^i for every $i \geq 2$.

The bigon β_{AB} has a left and a right boundary starting at A and B respectively. They both converge to the same point in $\partial\mathbb{H}^2$ that we call the *normal extremity* of the oriented edge (AB) and denote by ξ_{AB} .

Lemma 3.12. *The codes of the paths that follow the left and right boundaries of β_{AB} are the words v_R and u_L given by Table 1.*




FIGURE 7. The pair of spectacles S^f

Proof. We have to draw the corresponding bigon, and to describe its borders. This is depicted in Figure 8. The code is periodic, the period corresponding to one face in the case of r odd and two faces in the case of r even. \square

Let $S^f = (I_{A_0 \rightarrow B_0}^f, I_{B_0 \rightarrow A_0}^f)$ be the pair of spectacles such that the left extremity of $I_{A_0 \rightarrow B_0}^f$ and the right extremity of $I_{B_0 \rightarrow A_0}^f$ both coincide with the point $\xi_{A_0 B_0}$ (see Figure 7).

Lemma 3.13. *For every A in $\widehat{\mathcal{G}}_{p,q,r}$ and ξ on $\partial\mathbb{H}^2$, the path $\gamma_{A \rightarrow \xi}^{S^f}$ converges to ξ .*

Proof. Let (A^0, B^1, A^2, \dots) denote the consecutive vertices visited by $\gamma_{A \rightarrow \xi}^{S^f}$ and b_ξ be a Busemann function [Bus85] on \mathbb{H}^2 associated to ξ . We claim that b_ξ is decreasing when evaluated along every second point of $\gamma_{A \rightarrow \xi}^{S^f}$, that is, $b_\xi(A^0) > b_\xi(A^2) > b_\xi(A^4) > \dots$ holds. Indeed, one checks (see Figure 9) that for every i the spectacles $I_{B^{2i+1} \rightarrow A^{2i+2}}$ are included inside the interval of $\partial\mathbb{H}^2$ consisting of those directions that are closer to A^{2i+2} than to A^{2i} . \square




FIGURE 8. The code u_L of the right border of on infinite bigon is the periodic word $((a^{p-1}b)^{\frac{r-3}{2}}a^{p-1}b^2)^\infty$ when r is odd (on the left with $r = 5$), and $((a^{p-1}b)^{\frac{r-2}{2}}a^{p-2}(ba^{p-1})^{\frac{r-2}{2}}b^2)^\infty$ when r is even (on the right with $r = 6$).




FIGURE 9. For every ξ in the interval $I_{A_0 \to B_0} \cap I_{B_0 \to A_1}$ (approximately the interval between 12:00 and 1:30), the Busemann function b_ξ is smaller at A_1 than at A_0 : indeed $I_{A_0 \to B_0} \cap I_{B_0 \to A_1}$ is included in the half space defined by the perpendicular bisector of (A_0A_1) (dashed) and containing A_1 . Similarly for A_2 (1:30 to 3:30) and A_3 (3:30 to 6:00).

Now we have to see why for every η, ξ on $\partial\mathbb{H}^2$ there exists an admissible path that connects η to ξ .

Lemma 3.14. *The pair of spectacles S^f is accurate.*

Proof. We have to show that for every η, ξ on $\partial\mathbb{H}^2$ there exists an S^f -admissible path connecting η to ξ . Consider a sequence $(A_n)_{n \in \mathbb{N}}$ of points in the orbit of A_0 that converges to η and stays at a bounded distance from the hyperbolic geodesic connecting A_0 to η . By Lemma 3.13 all paths $\gamma_{A_n \rightarrow \xi}^{S^f}$ converge to ξ . Also every path $\gamma_{A_n \rightarrow \xi}^{S^f}$ is at a bounded distance from the geodesic connecting A_n to ξ . Therefore, every path $\gamma_{A_n \rightarrow \xi}^{S^f}$ is at a bounded distance from the geodesic that connects η to ξ . Since the orbit of A_0 contains only finitely many points in every ball of bounded radius, this implies that infinitely many paths $\gamma_{A_n \rightarrow \xi}^{S^f}$ go through the same point, hence ultimately coincide. By extraction, this yields an S^f -admissible path connecting η to ξ . \square

3.4. Uniqueness of the coding. The last missing point for proving Theorem B in the case $p > 2$ is to see when admissible paths fail to be unique. Denote the word $(a^{p-1}b)^{\frac{r-3}{2}}a^{p-2}b$ when r is odd and $(a^{p-1}b)^{\frac{r-2}{2}}a^{p-2}b(a^{p-1}b)^{\frac{r-4}{2}}a^{p-2}b$ when r is even by x_L , and the word $(b^{q-1}a)^{\frac{r-3}{2}}b^{q-2}a$ when r is odd and $(b^{q-1}a)^{\frac{r-2}{2}}b^{q-2}a(b^{q-1}a)^{\frac{r-4}{2}}b^{q-2}a$ when r is even by x_R . We then have (see Table 2) $w_L = x_L^\infty$ and $w_R = x_R^\infty$.

Lemma 3.15. *Assume r finite. Suppose that γ_1, γ_2 are two S^f -admissible paths connecting the same points on $\partial\mathbb{H}^2$. Then either γ_1, γ_2 coincide in a neighbourhood of ξ in which case their codes are of the form $(\dots x_L x_L x_L)w_1$ and $(\dots x_R x_R x_R)w_2$ where w_1 and w_2 are semi-infinite words that ultimately coincide, or γ, γ' are disjoint in which case their codes are x_L^∞ and x_R^∞ .*

Proof. First suppose that γ_1, γ_2 have no point in common. Since γ_1, γ_2 have the same extremities on $\partial\mathbb{H}^2$, they are separated by an infinite strip of faces of $\mathbb{H}^2 \setminus \widehat{\mathcal{G}}_{p,q,r}$, namely there exists faces $\dots, F^{-1}, F^0, F^1, \dots$, such that F^{i+1} is adjacent to F^i , that γ_1 is on the left of $\dots \cup F^{-1} \cup F^0 \cup F^1 \cup \dots$ and γ_2 on the right. Note that the width of the strip cannot exceed one face. Indeed, since all faces are $2r$ -gons with $r \geq 4$ whose vertices have angle at least $\pi/3$, one checks that in order to converge to the same point at infinity, one of the paths would have to use more than half of the sides of a face. This would contradict the admissibility of the path: indeed the code of such a path would contain at least $r/2$ consecutive blocks of the form $a^{p-1}b$ (or ab^{q-1}), and this is prohibited by the kneading sequences of Table 1.

Now, for every i , the intersection $F^i \cap F^{i+1}$ contains two vertices of $\widehat{\mathcal{G}}_{p,q,r}$ that we denote by A^i and B^i . Then γ_1 and γ_2 are two S^f -admissible paths connecting B^i and A^i to ξ . Since γ_1 does not go through A^i and γ_2 not through B^i , the point ξ belongs to $\partial\mathbb{H}^2 \setminus (I_{A^i \rightarrow B^i} \cup I_{B^i \rightarrow A^i})$ (the bottom gray interval on Figure 7) for every i . This forces $\gamma_1 \cap F^i \cap F^{i+1}$ to be in the orbit of A for every i and $\gamma_2 \cap F^i \cap F^{i+1}$ to be in the orbit of B .

Assume first r is odd. The previous remark implies that γ_1 and γ_2 both travel along $r-1$ sides of F^i , so that the faces F^{i-1} and F^{i+1} are opposite with respect to F^i . Therefore, the codes of γ_1 and γ_2 are x_L^∞ and x_R^∞ .

Next, assume r is even. The remark implies that γ_1 and γ_2 travel one along $r-2$ sides of F^i and the other along r sides. Similarly, in the face F^{i+1} , the two paths travel along $r-2$ and r sides of F^{i+1} . If the same path, say γ_1 , travels along r sides of F^i and F^{i+1} , then ξ actually belongs to the interval $I_{A^i \rightarrow B^i}$ so that γ_1 should have visited B^i , a

contradiction. Therefore γ_1 must travel along r sides of F^i and $r - 2$ of F^{i+1} , while γ_2 travels along $r - 2$ sides of F^i and r sides of F^{i+1} . By induction, the codes of γ_1 and γ_2 are x_L^∞ and x_R^∞ .

Now if γ_1, γ_2 have one point in common, by Lemma 3.5, they coincide after that point. By the same argument as above, the part between them is a semi-infinite chain of $2r$ -gons. The only possibility for such a chain is to be of the above form. \square

We can now conclude.

Proof of Theorem B in the case $p > 2$. We consider the pair of spectacles S^f introduced in Section 3.3. By Lemma 3.14 the pair S^f is accurate, meaning that for every η, ξ on $\partial\mathbb{H}^2$, there exists an S^f -admissible path connecting η to ξ . By Lemma 3.10, accurate paths are exactly the paths whose code verify the given inequalities. Finally, by Lemma 3.15, if η, ξ are lifts of the extremities of a periodic geodesic this S^f -admissible path is unique, except in the case of the words w_L and w_R . \square

3.5. Case $p = 2$. All definitions and lemmas of Section 3.1 and 3.2 and still valid when $p = 2$. Since faces of $\mathbb{H}^2 \setminus \widehat{\mathcal{G}}_{p,q,r}$ now have r sides and not $2r$, the definition of bigons (Definition 3.11) and the description of their boundaries (Lemma 3.12) need to be adapted.

We still consider the faces as $2r$ -gons (with angles at some of the vertices equal to π) and describe the corresponding boundaries. Up to switching between q and r , we can suppose $r \geq 5$. The kneading sequences in this case are given by Table 1 as shown by Figure 10.




FIGURE 10. An infinite bigon in the case $p = 2, r = 7$ on the left. The code u_L of the right border is the periodic word $((ab)^{\frac{r-3}{2}} ab^2)^\infty$ when r is odd (on the picture with $r = 7$) and $((ab)^{\frac{r-4}{2}} ab^2)^\infty$ when r is even. The code v_R of the left border is $b^{q-1}((ab^{q-1})^{\frac{r-5}{2}} ab^{q-2})^\infty$ when r is odd and $b^{q-1}((ab^{q-1})^{\frac{r-4}{2}} ab^{q-2})^\infty$ when r is even. The case $p = 2, r = 5$ on the right: the term $(ab^{q-1})^{\frac{r-5}{2}}$ disappears in v_R .

The proofs of Lemmas 3.13 and 3.14 need no special adaptation. Finally the proof of Lemma 3.15 can be translated to this case. The only modification is that for every i the

intersection $F^i \cap F^{i+1}$ consists of two vertices B_1^i and B_2^i that are both adjacent to some point A^i , and thus the point ξ belongs to $\partial\mathbb{H}^2 \setminus (I_{A^i \rightarrow B_1^i} \cup I_{A^i \rightarrow B_2^i})$. If r is odd this forces γ_1 to travel along $\frac{r-3}{2}$ sides of F^i and γ_2 along $\frac{r-1}{2}$ sides (or $\frac{r-1}{2}$ and $\frac{r-3}{2}$ sides respectively). At the next face F^{i+1} , the same argument as in the proof of Lemma 3.15 proves that the numbers have to alternate. Therefore the codes of γ_1 and γ_2 are those given by Table 2. If r is even the same proof also gives the codes of Table 2.

Finally the proof of Theorem B is now exactly the same that in the case $p > 2$.

4. THE TEMPLATE $\mathcal{T}_{p,q,r}$

We now introduce the main character of this story: the template $\mathcal{T}_{p,q,r}$ in $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$. The basic idea is that Theorem B gives a canonical way to distort $\mathbb{T}^1\mathbb{H}^2/G_{p,q,r}$ onto the 2-complex that is made of the fibers of the points of $\widehat{\mathcal{G}}_{p,q,r}/G_{p,q,r}$. However, this 2-complex is not a template because it branches along two circles, namely the fibers of the two vertices, instead of along intervals. The trick for proving Theorem A is then to replace the vertices of $\widehat{\mathcal{G}}_{p,q,r}$ by *roundabouts*, thus obtaining a template that has the desired properties. More precisely, we distort any path in $\widehat{\mathcal{G}}_{p,q,r}$ in a neighbourhood of each vertex. The crucial point is that since we make the distortion in \mathbb{H}^2 , we can choose to push the path to one side of the vertex or the other as we wish. We are therefore able to arrange all paths that pass through a vertex so that they turn around the vertex in a fixed direction (counterclockwise for any image of A_0 , clockwise for any image of B_0). Thanks to the admissibility of the original path, the transformed path will rotate by an angle at most $\frac{p-1}{p}2\pi$ around A_0 , and at most $\frac{q-1}{q}2\pi$ around B_0 .


For A, B two adjacent vertices of $\widehat{\mathcal{G}}_{p,q,r}$, we denote by B^+ the image of B by a $+2\pi/p$ -rotation around A and by A^+ the image of A by a $-2\pi/q$ -rotation around B .

Definition 4.1 (see Figure 11). Let $\mathcal{B}_{p,q,r}$ be the embedded oriented graph in $\mathbb{H}^2/G_{p,q,r}$ which is a bouquet of two oriented circles, one winding once around A_\downarrow and one winding minus one times around B_\downarrow . Let $\widetilde{\mathcal{B}}_{p,q,r}$ denote the lift of $\mathcal{B}_{p,q,r}$ in \mathbb{H}^2 . Every lift of a circle is called a *roundabout*. A *switch* is the contact point of two adjacent roundabouts. For A, B two adjacent vertices of $\widehat{\mathcal{G}}_{p,q,r}$, we denote by c_{AB} the arc of the roundabout around A that connects the edge (AB) to (AB^+) and by c_{BA} the arc of the roundabout around B that connects the edge (AB) to (A^+B) .

Every vertex of $\widehat{\mathcal{G}}_{p,q,r}$ is canonically associated to a roundabout of $\widetilde{\mathcal{B}}_{p,q,r}$, and every edge $\widehat{\mathcal{G}}_{p,q,r}$ to a switch of $\widetilde{\mathcal{B}}_{p,q,r}$.

Definition 4.2 (see Figure 12). Assume that $\gamma = (\dots, A^{-1}, B^0, A^1, \dots)$ is a simple path in $\widehat{\mathcal{G}}_{p,q,r}$. The associated *tango path* $\widetilde{\gamma}$ is the path in $\widetilde{\mathcal{B}}_{p,q,r}$ that is obtained by rotating around the roundabouts associated to the vertices of γ and changing at the switches that correspond to the edges of γ .

Define $\widetilde{\mathcal{T}}_{p,q,r}$ as the branched surface embedded in $\mathbb{T}^1\mathbb{H}^2$ that is made of those vectors of the form (x, v) , where x is a point on $\widetilde{\mathcal{B}}_{p,q,r}$ that belongs to an arc of the form c_{AB} (*resp.* c_{BA}) and v is a tangent vector that points into $I_{B \rightarrow A}^f$ (*resp.* $I_{A \rightarrow B}^f$). The surface $\widetilde{\mathcal{T}}_{p,q,r}$

FIGURE 11. The graph $\widetilde{\mathcal{B}}_{p,q,r}$.

is a surface made of ribbons of the form $c_{AB} \times I_{B \rightarrow A}^f$ (resp. $c_{BA} \times I_{A \rightarrow B}^f$) that branches in the fibers of the switches of $\widetilde{\mathcal{B}}_{p,q,r}$. We equip $\widetilde{\mathcal{T}}_{p,q,r}$ with a semi-flow, denoted by $\widetilde{\tau}_{p,q,r}$, whose orbits are, on every ribbon, of the form $c_{AB} \times \{*\}$ (resp. $c_{BA} \times \{*\}$). Since we are interested in the topology of orbits and not in the time, the speed is not relevant, we can arbitrarily decide that $\widetilde{\tau}_{p,q,r}$ travels along each branch at unit speed. In order to say that $\widetilde{\mathcal{T}}_{p,q,r}$ is a template, we need to check that ribbons are glued nicely.

Lemma 4.3. *Every branching arc of the surface $\widetilde{\mathcal{T}}_{p,q,r}$ is of the form $M \times I$, where M is a switch of the graph $\widetilde{\mathcal{B}}_{p,q,r}$ and, denoting the vertices of $\widetilde{\mathcal{B}}_{p,q,r}$ that surround M by A, B , the arc I in $\partial\mathbb{H}^2$ is the segment $I_{A \rightarrow B}^f \cup I_{B \rightarrow A}^f$.*

Proof. In the fiber of M four branches meet, namely $c_{AB}, c_{BA}, c_{A_B^-}$, and $c_{B_{A^-}}$. Their visual intervals are respectively $I_{B \rightarrow A}^f, I_{A \rightarrow B}^f, I_{B^- \rightarrow A}^f$, and $I_{A^- \rightarrow B}^f$. Figure 11 shows that all these segments are included in $I_{A \rightarrow B}^f \cup I_{B \rightarrow A}^f$. \square

Lemma 4.3 ensures that $\widetilde{\mathcal{T}}_{p,q,r}$ is indeed a template: it is made of ribbons, its branching arcs are segments, each branching segment has two incoming ribbons ($c_{A_B^-} \times I_{B^- \rightarrow A}^f$ and




FIGURE 12. The broken line represents a bi-infinite path in $\widehat{\mathcal{G}}_{p,q,r}$. The associated tango path in $\widetilde{\mathcal{B}}_{p,q,r}$ is thickened. All circles correspond to roundabouts.

$c_{B_{A^-}} \times I_{A^- \rightarrow B}^f$) that overlap and two outgoing ribbons ($c_{A_B} \times I_{B \rightarrow A}^f$ and $c_{B_A} \times I_{A \rightarrow B}^f$) that do not overlap. Along the branching segments, the vector fields induced by the different ribbons coincide. By construction, $\widetilde{\mathcal{T}}_{p,q,r}$ is $G_{p,q,r}$ -equivariant, so we can mod out.

Definition 4.4. The template $\mathcal{T}_{p,q,r}$ in $T^1\mathbb{H}^2/G_{p,q,r}$ is defined as the quotient $\widetilde{\mathcal{T}}_{p,q,r}/G_{p,q,r}$. It is equipped with the semi-flow $\tau_{p,q,r}$ that is the projection of $\widetilde{\tau}_{p,q,r}$.

We can now conclude.

Proof of Theorem A. First we check that $\mathcal{T}_{p,q,r}$ is embedded in the way described by Figure 1. Indeed, contracting the ribbons of $\widetilde{\mathcal{T}}_{p,q,r}$ into arcs bring $\mathcal{T}_{p,q,r}$ in a neighbourhood of the vector field v describes on Figure 2, and the arcs c_{A_B} , c_{B_A} wind once around the corresponding points A_\downarrow and B_\downarrow . For the framing of the ribbons of $\mathcal{T}_{p,q,r}$, notice that it is given by the direction of the fibers of $T^1\mathbb{H}^2/G_{p,q,r}$. The framing of Figure 1 is obtained by a rotation of 90 degrees, that is, by an isotopy. This proves that the template $\mathcal{T}_{p,q,r}$ is indeed embedded as in Figure 1.

Now, all we have to do is to describe a map, say $d_{p,q,r}$, from $T^1\mathbb{H}^2/G_{p,q,r}$ to $\mathcal{T}_{p,q,r}$ that will transport the orbits of the geodesic flow $\varphi_{p,q,r}$ onto the orbits of the semi-flow $\tau_{p,q,r}$ on the template, and check that its restriction to finite collection of periodic orbits is a topological equivalence and an isotopy. Actually, it is easier to do that in a $G_{p,q,r}$ -equivariant way in $T^1\mathbb{H}^2$.

Assume that η, ξ are two points in $\partial\mathbb{H}^2$ and denote by $g_{\eta \rightarrow \xi}$ the geodesics in \mathbb{H}^2 that connects them. By Theorem B, there also exists a unique admissible path $\gamma_{\eta \rightarrow \xi}^f$ in $\widehat{\mathcal{G}}_{p,q,r}$ that connects η and ξ . We denote it by $\widetilde{\gamma}_{\eta \rightarrow \xi}$ the associated tango path in $\widetilde{\mathcal{B}}_{p,q,r}$. The lift

of $g_{\eta \rightarrow \xi}$ in $T^1\mathbb{H}^2$ is the orbit $g_{\eta \rightarrow \xi} \times \{\xi\}$ of the geodesic flow $\varphi_{p,q,r}$. We now define $\widetilde{d}_{p,q,r}$ as the map that takes $g_{\eta \rightarrow \xi} \times \{\xi\}$ onto $\widetilde{\gamma}_{\eta \rightarrow \xi} \times \{\xi\}$, which is an orbit of $\widetilde{\tau}_{p,q,r}$. Of course, this can be done using an isotopy in the level $\mathbb{H}^2 \times \{\xi\}$ and in a $G_{p,q,r}$ -equivariant way. By Theorem B, the image of the map $\widetilde{d}_{p,q,r}$ is exactly the template $\widetilde{\mathcal{T}}_{p,q,r}$. Modding out the map $\widetilde{d}_{p,q,r}$ by $G_{p,q,r}$, we obtain the desired map $d_{p,q,r}$.

In order to check that we have an isotopy when we restrict to finite collections of periodic orbits, assume that γ, γ' are two arbitrary lifts in $T^1\mathbb{H}^2$ of two periodic orbits of $\varphi_{p,q,r}$. Then γ, γ' cannot point in the same direction in $\partial\mathbb{H}^2$, for otherwise they would become arbitrarily close, which is not possible for two periodic orbits. Therefore, the deformation $\widetilde{d}_{p,q,r}$ can be realized for these two orbits by two isotopies that live in different levels of $T^1\mathbb{H}^2$. These can be extended to a global isotopy of $T^1\mathbb{H}^2$. The same idea works for arbitrary (but finitely) many periodic orbits. \square

5. CONCLUDING REMARKS

5.1. Templates with two ribbons. In this article, we have given a construction of a template for some particular geodesic flows. One can wonder about the optimality of the result, that is, whether it can be extended to other geodesic flows on other surfaces or 2-dimensional orbifolds. It was proven in [BW83] that in this general case, there exists always a template, but with no control on the number of ribbons and the way they are embedded. Explicit constructions were given in [Deh15], but they always yield templates with more than two ribbons. Actually we have

Proposition 5.1. *Assume that G is a Fuchsian group such that \mathbb{H}^2/G is not a sphere with three cone points. Then there is no template with two ribbons that describes the isotopy classes of all periodic orbits of the geodesic flow on $T^1\mathbb{H}^2/G$.*

Proof. The set of isotopy classes that can be represented by a template with two ribbons is a submonoid with two generators of $\pi_1(T^1\mathbb{H}^2/G)$. Since the fundamental groups of all Fuchsian groups different from triangles groups have rank larger than 2, they cannot equal a monoid with two generators. \square

The natural task is then to wonder what are the next simplest templates and which groups they represent.

5.2. Space of codings. In Section 3 we have constructed a particular coding, relying on a particular pair of accurate spectacles. It would be interesting to understand which codings can be obtained in this way.

Question 5.2. Given p, q, r , what is the set of accurate spectacles for coding the periodic geodesics of $\mathbb{H}^2/G_{p,q,r}$?

More generally, given a generating set of $G_{p,q,r}$, a coding of periodic geodesics on $\mathbb{H}^2/G_{p,q,r}$ is a language. What we did is to describe one particular such language. Another example is given in [Pfe08] (it is not clear to us whether this coding can be obtained using an accurate pair of spectacles).

Question 5.3. Given a generating set of $G_{p,q,r}$, what is the set of those languages that encode the periodic geodesics of $\mathbb{H}^2/G_{p,q,r}$?

REFERENCES

- [AF91] ROY ADLER AND LEONARD FLATTO, Geodesic flows, interval maps, and symbolic dynamics, *Bull. Amer. Math. Soc.* **25** (1991), 229–334.
- [AGL08] AURÉLIEN ALVAREZ, ÉTIENNE GHYS, AND JOS LEYS, Dimensions, chapter 8 (2008). http://www.dimensions-math.org/Dim_CH7_E.htm
- [Ano67] DMITRI V. ANOSOV, Geodesic flows on closed Riemannian manifolds with negative curvature, *Proc. Steklov Inst. Math.* **90** (1967), 235pp.
- [BW83] JOAN S. BIRMAN AND ROBERT F. WILLIAMS, Knotted periodic orbits in dynamical systems—I: Lorenz’s Equations, *Topology* **22** (1983), 47–82. (erratum at www.math.columbia.edu/~jwb/bw-KP0-I-erratum.pdf)
- [BS79] RUFUS BOWEN AND CAROLINE SERIES, Markov maps associated with fuchsian groups, *Publ. Math. Inst. Hautes Études Sci.* **50** (1979), 153–170.
- [Bus85] HERBERT BUSEMANN, *The geometry of geodesics* (1985), Dover Books on Mathematics, Vol. 6.
- [CN08] ETHAN M. COVEN AND ZBIGNIEW H. NITECKI, On the genesis of symbolic dynamics as we know it, *Colloq. Math.* **110** (2008), 227–242.
- [dMvS93] WELLINGTON DE MELO AND SEBASTIAN VAN STRIEN, *One-Dimensional Dynamics* (1993), Springer, 605pp.
- [Deh15] PIERRE DEHORNOY, Geodesic flow, left-handedness, and templates, *Algebr. Geom. Topology* **15** (2015), 1525–1597. <http://arxiv.org/abs/1112.6296>
- [Deh15-2] PIERRE DEHORNOY, Which geodesic flows are left-handed ? <http://arxiv.org/abs/1501.02909>
- [Fri83] DAVID FRIED, Transitive Anosov flows and pseudo-Anosov maps, *Topology* **22** (1983), 299–303.
- [Ghy07] ÉTIENNE GHYS, Knots and dynamics, *Proc. Internat. Congress of Mathematicians I*, Eur. Math. Soc. (2007), 247–277.
- [Ghy09] ÉTIENNE GHYS, Right-handed vector fields and the Lorenz attractor, *Japan. J. Math.* **4** (2009), 47–61.
- [Had1898] JACQUES HADAMARD, Les surfaces à courbures opposées et leurs lignes géodésiques, *J. Math. Pures Appl.* **4** (1898), 27–74.
- [HS90] JOHN H. HUBBARD AND COLIN T. SPARROW, The classification of topologically expansive Lorenz maps, *Comm. Pure Appl. Math.* **43** (1990), 431–443.
- [Kat08] SVETLANA KATOK, Fuchsian groups, geodesic flows on surfaces of constant negative curvature and symbolic coding of geodesics, *Clay Math. Proc.* **10** (2008), 244–320.
- [Los13] JÉRÔME LOS, Volume entropy for surface groups via Bowen-Series like maps, to appear in *J. Topology*. <http://arxiv.org/abs/0908.3537>
- [Mon87] JOSÉ M. MONTESINOS-AMILIBIA, *Classical tessellations and three-manifolds*, Universitext, Springer (1987), 230pp.
- [MH38] MARSTEN MORSE AND GUSTAV A. HEDLUND, Symbolic Dynamics, *Amer. J. Math.* **60** (1938), 815–866.
- [Pfe08] MARKUS PFEIFFER, Automata and Growth Functions for the Triangle Groups, Diploma Thesis, Rheinisch-Westfälische Technische Hochschule Aachen. s
- [Pin11] TALÍ PINSKY, Templates for the geodesic flow, *Ergod. Theory Dynam. Systems* **34** (2014), 211–235. <http://arxiv.org/abs/1103.4499>
- [Pit11] VINCENT PIT, Codage du flot géodésique sur les surfaces hyperboliques de volume fini, PhD thesis, Bordeaux (2010).
- [Rat73] MARINA RATNER, Markov partitions for Anosov flows on n -dimensional manifolds, *Israel J. Math.* **15** (1973), 92–114.
- [Rol76] DALE ROLFSEN, *Knots and links*, Publish or Perish (1976), 439pp.
- [Ser81] CAROLINE SERIES, The infinite word problem and limit sets in Fuchsian groups, *Ergod. Theory Dynam. Systems* **1** (1981), 337–360.
- [Thu88] WILLIAM THURSTON, On the geometry and dynamics of diffeomorphisms of surfaces, *Bull. Amer. Math. Soc.* **19** (1988), 417–431.

UNIV. GRENOBLE ALPES, IF, F-38000 GRENOBLE, FRANCE; CNRS, IF, F-38000 GRENOBLE, FRANCE

E-mail address: pierre.dehornoy@ujf-grenoble.fr

URL: <http://www-fourier.ujf-grenoble.fr/~dehornop/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER BC, CANADA

E-mail address: tali@math.ubc.ca

URL: <http://www.math.ubc.ca/~tali/>