

TITLE (146 characters)

Cognition and Hippocampal Plasticity in the Mouse Is Altered by Monosomy of a Genomic Region Implicated in Down Syndrome

AUTHORS

Ignasi Sahún^{1,*}, Damien Marechal^{2,3,4,5*}, Patricia Lopes Pereira^{2,5}, Valérie Nalesso^{2,3,4,5}, Agnes Gruart⁶, José Maria Delgado Garcia⁶, Stylianos E. Antonarakis⁷, Mara Dierssen^{1,#} and Yann Hérault^{2,3,4,5,8,**}

Affiliations

¹Systems Biology Programme, Centre for Genomic Regulation (CRG); Universitat Pompeu Fabra (UPF); and Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Dr. Aiguader 88, E-08003 Barcelona, Spain.

²Institut de Génétique et de Biologie Moléculaire et Cellulaire, Illkirch, 1 rue Laurent Fries, 67404 Illkirch, France

³Centre National de la Recherche Scientifique, UMR7104, Illkirch, France

⁴Institut National de la Santé et de la Recherche Médicale, U964, Illkirch, France

⁵Université de Strasbourg, Illkirch, France.

⁶División de Neurociencias, Universidad Pablo de Olavide, Sevilla, Spain

⁷Department of Genetic Medicine and Development, University of Geneva Medical School, 1 Rue Michel-Servet, 1211 Geneva, Switzerland,

⁸Institut Clinique de la Souris, PHENOMIN, GIE CERBM, Illkirch, France

* Co-authorship

Joint leadership

** To whom correspondence should be addressed

Running head: gene dosage and cognition

[§] Present Adress : Transgenese et Archivage Animaux Modèles, TAAM, CNRS, UPS44, 3B rue de la Férollerie 45071 Orléans, France;

ABSTRACT

Down syndrome (DS) is due to increased copy number of human chromosome 21. The contribution of different genetic regions has been tested using mouse models. As shown previously, the *Abcg1-U2af1* genetic region contributes to cognitive defects in working and short-term recognition memory in Down syndrome mouse models. Here we analysed the impact of monosomy of the same genetic interval using a new mouse model, named Ms2Yah. We used several cognitive paradigms, and did not detect defects in the object recognition nor the Morris water maze tests. However, surprisingly, Ms2Yah mice displayed increased associative memory in a pure contextual fear conditioning test, and decreased social novelty interaction along with a larger long term potentiation recorded in the CA1 area following stimulation of Schaffer collaterals. Whole genome expression studies carried out on hippocampus showed that only the transcription of a small number of genes is affected, mainly from the genetic interval (*Cbs*, *Rsph1*, *Wdr4*), with a few additional ones, including the postsynaptic *Gabbr2*, *Gabbr1*, *Grid2p*, *Park2* and *Dlg1* and the components of the Ubiquitin mediated proteolysis (*Anapc1*, *Rnf7*, *Huwe1*, *Park2*). The *Abcg1-U2af1* region is undeniably encompassing dosage sensitive genes or elements whose change in copy number directly impact learning and memory, synaptic function and autistic related behavior.

Keywords

Aneuploidy, genetic dosage, mouse model, intellectual disabilities, trisomy 21

INTRODUCTION

Aneuploidy, defined as an abnormal number of copies of a genomic region, is a common cause of human genetic disorders associated with intellectual disabilities and autism (MEFFORD *et al.* 2012). Down syndrome (DS) is a chromosomal aneuploidy characterized by the presence of a supernumerary copy of the whole or part of human chromosome 21 (Hsa21) that gives rise to a syndromic intellectual disability. Although classically research has focused on the search of candidate genes, exciting new findings demonstrate that in addition to direct and indirect alteration of expression of Hsa21 and non-Hsa21 genes, alteration in copy number of functional, non-traditional genomic elements may impact the variability of the DS phenotype (LETOURNEAU *et al.* 2014) .

Several “dosage sensitive” regions, including genes and non-coding conserved elements, have been mapped across the length of Hsa21 and shown to be sufficient for induction of the complete phenotype of DS (KORBEL *et al.* 2009; LYLE *et al.* 2009). The results of such studies provide initial evidence supporting that some of the phenotypes of DS are not due to specific genes, but rather to the over/underexpression of a whole chromosomal domain or even to other lncRNA (UMLAUF *et al.* 2008). Until recently, the mouse models available have not explored the contribution of the gene-rich telomeric part of the Hsa21 that maps to the mouse chromosomes Mmu17, Mmu16 and Mmu10 and do not completely explain the complexity and variability of some phenotypes observed in DS patients. (REEVES *et al.* 1995; YU *et al.* 2010b; DUCHON *et al.* 2011). We previously demonstrated the contribution of trisomy of the *Abcg1–U2af1* genetic interval from the telomeric part of Hsa21 to DS, in a mouse model named Ts1Yah (LOPES PEREIRA *et al.* 2009). Ts1Yah mice

showed defects in novel object recognition, open-field and Y-maze tests, similar to other DS models, but the trisomy induced an improvement of the hippocampal-dependent spatial memory in the Morris water maze along with enhanced and longer lasting long-term potentiation *in vivo* in the hippocampus. Although DS induces several deleterious effects, Hsa21 also carries protective genes or regions, which may explain some “advantageous” phenotype (SUSSAN *et al.* 2008).

To analyze the dosage-sensitivity of this particular region, in the present study, we have generated the corresponding monosomy (Ms2Yah) model to the Ts1Yah and we have characterized various functional domains using behavioral and electrophysiological studies (LOPES PEREIRA *et al.* 2009). We used several cognitive paradigms (non-hippocampal dependent (object recognition, CS-freezing) and the hippocampal dependent (water maze, context freezing) tests) with locomotor (Rotarod) and social tests to evaluate defects observed in DS people. We expected to find an opposite phenotype to that observed in the Ts1Yah, but we could not detect alterations in the performance in neither the object recognition nor the Morris water maze test. However, surprisingly, monosomy of the *Abcg1-U2af1* genetic interval led to increased associative memory in a pure contextual fear conditioning test and altered social interaction along with a significant increase in the long term potentiation *in vivo*; to understand the possible underlying mechanisms in the hippocampus, we used a transcriptome approach and we observed a very light whole genome effect from the decrease in gene dosage associated with the *Abcg1-U2af1* region which highlights the contribution of genes from the interval to the learning and memory phenotypes. These changes were detected in the absence of abnormalities in the size or density of pyramidal neurons.

MATERIALS AND METHODS

Mouse line and Ethical Statement

The Ms2Yah mouse line carried a deletion of the region located downstream of *Abcg1* and upstream of *U2af1* and was bred on C57BL/6J for more than 10 generations (LOPES PEREIRA *et al.* 2009). Mice were handled with the agreement of the local ethical committee and in accordance with the European Council Guidelines for the Care and Use of Laboratory animals (accreditation 7320). They were housed under a 12h/12h light-dark cycle in ICS mouse facility in Illkirch (certificate C45-234-6) and shipped by authorized carriers to Barcelona for the behavioral assessment or to Sevilla for the *in vivo* LTP experiments. Mice were housed in standard cages with access to food and water ad libitum, under a controlled environment (temperature = 20 ± 1 °C; humidity = 60 %), and with a light/dark cycle of 12h (08:00 a.m. to 08:00 p.m.). All experimental protocols were performed under authorization (CEEA-PRBB; MDS-08-1060P1 and JMC-07-1001P1-MDS), and met the guidelines of the local (law 32/2007) and European regulations (EU directive 86/609, EU decree 2001-486) and Statement of Compliance with Standards for Use of Laboratory Animals by Foreign Institutions nº A5388-01, National Institutes of Health (USA). All investigators involved in animal experimentations have the appropriate training (Department de Medi Ambient i Habitatge, 214/1997). The CRG is authorized to work with genetically modified organisms (A/ES/05/I-13 and A/ES/05/14). All the behavioral testing was conducted in an isolated room by the same experimenter at the same time of the day (between 8:30 a.m. and 3:30 p.m.). The experimenter was blinded as to the genetic status of the animals. Dr Yann HERAULT, as the principal investigator in this study, was granted the accreditation 45-31 and 67-369 to perform the reported experiments with the collaborators Dr Dierssen and Dr Delgado-Garcia.

Mouse Genotyping

Genomic DNA was isolated from tail biopsies and the Ms2Yah allele was identified by a 3-primer PCR using one common Fwd primer (5'-CCAGCTGAAGATGGGTGTGTCTGC-3'), one Rev primer (5'-AGCCTCCCTGGGGACCTGAAA-3') specific for the wt allele and a Ms2Yah Rev (5'-AACGACCGAGCGCAGCGA-3') primer specific for the mutant. PCR reactions gave wt and Ms2Yah products of length 468bp and 272 bp respectively.

Behavioral analysis

All the experimental procedures, for motor and cognitive assessments have been previously described (BESSON *et al.* 2007; LOPES PEREIRA *et al.* 2009; DUCHON *et al.* 2011). 14 wild type (wt) and 13 Ms2Yah male littermates, were obtained from monosomic male crossed with wild-type females. They were 3 months old when they entered the test battery with the following order: 1-Rotarod; 2- Treadmill; 3-Novel Object Recognition; 4-Morris Water Maze; 5- Fear Conditioning; with 1 week between tests.

Briefly, motor performance was tested on the rotarod and treadmill tests, to evaluate motor coordination and learning. The experimental design consisted of two training trials on a rotarod (LE8500, Panlab- Harvard Apparatus, SL, Spain), at 4 rpm followed by consecutive 1 min trials at increasing FRS (4, 10, 14, 19, 24, and 34 rpm; inter-trial period 5 min), and one accelerating rod trial (4 to 40 rpm over 40-s). For each trial, the elapsed time until the mouse fell off the rod was recorded. The treadmill (LE8708, Panlab-Harvard Apparatus, SL, Spain) consisted of a belt (50 cm long and 20 cm wide) with adjustable speed (5 to 150 cm/s) and slope (0°–45°) and an electrified grid delivering a foot shock (0.6 mA) whenever the mice fell off the belt. Mice received eight 1 min trials on a single day session: two training trials (5

cm/s at 0°), and 6 test trials (5, 10, 20, 30, 40 and 50 cm/s at 20°). Number and duration of shocks were quantified.

In the novel object recognition task, a 10 min habituation session was performed in an open field (OF; 70 cm x 70 cm x 40 cm; TBS, SL, Spain) with two multi-colored interlocking plastic bricks (Tente; EXIN-LINES BROS S.A., Barcelona, Spain) with the same preference levels. The familiar ones were a simple colored columns and the unfamiliar was a colored abstract shape construction. 24 h later, a familiarization trial where mice were presented with a pair of identical objects for a maximum of 10 min was performed. Any investigative behaviour (head orientation, or sniffing) or deliberate contact with each object in a distance ≤ 2 cm is considered as exploration of the objects. Animals not exploring the objects for at least 20 s were discarded. After 1h, one of the familiar objects was changed to a new one, and the animals explored the OF for 5 min. Long term memory was operationally defined by the discrimination index (DI): time spent investigating the novel object minus the familiar one ($DI = (Novel\ Object\ Exploration\ Time / Total\ Exploration\ Time) - (Familiar\ Object\ Exploration\ Time / Total\ Exploration\ Time) \times 100$).

Animals were also tested in a visuo-spatial learning paradigm in the Morris Water Maze test (MWM) over 9 days (4 trials/session, 10-min inter-trial intervals). The water maze consisted of a circular pool (diameter, 1.50 m; height, 0.35 m; TBS, SL, Spain) filled with tepid water (24°C) opacified with non-toxic white ink. A white escape platform (8 cm diameter, height 24 cm) was located 1 cm below the water surface in a fixed position (NE quadrant, 25 cm away from the wall). The maze was surrounded by white curtains with 3 black patterns affixed to provide spatial cues. In a pre-training session the platform was visible in the center of the pool (day 1). In the five acquisition sessions (days 2-6), mice were introduced in the

pool through different locations (N, S, E, W) and were allowed to swim until they located the platform. Mice failing to find the platform within 60 s were placed on it for 20 s. A probe trial (day 7) in which the platform was removed was followed by a cued session (day 8) where the platform position was visible. Escape latencies, length of the swimming paths and swimming speed for each animal and trial were monitored and computed by a software tracking system SMART[®] (Panlab-Harvard Apparatus, SL, Spain) connected to a video camera placed above the pool, and Jtrack, an in-house designed software (ARQUÉ *et al.* 2008).

Finally in the pure contextual fear-conditioning paradigm, each mouse was placed in a test chamber (67 cm x 53 cm x 55 cm) inside a sound-attenuated compartment (Panlab-Harvard Apparatus, SL, Spain) with a weight transducer, two audio generators, an electrified grid and an incandescent light of 40W. In the conditioning session mice were allowed to explore freely the chamber for 2 min. Thereafter, an 8 kHz 100dB pure tone, which served as the conditioned stimulus (CS), was presented for 30 s, followed by a mild footshock (2 s, 0.2 mA), which served as the unconditioned stimulus (US; training phase). Fear testing was conducted 24 hr after conditioning during 2 min of free exploration and 3 min to exposure to the CS (test phase). Freezing time (s) was recorded and analyzed. The extinction was further evaluated by scoring freezing time in 2 min sessions over a 6 min period.

Social preference and discrimination was evaluated using a specific test with 3 successive and identical chambers (Stoelting, Dublin, Ireland). The protocol is similar to the one described by Moy and collaborators (Moy *et al.* 2004; Moy *et al.* 2008) and was described previously (DUCHON *et al.* 2011) In the habituation, tested mice were allowed to explore freely the three chambers for 10 min starting from the intermediate compartment and with the two side and opposite chambers having empty wire cage. In the second phase,

the test mouse was placed in the central box, while an unfamiliar mouse (stranger 1) was now put in one of the wire cages in a random and balanced manner. The doors were re-opened and the test mouse was allowed to explore the 3 chambers for 10 min. Time spent in each of the chambers, the number of entries into each chamber and the time spent sniffing each wire cage were recorded for social preference. In a third phase, the social discrimination was evaluated with a new stranger mouse (stranger 2) placed into the empty wire cage and the test mouse was allowed to explore again the entire arena for 6 min, having the choice between the first, already-investigated mouse (stranger 1) and the novel unfamiliar mouse (stranger 2). The same measures were taken as for the social preference. The olfactory tests, was described previously (DUCHON *et al.* 2011).

Electrophysiology

Another experiment in which we found interesting results in Ts1Yah was the activity of hippocampal synapses; change was obviously expected in Ms2Yah correlatively with *Abcg1-U2af1* dosage diminution. Six month old control and mutant males (n = 10 per group) were anesthetized with 0.8-3% halothane at a flow rate of 1 L/min oxygen. Animals were implanted with bipolar stimulating electrodes aimed at the right Schaffer collateral-commissural pathway of the dorsal hippocampus (2 mm lateral and 1.5 mm posterior to Bregma; depth from brain surface, 1.0-1.5 mm; Paxinos & Franklin, 2001) and with two recording electrodes aimed at the CA1 area (1.2 mm lateral and 2.2 mm posterior to Bregma; depth from brain surface, 1.0-1.5 mm). The final position of hippocampal electrodes was determined using the field potential depth profile evoked by paired (40 ms interval) pulses presented at the Schaffer collateral pathway as a guide. A bare silver wire (0.1 mm) was affixed to the skull as a ground. The four wires were connected to a 4-pin socket and the

socket was fixed to the skull with the help of two small screws and dental cement (see Gruart et al. (2006) for details of this chronic preparation).

Recording and stimulation procedures

Field excitatory post-synaptic potentials (fEPSPs) were recorded with Grass P511 differential amplifiers. Electrical stimulus presented to Schaffer collaterals consisted of 100 μ s, square, biphasic pulses presented alone, paired, or in trains. Stimulus intensities ranged from 0.02 to 500 μ A for the construction of the input/output curves. For paired pulse facilitation, the stimulus intensity was set well below the threshold for evoking a population spike, usually 35% of the intensity necessary for evoking a maximum fEPSP response (GUREVICIENE *et al.* 2004). Paired pulses were presented at six (10, 20, 40, 100, 200, and 500) different pulse intervals. The stimulus intensity was also set at 35% of its asymptotic value for LTP induction. For LTP induction, each animal was presented with a high frequency stimulation (HFS) protocol consisting of five trains (200 Hz, 100 ms) of pulses at a rate of 1/s. This protocol was presented 6 times in total, at intervals of 1 min. The 100 μ s, square, biphasic pulses used to evoke LTP were applied at the same intensity used for the single pulse presented following CS presentation. Further details of this chronic preparation can be found in previous work (GRUART *et al.* 2006).

Histological procedures and cresyl violet / Luxol blue staining

Animals were anesthetized with pentobarbital (25 mg/kg) and intracardially perfused with sterile PBS (0.01 M) followed by 4% paraformaldehyde (PFA). Then brains were dissected and kept in 4% PFA solution at 4°C (overnight). Brains were dehydrated, embedded in paraffin and 5-7 μ m coronal sections were processed using microtome (Leica).

Sections were selected between Bregma + 2.00 and + 2.50 mm. Deparaffinized sections were colored using classical procedure for Luxol blue and cresyl violet. Slides were dried over night at RT and then mounted with pertex™. The sections were imaged in brightfield using the NanoZoomer Digital Pathology-2.0 HT (C9600; Hamamatsu Photonics, K.K., Japan). The CA1, CA2-3 and Dentate Gyrus areas were estimated using the NDP.view software. On the same sections, relative neuronal density was estimated by counting the cresyl violet positive cells in 3 randomized fields at magnification x40 for each hippocampal area. For each individual, 5 sections were used and the mean was calculated with 5 animals per genotype (wt and Ms2Yah).

Expression profiling analysis

Hippocampi were isolated from 5 male mice at 3 month of age and flash frozen. Total RNA was extracted using RNeasy mini kit extraction (QIAGEN) according to the manufacturer's instructions. The quality of all RNA samples was controlled using an Agilent 2100 Bioanalyzer (Agilent Technologies). Total RNA was converted to cDNA using Superscript II (Invitrogen). Hybridization was carried out on Affymetrix Mouse Gene 1.0 ST. Raw data of fluorescence was then normalized with the Robust Multiarray Average (RMA) algorithm developed by Irizarry *et al.* (IRIZARRY *et al.* 2003) and we obtained 35,557 probes RMA values. We fixed the background with RMA signals over 4.36 and selected 24697 probes. Values were log transformed and finally we calculated the fold change between the two groups for analysis. Hierarchical clustering was carried out with Cluster3.0 software using Euclidian distances to calculate the distances between the genes and between the samples. Functional annotation was performed using Databases as the known mammalian phenotypes database available at the Mouse Genome Informatics. The microarray data were submitted to the

Array Express Home under the accession number (E-MTAB-988). We applied selection filters on the raw data provided by RMA algorithm to extract relevant differences between our 2 groups. The final data file retained recapitulates all the genes which had a statistical difference (t-test $P < 0.05$) between wt and Ms2Yah, without unknown transcript (no mRNA accession number) and multiple probes for a single gene. We estimated the false positive rate, with $n_1 = \alpha \times N$ (total genes selected) and $n_2 =$ number of values $< \alpha$; α is the threshold P -value used in the student t-test. We applied this to the first batch of probes after background threshold. For $\alpha = 0.05$, we had $n_1 = 80$ and $n_2 = 171$, so the allowed error is $n_1/n_2 = 0.46$.

For the QRT-PCR experiments we considered 14 genes mapped on Mmu17, between the *Abcg1* and *U2af1* genes (mm9 database ENSEMBL); assays for the *Umodl1*, *Abcg1*, *U2af1* and *Cryaa* genes were also designed to check the expression of genes located at the borders of the interval. In addition, we included assays for 17 genes located outside the region of interest. Among these, three genes for each tissue were selected for normalization and the remaining genes were used as additional controls. Selection of normalization genes was performed with GeNorm software (VANDESOMPELE *et al.* 2002). To assess the differences in gene expression values between monosomic and wt samples, we performed the student t test.

RESULTS

General behavior and Motor learning

The Ms2Yah were generated like the Ts1Yah mice (LOPES PEREIRA *et al.* 2009) after chromosomal engineering in embryonic stem cells. Ms2Yah individuals are heterozygous for the deletion of the *Abcg1-U2af1* region encompassing 16 genes. The carrier mouse were obtained and appeared normal without any gross abnormal phenotypes or stereotypic behavior in our standard battery of test. However to assess the motor learning and behavior we compared the Ms2Yah mice with controls using the rotarod and the treadmill paradigms. No significant differences between genotypes were detected in the rotarod test (Figure 1A). The latency to fall off the rod was similar both at constant speeds (4, 10, 14, 19, 24 and 34 rpm) and in the acceleration test (4-40 rpm in 40s). However, in treadmill test, we observed a worse performance in the last session (ANOVA $F_{(1,26)}=4.980$ $P=0.035$; Figure 1B), in which Ms2Yah mice receive a high number of shocks than wild type mice, suggesting a poorer performance on high task demands. The locomotion is not affected in mutant mice, but it appears that Ms2Yah have difficulties to maintain their effort for a long period. No coordination and motor learning impairments were found but muscle activity might be affected.

Cognitive tests

We then investigated whether the decrease in copy number of the *Abcg1-U2af1* region has an impact on cognitive functions as observed for the Ts1Yah mice (LOPES PEREIRA *et al.* 2009). No differences were observed between genotypes in object recognition test (ANOVA $F_{(1,26)}=0.015$ $P=0.903$), nor in the Morris water maze, during the acquisition and

cued sessions (Repeated Measures ANOVA $F_{(1,26)}=0.070$ $P=0.794$). Mice of both genotypes reached the platform with similar latencies (Repeated Measures ANOVA $F_{(1,26)}=0.010$ $P=0.923$) and no differences in swimming speed were detected. In the removal session, no differences were observed in the preference for the trained quadrant (ANOVA $F_{(1,26)}=0.013$ $P=0.909$). In the reversal session no differences in learning the new platform position were observed between genotypes (ANOVA $F_{(1,26)}=0.040$ $P=0.842$). However, in the pure contextual fear conditioning test a tendency to increase the freezing time in test session in Ms2Yah mice was found (ANOVA $F_{(1,26)}=2.963$ $P<0.05$). This increase is significantly higher in Ms2Yah than in wild type mice (Repeated measures $F_{(1,26)}=5.680$ $P=0.025$) with respect to the freezing in training sessions (Figure 2).

Social interest and social discrimination

We evaluated the social behavior of Ms2Yah mice using the three compartment test (Moy *et al.* 2004; Moy *et al.* 2008; DUCHON *et al.* 2011). The wild type and monosomic animals explored similarly the 3 chambers (ANOVA “genotype x chambers”, $F_{(2,36)}=2.588$ $P=0.089$), excluding a place preference in the arena. During the social interest session, control and Ms2Yah mice displayed strong preference for a congener mouse and spent more time exploring the chamber with the congener compared to the empty cage chamber (Two ways ANOVA “congener (chamber 1) versus empty cage (chamber 2)” $F_{(1,24)}=14.17$ $P<0.001$; post-hoc Tuckey “congener (chamber 1) versus empty cage (chamber 2)”, wt $P=0.016$; Ms2Yah $P=0.012$; Figure 3A). Likewise, the percentage of time spent by the control littermate sniffing at the wire cage containing the mouse is increased versus the empty one (ANOVA “chamber1 versus chamber2” $F_{(1,24)}=21.78$ $P<0.001$; post-hoc Tuckey “Stranger1 versus

empty cage”, wt $P < 0.001$; Ms2Yah $P = 0.013$; Figure 3B). We conclude that there was no interference for social interaction due to the monosomic condition.

During the social discrimination there was no difference in the time spent in the chamber with the newly introduced congener (chamber 2) for both groups (ANOVA “familiar (chamber 1) versus novel congener (chamber 2)” $F_{(1,24)} = 0.001$ $P = 0.923$; post-hoc Tuckey “familiar versus novel”, wt $P = 0.167$; Ms2Yah $P = 0.131$; Figure 3C). The tendency for social discrimination was certainly due to a slight difference of the control and Ms2Yah group in exploring the chamber 2 (ANOVA “genotype x chamber”, $F_{(1,24)} = 4.467$ $P = 0.045$); the wt and Ms2Yah displayed different exploratory patterns, confounding the altered pattern of the monosomics. Similarly, the percentage of time spent sniffing the new congener in chamber 2 was affected (ANOVA “Stranger1 chamber versus Stranger2 chamber” $F_{(1,24)} = 6.690$ $P = 0.016$) (Figure 3C) but the wt spent more time exploring the new congener than the Ms2Yah mice (Post-hoc Tuckey “stranger 1 cage versus stranger 2 cage”, wt $P = 0.045$; Ms2Yah $P = 0.08$; time of sniffing the new congener: wt 1.91 ± 0.27 and Ms2Yah: 1.44 ± 0.31). Then we checked that the Ms2Yah have no alteration in odor recognition (Student t-test “Trial 4 versus Test session” wt $P = 0.045$ and Ms2Yah $P = 0.024$; figure 3E). This experiment demonstrated that Ms2Yah animals had a lower interaction with new congeners.

Input/output curves and paired-pulse facilitation

Since overexpression of genes located in the *Abcg1-U2af1* region led to increased long-term potentiation (LTP) in the hippocampal CA3-CA1 synapse (LOPES PEREIRA *et al.* 2009), we explored the effect of the monosomy on this phenotype. In our current experiment, both wild type (Wt) and transgenic (Ms2Yah) mice presented similar increases in the slope of fEPSP evoked at the CA1 area following the presentation of singles pulses of increasing

intensity at the ipsilateral Schaffer collaterals (Figure 4). In all cases, these relationships were fitted by sigmoid curves ($r > 0.9$, $P < 0.0001$), thus suggesting normal functioning of the CA3-CA1 synapse in the two genotypes.

As illustrated in Figure 4B, both wild type and monosomic animals presented a significant [$F_{(12,135)} = 56.126$, $P < 0.001$] increase of the response to the 2nd pulse at short (20-100 ms) time intervals. No significant differences between groups were observed at any of the selected (10, 20, 40, 100, 200, and 500 ms) intervals [$F_{(12,135)} = 0.735$, $P = 0.540$], thus suggesting normal short term hippocampal plasticity.

Comparison of LTP evoked in alert behaving mice

For the LTP study, and in order to obtain a baseline, animals were stimulated every 20 s for ≥ 15 min at Schaffer collaterals (Figure 4C). After a stable baseline was obtained, the experimental animal was presented with the HFS protocol (see Methods). After HFS, the same single stimulus used to generate baseline records was presented at the initial rate (3/min) for another 30 min. Recording sessions were repeated 24 and 48 h later for 15 min each (Figure 4C). Using this HFS and recording protocols, both experimental groups presented a significant LTP for the first 24 h [$P < 0.001$; $F_{(24,216)} = 10.785$], but with some interesting differences between them. Thus, the LTP response presented by Ms2Yah mice was significantly larger than that presented by the Ms2Wt group for most part of the recording period ($P < 0.01$).

Morphological analysis of hippocampus

We measured hippocampal areas (whole and subregions): the mean of entire hippocampus for wild type was 2.01 and 1.92 mm² for the mutant and the selected region,

CA1 stretched to $0,88 \pm 0.09 \text{ mm}^2$ (wt) and $0,83 \pm 0.07 \text{ mm}^2$ (Ms2Yah), CA2-3 $0,54 \pm 0.07 \text{ mm}^2$ (wt) and $0,51 \pm 0.06 \text{ mm}^2$ (Ms2Yah) with the dentate gyrus with $0,59 \pm 0.02 \text{ mm}^2$ (wt) and $0.58 \pm 0.03 \text{ mm}^2$ (Ms2Yah). The cell count was performed to evaluate the neuronal density in hippocampal subregions, by counting cresyl violet stained cells: in CA1, we calculated 1140 cells/ mm^2 in the wt similar to 1199 cells/ mm^2 in Ms2Yah; in CA2-3 region we quantified 1225 cells/ mm^2 in wild type compared to 1245 in mutant group. Finally, in the dentate gyrus area, the density was evaluated to be 1469 cells/ mm^2 in the wt and to 1565 cells/ mm^2 in the Ms2Yah. A student t test was performed and we did not observe any size difference in the hippocampus nor neuronal density. There is neither signs of volume changes nor layer disorganization. We concluded that Ms2Yah did not show abnormal morphology of hippocampus.

Whole Genome expression analysis

To better characterize the mechanisms underlying the behavioral and LTP phenotype observed in the *Abcg1-U2af1* monosomy, we carried out whole genome expression analysis on adult wt and Ms2Yah hippocampi using Affymetrix Gene Chip technology (RAVEAU *et al.* 2012). 24,697 transcripts were found expressed in the hippocampi over our threshold background of which 12009 known with an mRNA accession number. After normalization, we only found a subset of 80 genes that were statistically misexpressed with a fold change higher than 1.20 or lower than 0.80 (table 1). We also performed a cluster analysis showing a clear discrimination between wt and Ms2Yah samples with 2 groups of genes (Figure 5) and The first group corresponds to 33 down-regulated genes, with as expected *Cbs* (Fold change (FC)=0.69 with $P=0.02$) and *Rsph1/Tsga2* (FC=0.80 with $P=0.02$) from the monosomic

interval. The second group encompassed a series of 139 upregulated genes. Most of the genes were located outside the aneuploid regions, indicating a small genome-wide effect on gene expression in the Ms2Yah adult hippocampi. We analyzed the enrichment of functional annotation, using DAVID (HUANG *et al.* 2009) and found an enrichment for postsynaptic cell membrane proteins encoded by *Gabrr2*, *Gabbr1*, *Grid2p*, *Park2* and *Dlg1* (Fold enrichment=13.75; Pvalue= 7.46E-04; FDR= 0.87) and with members of the Ubiquitin mediated proteolysis (*Anapc1*, *Rnf7*, *Huwe1*, *Park2*; Fold enrichment=7.67; Pvalue= 0.012; FDR= 11.13).

To confirm the expression pattern detected in the whole genome approach, we performed quantitative RTPCR on RNA isolated from monosomic hippocampi and their respective controls. All methods from RNA extraction to cDNA quantification have been already described in previous work (LOPES PEREIRA *et al.* 2009; RAVEAU *et al.* 2012). We focused on the *Abcg1-U2af1* genes plus gene located either at the borders, and a few genes among Mmu10 and 16 homologous regions. Genes expressed from the monosomic interval were down regulated as anticipated, with a strong and significant decrease for *Ubash3a*, *Pde9a*, *Wdr4*, *Ndufv3*, *Pknox1* and *Cbs* (Figure 6) whereas the expression a subset of genes, located at the bordering regions (*Umodl1*, *Abcg1*, *U2af1*, *Cryaa*) or outside in the genome in regions homologous to human chromosome or elsewhere, was not affected. We conclude that limited change in the genetic expression was induced by the monosomy of the *Abcg1-U2af1* interval.

DISCUSSION

We evaluated here, the cognitive and electrophysiological phenotypes of a mouse model monosomic for the *Abcg1–U2af1* genetic interval. This model helps us to evaluate the contribution of not only the 16 encompassed coding genes but also other functional elements within this chromosomal region towards cognition. Previous studies of trisomic mice for the same region (Ts1Yah) showed a complex cognitive phenotype that combined alteration in specific domains that are affected in well characterized Down syndrome (DS) models such as non-associative learning and short-term spatial and recognition memory (LOPES PEREIRA *et al.* 2009). Similar results were obtained for a model encompassing the same region (YU *et al.* 2010b). However, trisomy of the interval improved the performance in spatial learning, along with higher and long-lasting LTP, suggesting improved hippocampal function. Thus, we concluded that, whatever the copy number, the *Abcg1-U2af1* region is necessary to maintain normal LTP activation in hippocampus.

The Df(17)1Yey/+ monosomic mouse model, encompassing the *Abcg1-Rpr1b* interval, displayed reduced swimming speed in the Morris water maze and decreased freezing time in the contextual fear conditioning paradigm (YU *et al.* 2010a). In the present report, the Ms2Yah, mice showed slightly different phenotypes: a mild motor defect found in the treadmill; no spatial or recognition memory impairment with no changes in the swimming activity; and improvement in pure contextual fear associative memory along with increased LTP, and a light sociability discrimination defect. The phenotypic differences observed in the two models certainly result from the size of the deficiency: shorter in the Ms2Yah with 4 more disomic genes namely *Cryaa*, *Sik1*, *Hsf2bp* and *Rrp1b* compared with the Df(17)1Yey.

According to the Allen brain atlas, *Cryaa*, *Sik1* and *Rrp1b* are expressed in the adult mouse brain, including the hippocampus (all three), the cortex (*Sik1* and *Rrp1b*) and the cerebellum (*Rrp1b*). *Cryaa* is more likely a molecular chaperone, stabilizing other partners, but with yet unknown brain function (FAN *et al.* 2014). *Sik1* is a HDAC kinase contributing to the CREB pathway, with a wide impact on Brain function (BERDEAUX *et al.* 2007; TAKEMORI *et al.* 2007) and *Rrp1b*, a chromatin-associated factor, a substrate of the MAPK14 (VARJOSALO *et al.* 2013) are more promising candidate.

Interestingly the Ms2Yah is not mirroring the Ts1Yah phenotypes with no changes in the MWM and the Novel object recognition (LOPES PEREIRA *et al.* 2009). Short-term plastic processes are not affected in Ms2Yah animals, since the synaptic facilitation evoked by the presentation of a pair of pulses, a typical presynaptic short-term plastic property of the hippocampal CA3-CA1 synapse, which has been related to the process of neurotransmitter release was not affected in monosomic mice. However, similar to what was previously described in the homologous trisomy, Ms2Yah mice present larger and longer-lasting LTPs than their respective littermate wild types, thus suggesting a dissociation of hippocampal function between spatial memory, novelty discrimination and fear memory in the Ms2Yah model. Comparing those two functions affected, we could hypothesize that spatial and working memory are related to short term plasticity because it is associated only with trisomic mouse model, but associative memory, evaluated by fear conditioning, implicated more neuronal circuits than ones structuring around the hippocampus. The high LTP levels recorded are only present in mice models restricted to syntenic region between Mmu17 and Hsa21. This means that the role of those few common genes could have an underestimated impact on Down syndrome cognition disabilities.

We further explored the molecular origins of the monosomic phenotypes. The transcriptomic study performed comparing the expression profile in wild type and mutant hippocampi points to the extreme sensitivity of the modifications generated by the deletion. On enhancing the stringence and the power of the statistical tests, the analysis highlighted 80 misregulated coding genes (Table 1). Some of them are involved in cognitive function such as *Gabrr1*, *Gabrr2*, *Grid2p*, *Park2* and *Dlg1* encoding postsynaptic cell membrane proteins. *Grid2ip* is an interacting protein of delta subunit of glutamate receptor expressed in the cerebellum and in the hippocampus (MIYAGI *et al.* 2002; MATSUDA *et al.* 2006). Inactivation of *Grid2ip* facilitates LTD in the cerebellum with more sensitivity to calcium response compared to wt (TAKEUCHI *et al.* 2008). Interestingly, *Grid2ip* is a partner of the GluRd2 which is involved in motor learning and motor coordination (KASHIWABUCHI *et al.* 1995). *Grid2ip* could be a good candidate to explore change in the excitability of the neurons and the motor function defects observed in the Ms2Yah mice. We detected also variations of *Park2*, a regulator of the dopamine production in the striatum, whose loss decreases synaptic excitability (GOLDBERG *et al.* 2003). Here, *Park2* was overexpressed in Ms2Yah and may contribute to the enhanced LTP. *Dlg1* the Discs, large homolog 1 multidomain scaffolding protein is involved in synaptogenesis and postsynaptic signaling and potentially involved in brain pathologies (DE BARTOLOMEIS and FIORE 2004; NITHIANANTHARAJAH *et al.* 2013). Interestingly *Dlg1* and *Pak2* were recently found deleted in autism (QUINTERO-RIVERA *et al.* 2010). However both *Gabrr1* and *Gabrr2*, or *Park2* and *Dlg1*, are located at proximity respectively on mouse chromosome 4 or 16 and are coregulated in our Ms2Yah hippocampi. In addition *Sox2* a well-known transcription factor controlling neural stem cells is downregulated in the Ms2Yah brain (WEGNER and STOLT 2005; SHI *et al.* 2008).

From the *Abcg1-U2af1* deleted genetic interval, 6 out of 8 genes expressed in the hippocampus, namely *Ubash3a*, *Pde9a*, *Wdr4*, *Ndufv3*, *Pknox1* and *Cbs* were found downregulated compared to control. Little is known about the functions of *Ubash3a* and *Wdr4* and *Pknox1* in the brain, whereas *Pde9a* encodes a phosphodiesterase specifically expressed in hippocampus, whose inhibition increases early LTP after stimulation (KROKER *et al.* 2012). Thus *Pde9a*, with *Grid2ip*, is a candidate for the enhanced LTP observed in vivo. The *Ndufv3* dehydrogenase is a member of a family that contribute to bipolar disorders or depression (BEN-SHACHAR and KARRY 2008) but the role of this particular protein is actually unknown. *Cbs* is mutated in the intellectual disability homocystinuria. *Cbs* could impact the neuronal function through various mechanisms including the synthesis of H₂S, a new gaseous neurotransmitter (KIMURA 2002; CHEN *et al.* 2004).

During the social interest and discrimination test, we found an unexpected phenotype with the Ms2Yah mice showing reduced interaction with a new congener. This behavior is not due to a deficiency in odor system but suggests memory impairment. Interestingly the *Gabrr1* and *Gabrr2* genes are associated with several cognitive defects including schizophrenia (GREEN *et al.* 2010), and autism (MA *et al.* 2005). Their altered expression observed in the Ms2Yah hippocampi could be linked to the abnormal phenotypes observed in the fear conditioning and the social interaction. With this lack of interest for a novel individual, the Ms2Yah mice develop a specific autistic phenotype. Several Autistic Spectrum Disorders (ASD) are linked with copy number variants (MORENO-DE-LUCA *et al.* 2010; SANDERS *et al.* 2011; LUO *et al.* 2012). In the Autism database (BASU *et al.* 2009), we found a few duplications and one deletion in patients that included segments from the *Abcg1-U2af1* region (JACQUEMONT *et al.* 2006; MARSHALL *et al.* 2008; PINTO *et al.* 2010; LEVY *et al.* 2011; SANDERS *et al.* 2011). Thus the Ms2Yah mice might represent a model for autism

with several functions affected in the brain without any noticeable cellular defects, but with changes in the expression of genes involved in the postsynaptic functions without any stereotypies detected so far.

Partial deletions of 21q are rare and these patients display a highly variable phenotype depending on the size and position of the deletion. A review of the literature identified 38 cases with pure 21q deletions. Partial monosomy 21 was recently segregated into three regions associated with variable clinical severity. Terminal deletions (21q22.2-ter and 21q22.3-ter), or interstitial hemizygous loss of 16.4 Mb (21q21.1-q22.11) gave rise to relatively mild phenotypes while other 21q deletions produce a greater number of dysmorphic features and some major malformations related to genes in the interval (LINDSTRAND *et al.* 2010; ROBERSON *et al.* 2011). The combined information of the 38 available cases suggests that the *ITSN1* gene is involved in severe mental retardation in patients with 21q deletion. In addition, a critical region of 0.56 Mb containing four genes, *KCNE1*, *DSCR1*, *CLIC6* and *RUNX1*, is associated with severe congenital heart defects, and deletions of the most proximal 15-17 Mb of 21q is associated with mild or no cognitive impairment, but may lead to problems with balance and motor function. Several cases of chromosomal aberration encompassing the Human chromosome 21q22.3 up to the qter regions have been reported (LYLE *et al.* 2009; HANNACHI *et al.* 2011; MELIS *et al.* 2011; ROBERSON *et al.* 2011) with mild phenotypes. In the ECARUCA database (www.ecaruca.net), 8 of 16 cases with complex aberration encompassing the 22q3 region of Hsa21 displayed mental retardation (FERRANTE *et al.* 1983; MCGINNISS *et al.* 1992; FEENSTRA *et al.* 2006). In our previous analysis of the monosomy of the most telomeric part of the human chromosome 21, we found no phenotypes attached to the deletion of the *Cstb-Prmt2* genetic interval (DUCHON *et al.* 2011) similar to the mild effect of the 21qter monosomy in human. With this new report, the

Abcg1-U2af1 region should be considered as an important genetic interval, containing key dosage sensitive elements controlling neurological phenotypes. The Ms2Yah mice should be further explored as a model of Hsa21 terminal deletion for which little consequence have been described so far (LYLE *et al.* 2009; LINDSTRAND *et al.* 2010; ROBERSON *et al.* 2011). On the opposite, increase in the *Abcg1-U2af1* region dosage may be determinant for the social behavior observed in Down syndrome.

ACKNOWLEDGEMENTS

We thank members of the research group, of the IGBMC laboratory, of the ICS and of the AnEUploidy consortium for their helpful comments (www.aneuploidy.org). We are grateful to the animal care-takers of the CNRS UPS44 TAAM unit and of the ICS. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The project was supported by the French National Centre for Scientific Research (CNRS), the French National Institute of Health and Medical Research (INSERM), the University of Strasbourg and the “Centre Europeen de Recherche en Biomedecine”, the “Fondation Jerome Lejeune”, Koplowitz, FRAXA and AFM Foundations, Spanish Ministry of Economy (SAF2010-16427, SAF2007-31093-E, FIS-PI 082038), Marató TV3, and the European commission (AnEUploidy project (LSHG-CT-2006-037627) to YH and MD. The laboratory of MD is supported by DURSI (Grups consolidats 09 2009SGR1313).

FIGURES LEGENDS

Figure 1: Locomotor and behavioral assessment of the Ms2Yah monosomic model for the Abcg1-U2af1 region. (A) Rotarod test. No differences were observed between genotypes in the two training sessions prior to the test. In addition, no differences were observed between both genotypes in the latency of falling from the rotating rod. All mice acquired the criteria to perform the task at the same time. (B) Treadmill test. In the treadmill test, all mice showed a normal pattern of locomotion, without apparent alterations. But in the last trial of the test, at the highest demanding speed of 50 rpm, significant difference between genotypes was showed (ANOVA $F_{(1,26)}= 4,980$ $P=0,035$). Ms2Yah mice received significantly more shocks than control wild type mice.

Figure 2: Altered pure contextual fear conditioning in the Ms2Yah mice

(A) When we analyzed the total freezing time during the test phase, no differences between both genotypes were observed, beyond a non-significant trend to increase the percentage of freezing in Ms2Yah mice. However, when we analyzed the freezing during the training phase compared to the test phase, a significant difference between both genotypes was observed (Repeated measures $F_{(1,26)}=5.680$ $P=0.025$), being much higher in Ms2Yah mice respect to the wild-type mice one. (B) we then further analysed the extinction of contextual fear memory by scoring freezing time in 2 min session over 6 min but did not detect any significant changes.

Figure 3: Social novelty performance is altered in the Ms2Yah mice. Percentage of time spent in the different chambers (A,C) and of time spent sniffing the cages (B,D). In A and B,

chamber 1 contained a mouse (familiar) and the chamber 2 is empty. The wt mice showed a strong preference for the chambers and the cage containing an unfamiliar mouse while monosomic mice showed no significant interaction with its congener. In C and D, a new mouse (Novel) was placed in the chamber 2. Here the wt mice had a preference for the chambers or spend more time sniffing the cage with the novel individual whereas the Ms2Yah mice failed to interact with the new mouse and spend more time in the chamber with the familiar individual. In the last figure E, odor learning and discrimination was controlled to exclude olfactory defect in the Ms2Yah mice. Data shown are mean+ S.E.M. for each strain. Statistical analyses are based on student t test (**p<0,001; *p>0,01; *p>0,05)

Figure 4: Changes in Long Term Potentiation in Ms2Yah mice compared to wild type

littermates. (A) Input/output curves for the CA3-CA1 synapse. A single (100 μ s, biphasic) pulse was presented to Schaffer collaterals at increasing intensities (in mA) while recording the evoked fEPSP at the CA1 area for wild type (Ms2Wt, dark blue squares) and transgenic (Ms2Ysh, light blue squares) Ms2 mice. Some fEPSPs collected from the two types of mouse are illustrated at the right. Ms2Yah mice presented input/output curves similar to those presented by Ms2Wt animals. (B) Paired-pulse facilitation. The data shown are mean \pm SEM slopes of the second fEPSP expressed as a percentage of the first for six (10, 20, 40, 100, 200, 500) interstimulus intervals. The two groups of mice presented similar paired-pulse facilitation at intervals of 20-100 ms. (C) The upper panel illustrates examples of fEPSPs collected from selected Ms2Wt (dark blue square) and Ms2Yah (light blue square) animals before (baseline) and after (days 1-3) high-frequency stimulation (HFS) of Schaffer collaterals. The bottom graph illustrates the time course of LTP evoked in the CA1 area (fEPSP mean \pm SEM) following HFS for Ms2Wt and Ms2Yah mice. The HFS was presented

after 15 min of baseline recordings, at the time marked by the dashed line. The fEPSP is given as a percentage of the baseline (100%) slope. Although the two groups presented a significant increase (ANOVA, two-tailed) in fEPSP slope following HFS when compared with baseline records, values collected from the MS2Yah group were significantly (* $P < 0.01$; $F_{(24,216)} = 10.785$) larger than those collected from Ms2Wt mice at the indicated times.

Figure 5: Clustering derived from statistically deregulated genes in Ms2Yah hippocampi.

Whole genome microarray expression analysis was performed on RNA isolated from whole adult mouse hippocampi and statistically deregulated genes were assessed using GeneSpring software. Under-expression and over-expression are represented in green and red respectively, expression levels were calculated by comparison to mean expression level of all arrays for each gene. All the selected genes were selected through significant differentiated genes (p value < 0.05) and with an up/down cut off on the fold change ($0,8 < x$ for the group 1 and $x > 1,2$ for the group 2).

Figure 6: Quantitative PCR expression study focusing on genes located in the *Abcg16U2af1* regions and on genes homologous to *Hsa21* located in additional genetic interval. We

tested Ms2Yah region genes, borders and relevant genes on the mouse chromosome 10 and 16. These results confirm the transcriptome analysis showing underexpressed genes from the *Abcg1-U2af1* region compared to all the other genes tested.

TABLES

Group	Gene Name	Chr	wt	Ms2Yah	FC	Test
			mean±sem	mean±sem		Post Hoc
1	<i>Cldn5</i>	16	0.77 ± 0.15	0.52 ± 0.09	0.68	*
1	<i>Cbs</i>	17	0.9 ± 0.12	0.62 ± 0.05	0.69	*
1	<i>Tmem234</i>	4	0.84 ± 0.1	0.63 ± 0.07	0.75	*
1	<i>Rps2</i>	17	0.98 ± 0.08	0.74 ± 0.07	0.76	*
1	<i>Fam40b/Strip2</i>	6	0.95 ± 0.06	0.72 ± 0.08	0.76	*
1	<i>Tek</i>	4	0.85 ± 0.1	0.66 ± 0.14	0.77	**
1	<i>Ggcx</i>	6	0.99 ± 0.04	0.76 ± 0.09	0.77	*
1	<i>Sox2</i>	3	0.86 ± 0.11	0.67 ± 0.15	0.78	*
1	<i>Hist1h1c</i>	13	1.05 ± 0.15	0.82 ± 0.08	0.78	*
1	<i>Hyou1</i>	9	0.89 ± 0.08	0.69 ± 0.09	0.78	*
1	<i>Copg</i>	6	0.9 ± 0.07	0.7 ± 0.08	0.78	**
1	<i>Slco1c1</i>	6	0.95 ± 0.1	0.74 ± 0.12	0.79	*
1	<i>Gm12696</i>	4	0.71 ± 0.13	0.56 ± 0.12	0.79	*
1	<i>Elk1</i>	X	0.9 ± 0.08	0.71 ± 0.06	0.79	*
1	<i>Gm11599</i>	1	1.02 ± 0.07	0.8 ± 0.06	0.79	*
1	<i>Spsb3</i>	17	0.93 ± 0.04	0.73 ± 0.09	0.79	*
1	<i>Slc39a1</i>	3	1.04 ± 0.09	0.82 ± 0.15	0.79	*
1	<i>2810006K23Rik</i>	5	0.92 ± 0.06	0.73 ± 0.1	0.79	*
1	<i>Tbl3</i>	17	1.07 ± 0.12	0.85 ± 0.07	0.79	*
1	<i>Ms13l2</i>	10	0.96 ± 0.04	0.76 ± 0.12	0.8	*
1	<i>Slc37a3</i>	6	0.88 ± 0.05	0.7 ± 0.12	0.8	*
1	<i>Smyd5</i>	6	0.93 ± 0.09	0.74 ± 0.09	0.8	*
1	<i>Hmcn1</i>	1	0.87 ± 0.08	0.7 ± 0.09	0.8	***
1	<i>Anapc1</i>	2	0.97 ± 0.05	0.77 ± 0.17	0.8	*
2	<i>Dlg1</i>	16	0.87 ± 0.08	1.04 ± 0.1	1.2	*
2	<i>Kcnn3</i>	3	1.04 ± 0.05	1.25 ± 0.13	1.2	*
2	<i>B430305J03Rik</i>	3	1.08 ± 0.07	1.29 ± 0.1	1.2	*
2	<i>Trp53inp1</i>	4	1.12 ± 0.13	1.35 ± 0.15	1.2	*
2	<i>Gdpd2</i>	X	0.95 ± 0.04	1.14 ± 0.07	1.21	*
2	<i>Gabrr2</i>	4	1.28 ± 0.13	1.55 ± 0.09	1.21	*
2	<i>Gm7820</i>	X	1.09 ± 0.04	1.31 ± 0.12	1.21	*
2	<i>Kitl</i>	10	0.88 ± 0.07	1.06 ± 0.08	1.21	*
2	<i>Al836003</i>	15	1.05 ± 0.03	1.26 ± 0.1	1.21	*
2	<i>Cav3</i>	6	0.97 ± 0.04	1.17 ± 0.1	1.21	*
2	<i>S100a10</i>	3	0.91 ± 0.07	1.1 ± 0.1	1.21	*
2	<i>Vma21</i>	X	1.08 ± 0.15	1.31 ± 0.03	1.21	*
2	<i>Figf</i>	X	0.93 ± 0.15	1.13 ± 0.12	1.22	*
2	<i>2210404J11Rik</i>	17	0.98 ± 0.07	1.2 ± 0.13	1.23	*
2	<i>Trhr</i>	15	0.66 ± 0.14	0.81 ± 0.09	1.23	*
2	<i>Tsc22d3</i>	X	1.23 ± 0.26	1.52 ± 0.19	1.23	*
2	<i>Gm4876</i>	6	0.91 ± 0.09	1.12 ± 0.11	1.24	*
2	<i>Arrdc2</i>	8	1.19 ± 0.11	1.47 ± 0.21	1.24	*
2	<i>Cpsf7</i>	19	0.85 ± 0.11	1.06 ± 0.07	1.24	*
2	<i>Cntn6</i>	6	0.92 ± 0.08	1.14 ± 0.13	1.24	*
2	<i>Dclre1b</i>	3	1.05 ± 0.07	1.31 ± 0.16	1.25	**

2	<i>Cacna1g</i>	11	0.92 ± 0.06	1.16 ± 0.14	1.26	*
2	<i>Huwe1</i>	X	1.1 ± 0.11	1.39 ± 0.13	1.26	*
2	<i>Fam70a</i>	X	0.92 ± 0.09	1.16 ± 0.12	1.26	*
2	<i>Gm129</i>	3	1.12 ± 0.26	1.41 ± 0.24	1.26	*
2	<i>Gabrr1</i>	4	1.03 ± 0.05	1.31 ± 0.05	1.27	**
2	<i>Gm9801</i>	7	0.95 ± 0.08	1.21 ± 0.14	1.27	*
2	<i>Anapc1</i>	2	0.95 ± 0.15	1.2 ± 0.08	1.27	*
2	<i>Thpo</i>	16	1.03 ± 0.1	1.31 ± 0.1	1.27	*
2	<i>Park2</i>	17	1.21 ± 0.14	1.55 ± 0.08	1.28	**
2	<i>Gm5148</i>	3	0.96 ± 0.08	1.22 ± 0.13	1.28	*
2	<i>Rnf7</i>	9	0.96 ± 0.04	1.23 ± 0.13	1.28	*
2	<i>Bmp2</i>	2	0.99 ± 0.06	1.26 ± 0.17	1.28	*
2	<i>Cit</i>	5	0.92 ± 0.06	1.18 ± 0.09	1.28	*
2	<i>Uevld</i>	7	0.86 ± 0.07	1.11 ± 0.14	1.28	*
2	<i>Usp54</i>	14	1.12 ± 0.09	1.44 ± 0.25	1.29	*
2	<i>Rnu1b1</i>	3	1.01 ± 0.02	1.31 ± 0.09	1.29	**
2	<i>1700048020Rik</i>	9	1.07 ± 0.03	1.39 ± 0.13	1.29	*
2	<i>Bruno15</i>	4	1 ± 0.11	1.3 ± 0.15	1.3	*
2	<i>Elt1</i>	3	1.08 ± 0.12	1.42 ± 0.31	1.32	*
2	<i>Carhsp1</i>	16	1.06 ± 0.09	1.4 ± 0.17	1.32	*
2	<i>Dhrs1</i>	14	0.98 ± 0.12	1.31 ± 0.14	1.33	*
2	<i>Mybl1</i>	1	1.07 ± 0.11	1.44 ± 0.22	1.35	*
2	<i>Dnahc3</i>	7	0.98 ± 0.07	1.33 ± 0.25	1.36	*
2	<i>Mir692-1</i>	17	0.8 ± 0.14	1.09 ± 0.11	1.36	*
2	<i>Gpr4</i>	7	0.89 ± 0.09	1.22 ± 0.14	1.37	*
2	<i>mIRNAAL589870</i>	UN	1.08 ± 0.13	1.49 ± 0.16	1.37	**
2	<i>Rin3</i>	12	1.06 ± 0.04	1.48 ± 0.25	1.39	*
2	<i>Camk2d</i>	3	0.97 ± 0.06	1.36 ± 0.2	1.4	*
2	<i>Erdr1</i>	UN	1.13 ± 0.14	1.59 ± 0.07	1.41	*
2	<i>Grid2ip</i>	5	0.84 ± 0.08	1.19 ± 0.1	1.41	*
2	<i>Taf1d</i>	9	1.09 ± 0.17	1.56 ± 0.1	1.43	**
2	<i>Ccl28</i>	UN	0.81 ± 0.19	1.16 ± 0.1	1.44	*
2	<i>Plekhf1</i>	7	1.39 ± 0.25	2.05 ± 0.32	1.47	**
2	<i>Meg3</i>	12	0.85 ± 0.17	1.25 ± 0.15	1.48	*
2	<i>3110052M02Rik</i>	17	1.05 ± 0.16	1.59 ± 0.04	1.52	**


Table 1: Genes deregulated in the Ms2Yah mice hippocampi. Group of genes down- (1) or u- (2) regulated for which the statistical test is below 0.05 are listed here. “Chr” for chromosome and “FC” for Fold Change. The post Hoc statistical test used is a bilateral t-test. (P values: ***: P<0.001; **: P<0.01; *: P<0.05).

REFERENCES

- Arqué, G., V. Fotaki, D. Fernández, M. Martínez de Lagrán, M. Arbonés *et al.*, 2008 Impaired spatial learning strategies and novel object recognition in mice haploinsufficient for the dual specificity tyrosine-regulated kinase-1A (Dyrk1A). *PLoS One* 3: e2575.
- Basu, S. N., R. Kollu and S. Banerjee-Basu, 2009 AutDB: a gene reference resource for autism research. *Nucleic Acids Res* 37: D832-836.
- Ben-Shachar, D., and R. Karry, 2008 Neuroanatomical Pattern of Mitochondrial Complex I Pathology Varies between Schizophrenia, Bipolar Disorder and Major Depression. *Plos One* 3.
- Berdeaux, R., N. Goebel, L. Banaszynski, H. Takemori, T. Wandless *et al.*, 2007 SIK1 is a class II HDAC kinase that promotes survival of skeletal myocytes. *Nat Med* 13: 597-603.
- Besson, V., V. Brault, A. Duchon, D. Togbe, J.-C. Bizot *et al.*, 2007 Modeling the monosomy for the telomeric part of human chromosome 21 reveals haploinsufficient genes modulating the inflammatory and airway responses. *Hum Mol Genet* 16: 2040-2052.
- Chen, X. L., K. H. Jhee and W. D. Kruger, 2004 Production of the neuromodulator H₂S by cystathionine beta-synthase via the condensation of cysteine and homocysteine. *Journal of Biological Chemistry* 279: 52082-52086.
- de Bartolomeis, A., and G. Fiore, 2004 Postsynaptic density scaffolding proteins at excitatory synapse and disorders of synaptic plasticity: implications for human behavior pathologies. *Int Rev Neurobiol* 59: 221-254.
- Duchon, A., S. Pothion, V. Brault, A. J. Sharp, V. L. J. Tybulewicz *et al.*, 2011 The telomeric part of the human chromosome 21 from *Cstb* to *Prmt2* is not necessary for the locomotor and short-term memory deficits observed in the Tc1 mouse model of Down syndrome. *Behavioural Brain Research* 217: 271-281.
- Fan, Q., L. Z. Huang, X. J. Zhu, K. K. Zhang, H. F. Ye *et al.*, 2014 Identification of proteins that interact with alpha A-crystallin using a human proteome microarray. *Mol Vis* 20: 117-124.
- Feenstra, I., J. Fang, D. A. Koolen, A. Siezen, C. Evans *et al.*, 2006 European Cytogeneticists Association Register of Unbalanced Chromosome Aberrations (ECARUCA); an online database for rare chromosome abnormalities. *Eur J Med Genet* 49: 279-291.
- Ferrante, E., P. Vignetti, M. Antonelli, L. Bruni, S. Bertasi *et al.*, 1983 Partial monosomy for a 21 chromosome. Report of a new case of r(21) and review of the literature. *Helv Paediatr Acta* 38: 73-80.
- Goldberg, M. S., S. M. Fleming, J. J. Palacino, C. Cepeda, H. A. Lam *et al.*, 2003 Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons. *Journal of Biological Chemistry* 278: 43628-43635.
- Green, E. K., D. Grozeva, V. Moskvina, M. L. Hamshere, I. R. Jones *et al.*, 2010 Variation at the GABA(A) Receptor Gene, Rho 1 (GABRR1) Associated With Susceptibility to Bipolar Schizoaffective Disorder. *American Journal of Medical Genetics Part B-Neuropsychiatric Genetics* 153B: 1347-1349.
- Gruart, A., M. D. Munoz and J. M. Delgado-Garcia, 2006 Involvement of the CA3-CA1 synapse in the acquisition of associative learning in behaving mice. *Journal of Neuroscience* 26: 1077-1087.
- Gureviciene, I., S. Ikonen, K. Gurevicius, A. Sarkaki, T. van Groen *et al.*, 2004 Normal induction but accelerated decay of LTP in APP+PS1 transgenic mice. *Neurobiology of Disease* 15: 188-195.
- Hannachi, H., S. Mougou-Zerelli, I. BenAbdallah, N. Mama, I. Hamdi *et al.*, 2011 Clinical and molecular characterization of a combined 17p13.3 microdeletion with partial monosomy 21q21.3 in a 26-year-old man. *Cytogenet Genome Res* 135: 102-110.
- Huang, D. W., B. T. Sherman and R. A. Lempicki, 2009 Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 4: 44-57.
- Irizarry, R. A., B. Hobbs, F. Collin, Y. D. Beazer-Barclay, K. J. Antonellis *et al.*, 2003 Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249-264.

- Jacquemont, M. L., D. Sanlaville, R. Redon, O. Raoul, V. Cormier-Daire *et al.*, 2006 Array-based comparative genomic hybridisation identifies high frequency of cryptic chromosomal rearrangements in patients with syndromic autism spectrum disorders. *J Med Genet* 43: 843-849.
- Kashiwabuchi, N., K. Ikeda, K. Araki, T. Hirano, K. Shibuki *et al.*, 1995 Impairment of motor coordination, Purkinje cell synapse formation, and cerebellar long-term depression in GluR delta 2 mutant mice. *Cell* 81: 245-252.
- Kimura, H., 2002 Hydrogen sulfide as a neuromodulator. *Molecular Neurobiology* 26: 13-19.
- Korbel, J. O., T. Tirosh-Wagner, A. E. Urban, X. N. Chen, M. Kasowski *et al.*, 2009 The genetic architecture of Down syndrome phenotypes revealed by high-resolution analysis of human segmental trisomies. *Proceedings of the National Academy of Sciences of the United States of America* 106: 12031-12036.
- Kroker, K. S., G. Rast, R. Giovannini, A. Marti, C. Dorner-Ciossek *et al.*, 2012 Inhibition of acetylcholinesterase and phosphodiesterase-9A has differential effects on hippocampal early and late LTP. *Neuropharmacology* 62: 1964-1974.
- Letourneau, A., F. A. Santoni, X. Bonilla, M. R. Sailani, D. Gonzalez *et al.*, 2014 Domains of genome-wide gene expression dysregulation in Down's syndrome. *Nature* 508: 345-350.
- Levy, D., M. Ronemus, B. Yamrom, Y. H. Lee, A. Leotta *et al.*, 2011 Rare de novo and transmitted copy-number variation in autistic spectrum disorders. *Neuron* 70: 886-897.
- Lindstrand, A., H. Malmgren, S. Sahlen, J. Schoumans, A. Nordgren *et al.*, 2010 Detailed molecular and clinical characterization of three patients with 21q deletions. *Clinical Genetics* 77: 145-154.
- Lopes Pereira, P., L. Magnol, I. Sahún, V. Brault, A. Duchon *et al.*, 2009 A new mouse model for the trisomy of the Abcg1-U2af1 region reveals the complexity of the combinatorial genetic code of down syndrome. *Hum Mol Genet* 18: 4756-4769.
- Luo, R., S. J. Sanders, Y. Tian, I. Voineagu, N. Huang *et al.*, 2012 Genome-wide transcriptome profiling reveals the functional impact of rare de novo and recurrent CNVs in autism spectrum disorders. *Am J Hum Genet* 91: 38-55.
- Lyle, R., F. Bena, S. Gagos, C. Gehrig, G. Lopez *et al.*, 2009 Genotype-phenotype correlations in Down syndrome identified by array CGH in 30 cases of partial trisomy and partial monosomy chromosome 21. *European Journal of Human Genetics* 17: 454-466.
- Ma, D. Q., P. L. Whitehead, M. M. Menold, E. R. Martin, A. E. Ashley-Koch *et al.*, 2005 Identification of significant association and gene-gene interaction of GABA receptor subunit genes in autism. *American Journal of Human Genetics* 77: 377-388.
- Marshall, C. R., A. Noor, J. B. Vincent, A. C. Lionel, L. Feuk *et al.*, 2008 Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet* 82: 477-488.
- Matsuda, K., S. Matsuda, C. M. Gladding and M. Yuzaki, 2006 Characterization of the delta 2 glutamate receptor-binding protein delphilin - Splicing variants with differential palmitoylation and an additional PDZ domain. *Journal of Biological Chemistry* 281: 25577-25587.
- McGinniss, M. J., H. H. Kazazian, G. Stetten, M. B. Petersen, H. Boman *et al.*, 1992 Mechanisms of ring chromosome formation in 11 cases of human ring chromosome 21. *Am J Hum Genet* 50: 15-28.
- Mefford, H. C., M. L. Batshaw and E. P. Hoffman, 2012 Genomics, intellectual disability, and autism. *N Engl J Med* 366: 733-743.
- Melis, D., R. Genesio, G. Cappuccio, V. MariaGinocchio, R. D. Casa *et al.*, 2011 Mental retardation, congenital heart malformation, and myelodysplasia in a patient with a complex chromosomal rearrangement involving the critical region 21q22. *Am J Med Genet A* 155A: 1697-1705.
- Miyagi, Y., T. Yamashita, M. Fukaya, T. Sonoda, T. Okuno *et al.*, 2002 Delphilin: a novel PDZ and formin homology domain-containing protein that synaptically colocalizes and interacts with glutamate receptor delta 2 subunit. *Journal of Neuroscience* 22: 803-814.
- Moreno-De-Luca, D., J. G. Mulle, E. B. Kaminsky, S. J. Sanders, S. M. Myers *et al.*, 2010 Deletion 17q12 is a recurrent copy number variant that confers high risk of autism and schizophrenia. *Am J Hum Genet* 87: 618-630.
- Moy, S. S., J. J. Nadler, A. Perez, R. P. Barbaro, J. M. Johns *et al.*, 2004 Sociability and preference for social novelty in five inbred strains: an approach to assess autistic-like behavior in mice. *Genes Brain Behav* 3: 287-302.

- Moy, S. S., J. J. Nadler, N. B. Young, R. J. Nonneman, S. K. Segall *et al.*, 2008 Social approach and repetitive behavior in eleven inbred mouse strains. *Behav Brain Res* 191: 118-129.
- Nithianantharajah, J., N. H. Komiyama, A. McKechnie, M. Johnstone, D. H. Blackwood *et al.*, 2013 Synaptic scaffold evolution generated components of vertebrate cognitive complexity. *Nat Neurosci* 16: 16-24.
- Pinto, D., A. T. Pagnamenta, L. Klei, R. Anney, D. Merico *et al.*, 2010 Functional impact of global rare copy number variation in autism spectrum disorders. *Nature* 466: 368-372.
- Quintero-Rivera, F., P. Sharifi-Hannauer and J. A. Martinez-Agosto, 2010 Autistic and psychiatric findings associated with the 3q29 microdeletion syndrome: case report and review. *Am J Med Genet A* 152A: 2459-2467.
- Raveau, M., J. M. Lignon, V. Nalesso, A. Duchon, Y. Groner *et al.*, 2012 The App-Runx1 Region Is Critical for Birth Defects and Electrocardiographic Dysfunctions Observed in a Down Syndrome Mouse Model. *Plos Genetics* 8.
- Reeves, R. H., N. G. Irving, T. H. Moran, A. Wohn, C. Kitt *et al.*, 1995 A MOUSE MODEL FOR DOWN-SYNDROME EXHIBITS LEARNING AND BEHAVIOR DEFICITS. *Nature Genetics* 11: 177-184.
- Roberson, E. D., E. S. Wohler, J. E. Hoover-Fong, E. Lisi, E. L. Stevens *et al.*, 2011 Genomic analysis of partial 21q monosomies with variable phenotypes. *Eur J Hum Genet* 19: 235-238.
- Sanders, S. J., A. G. Ercan-Sencicek, V. Hus, R. Luo, M. T. Murtha *et al.*, 2011 Multiple recurrent de novo CNVs, including duplications of the 7q11.23 Williams syndrome region, are strongly associated with autism. *Neuron* 70: 863-885.
- Shi, Y., G. Sun, C. Zhao and R. Stewart, 2008 Neural stem cell self-renewal. *Crit Rev Oncol Hematol* 65: 43-53.
- Sussan, T. E., A. N. Yang, F. Li, M. C. Ostrowski and R. H. Reeves, 2008 Trisomy represses Apc(Min)-mediated tumours in mouse models of Down's syndrome. *Nature* 451: 73-U10.
- Takemori, H., J. Kajimura and M. Okamoto, 2007 TORC-SIK cascade regulates CREB activity through the basic leucine zipper domain. *FEBS J* 274: 3202-3209.
- Takeuchi, T., G. Ohtsuki, T. Yoshida, M. Fukaya, T. Wainai *et al.*, 2008 Enhancement of Both Long-Term Depression Induction and Optokinetic Response Adaptation in Mice Lacking Delphilin. *Plos One* 3.
- Umlauf, D., P. Fraser and T. Nagano, 2008 The role of long non-coding RNAs in chromatin structure and gene regulation: variations on a theme. *Biological Chemistry* 389: 323-331.
- Vandesompele, J., K. De Preter, F. Pattyn, B. Poppe, N. Van Roy *et al.*, 2002 Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* 3.
- Varjosalo, M., S. Keskitalo, A. Van Drogen, H. Nurkkala, A. Vichalkovski *et al.*, 2013 The protein interaction landscape of the human CMGC kinase group. *Cell Rep* 3: 1306-1320.
- Wegner, M., and C. C. Stolt, 2005 From stem cells to neurons and glia: a Soxist's view of neural development. *Trends Neurosci* 28: 583-588.
- Yu, T., S. Clapcote, Z. Li, C. Liu, A. Pao *et al.*, 2010a Deficiencies in the region syntenic to human 21q22.3 cause cognitive deficits in mice. *Mamm Genome* 21: 258-267.
- Yu, T., Z. Y. Li, Z. P. Jia, S. J. Clapcote, C. H. Liu *et al.*, 2010b A mouse model of Down syndrome trisomic for all human chromosome 21 syntenic regions. *Human Molecular Genetics* 19: 2780-2791.

A**B**