

Safyan Ahmad; Shamsa Kanwal; Talat Firdous
Cohen-Macaulay modifications of the vertex cover ideal of a graph

Czechoslovak Mathematical Journal, Vol. 68 (2018), No. 3, 843–852

Persistent URL: <http://dml.cz/dmlcz/147372>

Terms of use:

© Institute of Mathematics AS CR, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

COHEN-MACAULAY MODIFICATIONS OF THE
VERTEX COVER IDEAL OF A GRAPH

SAFYAN AHMAD, SHAMSA KANWAL, Lahore, TALAT FIRDOUS, Gujrat

Received January 15, 2017. Published online June 7, 2018.

Abstract. We study when the modifications of the Cohen-Macaulay vertex cover ideal of a graph are Cohen-Macaulay.

Keywords: monomial ideal, minimal vertex cover, polarization of ideal, chordal graph

MSC 2010: 13A02, 13D02, 13D25, 13P10

1. INTRODUCTION

Let K be a field and $S = K[x_1, \dots, x_n]$ the polynomial ring in n variables over K , and let I be a squarefree Cohen-Macaulay monomial ideal in S . We will denote the unique minimal system of monomial generators of I by $G(I)$. Let $G(I) = \{u_1, \dots, u_m\}$, then we call a monomial ideal J a modification of I , if $G(J) = \{v_1, \dots, v_m\}$ and $\text{supp}(u_i) = \text{supp}(v_i)$ for all i . By support of a monomial u we mean the set $\text{supp}(u) = \{i : x_i \text{ divides } u\}$. A monomial ideal J is called a *trivial modification* of I , if there exist nonnegative integers a_1, \dots, a_n such that J is obtained from I by the substitutions $x_i \mapsto x_i^{a_i}$ for all i . Obviously, if J is a trivial modification of I , then J is Cohen-Macaulay as $J = \varphi(I)S$ where $\varphi: S \rightarrow S$ is a flat K -algebra homomorphism with $\varphi(x_i) = x_i^{a_i}$ for all i .

Let G be a simple connected graph on the vertex set $V(G) = \{v_1, \dots, v_n\}$ with the edge set $E(G)$. The vertex cover ideal I_G associated to G is the ideal generated by all monomials of the form $\prod_{x_i \in C} x_i$ for all minimal vertex covers C of G . Recall that by a minimal vertex cover we mean a subset $C \subset V(G)$ such that every edge has at least one vertex in C and no proper subset of C has the same property,

$$I_G = \bigcap_{\{v_i, v_j\} \in E(G)} (x_i, x_j).$$

Dually one defines the edge ideal

$$I(G) = (x_i x_j : \{v_i, v_j\} \in E(G))$$

Let Δ_G be the simplicial complex whose Stanley-Reisner ideal I_{Δ_G} coincides with $I(G)$. Then $I_G = I_{\Delta_G^\vee}$, where Δ_G^\vee is the Alexander dual of Δ_G .

Recall that a graph G is chordal if each cycle in G of length greater than 3 has a chord and the complementary graph \overline{G} of G is the graph with $V(\overline{G}) = V(G)$ and $E(\overline{G}) = \{\{v_i, v_j\} : \{v_i, v_j\} \notin E(G)\}$. By using the Alexander duality and results by Eagon-Reiner [4] as well as Fröberg [5] we immediately obtain the following statement.

Proposition 1.1 ([1]). *The ideal I_G is Cohen-Macaulay if and only if the complementary graph \overline{G} is chordal.*

The purpose of this paper is to complement the results presented in the paper [1]; related questions have been studied in [2], [3] and [7].

Let us first review the concept of polarization. Given a monomial

$$u = x_1^{a_1} x_2^{a_2} \dots x_n^{a_n}$$

we define the following monomial in a new set of variables:

$$u^p = \prod_{i=1}^n \prod_{j=1}^{a_i} x_{ij}.$$

Now let $I \subset S$ be an arbitrary monomial ideal with the minimal set of monomial generators $\{u_1, \dots, u_m\}$. Then we set

$$I^p = (u_1^p, \dots, u_m^p).$$

This ideal is called the *polarization* of I . If we choose an arbitrary set $\{v_1, \dots, v_r\}$ of monomial generators of I , then we have

$$I^p = I^p R = (v_1^p, \dots, v_r^p) R,$$

where R is the polynomial ring over K in the variables which are needed to polarize the monomials v_i . We will also need the following rule: Suppose $I = I_1 \cap I_2 \cap \dots \cap I_r$ where each I_j is a monomial ideal, then

$$(1.1) \quad I^p = I^p R = (I_1^p R \cap I_2^p R \cap \dots \cap I_r^p R),$$

where R is again the polynomial ring over K in the variables which are needed to polarize all the monomials involved.

Proposition 1.2 ([6], Corollary 1.6.3). *Let I be a monomial ideal. The following condition are equivalent:*

- ▷ I is Cohen-Macaulay.
- ▷ I^p is Cohen-Macaulay.

We need some preparation to formulate the main results.

2. MODIFICATIONS OF FIRST TYPE

Let G be a simple connected graph with vertex set $V(G)$ and edge set $E(G)$ and let I_G be the vertex cover ideal of G ,

$$I_G = \bigcap_{\{v_i, v_j\} \in E(G)} (x_i, x_j).$$

In this section we consider those modifications where we take powers of both variables in the prime ideal in I_G corresponding to some edge, in the primary decomposition of I_G , i.e. of the form

$$J = (x_{a_i}^m, x_{b_i}^m) \cap \left(\bigcap_{\{a_j, b_j\} \in E(G), i \neq j} (x_{a_j}, x_{b_j}) \right).$$

Definition 2.1. For $\{a, b\} \in E(G)$ and $m \in \mathbb{Z}$, $m > 1$, we define a new graph $\overline{G_{m,ab}}$ with vertex set $V(\overline{G_{m,ab}}) = V(G) \cup \{a_{11}, \dots, a_{1m-1}, b_{11}, \dots, b_{1m-1}\}$, where $a_{1i}, b_{1i} \notin V(G)$ for all $i = 1, \dots, m-1$ and edge set

$$\begin{aligned} E(\overline{G_{m,ab}}) = E(G) \cup & \left(\bigcup_{i=1}^{m-1} \{\{a_{1i}, v\}, \{u, b_{1i}\} : u, v \in V(G), u \neq a, v \neq b\} \right) \\ & \cup \left(\bigcup_{i \neq j, i, j=1}^{m-1} \{a_{1i}, a_{1j}\} \right) \cup \left(\bigcup_{i \neq j, i, j=1}^{m-1} \{b_{1i}, b_{1j}\} \right) \end{aligned}$$

With the above notation, we have the following lemma.

Lemma 2.2. *Let G be a graph with $|V(G)| \geq 4$ and $\{a, b\} \in E(G)$. If there exist $c, d \in V(G) \setminus \{a, b\}$ with $\{c, d\} \in E(G)$ then $\overline{G_{m,ab}}$ contains minimal cycles $\{a_{1i}, c, b_{1i}, d\}$ for all $i = 1, \dots, m-1$.*

Proof. Suppose there exist $c, d \in V(G) \setminus \{a, b\}$ with $\{c, d\} \in E(G)$. By the definition of $\overline{G_{m,ab}}$, we know that

$$E(\overline{G_{m,ab}}) = E(\overline{G}) \cup \left(\bigcup_{i=1}^{m-1} \{\{a_{1i}, v\}, \{u, b_{1i}\} : u, v \in V(G), u \neq a, v \neq b\} \right) \\ \cup \left(\bigcup_{i \neq j, i, j=1}^{m-1} \{a_{1i}, a_{1j}\} \right) \cup \left(\bigcup_{i \neq j, i, j=1}^{m-1} \{b_{1i}, b_{1j}\} \right).$$

As $\{c, d\} \in E(G)$, so $\{c, d\} \notin E(\overline{G_{m,ab}})$. Also by the definition of $\overline{G_{m,ab}}$ it is clear that $\{a_{1i}, b_{1j}\} \notin E(\overline{G_{m,ab}})$ for all $i, j = 1, \dots, m-1$ and $\{a_{1i}, c\}, \{a_{1i}, d\}, \{c, b_{1i}\}, \{d, b_{1i}\} \in E(\overline{G_{m,ab}})$ for all $i = 1, \dots, m-1$. Using all these facts, we have minimal cycles $\{a_{1i}, c, b_{1i}, d\}$ in $\overline{G_{m,ab}}$ for all $i = 1, \dots, m-1$. \square


Another observation regarding $\overline{G_{m,ab}}$ is recorded as the following lemma.

Lemma 2.3. *Let G be a graph with $|V(G)| \geq 4$. If \overline{G} is chordal then $\overline{G_{m,ab}}$ has no minimal cycle of length not less than 5.*


Proof. Since \overline{G} is chordal, all its minimal cycles have length 3. Suppose that $\overline{G_{m,ab}}$ contains a minimal cycle C of length not less than 5; as \overline{G} is chordal it follows that $V(C) \not\subset V(\overline{G})$. Thus there exists $v \in V(C)$ such that $v \notin V(\overline{G})$ and by the definition of $\overline{G_{m,ab}}$, $v \in \{a_{11}, \dots, a_{1m-1}, b_{11}, \dots, b_{1m-1}\}$.

If $v = a_{1i}$ for some $i = 1, \dots, m-1$, we know from the definition that a_{1i} is adjacent to every vertex in $\overline{G_{m,ab}}$ except $b, b_{11}, \dots, b_{1m-1}$. Thus C must contain the edge formed by b and b_{1t} for some $t \in \{1, \dots, m-1\}$; note that $b_{1k} \notin V(C)$ for $k \neq t$ because $\{b, b_{1k}\}, \{b_{1k}, b_{1j}\} \in E(\overline{G_{m,ab}})$ for all $k \neq j; k, j = 1, \dots, m-1$.

Similar reasoning shows that C also contains a and so $\{a, a_{1i}\}, \{b, b_{1t}\} \in E(C)$. Now for any $l \in V(C) \setminus \{a, a_{1i}, b, b_{1t}\}$, we have $\{l, a_{1i}\}, \{l, b_{1t}\} \in E(\overline{G_{m,ab}})$. Thus the cycle must be of the following form:



As $\{a, b\} \notin E(\overline{G_{m,ab}})$ and this is a cycle, we must have at least one more vertex in this cycle, say o ,



where $o \notin \{a, b, a_{1i}, b_{1t}, l\}$.

But then by the definition of $\overline{G_{m,ab}}$, we must have $\{a_{1i}, o\}, \{b_{1t}, o\} \in E(\overline{G_{m,ab}})$, thus no such cycle of length not less than 5 exists in $\overline{G_{m,ab}}$. The case when $v = b_{1i}$ for any $i = 1, \dots, m - 1$, can be proved along the same lines. \square

Proposition 2.4. *Let \overline{G} be a chordal graph with $|V(\overline{G})| \geq 4$, then $\overline{G_{m,ab}}$ is not chordal if and only if there exist $\{c, d\} \in E(G)$ with $c, d \in V(G) \setminus \{a, b\}$.*

Proof. If there exist $\{c, d\} \in E(G)$ with $c, d \in V(G) \setminus \{a, b\}$, Lemma 2.2 guarantees that $\overline{G_{m,ab}}$ contains at least one minimal 4-cycle through c and d , thus $\overline{G_{m,ab}}$ is not chordal.

Conversely if $\overline{G_{m,ab}}$ is not chordal and \overline{G} is chordal, Lemma 2.3 ensures that $\overline{G_{m,ab}}$ contains a 4 cycle, say $\{p, q, r, s\}$. As \overline{G} is chordal, some of these vertices do not belong to $V(\overline{G})$. Moreover, $\{q, s\}$ does not belong to $E(\overline{G})$, without loss of generality, we may assume that $p = a_{1i}$ for some $i = 1, \dots, m - 1$, then neither q nor s are b . Since $\{a_{1i}, r\} \notin E(\overline{G_{m,ab}})$ we have that $r = b$ or $r = b_{1j}$ for some $j = 1, \dots, m - 1$. By the definition of $\overline{G_{m,ab}}$, we have $q \neq a, s \neq a$ so that $\{q, s\}$ is the requested edge, as desired. \square

Theorem 2.5. *Let G be a simple connected graph and let*

$$I_G = \bigcap_{\{a_i, b_i\} \in E(G)} (x_{a_i}, x_{b_i})$$

be the Cohen-Macaulay vertex cover ideal of G . Then

$$J = (x_{a_i}^m, x_{b_i}^m) \cap \left(\bigcap_{\{a_j, b_j\} \in E(G), i \neq j} (x_{a_j}, x_{b_j}) \right)$$

is Cohen-Macaulay for no $m > 1$ if and only if there exist an edge $\{c, d\} \in E(G)$ such that $c, d \notin \{a_i, b_i\}$.

Proof. The ideal I_G is Cohen-Macaulay if and only if \overline{G} is chordal. Using primary decomposition and polarization we can observe that the ideal J will be Cohen-Macaulay if and only if the graph $\overline{G_{m,a_i b_i}}$ is chordal. But by Proposition 2.4, $\overline{G_{m,a_i b_i}}$ is not chordal if and only if there exist $\{c, d\} \in E(G)$ with $c, d \in V(G) \setminus \{a_i, b_i\}$. Thus the ideal J will not be Cohen-Macaulay if and only if there exist $\{c, d\} \in E(G)$ with $c, d \in V(G) \setminus \{a_i, b_i\}$, completing the proof. \square

Corollary 2.6. *If $\{a_i, b_i\}$ is a minimal vertex cover of G , then the ideal*


$$J = (x_{a_i}^m, x_{b_i}^m) \cap \left(\bigcap_{\{a_j, b_j\} \in E(G), i \neq j} (x_{a_j}, x_{b_j}) \right)$$

is Cohen-Macaulay for all $m \in \mathbb{Z}^+$.

Proof. As $\{a_i, b_i\}$ is a minimal vertex cover of G , there exists no edge $\{c, d\} \in E(G)$ such that $c, d \notin \{a_i, b_i\}$, so that J is Cohen-Macaulay. \square

Example 2.7.

(1) Consider the graph G with the vertex set $V(G) = \{v_1, v_2, v_3, v_4\}$ and edge set $E(G) = \{\{v_1, v_2\}, \{v_2, v_3\}, \{v_3, v_4\}\}$:



Here the vertex cover ideal will be

$$I_G = (x_1, x_2) \cap (x_2, x_3) \cap (x_3, x_4).$$

Now, there exists an edge $\{v_3, v_4\}$ such that $v_3, v_4 \notin \{v_1, v_2\}$, thus Theorem 2.5 guarantees that the ideal

$$J = (x_1^m, x_2^m) \cap (x_2, x_3) \cap (x_3, x_4)$$

will not be Cohen-Macaulay for any $m > 1$. The same is true with $\{v_1, v_2\}$ and $\{v_3, v_4\}$ interchanged. On the other hand the edge $\{v_2, v_3\}$ is a minimal vertex cover for this graph so by Theorem 2.5, the ideal

$$K = (x_1, x_2) \cap (x_2^m, x_3^m) \cap (x_3, x_4)$$

is Cohen-Macaulay for all choices of $m \in \mathbb{Z}^+$.

(2) Let us consider the graph G with $V(G) = \{v_1, v_2, v_3, v_4, v_5\}$ and $E(G) = \{\{v_1, v_2\}, \{v_2, v_3\}, \{v_2, v_5\}, \{v_3, v_4\}\}$. Then

$$I_G = (x_1, x_2) \cap (x_2, x_3) \cap (x_3, x_4) \cap (x_2, x_5).$$

By Theorem 2.5, all the following ideals are not Cohen-Macaulay for any $m > 1$,

$$\begin{aligned} J_1 &= (x_1^m, x_2^m) \cap (x_2, x_3) \cap (x_3, x_4) \cap (x_2, x_5); \\ J_2 &= (x_1, x_2) \cap (x_2, x_3) \cap (x_3^m, x_4^m) \cap (x_2, x_5); \\ J_3 &= (x_1, x_2) \cap (x_2, x_3) \cap (x_3, x_4) \cap (x_2^m, x_5^m). \end{aligned}$$

On the other hand, the ideal

$$J_4 = (x_1, x_2) \cap (x_2^m, x_3^m) \cap (x_3, x_4) \cap (x_2, x_5)$$

is Cohen-Macaulay for all choices of $m > 1$, $m \in \mathbb{Z}$ because $\{v_2, v_3\}$ is a minimal vertex cover of G .

(3) Consider the graph with vertex set $\{v_1, \dots, v_n\}$ and edge set $\{\{v_1, v_i\}: 2 \leq i \leq n\}$, the so called *bouquet graph*. Then

$$J = \bigcap_{\{v_1, v_i\} \in E(G)} (x_1^{m_i}, x_i^{m_i})$$

is Cohen-Macaulay for all $m_i > 1$, $m_i \in \mathbb{Z}$.

(4) Finally, for the graph K_3 , all its edges are minimal vertex covers of K_3 . Thus the ideal

$$J = (x_1^l, x_2^l) \cap (x_2^m, x_3^m) \cap (x_3^n, x_1^n)$$

is Cohen-Macaulay for all choices of $l, m, n \in \mathbb{Z}^+$.

3. MODIFICATIONS OF SECOND TYPE

In this section we consider the modifications of the form

$$J = (x_{a_i}^m, x_{b_i}) \cap \left(\bigcap_{\{a_j, b_j\} \in E(G), i \neq j} (x_{a_j}, x_{b_j}) \right).$$

We recall that a subset $T \subset V(G)$ is called an *independent set* of G , if for all $v_i, v_j \in T$ it holds that $\{v_i, v_j\} \notin E(G)$. An independent set T is called *maximal*, if it is not a proper subset of any independent set, see [8]. The set of vertices adjacent to v_i will be denoted by $N_G(v_i)$. In [1], the first author proved the following result:

Theorem 3.1. *Suppose that I_G is Cohen-Macaulay. Let $W = \{v_{i_1}, \dots, v_{i_r}\}$ be a set of pairwise distinct vertices of G with the property that each v_{i_k} belongs to exactly one maximal independent set T_k of G , where $T_k \neq T_l$ for $k \neq l$. For each*

$v_{i_k} \in W$ choose a nonempty subset $A_k \subset N_G(v_{i_k})$ with the property that, if some $v_{i_k} \in A_j$ then $v_{i_j} \notin A_k$ and $A_k \cap A_l = \emptyset$ for $k \neq l$, and let

$$J = \bigcap_{\{v_i, v_j\}} (x_i, x_j) \cap \bigcap_{k=1}^r \bigcap_{v_j \in A_k} (x_{i_k}, x_j^{a_j}),$$

where the first intersection is taken over all edges $\{v_i, v_j\}$ different from the edges $\{v_{i_k}, v_j\}$ with $v_j \in A_k$, and where each a_j is a positive integer. Then J is Cohen-Macaulay.

We will now prove the converse.

Definition 3.2. Let G be a graph, $v \in V(G)$ and w a new vertex not belonging to $V(G)$. We let G_v be the graph with $V(G_v) = V(G) \cup \{w\}$ and $E(G_v) = E(G) \cup \{u, w\} : u \in V(G), u \neq v\}$.

Let $c(G)$ be the maximum length of a chord-less cycle in G .

Lemma 3.3 ([1], Lemma 3.2). *Suppose G is chordal. Then $c(G_v) \leq 4$.*

Definition 3.4. Let $v \in V(G)$ and $N_G(v) = \{v_1, \dots, v_r\}$. Then we define $c_G(v)$ to be the cardinality of the set

$$\{\{v_i, v_j\} : \{v_i, v_j\} \notin E(G); 1 \leq i < j \leq r\},$$

and call $c_G(v)$ the cycle number of v in G .

Remark 3.5. Note that $c_G(v) = 0$ if and only if the restriction of G to the vertex set $\{v\} \cup N_G(v) \subset V(G)$ is a clique in G (a complete subgraph of G). Observe that if $\{v\} \cup N_G(v)$ is a clique, it is indeed a maximal clique in $V(G)$, since it contains all neighbors of v .

Let us see an immediate consequence of Lemma 3.3.

Lemma 3.6. *G_v is chordal if and only if $c_G(v) = 0$.*

Now we are ready to state and prove our main theorem of this section.

Theorem 3.7. *Let G be a graph such that the vertex cover ideal*

$$I_G = \bigcap_{\{v_i, v_j\} \in E(G)} (x_i, x_j)$$

is Cohen-Macaulay. If for some $v_i \in V(G)$ there exists $v_k, v_l \in V(G)$ with $\{v_k, v_l\} \in E(G)$ and $\{v_i, v_k\}, \{v_i, v_l\} \notin E(G)$, then for any $v_m \in N_G(v_i)$

$$J = \bigcap_{\{v_i, v_j\} \in E(G); j \neq m} (x_i, x_j) \cap (x_i, x_m^n)$$

is not Cohen-Macaulay, for any $n \geq 2$.

Proof. It is enough to prove the theorem for $n = 2$. Since, by assumption, I_G is Cohen-Macaulay, it follows that \overline{G} is chordal. Suppose for some $v_i \in V(G)$ there exists $v_k, v_l \in V(G)$ such that $\{v_k, v_l\} \in E(G)$ with $\{v_i, v_k\}, \{v_i, v_l\} \notin E(G)$. Then $\{v_k, v_l\} \notin E(\overline{G})$ and $\{v_i, v_k\}, \{v_i, v_l\} \in E(\overline{G})$.

As $\{v_i\} \cup N_{\overline{G}}(v_i)$ is not a clique in \overline{G} , $c_{\overline{G}}(v_i) \neq 0$ and \overline{G}_{v_i} is not chordal.

Since

$$J = \bigcap_{\{v_i, v_j\} \in E(G); j \neq m} (x_i, x_j) \cap (x_i, x_m^2)$$

we have

$$J^p = \bigcap_{\{v_i, v_j\} \in E(G); j \neq m} (x_i, x_j) \cap (x_i, x_m) \cap (x_i, w) = \bigcap_{\{v_i, v_j\} \in E(G)} (x_i, x_j) \cap (x_i, w).$$


Let H be the graph obtained from G by adding a whisker with vertex i . Then $J^p = I_H$, where H is the complementary graph of \overline{G}_{v_i} ; this implies that J^p is not Cohen-Macaulay and hence J is not Cohen-Macaulay. \square

Now we will formulate a complete example to demonstrate the result.

Example 3.8. Consider the graph shown in the figure:



The vertex cover ideal associated to this graph is $I_G = (x_1, x_2) \cap (x_2, x_3) \cap (x_3, x_4)$. The complementary graph \overline{G} of G is



As this graph is chordal, the ideal I_G is Cohen-Macaulay. Now, $\{v_3, v_4\} \in E(G)$ and $\{v_1, v_3\}, \{v_1, v_4\} \notin E(G)$. Moreover, $N_G(v_1) = \{v_2\}$, whence by Theorem 3.7, the ideal

$$I_G = (x_1, x_2^n) \cap (x_2, x_3) \cap (x_3, x_4)$$

is not Cohen-Macaulay for any n greater than 1.

Acknowledgements. We gratefully acknowledge the referee for his comments and suggestions to improve the presentation of the paper and for pointing out grammatical mistakes.

References

- [1] *S. Ahmad*: Cohen-Macaulay intersections. Arch. Math. *92* (2009), 228–236. [zbl](#) [MR](#) [doi](#)
- [2] *S. Ahmad, M. Naeem*: Cohen-Macaulay monomial ideals with given radical. J. Pure Appl. Algebra. *214* (2010), 1812–1817. [zbl](#) [MR](#) [doi](#)
- [3] *S. Ahmad, M. Naeem*: Classes of simplicial complexes which admit non-trivial Cohen-Macaulay modifications. Stud. Sci. Math. Hung. *52* (2015), 423–433. [zbl](#) [MR](#) [doi](#)
- [4] *J. A. Eagon, V. Reiner*: Resolutions of Stanley-Reisner rings and Alexander duality. J. Pure Appl. Algebra *130* (1998), 265–275. [zbl](#) [MR](#) [doi](#)
- [5] *R. Fröberg*: Rings with monomial relations having linear resolutions. J. Pure Appl. Algebra *38* (1985), 235–241. [zbl](#) [MR](#) [doi](#)
- [6] *J. Herzog, T. Hibi*: Monomial Ideals. Graduate Texts in Mathematics 260. Springer, London, 2011. [zbl](#) [MR](#) [doi](#)
- [7] *J. Herzog, Y. Takayama, N. Terai*: On the radical of a monomial ideal. Arch. Math. *85* (2005), 397–408. [zbl](#) [MR](#) [doi](#)
- [8] *R. H. Villarreal*: Monomial Algebras. Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, 2015. [zbl](#) [MR](#) [doi](#)

Authors' addresses: Safyan Ahmad, Shamsa Kanwal, Abdus Salam School of Mathematical Sciences, GC University Lahore, Lahore, Pakistan, e-mail: safyank@gmail.com, lotus_zone16@yahoo.com; Talat Firdous, University of Gujrat, Gujrat, Pakistan, e-mail: talatfirdous25@gmail.com.