

Collapse and dispersal of a homogeneous spin fluid in Einstein–Cartan theory

M. Hashemi, S. Jalalzadeh^a, A. H. Ziaie

Department of Physics, Shahid Beheshti University, G. C., Evin, Tehran 19839, Iran

Received: 16 July 2014 / Accepted: 6 January 2015 / Published online: 4 February 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract In the present work, we revisit the process of gravitational collapse of a spherically symmetric homogeneous dust fluid which is described by the Oppenheimer–Snyder (OS) model (Oppenheimer and Snyder in *Phys Rev D* 56:455, 1939). We show that such a scenario would not end in a spacetime singularity when the spin degrees of freedom of fermionic particles within the collapsing cloud are taken into account. To this purpose, we take the matter content of the stellar object as a homogeneous Weysenhoff fluid. Employing the homogeneous and isotropic FLRW metric for the interior spacetime setup, it is shown that the spin of matter, in the context of a negative pressure, acts against the pull of gravity and decelerates the dynamical evolution of the collapse in its later stages. Our results show a picture of gravitational collapse in which the collapse process halts at a finite radius, whose value depends on the initial configuration. We thus show that the spacetime singularity that occurs in the OS model is replaced by a non-singular bounce beyond which the collapsing cloud re-expands to infinity. Depending on the model parameters, one can find a minimum value for the boundary of the collapsing cloud or correspondingly a threshold value for the mass content below which the horizon formation can be avoided. Our results are supported by a thorough numerical analysis.

1 Introduction

One of the most important questions in a gravitational theory, such as general relativity (GR), and relativistic astrophysics is the gravitational collapse of a massive star under its own gravity at the end of its life cycle. A process in which a sufficiently massive star undergoes a continual gravitational collapse on exhausting its nuclear fuel, without achieving an equilibrium state [2]. According to the singularity theorems in GR [3–5], the spacetimes describing the solutions

of the Einstein field equations in a typical scenario of the collapse would inevitably admit singularities.¹ These theorems are based on three main assumptions under which the existence of a spacetime singularity is foretold in the form of geodesic incompleteness in the spacetime. The first premise is in the form of a suitable causality condition that ensures a physically reasonable global structure of the spacetime. The second premise is an energy condition that requires the positivity of the energy density at the classical regime as seen by a local observer. The third one requires that gravity be so strong that trapped surface² formation must occur during the dynamical evolution of a continual gravitational collapse.

The first detailed treatment of the gravitational collapse of a massive star, within the framework of GR, was published by Oppenheimer and Snyder [1]. They concluded that gravitational collapse of a spherically symmetric homogeneous dust cloud would end in a black hole. Such a black hole is described by the presence of a horizon which covers the spacetime singularity. This scenario provides the basic motivation for the physics of black holes and the cosmic censorship conjecture (CCC) [8–11]. This conjecture states that the spacetime singularities that develop in a gravitational scenario of the collapse are necessarily covered by the event horizons, thus ensuring that the collapse end-product is a black hole only. As no proof, nor a stringent mathematical formulation of the CCC has been available so far, a great deal of effort has been made in the past decades to perform a detailed study of several collapse settings in GR, in order to extend our understanding of this phenomenon.

While black hole physics has given rise to interesting theoretical as well as astrophysical progress, it is necessary, how-

¹ These are the spacetime events where the metric tensor is undefined or is not suitably differentiable, the curvature scalars and densities are infinite and the existing physical framework would break down [6].

² A trapped surface is a closed two-surface on which both in-going as well as out-going light signals normal to it are necessarily converging [7].

^a e-mails: s.jalalzadeh@gmail.com; sjalalzadeh@sbu.ac.ir

ever, to investigate more realistic collapse settings in order to put black hole physics on a firm status. This is because the OS model is rather idealized and pressures as well as inhomogeneities within the matter distributions would play an important role in the collapse dynamics of any realistic stellar object. It is therefore of significant importance to broaden the study of gravitational collapse to more realistic models in order to deal with this question: what ways are there for possible departures in final outcomes, as opposed to the homogeneous dust cloud collapse? Within this context, several gravitational collapse settings have been investigated over the past years which represent the occurrence of naked singularities.³ Work along this line has been reported in the literature within a variety of models; among them we quote the role of inhomogeneities within the matter distribution on the final fate of gravitational collapse [12–22], collapse of a perfect fluid with heat conduction [23–26], the effects of shear on the collapse end-product [27–30], and the collapse process in the context of different gravitational theories [31, 32] (see also [33, 34] for recent reviews).

On the other hand, though GR has emerged as a highly successful theory of gravitation, it suffers from the occurrence of spacetime singularities under physically reasonable conditions. It is therefore plausible to seek for the alternative theories of gravitation whose geometrical attributes are not present in GR. This allows for the inclusion of more realistic matter fields within the structure of stellar objects, in order to cure the singularity problem. In this regard, since the realistic stars are made up of fermions, it would be difficult to reject the role of intrinsic angular momentum (spin) of fermions in collapse studies. As we shall see, the inclusion of spin of fermions and thus its possible effects on the collapse dynamics could be of significant importance, specially at the late stages of the collapse setting where these effects could go against the gravitational attraction to ultimately balance it. In such a scenario the collapse may no longer terminate in a spacetime singularity and instead is replaced by a bounce, a point at which the contraction of matter cloud stops and an expanding phase begins. However, if the spin effects are explicitly present, then GR will no longer be the relevant theory to describe the collapse dynamics. In GR, the energy–momentum tensor couples to the metric, while in the presence of fermions, it is expected that the intrinsic angular momentum is coupled to a geometrical quantity related to the rotational degrees of freedom in the spacetime, the so called spacetime torsion. This obviously is not possible in the ambit of GR so that one is forced to modify the theory in order to introduce torsion and relate it to the spin degrees

of freedom of fermions. This point of view suggests a spacetime manifold which is non-Riemannian. One such framework, within which the inclusion of spin effects of fermions can be worked out and thus will allow non-trivial dynamical consequences to be extracted is the Einstein–Cartan (EC) theory [35–39]. Within this context, many cosmological models have been found in which the unphysical big bang singularity is replaced with a bounce at a finite amount the scalar factor [40–54]. From another perspective, the research of the recent years has shown that in the final stages of a typical scenario of the collapse where a high energy regime governs, the effects of quantum gravity would regularize the singularity that happens in the classical model [55, 56]. In cosmological settings, it is shown that non-perturbative quantum geometric effects in loop quantum cosmology would replace the classical singularity by a quantum bounce in the high energy regime where the loop quantum modifications are dominant [57–61]. However, since the full quantum theory of gravity has not yet been discovered, investigating the repulsive spin effects of fermions, which is more physically reasonable and confirmed observationally, on the final state of collapse could be well motivated. The organization of this paper is as follows: In Sect. 2 we give a brief review on the field equations in EC theory and the phenomenological Weysenhoff model. In Sect. 3, we study the collapse dynamics in the presence of spin effects and the possibility of singularity removal. Finally, our conclusions are drawn in Sect. 4.

2 Einstein–Cartan theory

As we know, the dynamics of the gravitational field, i.e., the metric field, in GR is described by the Hilbert–Einstein action with the Lagrangian which is linear in curvature scalar. Contrary to GR, in the gravity with torsion there is a considerable freedom in constructing the dynamical scheme, since one can define much more invariants from torsion and curvature tensors. There are two most attractable classes of models, namely: the EC theory and the quadratic theories. In this work we are interested in the former one for which the action integral is given by

$$S = \int d^4x \sqrt{-g} \left\{ \frac{-\hat{R}}{2\kappa} + \mathcal{L}_m \right\}, \quad (1)$$

where $\kappa = 8\pi G$ (we set $c = 1$) is the gravitational coupling constant, \mathcal{L}_m is the Lagrangian for the matter fields and \hat{R} is the EC curvature scalar constructed out of the general asymmetric connection $\hat{\Gamma}^\alpha_{\mu\nu}$, i.e., the connection of the Riemann–Cartan manifold. The torsion tensor $T^\alpha_{\mu\nu}$ is defined as the antisymmetric part of the affine connection, given by

$$T^\mu_{\alpha\beta} = \frac{1}{2} [\hat{\Gamma}^\mu_{\alpha\beta} - \hat{\Gamma}^\mu_{\beta\alpha}]. \quad (2)$$

³ In this case, the horizons are delayed or failed to form during collapse, as governed by the internal dynamics of the collapsing object. Then the scenario where the super-dense regions are visible to external observers occurs, and a visible naked singularity forms [2].

From the metricity condition, $\hat{\nabla}_\alpha g_{\mu\nu} = 0$ we can find the affine connection as

$$\hat{\Gamma}^\mu_{\alpha\beta} = \left\{ \begin{matrix} \mu \\ \alpha\beta \end{matrix} \right\} + K^\mu_{\alpha\beta}, \tag{3}$$

where the first part is the Christoffel symbols and the second part is the contorsion tensor defined as

$$K^\mu_{\alpha\beta} = T^\mu_{\alpha\beta} + T_{\alpha\beta}{}^\mu + T_{\beta\alpha}{}^\mu. \tag{4}$$

Extremizing the total action with respect to the contorsion tensor gives the Cartan field equation as

$$T^\alpha_{\mu\beta} - \delta^\alpha_\beta T^\gamma_{\mu\gamma} + \delta^\alpha_\mu T^\gamma_{\beta\gamma} = -\frac{1}{2}\kappa\tau_{\mu\beta}{}^\alpha, \tag{5}$$

where $\tau^{\mu\alpha\beta} = 2(\delta\mathcal{L}_m/\delta K_{\mu\alpha\beta})/\sqrt{-g}$ is the spin angular momentum tensor [38, 39]. It is worth noting that the equation governing the torsion tensor is an equation of pure algebraic type, i.e., the torsion is not allowed to propagate beyond the matter distribution as a torsion wave or through any interaction of non-vanishing range [38, 39]; therefore it can be only nonzero inside material bodies. Varying the total action with respect to the metric tensor leads to the Einstein field equation with additional terms on the curvature side that are quadratic in the torsion tensor [62–64]. Substituting for the torsion tensor, from Eq. (5), into these terms we get the combined field equations [38, 39, 53, 54, 62–66]:

$$G_{\mu\beta}(\{\}) = \kappa(\mathcal{T}_{\mu\beta} + \Sigma_{\mu\beta}), \tag{6}$$

where $\mathcal{T}_{\mu\beta} = 2(\delta\mathcal{L}_m/\delta g^{\mu\beta})/\sqrt{-g}$ is the dynamical energy-momentum tensor and $\Sigma_{\mu\beta}$ can be considered as representing the contribution of an effective spin–spin interaction [38, 39], i.e., the product terms

$$\begin{aligned} \Sigma_{\mu\beta} = & \frac{1}{2}\kappa \left[\tau_{\mu\alpha}{}^\alpha \tau_{\beta\gamma}{}^\gamma - \tau_{\mu}{}^{\alpha\gamma} \tau_{\beta\gamma\alpha} - \tau_{\mu}{}^{\alpha\gamma} \tau_{\beta\alpha\gamma} \right. \\ & + \frac{1}{2}\tau^{\alpha\gamma}{}_\mu \tau_{\alpha\gamma\beta} + \frac{1}{4}g_{\mu\beta} \left(2\tau_{\alpha\gamma\epsilon} \tau^{\alpha\epsilon\gamma} \right. \\ & \left. \left. - 2\tau_{\alpha\gamma}{}^\gamma \tau^{\alpha\epsilon}{}_\epsilon + \tau^{\alpha\gamma\epsilon} \tau_{\alpha\gamma\epsilon} \right) \right]. \tag{7} \end{aligned}$$

It now is obvious that the second term on the right hand side of (6), represents a correction (though very weakly at ordinary densities as this term carries a factor κ^2) to the dynamical energy-momentum tensor, which takes into account the spin contributions to the geometry of the manifold.⁴ However, the spin corrections are significant only at the late stages of gravitational collapse of a compact object, where superdense regions of extreme gravity are involved. Therefore, we have a good motivation to investigate the collapse process of material fluid sources which are endowed with spin. Let us now apply Eq. (6) to estimate the influence of spin in the case of the Weyssenhoff fluid, which generalizes the perfect fluid of GR to the case of non-vanishing spin. This model of the

fluid was first studied by Weyssenhoff and Raabe [67–69] and extended by Obukhov and Korotky in order to build cosmological models based on the EC theory [70]. In the model presented in this paper we employ an ideal Weyssenhoff fluid which is considered as a continuous medium whose elements are characterized by the intrinsic angular momentum (spin) of particles. In this model the spin density is described by the second-rank antisymmetric tensor $S_{\mu\nu} = -S_{\nu\mu}$. The spin tensor for the Weyssenhoff fluid is then postulated to be

$$\tau_{\mu\nu}{}^\alpha = S_{\mu\nu}U^\alpha, \tag{8}$$

where U^α is the four-velocity of the fluid element. The Frenkel condition which arises by varying the Lagrangian of the sources [70] requires $S^{\mu\nu}U_\nu = 0$. This condition further restricts the torsion tensor to be traceless.

From the microscopical viewpoint, a randomly oriented gas of fermions is the source for the spacetime torsion. However, we have to treat this issue from a macroscopic perspective, which means we need to perform suitable spacetime averaging. In this respect, the average of the spin density tensor vanishes, $\langle S_{\mu\nu} \rangle = 0$ [38, 39, 71]. But even with this term vanishing at macroscopic level, the square of the spin density tensor $S^2 = \frac{1}{2}\langle S_{\mu\nu}S^{\mu\nu} \rangle$ contributes to the total energy-momentum tensor. Taking these considerations into account, the relations (6)–(8) then give the Einstein field equation with spin correction terms [38, 39, 65, 66]

$$G_{\mu\nu} = \kappa \left(\rho + p - \frac{\kappa}{2}S^2 \right) U_\mu U_\nu - \kappa \left(p - \frac{\kappa}{4}S^2 \right) g_{\mu\nu}, \tag{9}$$

where ρ and p are the usual energy density and pressure of the perfect fluid satisfying a barotropic equation of state $p = w\rho$. Thus, the EC equation for such a spin fluid would be equivalent to the Einstein equation for a perfect fluid with the effective energy density $\rho_{\text{eff}} = \rho - \frac{\kappa}{4}S^2$ and effective pressure $p_{\text{eff}} = p - \frac{\kappa}{4}S^2$. This estimate shows that the contribution of the spin of the fermions to the gravitational interaction is negligible in the case of normal matter densities (e.g., in the early stages of the collapse process) while in the late stages of the collapse, where one encounters ultra-high energy densities, it is the spin contribution that decides the final fate of the scenario of the collapse. This is the subject of our next discussion.

3 Spin effects on the collapse dynamics and singularity removal

The study of gravitational collapse of a compact object and its importance in relativistic astrophysics was initiated since the work of Datt [72] and OS [1] (see also [73] for a pedagogical discussion). This model, which simplifies the complexity of such an astrophysical scenario, describes the process of gravitational collapse of a homogeneous dust cloud with no rotation and internal stresses in the framework of GR. Assum-

⁴ We note that if the spin is switched off, the field Eq. (6) reduces to the ordinary Einstein’s field equation.

ing the interior geometry of the collapsing object to be that of the FLRW metric, they investigated the dynamics of the continuous gravitational collapse of such a matter distribution under its own weight and showed that for an observer comoving with the fluid, the radius of the star crushes to zero size and the energy density diverges in a finite proper time. For this idealized model, which showed that a black hole is developed as the collapse end-state, the only evolving portion of the spacetime is the interior of the collapsing object, while the exterior spacetime remains like that of the Schwarzschild solution with a dynamical boundary. However, in more realistic scenarios, the dynamical evolution of a collapse setting would be significantly different in the later stages of the collapse where the inhomogeneities are introduced within the densities and pressures. These effects could alter the dynamics of the horizons and, consequently, the scenario describing the fate of collapse [74].

The spin effects within more realistic stellar collapse models could have considerable effects on the collapse dynamics as we shall see in this section. In order to deal with this purpose, the matter content of the collapsing object is taken as a homogeneous and isotropic Weyssenhoff fluid that collapses under its own gravity. We then parametrize the interior line element as

$$ds^2 = dt^2 - \frac{a(t)^2 dr^2}{1 - kr^2} - R^2(t, r) d\Omega^2, \tag{10}$$

where $R(t, r) = ra(t)$ is the physical radius of the collapsing star with $a(t)$ is the scale factor, k is a constant that is related to the curvature of spatial metric, and $d\Omega^2$ is the standard line element on the unit two-sphere. The field equations then read

$$\begin{cases} \left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho - \frac{(4\pi G)^2}{3}S^2, \\ \ddot{a} = -\frac{4\pi G}{3}(\rho + 3p) + \frac{2}{3}(4\pi G)^2 S^2. \end{cases} \tag{11}$$

The contracted Bianchi identities give rise to the continuity equation:

$$\dot{\rho}_{\text{eff}} = -3\frac{\dot{a}}{a}(\rho_{\text{eff}} + p_{\text{eff}}), \tag{12}$$

whence we have

$$\dot{\rho} = -3\frac{\dot{a}}{a}(\rho + p), \quad (S^2)^\cdot = -6\frac{\dot{a}}{a}S^2. \tag{13}$$

The first parts of the above equations give

$$\rho = \rho_i \left(\frac{a}{a_i}\right)^{-3(1+w)}, \tag{14}$$

where ρ_i is the initial energy density profile and a_i is the initial value of the scale factor at the initial epoch. A suitable averaging procedure leads to the following relation between the spin squared and energy densities [75]:

$$S^2 = \frac{\hbar^2}{8} A_w^{-\frac{2}{1+w}} \rho^{\frac{2}{1+w}}, \tag{15}$$

where A_w is a dimensional constant that depends on the equation of state parameter. It should be noticed that substituting (14) into the above expression leads to $S^2 \propto a^{-6}$, which is nothing but the solution of the second part of (13). The field equations can then be re-written as

$$\begin{cases} \left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = 2Ca^{-3(1+w)} - Da^{-6}, \\ \ddot{a} = -C(1 + 3w)a^{-3(1+w)} + 2Da^{-6}, \end{cases} \tag{16}$$

where $C = \frac{4\pi G}{3}\rho_i a_i^{3(1+w)}$ and $D = \frac{(4\pi G)^2}{24}\hbar^2 A_w^{-\frac{2}{1+w}} \rho_i^{\frac{2}{1+w}} a_i^6$. Next we proceed to study the collapse evolution for different values of the spatial curvature. We assume that the star begins its contraction phase from a stable situation, i.e. $\dot{a}(t_i) = 0$, where t_i is the initial time at which the collapse commences. Thus, from the first part of (16) we find

$$k = \left[\frac{2C}{D} - a_i^{3(w-1)}\right] D a_i^{-(1+3w)}. \tag{17}$$

Depending on the sign of the expression in double brackets, the constant k may be either positive, negative or zero. Therefore we may write

$$\begin{cases} k > 0 & \frac{2C}{D} > a_i^{3(w-1)}, \\ k \leq 0 & \frac{2C}{D} \leq a_i^{3(w-1)}. \end{cases} \tag{18}$$

Let us consider the dust fluid ($w = 0$) for which the solution of ($k = 0$) clearly represents an expanding solution. For the case ($k < 0$) the collapse velocity is non-real which is physically implausible [76]. Thus the only remaining case is $k > 0$ for which we are to investigate the collapse dynamics for large and small values of the scale factor, i.e., the early and late stages of the collapse process, respectively.

In the early stages of the collapse, the spin contribution is negligible and thus the first part of (16) can be approximated by

$$\dot{a}^2 \cong -k + \frac{2C}{a}. \tag{19}$$

Performing the transformation $ad\xi = \sqrt{k}dt$ we get the solution

$$\begin{cases} a(\xi) = \frac{C}{k}(1 \pm \cos(\xi)), \\ t(\xi) = \frac{C}{k^{\frac{3}{2}}}(\xi \pm \sin(\xi)) + t_i, \end{cases} \tag{20}$$

where according to Eq. (19) $a_i \cong 2C/k$.

For the collapse evolution where the scale factor has become small enough and the spin effects are dominant, the




Fig. 1 Time behavior of the scale factor (*left panel*) and the collapse velocity (*right panel*) for $C = 1.11$, $a_i = 1$, and $\dot{a}(t_i) = 0$, $D = 0.08$ (*full curve*), and $D = 0.0$ (*dashed curve*)

k term in the first Friedmann equation (16) can be neglected compared to the rest. We then get

$$\dot{a}^2 \cong \frac{2C}{a} - \frac{D}{a^4}, \tag{21}$$

for which the solution is given by

$$a(t) = \left\{ a_s^3 + \frac{9}{2}C(t - t_s)^2 - \sqrt{18C}(t - t_s)\sqrt{a_s^3 - \frac{D}{2C}} \right\}^{\frac{1}{3}}, \tag{22}$$

where $t_s > t_i$ represents the time at which the small scale factor regime starts at a finite value of the scale factor $a_s < a_i$. The solution (22) exhibits a bounce at a finite time, say $t = t_b$, where the collapse halts ($\dot{a}(t_b) = 0$) at a minimum value of the scale factor given by

$$a_{\min} = \left[\frac{D}{2C} \right]^{\frac{1}{3}} = \left[\frac{\pi G \hbar^2 \rho_i}{4A_0^2} \right]^{\frac{1}{3}} a_i. \tag{23}$$

We note that for $a > a_{\min}$, Eq. (22) is always real.

For a physically reasonable collapse setting the weak energy condition (WEC) must be satisfied. This condition states that for any non-spacelike vector field $T_{\alpha\beta}^{\text{eff}} V^\alpha V^\beta \geq 0$, which for our model amounts to $\rho_{\text{eff}} \geq 0$ and $\rho_{\text{eff}} + p_{\text{eff}} \geq 0$. The first inequality suggests that $\rho \geq 2\pi GS^2$, while the latter implies $\rho \geq 4\pi GS^2$. The first inequality with the use of (15) gives

$$\rho \leq \frac{4A_0^2}{\pi G \hbar^2} = \rho_i \left(\frac{a_i}{a_{\min}} \right)^3, \tag{24}$$

whereby considering (14) we arrive at $a/a_{\min} \geq 1$. Since the scale factor never reaches the values smaller than a_{\min} , this inequality is always held implying the satisfaction of positive energy density condition. Moreover, the second inequality with similar calculations for dust tells us $a/a_{\min} \geq 2^{\frac{1}{3}}$. This means that in the later stages of the collapse as governed by a spin dominated regime, WEC is violated. Such a violation

of the weak energy condition can be compared to the models where the quantum effects in the scenario of the collapse have been discussed [77]. In brief, we have WEC violation for the following interval:

$$a_{\min} \leq a < 2^{\frac{1}{3}} a_{\min}. \tag{25}$$

3.1 Numerical analysis

In order to get a better understanding of the situation we perform a numerical simulation for the time behavior of the scale factor, the collapse velocity, its acceleration and Kretschmann scalar, by solving the second part of (16) numerically and taking the first part as the initial constraint. The left panel of Fig. 1 shows that if the spin effects are neglected the collapse process terminates in a spacetime singularity (dashed curve) where the scale factor vanishes at a finite amount of comoving time. As the full curve shows, the scale factor begins its evolution from its initial value but deviates from the singular curve as the collapse advances. It then reaches a minimum value (a_{\min}) at the bounce time, $t = t_b$, after which the collapsing phase turns into an expansion. The scale factor never vanishes and hence the spacetime is regular throughout the contracting and expanding phases.

The diagram for the speed of collapse indeed verifies such a behavior (see the full curve in the right panel of Fig. 1) where the collapse begins at rest with the speed changing its sign from negative to positive values at the bounce time. The behavior of the collapse acceleration gives us more interesting results. We see that \ddot{a} changes its sign at two inflection points (see the left panel of Fig. 2) in such a way that for $t < t_{1\text{inf}}$ the collapse undergoes an accelerated contracting phase ($\ddot{a} < 0$ and $\dot{a} < 0$). For the time interval $t_{1\text{inf}} < t < t_b$ the collapse process experiences a decelerated contracting phase where $\ddot{a} < 0$ and $\dot{a} < 0$. After the bounce occurs, the scenario enters an inflationary expanding regime till the second inflection point is reached, i.e., $\ddot{a} > 0$ and $\dot{a} > 0$ for $t_b < t < t_{2\text{inf}}$. Finally a decelerating expanding phase




Fig. 2 Time behavior of collapse acceleration (*left panel*) and Kretschmann scalar (*right panel*) for $C = 1.11$, $a_i = 1$, and $\dot{a}(t_i) = 0$, $D = 0.08$ (*full curve*), and $D = 0.0$ (*dashed curve*)

commences once the collapse acceleration passes through its second inflection point (the same cosmological scenario has been discussed in [71]). The collapsing cloud then disperses at later times. Concomitantly, the Kretschmann scalar increases toward a maximum value but remains finite during the whole process of contraction and expansion (see the full curve in the right panel of Fig. 2) signaling the avoidance of a spacetime singularity.

Now, what would happen to the formation of an apparent horizon during the entire evolution of the collapsing cloud? Especially we may ask: is the bounce visible or not? In order to answer this question we proceed by recasting the metric (10) into the double-null form:

$$ds^2 = -2d\zeta^+d\zeta^- + R^2d\Omega^2, \tag{26}$$

with the null one-forms defined as

$$\begin{aligned} d\zeta^+ &= -\frac{1}{\sqrt{2}} \left[dt - \frac{a}{\sqrt{1-kr^2}} dr \right], \\ d\zeta^- &= -\frac{1}{\sqrt{2}} \left[dt + \frac{a}{\sqrt{1-kr^2}} dr \right]. \end{aligned} \tag{27}$$

From the above expressions we can easily find the null vector fields as

$$\begin{aligned} \partial_+ &= \frac{\partial}{\partial \zeta^+} = -\sqrt{2} \left[\partial_t - \frac{\sqrt{1-kr^2}}{a} \partial_r \right], \\ \partial_- &= \frac{\partial}{\partial \zeta^-} = -\sqrt{2} \left[\partial_t + \frac{\sqrt{1-kr^2}}{a} \partial_r \right]. \end{aligned} \tag{28}$$

The condition for the radial null geodesics, $ds^2 = 0$, leaves us with the two kinds of null geodesics characterized by $\zeta^+ = \text{constant}$ and $\zeta^- = \text{constant}$. The expansions along these two congruences are given by

$$\theta_{\pm} = \frac{2}{R} \partial_{\pm} R. \tag{29}$$

In a spherically symmetric spacetime, the Misner–Sharp quasi-local mass which is the total mass within the radial




Fig. 3 Time behavior of the apparent horizon curve for $a_i = 1$ and $\dot{a}(t_i) = 0$, $C = 1.11$, $D = 0.08$ (*full curve*), and $C = 1.11$, $D = 0$ (*dashed curve*)

coordinate r at the time t is defined as [78–81]

$$\begin{aligned} m(t, r) &= \frac{R(t, r)}{2} (1 + g^{\mu\nu} \partial_{\mu} R(t, r) \partial_{\nu} R(t, r)) \\ &= \frac{R(t, r)}{2} \left(1 + \frac{R(t, r)^2}{2} \theta_+ \theta_- \right). \end{aligned} \tag{30}$$

Therefore, it is the ratio $2m(t, r)/R(t, r) = \dot{R}(t, r)^2 + kr^2$ that controls the formation or otherwise of the trapped surfaces so that the apparent horizon defined as the outermost boundary of the trapped surfaces is given by the condition $\theta_+ \theta_- = 0$ or equivalently $2m(t, r_{ah}(t)) = R(t, r_{ah}(t))$. The equation for the apparent horizon curve then reads

$$R(t, r_{ah}(t))^{-2} = \left(\frac{\dot{a}(t)}{a(t)} \right)^2 + \frac{k}{a(t)^2}, \tag{31}$$

or by the virtue of the first part of 16

$$r_{ah}(a(t)) = \left[\frac{2C}{a(t)} - \frac{D}{a(t)^4} \right]^{-\frac{1}{2}}, \tag{32}$$

which shows the time at which the shell labeled by r becomes trapped. Figure 3 shows the time behavior of the apparent

horizon where we see that, in the presence of spin effects, the apparent horizon decreases (see the full curve) for a while to a minimum value, where the first inflection point is reached ($\ddot{a}(t_{\text{inf}}) = 0$). It then increases to a finite maximum at the bounce time and converges again in the post-bounce regime to the same minimum value where the acceleration vanishes for the second time. The apparent horizon goes to infinity at later times. In order to find the minimum value for the apparent horizon curve we can easily extremize Eq. (31) to get

$$a_\star = \left(\frac{2D}{C}\right)^{\frac{1}{3}}. \tag{33}$$

Therefore there exists a minimum radius

$$r_{\text{min}} = r_{\text{AH}}(a_\star) = \frac{1}{\sqrt{2}a_i} \left(\frac{\hbar}{\pi G \rho_i A_0}\right)^{\frac{1}{3}}, \tag{34}$$

for which if the boundary of the collapsing object is taken as $r_b < r_{\text{min}}$, the apparent horizon does not develop throughout the collapsing and expanding phases and therefore the bounce is uncovered. From the viewpoint of Eq. (31), we can also deduce that since the collapse velocity is bounded the apparent horizon never converges to zero, implying that there exists a minimum radius (the dashed red curve) below which no horizon would form to meet the boundary of the collapsing object. However, the apparent horizon does not diverge at the bounce since $k > 0$; and to which extent it could grow depends on the initial configuration of the stellar object. When the spin effects are absent, the apparent horizon decreases monotonically (see the dashed curve) to finally cover the resulting singularity. There cannot be found any minimum for the boundary of the collapsing cloud so that the formation of the horizon can be prevented.

The existence of a minimum value for the boundary implies that there exists a minimum value for the total mass contained within the collapsing cloud. Let us be more precise. Using Eq. (30) we can re-write the dynamical interior field equations (16) as

$$\begin{aligned} \partial_r m(t, r) &= 4\pi G \rho_{\text{eff}} R(t, r)^2 \partial_r R(t, r), \\ \partial_t m(t, r) &= -4\pi G p_{\text{eff}} R(t, r)^2 \partial_t R(t, r), \end{aligned} \tag{35}$$

whereby integration of the first part gives

$$m(t, r) = \frac{4\pi G}{3} r^3 \rho_i a_i^3 \left[1 - \frac{\pi G \hbar^2}{4A_0^2} \rho_i \left(\frac{a_i}{a(t)}\right)^3 \right]. \tag{36}$$

The above expression together with (33) and (34) gives the threshold mass confined within the radius r_{min} as

$$m_\star = m(a_\star, r_{\text{min}}) = \frac{\hbar}{\sqrt{2}A_0}. \tag{37}$$

Thus, if the total mass is chosen so that $m < m_\star$, there would not exist enough mass within the collapsing cloud at the later

stages of the collapse to get the light trapped and, as a result, formation of the apparent horizon is avoided.

Furthermore, the time derivative of the mass function, using the second part of (35),

$$\partial_t m(t, r) = \frac{\pi^2 G^2 \hbar^2 \rho_i^2 a_i^6}{A_0^2 a(t)^4} r^3 \dot{a}(t), \tag{38}$$

is negative throughout the contracting phase. This may be interpreted as if some mass may be thrown away from the stellar object till the bounce time is approached. At this time, $m(a_{\text{min}}, r) = 0$, which can be imagined as the whole evaporation of the collapsing cloud. After this time, when the expanding phase begins, the ejected mass may be regained since $\dot{m}|_{(t>t_b)} > 0$. We note that such a behavior is due to the homogeneity of the model since all the shells of matter collapse or expand simultaneously. For the case of dust fluid considered here, the exterior region of the star can be modeled by a Schwarzschild spacetime since the spin effects are negligible at the early stages of the collapse. However, as the collapse advances, the mass profile is no longer constant due to the presence of a negative pressure, which originates from the spin contribution. Hence, at the very late stages of the collapse, the Schwarzschild spacetime may not be a suitable candidate for the matching process and instead the interior region should be smoothly matched to the exterior generalized Vaidya metric [82,83]. Let us consider the exterior spacetime in retarded null coordinates:

$$ds_{\text{out}}^2 = f(u, r_v) du^2 + 2du dr_v - r_v^2 (d\theta^2 + \sin^2 \theta d\phi^2), \tag{39}$$

where $f(u, r_v) = 1 - 2M(r_v, u)/r_v$ with $M(r_v, u)$ being the Vaidya mass. We label the exterior coordinates as $\{X_{\text{out}}^\mu\} \equiv \{u, r_v, \theta, \phi\}$ where u is the retarded null coordinate labeling different shells of radiation and r_v is the Vaidya radius. The above metric is to be matched through the timelike hypersurface Σ given by the condition $r = r_b$ to the interior line element given by (10). The interior coordinates are labeled as $\{X_{\text{in}}^\mu\} \equiv \{t, r, \theta, \phi\}$. The induced metrics from the interior and exterior spacetimes close to Σ then read

$$ds_{\Sigma_{\text{out}}}^2 = dt^2 - a^2(t) r_b^2 (d\theta^2 + \sin^2 \theta d\phi^2) \tag{40}$$

and

$$ds_{\Sigma_{\text{out}}}^2 = [f(u(t), r_v) \dot{u}^2 + 2\dot{r}_v \dot{u}] dt^2 - r_v^2(t) (d\theta^2 + \sin^2 \theta d\phi^2). \tag{41}$$

Matching the induced metrics give

$$f \dot{u}^2 + 2\dot{r}_v \dot{u} = 1, \quad r_v(t) = r_b a(t). \tag{42}$$

The unit vector fields normal to the interior and exterior hypersurfaces can be obtained as

$$n_{\mu}^{\text{in}} = \left[0, \frac{a(t)}{\sqrt{1-kr^2}}, 0, 0 \right],$$

$$n_{\mu}^{\text{out}} = \frac{1}{[f(u, r_v)\dot{v}^2 + 2\dot{r}_v\dot{u}]^{\frac{1}{2}}} [-\dot{r}_v, \dot{u}, 0, 0]. \quad (43)$$

The extrinsic curvature tensors for the interior and exterior spacetimes are given by

$$K_{ab}^{\text{in}} = -n_{\mu}^{\text{in}} \left[\frac{\partial^2 X_{\text{in}}^{\mu}}{\partial y^a \partial y^b} + \hat{\Gamma}_{\nu\sigma}^{\mu \text{in}} \frac{\partial X_{\text{in}}^{\nu}}{\partial y^a} \frac{\partial X_{\text{in}}^{\sigma}}{\partial y^b} \right] \quad (44)$$

and

$$K_{ab}^{\text{out}} = -n_{\mu}^{\text{out}} \left[\frac{\partial^2 X_{\text{out}}^{\mu}}{\partial y^a \partial y^b} + \{\mu\}_{\nu\sigma}^{\text{out}} \frac{\partial X_{\text{out}}^{\nu}}{\partial y^a} \frac{\partial X_{\text{out}}^{\sigma}}{\partial y^b} \right], \quad (45)$$

respectively, where $y^a = \{t, \theta, \phi\}$ are coordinates on the boundary. We note that in computing the components of the extrinsic curvature of the interior spacetime, the general affine connection should be utilized. However, from Eqs. (3–5) together with (8), we see that the affine connection is finally obtained linearly with respect to the spin density tensor. Therefore, by a suitable spacetime averaging, only the Christoffel symbols would remain to be used in (44). The non-vanishing components of the extrinsic curvature tensors then read

$$K_{tt}^{\text{in}} = 0, \quad K_{\theta}^{\theta \text{in}} = K_{\phi}^{\phi \text{in}} = \frac{\sqrt{1-kr_b^2}}{r_b a(t)},$$

$$K_{tt}^{\text{out}} = -\frac{\dot{u}^2 [ff_{,r_v}\dot{u} + f_{,u}\dot{u} + 3f_{,r_v}\dot{r}_v] + 2(\dot{u}\ddot{r}_v - \dot{r}_v\ddot{u})}{2(f\dot{u}^2 + 2\dot{r}_v\dot{u})^{\frac{3}{2}}},$$

$$K_{\theta}^{\theta \text{out}} = K_{\phi}^{\phi \text{out}} = \frac{f\dot{u} + \dot{r}_v}{r_v \sqrt{f\dot{u}^2 + 2\dot{r}_v\dot{u}}}. \quad (46)$$

Matching the components of extrinsic curvatures on the boundary give

$$f\dot{u} + \dot{r}_v = \sqrt{1-kr_b^2}, \quad (47)$$

$$\dot{u}^2 [ff_{,r_v}\dot{u} + f_{,u}\dot{u} + 3f_{,r_v}\dot{r}_v] + 2(\dot{u}\ddot{r}_v - \dot{r}_v\ddot{u}) = 0. \quad (48)$$

A straightforward but lengthy calculation reveals that (48) results in $f(r_v, u) = f(r_v)$ on the boundary [84]. Furthermore from (47) and the first part of (42) we get

$$\dot{r}_v = -(1 - f - kr_b^2)^{\frac{1}{2}}, \quad (49)$$

whence using the second part of (42) we readily arrive at the following equality:

$$M(r_v) = m(t, r_b). \quad (50)$$

Thus from the exterior viewpoint, Eq. (37) implies that there can be found a mass threshold so that, for the mass distributions below such a threshold, the apparent horizon would fail to intersect the surface boundary of the collapsing cloud. Moreover, in view of (36), we observe that as the scale factor increases in the post-bounce regime, the second term

decreases and vanishes at late times. This leaves us with a Schwarzschild exterior spacetime with a constant mass.

4 Concluding remarks

We studied the process of gravitational collapse of a massive star whose matter content is a homogeneous Weyssenhoff fluid in the context of EC theory. Such a fluid is considered as a perfect fluid with spin correction terms that stem from the presence of intrinsic angular momentum of fermionic particles within a real star. The main objective of this paper was to show that, contrary to the OS model, if the spin contributions of the matter sources are included in the gravitational field equations, the scenario of the collapse does not necessarily end in a spacetime singularity. The spin effects can be negligible at the early stages of the collapse, while as the collapse proceeds, these effects would play a significant role in the final fate of the scenario of the collapse. This situation can be compared to the singularity removal for a FLRW spacetime in the very early universe, as we go *backwards in time* [40–54]. We showed that in contrast to the homogeneous dust collapse which leads inevitably to the formation of a spacetime singularity, the occurrence of such a singularity is avoided and instead a bounce occurs at the end of the contracting phase. The whole evolution of the star experiences four phases two of which are in the contracting regime and the other two ones are in the post-bounce regime. While, in the homogeneous dust case without spin correction terms, the singularity is necessarily dressed by an event horizon, the formation of such a horizon can always be prevented by suitably choosing the surface boundary of the collapsing star. This signals that there exists a critical threshold value for the mass content, below which no horizon would form. The same picture can be found in [85] where the non-minimal coupling of gravity to fermions is allowed. Besides the model presented here, non-singular scenarios have been reported in the literature within various models such as $f(R)$ theories of gravity in the Palatini [86] and metric [87] formalisms, non-singular cosmological settings in the presence of a spinning fluid in the context of EC theory [88], bouncing scenarios in brane models [89–93], and modified Gauss–Bonnet gravity [94] (see also [95] for recent review). While the spacetime singularities could generically occur as the end-product of a continual gravitational collapse, it is widely believed that in the very final stages of the collapse where the scales are comparable to the Planck length and extreme gravity regions are dominant, quantum corrections could generate a strong negative pressure in the interior of the cloud to finally resolve the classical singularity [96–107]. Finally, as we come near the end of this paper we should point out that quantum effects due to particle creation could possibly avoid the cosmological [108–111] as well as astrophysical singularities [112].

Acknowledgments The authors would like to sincerely thank the anonymous referee for constructive and helpful comments to improve the original manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

Funded by SCOAP³ / License Version CC BY 4.0.

References

- J. Oppenheimer, H. Snyder, Phys. Rev. D **56**, 455 (1939)
- P.S. Joshi, *Gravitational Collapse and Space-Time Singularities*. (Cambridge University Press, London, 2007)
- R. Penrose, Phys. Rev. Lett. **14**, 57 (1965)
- S.W. Hawking, Proc. R. Soc. Lond. A **300**, 187 (1967)
- S.W. Hawking, R. Penrose, Proc. R. Soc. Lond. A **314**, 529 (1970)
- S.W. Hawking, G.F.R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, 1973)
- V.P. Frolov, I.D. Novikov, *Black Hole Physics: Basic Concepts and New Developments*. Fundamental Theories of Physics (Springer, Berlin, 1998)
- R. Penrose, Riv. Nuovo Cimento **1**, 252 (1969)
- R. Penrose, in *General relativity, an Einstein Centenary Survey*, ed. by S.W. Hawking, W. Israel (Cambridge University Press, Cambridge, 1973)
- R. Penrose, in *Gravitational Radiation and Gravitational Collapse*, ed. by C. DeWitt-Morette. IAU Symposium, vol. 64 (Reidel, Dordrecht, 1974)
- R.M. Wald. [arXiv:gr-qc/9710068](https://arxiv.org/abs/gr-qc/9710068)
- D.M. Eardley, L. Smarr, Phys. Rev. D **19**, 2239 (1979)
- D. Christodoulou, Commun. Math. Phys. **93**, 171 (1984)
- R.P.A.C. Newman, Class. Quantum Gravity **3**, 527 (1986)
- B. Waugh, K. Lake, Phys. Rev. D **38**, 1315 (1988)
- G. Grillo, Class. Quantum Gravity **8**, 739 (1991)
- I.H. Dwivedi, P.S. Joshi, Class. Quantum Gravity **9**, L69 (1992)
- T.P. Singh, P.S. Joshi, Class. Quantum Gravity **13**, 559 (1996)
- P.S. Joshi, I.H. Dwivedi, Phys. Rev. D **47**, 5357 (1993)
- P.S. Joshi, T.P. Singh, Phys. Rev. D **51**, 6778 (1995)
- L. Herrera, A. Di Prisco, J.L. Hernandez-Pastora, N.O. Santos, Phys. Lett. A **237**, 113 (1998). [arXiv:gr-qc/9711002](https://arxiv.org/abs/gr-qc/9711002)
- P.S. Joshi, D. Malafarina, Gen. Relativ. Gravit. **45**, 305 (2012). [arXiv:1105.4336](https://arxiv.org/abs/1105.4336)
- O. Bergmann, Phys. Lett. A **82**, 383 (1981)
- S.R. Maiti, Phys. Rev. D **25**, 2518 (1982)
- A.K. Sanyal, D. Ray, J. Math. Phys. **25**, 1975 (1984)
- B. Modak, J. Astrophys. Astron. **5**, 317 (1984)
- P.S. Joshi, R. Goswami, N. Dadhich. [arXiv:gr-qc/0308012](https://arxiv.org/abs/gr-qc/0308012)
- S.M.C.V. Goncalves, Phys. Rev. D **69**, 021502(R) (2004). [arXiv:gr-qc/0312123](https://arxiv.org/abs/gr-qc/0312123)
- P.S. Joshi, N. Dadhich, R. Maartens, Phys. Rev. D **65**, 101501 (2002). [arXiv:gr-qc/0109051](https://arxiv.org/abs/gr-qc/0109051)
- F.C. Mena, B.C. Nolan, R. Tavakol, Phys. Rev. D **70**, 084030 (2004). [arXiv:gr-qc/0405041](https://arxiv.org/abs/gr-qc/0405041)
- A.H. Ziaie, K. Atazadeh, S.M.M. Rasouli, Gen. Relativ. Gravit. **43**, 2943 (2011). [arXiv:1106.5638](https://arxiv.org/abs/1106.5638)
- N. Bedjaoui, P.G. LeFloch, J.M. Martin-Garcia, J. Novak, Class. Quantum Gravity **27**, 245010 (2010). [arXiv:1008.4238](https://arxiv.org/abs/1008.4238)
- P.S. Joshi, Pramana **55**, 529 (2000). [arXiv:gr-qc/0006101](https://arxiv.org/abs/gr-qc/0006101)
- P.S. Joshi, D. Malafarina, Int. J. Mod. Phys. D **20**, 2641 (2011). [arXiv:1201.3660](https://arxiv.org/abs/1201.3660)
- D.Z. Freedman, A. Van Proeyen, *Supergravity* (Cambridge University Press, Cambridge, 2012)
- T. Ortin, *Gravity and Strings* (Cambridge University Press, Cambridge, 2004)
- M. Blagojevic, F.W. Hehl, *Gauge Theories of Gravitation a Reader with Commentaries* (World Scientific Publishing Co., Inc, Singapore; reprint edition Imperial College Press, London, 2013)
- F.W. Hehl, P. Von der Heyde, G.D. Kerlick, J.M. Nester, Rev. Mod. Phys. **48**, 393 (1976)
- F.W. Hehl, P. Von der Heyde, G.D. Kerlick, Phys. Rev. D **10**, 1066 (1974)
- W. Kopczynski, Phys. Lett. A **39**, 219 (1972)
- W. Kopczynski, Phys. Lett. A **43**, 63 (1973)
- J. Stewart, P. Hajicek, Nature (Phys. Sci.) **242**, 7 (1973)
- J. Tafel, Phys. Lett. A **45**, 341 (1973)
- A. Trautman, Nature, (Phys. Sci.) **242**, 7 (1973)
- A.K. Raychaudhuri, Phys. Rev. D **12**, 952 (1975)
- G.D. Kerlick, Ann. Phys. **99**, 127 (1976)
- B. Kuchowicz, Astrophys. Space Sci. **39**, 157 (1976)
- B. Kuchowicz, Gen. Relativ. Gravit. **9**, 511 (1978)
- M. Gasperini, Gen. Relativ. Gravit. **30**, 1703 (1998). [arXiv:gr-qc/9805060](https://arxiv.org/abs/gr-qc/9805060)
- N.J. Poplawski, Gen. Relativ. Gravit. **44**, 1007 (2012). [arXiv:1105.6127](https://arxiv.org/abs/1105.6127)
- B. Vakili, S. Jalalzadeh, Phys. Lett. B **28**, 726 (2013). [arXiv:1308.2877](https://arxiv.org/abs/1308.2877)
- J.-A. Lu. [arXiv:1404.4440](https://arxiv.org/abs/1404.4440) [gr-qc]
- N.J. Poplawski, Phys. Rev. D **85**, 107502 (2012). [arXiv:1111.4595](https://arxiv.org/abs/1111.4595)
- N.J. Poplawski, Astron. Rev. **8**, 108 (2013). [arXiv:1106.4859](https://arxiv.org/abs/1106.4859)
- M. Bojowald, R. Goswami, R. Maartens, P. Singh, Phys. Rev. Lett. **95**, 091302 (2005). [arXiv:gr-qc/0503041](https://arxiv.org/abs/gr-qc/0503041)
- C. Bambi, D. Malafarina, L. Modesto, Eur. Phys. J. C **74**, 2767 (2014). [arXiv:1306.1668](https://arxiv.org/abs/1306.1668)
- M. Bojowald, Phys. Rev. Lett. **86**, 5227 (2001). [arXiv:gr-qc/0102069](https://arxiv.org/abs/gr-qc/0102069)
- M. Bojowald, Nat. Phys. **3**, 523 (2007)
- P. Singh, K. Vandersloot, G.V. Vereshchagin, Phys. Rev. D **74**, 043510 (2006). [arXiv:gr-qc/0606032](https://arxiv.org/abs/gr-qc/0606032)
- T. Cailleteau, A. Cardoso, K. Vandersloot, D. Wands, Phys. Rev. Lett. **101**, 251302 (2008). [arXiv:0808.0190](https://arxiv.org/abs/0808.0190)
- A. Ashtekar, D. Sloan, Phys. Lett. B **694**, 108 (2010). [arXiv:0912.4093](https://arxiv.org/abs/0912.4093)
- V. De Sabbata, M. Gasperini, *Introduction to Gravitation* (World Scientific, Singapore, 1986)
- V. De Sabbata, C. Sivaram, *Spin and Torsion in Gravitation* (World Scientific, Singapore, 1994)
- V. De Sabbata, C. Sivaram, Astrophys. Space Sci. **165**, 51 (1990)
- N.J. Poplawski, Phys. Lett. B **694**, 181 (2010). [arXiv:1007.0587](https://arxiv.org/abs/1007.0587)
- N.J. Poplawski. [arXiv:1410.3881](https://arxiv.org/abs/1410.3881) [gr-qc]
- J. Weyssenhoff, A. Raabe, Acta Phys. Pol. **9**, 7 (1947)
- J.R. Ray, L.L. Smalley, Phys. Rev. D **27**, 1383 (1983)
- G.A. Maugin, Ann. Inst. Henri Poincare **20**, 41 (1974)
- Y.N. Obukhov, V.A. Korotky, Class. Quantum Gravity **4**, 1633 (1987)
- M. Gasperini, Phys. Rev. Lett. **56**, 26 (1986)
- B. Datt, Z. Physik. **108**, 314 (1938)
- A. Fabbri, J. Navarro-Salas, *Modeling Black Hole Evaporation* (ICP-World Scientific, London, 2005)
- P.S. Joshi, D. Malafarina, R.V. Saraykar, Int. J. Mod. Phys. D **21**, 1250066 (2012)
- I.S. Nurgaliev, W.N. Ponomarev, Phys. Lett. B **130**, 6 (1983)
- Y. Nambu, M. Sasaki, Prog. Theor. Phys. **79**, 1 (1988)
- C. Bambi, D. Malafarina, L. Modesto, Phys. Rev. D **88**, 044009 (2013). [arXiv:1305.4790](https://arxiv.org/abs/1305.4790)
- C.W. Misner, D.H. Sharp, Phys. Rev. **136**, B571 (1964)
- S.A. Hayward, Phys. Rev. D **53**, 1938 (1996)
- S.A. Hayward, Phys. Rev. D. **49**, 6467 (1994)
- T. Koike, H. Onozawa, M. Siino. [arXiv:gr-qc/9312012](https://arxiv.org/abs/gr-qc/9312012)

82. N.O. Santos, Mon. Not. R. Astron. Soc. **216**, 403 (1985)
83. W.B. Bonnor, A.K.G. de Oliveira, N.O. Santos, Phys. Rep. **181**, 269 (1989)
84. B.K. Tippett, V. Husain, Phys. Rev. D **84**, 104031 (2011). [arXiv:1106.1118](#)
85. C. Bambi, D. Malafarina, A. Marciano, L. Modesto, Phys. Lett. B **734**, 27 (2014). [arXiv:1402.5719](#)
86. C. Barragan, G.J. Olmo, H. Sanchis-Alepuz, Phys. Rev. D **80**, 024016 (2009). [arXiv:0907.0318](#)
87. N. Paul, S. N. Chakrabarty, K. Bhattacharya, JCAP **1410**, 10,009 (2014). [arXiv:1405.0139](#)
88. S.D. Brechet, M.P. Hobson, A.N. Lasenby, Class. Quantum Gravity **24**, 6329 (2007). [arXiv:0706.2367](#)
89. S. Mukherji, M. Peloso, Phys. Lett. B **547**, 297 (2002). [arXiv:hep-th/0205180](#)
90. E.M. Prodanov, Phys. Lett. B **530**, 210 (2002). [arXiv:hep-th/0103151](#)
91. Y. Shtanov, V. Sahni, Phys. Lett. B **557**, 1 (2003). [arXiv:gr-qc/0208047](#)
92. R. Maier, I.D. Soares, E.V. Tonini, Phys. Rev. D **79**, 023522 (2009). [arXiv:1212.3766](#)
93. R. Maier, N. Pinto-Neto, I.D. Soares, Phys. Rev. D **87**, 043528 (2013). [arXiv:1301.5250](#)
94. K. Bamba, A.N. Makarenko, A.N. Myagky, S.D. Odintsov, Phys. Lett. B **732**, 349 (2014). [arXiv:1403.3242](#)
95. M. Novello, S.E.P. Bergliaffa, Phys. Rep. **463**, 127 (2008). [arXiv:0802.1634](#)
96. V.P. Frolov, G.A. Vilkovisky, Phys. Lett. B **106**, 307 (1981)
97. P. Singh, A. Toporensky, Phys. Rev. D **69**, 104008 (2004). [arXiv:gr-qc/0312110](#)
98. V. Husain, O. Winkler, Class. Quantum Gravity **22**, L127 (2005). [arXiv:gr-qc/0410125](#)
99. R. Goswami, P.S. Joshi, P. Singh, Phys. Rev. Lett. **96**, 031302 (2006). [arXiv:gr-qc/0506129](#)
100. E. Greenwood, D. Stojkovic, JHEP **0806**, 042 (2008). [arXiv:0802.4087](#)
101. M. Bojowald, G.M. Paily, Class. Quantum Gravity **29**, 242002 (2012). [arXiv:1206.5765](#)
102. A. Kreienbuehl, T. Pawlowski, Phys. Rev. D **88**, 043504 (2013). [arXiv:1302.6566](#)
103. Y. Tavakoli, J. Marto, A. Dapor, Int. J. Mod. Phys. D **23**, 1450061 (2014). [arXiv:1303.6157](#)
104. Y. Tavakoli, J. Marto, A. Dapor, Springer Proc. Math. Stat. **60** (2014). doi:10.1007/978-3-642-40157-2
105. C. Rovelli, F. Vidotto. [arXiv:1401.6562](#)
106. Y. Zhang, Y. Zhu, L. Modesto, C. Bambi. [arXiv:1404.4770](#)
107. Y. Liu, D. Malafarina, L. Modesto, C. Bambi, Phys. Rev. D **90**, 044040 (2014). [arXiv:1405.7249](#)
108. L. Parker, S.A. Fulling, Phys. Rev. D **7**, 2357 (1973)
109. T.P. Singh, Gen. Relativ. Gravit. **30**, 1563 (1998)
110. J. Haro. [arXiv:1011.4772](#)
111. M.J.S. Houndjo, A.V. Monwanou, J.B. Chabi Orou, Int. J. Mod. Phys. D **20**, 2449 (2011). [arXiv:1105.3165](#)
112. Laura Mersini-Houghton, Phys. Lett. B **738**, 61 (2014). [arXiv:1406.1525](#)