



REVIEW ARTICLE

Colorectal cancer in inflammatory bowel disease: Results of the 3rd ECCO pathogenesis scientific workshop (I)



Shaji Sebastian^{a,*}, Vincent Hernández^b, Pär Myrelid^c, Revital Kariv^d, Epameinondas Tsianos^e, Murat Toruner^f, Marc Marti-Gallostra^g, Antonino Spinelli^h, Andrea E. van der Meulen-de Jongⁱ, Elif Saritas Yuksel^j, Christoph Gasche^k, Sandro Ardizzone^l, Silvio Danese^{m,**}

^a Hull & East Yorkshire Hospitals NHS Trust, Hull York Medical School, Hull, United Kingdom

^b Gastroenterology Department, Complejo Hospitalario Universitario de Vigo, Vigo, Spain

^c Division of Surgery, Department of Clinical and Experimental Medicine, Faculty of Health Sciences, Linköping University, County Council of Östergötland, Linköping, Sweden

^d Service for Gastrointestinal Malignancies, Department of Gastroenterology & Liver Disease, Tel Aviv Sourasky Medical Center, Israel

^e University of Ioannina, 1st Division of Internal Medicine and Hepato-Gastroenterology Unit, Greece

^f Ankara University Medical School, Ibni Sina Hospital, Division of Gastroenterology, Ankara, Turkey

^g Department of Colorectal Surgery, University Hospital of Valle de Hebron, Barcelona, Spain

^h Dipartimento e Cattedra di Chirurgia Generale, Istituto Clinico Humanitas IRCCS, Università degli Studi di Milano, Rozzano, Milano, Italy

ⁱ Leiden University Medical Center, Department of Gastroenterology, Leiden, The Netherlands

^j Department of Gastroenterology, Katip Celebi University, Ataturk Research and Teaching Hospital, Izmir, Turkey

^k Christian Doppler Laboratory on Molecular Cancer Chemoprevention, Division of Gastroenterology, Medical University of Vienna, Vienna, Austria

^l Chair of Gastroenterology, "L. Sacco" University Hospital, Milan, Italy

^m Department of Gastroenterology, Istituto Clinico Humanitas, Milan, Italy

Received 23 March 2013; accepted 5 April 2013

Abbreviations: AOM, Azoxymethane; IBD, Inflammatory bowel disease; UC, Ulcerative colitis; CD, Crohn's disease; CRC, Colorectal cancer; CAC, Colitis-associated cancer; CIN, Chromosomal instability; IL, Interleukins; DSS, Dextran sodium sulphate; IC, Indeterminate colitis; PSC, Primary sclerosing cholangitis; OR, Odds ratio; SIR, Standardised incidence ratio; RR, Relative risk; MRR, Mortality rate ratio; TLR, Toll-like receptor; NF- κ B, Nuclear factor- κ B; TGF β , Transforming growth factor- β ; MSI, Microsatellite instability; TNF α , Tumour necrosis factor- α ; 5-ASA, 5-amino salicylic acid.

* Correspondence to: S. Sebastian, Hull & East Yorkshire Hospitals NHS Trust, Hull York Medical School, Anlaby Road, Hull, United Kingdom.

** Correspondence to: S. Danese, IBD Center, Department of Gastroenterology, Istituto Clinico Humanitas, Via Manzoni 56, 20089 Rozzano, Milan, Italy.

E-mail addresses: Shaji.Sebastian@hey.nhs.uk (S. Sebastian), sdanese@hotmail.com (S. Danese).

KEYWORDS

Colorectal cancer;
Epidemiology;
Chemoprevention;
Pathogenesis;
Outcomes

Abstract

Epidemiological studies demonstrate an increased risk of colorectal cancer in patients with inflammatory bowel disease (IBD). A detailed literature review was conducted on epidemiology, risk factors, pathophysiology, chemoprevention and outcomes of colorectal cancer (CRC) in IBD as part of the 3rd ECCO scientific pathogenesis workshop.

Crown Copyright © 2013 Published by Elsevier B.V. on behalf of European Crohn's and Colitis Organisation. All rights reserved.

Contents

1. Introduction	6
2. Epidemiology of colorectal cancer in IBD	6
2.1. Methodological considerations	6
2.2. Epidemiology of colorectal cancer in ulcerative colitis	7
2.3. Epidemiology of colorectal cancer in Crohn's disease	7
2.4. Colorectal cancer in indeterminate colitis	7
3. Risk factors for CRC in patients with IBD	7
3.1. Duration and extent	8
3.2. Age of onset	8
3.3. Primary sclerosing cholangitis	8
3.4. Other risk factors	8
4. The pathophysiology of colitis-associated cancer	8
4.1. Regulatory signals in colitis-associated cancer (CAC)	8
4.1.1. Toll-like receptors (TLRs)	8
4.1.2. Cytokines	8
4.1.3. Chemokines	8
4.1.4. Negative regulators	9
4.2. Molecular pathways of CAC	10
4.3. The field effect	10
5. Outcome of colorectal cancer in IBD	10
6. Chemoprevention of colorectal cancer in IBD	11
6.1. 5-Aminosalicylates	11
6.2. Ursodeoxycholic acid	12
6.3. Immunomodulators	12
6.4. Folic acid	13
6.5. Statins	13
6.6. Chemoprevention in CD	13
7. Conclusions and areas for future research	13
Conflict of interest	13
References	14

1. Introduction

We conducted a detailed literature review on epidemiology, risk factors, pathophysiology, chemoprevention and outcomes of colorectal cancer (CRC) in inflammatory bowel disease (IBD) as part of the 3rd ECCO scientific pathogenesis workshop. The results of a literature search are presented here.

2. Epidemiology of colorectal cancer in IBD**2.1. Methodological considerations**

IBD is associated with an increased risk of CRC, which is thought to be primarily related to long-standing chronic

inflammation.¹ However, the magnitude of the risk has been difficult to estimate, as many factors may bias study results² including patient selection, number of patients, completeness of case recruitment and ascertainment and duration of follow-up.^{2,3}

Early studies from tertiary referral centres reported a high risk of CRC, but as more severe cases were included, they tended to overestimate the risk. In contrast, population-based studies covering defined geographical areas report a more conservative risk. However, these studies probably included patients with limited and less severe disease, and may have underestimated the risk. In addition, geographical differences have been described, with an increased incidence of IBD-related CRC reported in the USA and in the UK compared with Scandinavia.⁴ Similarly, the 10-year follow-up of the European

Collaborative IBD inception cohort showed that Northern European centres had higher rates of IBD-related CRC than Southern centres (prevalence 0.9% vs. 0.25%, respectively; $p = 0.17$),⁵ although this North-South gradient was not apparent at 15 year follow-up.⁶ Similarly, studies from Asian countries have shown important differences between countries.^{7–10}

2.2. Epidemiology of colorectal cancer in ulcerative colitis

In 2001, a meta-analysis of cohort and case-controlled studies⁴ on the risk of CRC in UC estimated an incidence rate of 3/1000 person years disease duration (PYD), an annual risk of 0.3% and a cumulative risk of 18.4% after 30 years of disease. The incidence rate progressively increased each decade from 2/1000 PYD to 7/1000 for the second and 12/1000 for the third decade.

More recent population-based studies and meta-analyses have shown that the risk is lower than previously described or even similar to that of the general population.^{4,11–22} (Table 1). This could be explained by methodological aspects (hospital-based vs. population-based studies), by a true decrease in risk due to better disease control, higher colectomy rates, use of drugs with chemoprotective effects or more widespread endoscopic surveillance in high-risk patients.²

Castañó-Milla et al.²¹ reported an overall incidence rate of 1.67/1000 PYD and incidence rates per decade were estimated at 1.01/1000, 3.75/1000 and 5.85/1000 PYD for the first, second and third decades, respectively. In a meta-analysis of prospective population-based studies, Jess et al.,²² found that an average of 1.6% of patients with UC were diagnosed with CRC during the first 14 year follow-up, and the estimated standardised incidence ratio (SIR) was 2.39 (2.1–2.7).

Recent time-trend studies also demonstrate a decreasing risk of CRC in UC patients. Jess et al.¹⁹ found that the relative risk (RR) decreased from 1.34 between 1979 and 1988 to 0.57 between 1999 and 2008. In a meta-analysis by Castañó-Milla et al.,²¹ the incidence rate was found to decrease from 4.29/1000 PYD in studies published in the 1950s to 1.09/1000 PYD in the studies published between 2000 and 2011.

2.3. Epidemiology of colorectal cancer in Crohn's disease

The role of Crohn's disease (CD) as a risk factor of CRC is controversial and, compared with UC, the risk is modest. In a meta-analysis of population-based studies in CD, Jess et al.²³ estimated a pooled SIR for CRC of 1.9 (1.4–2.5). Canavan et al.²⁴ found a RR of 2.5 (1.3–4.7) among all CD patients, with a higher risk for CD-colitis (RR 4.5 (1.3–14.9)); this study estimated a cumulative risk of 2.9% at 10 years, 5.6% at 20 years and 8.3% at 30 years. A similar risk (RR 2.44 (1.56–3.82)) was reported in a meta-analysis by von Roon et al.²⁵ Laukoetter et al.²⁶ reported an incidence of 0.5/1000 PYD, a 2- to 3-fold increase over the incidence of CRC in an age-matched background population.

Friedman et al. found dysplasia (low or high grade) or CRC in 18.5% of patients, and after a negative screening colonoscopy the probability of developing HGD or CRC was 7% by the 10th surveillance examination.²⁷ Basseri et al. detected LGD, HGD or CRC in 5.6% of patients,²⁸ but in this cohort the number of patients with extensive colitis was lower than in Friedman's (55% vs. 90%).

2.4. Colorectal cancer in indeterminate colitis

Only one study evaluating the epidemiology of CRC in indeterminate colitis (IC) was identified.²⁹ Stewenius et al., in 1995, followed an inception cohort of 471 UC patients and 100 IC patients for a mean of 14.8 years. Three of 100 patients with IC developed CRC, an incidence of 2.4/1000 PYD (SIR 8.6, 95% CI 1.8–25.1), compared with an incidence of 1.4/1000 PYD in patients with UC.

3. Risk factors for CRC in patients with IBD

Reported risk factors for CRC include extensive disease,^{4,30} young age at diagnosis,^{4,31} family history of CRC,³² co-existing primary sclerosing cholangitis (PSC)³³ and persistent inflammation of the colon.^{34–36}

Table 1 CRC risk in UC patients. Population-based studies.

	Annual incidence	Cumulative incidence	Risk (95% CI)
Palli ¹¹	0.12%		1.79 (0.85–3.28)
Bernstein ¹²	Colon 0.16%		Colon, RR 2.75 (1.91–3.97)
	Rectum 0.06%		Rectum, RR 1.90 (1.05–3.43)
Winther ¹³	0.06%	2.1% at 30 years	SIR 1.05 (0.56–1.79)
Jess ¹⁴	0.10%	2% at 25 years	SIR 1.1 (0.4–2.4)
Lakatos ¹⁵	0.15%	7.5% at 30 years	SIR 1.74 (1.01–3.0)
Jess ¹⁶			SIR 3.1 (0.1–17)*
Soderlund ¹⁷		3.5% at 30 years**	SIR 2.7 (2.3–3.2)
Jess ¹⁹	0.051%		RR 1.07 (0.95–1.21)

RR: Relative risk. SIR: Standardised incidence ratio.

* Danish EC-IBD Cohort.

** Estimated from Kaplan–Meier curve.

3.1. Duration and extent

A meta-analysis by Eaden⁴ shows that in UC patients the risk of CRC becomes appreciable after 10 years, and it is assumed that risk increases with disease duration. Guidelines recommend that CRC screening be started 6–10 years after onset of symptoms in extensive colitis.^{37–39} However, CRC may develop earlier, with some patients diagnosed simultaneously with CRC and IBD, and about 9–15% presenting with CRC with a disease duration of <6 years.⁴⁰ However, data from the St Mark's Hospital London surveillance cohort report that the incidence of CRC remained constant or decreased with increasing disease duration.⁴¹ In a Swedish cohort of 7607 IBD patients followed from 1960 to 2004, the incidence of CRC (adjusted for the type and extent of IBD, sex, age and time since diagnosis) did not show any statistically significant trend over successive calendar periods of follow-up.¹⁷

The risk of CRC has been found to be higher in extensive UC than in left-sided UC, while in proctitis it is similar to that of the general population,³⁰ and this has been confirmed in a recent meta-analysis.²² However, Soderlund et al.,¹⁷ analysing an original Scandinavian cohort,³⁰ found that the RR of cancer in patients with proctitis was 1.7 (95% CI, 1.2–2.4) compared with that of the general population. Some authors suggested a possible proximal extension of the disease in patients with an initial rectal involvement.⁴² In Crohn's colitis, the risk of CRC is similar to that of UC if the extension and duration are comparable.^{43–46}

3.2. Age of onset

In the original meta-analysis by Eaden et al., young age at onset of IBD showed a trend towards a slightly increased risk of CRC, and the incidence rate of CRC in UC diagnosed in childhood was higher than the incidence in adults⁴; this has been confirmed in a more recent meta-analysis.²² In CD, young age has also been associated with an increased risk of CRC.³¹ However, this has not been confirmed in other studies, with some even suggesting that CRC may develop earlier in patients diagnosed with IBD at older ages.^{47–49}

3.3. Primary sclerosing cholangitis

PSC is a recognized risk factor for CRC in patients with colitis, with a 4-fold increase in risk compared with UC patients without PSC.^{15,19,33} Dysplasia or CRC have also been found to appear soon after the diagnosis of both diseases in some patients (21.5 per 100 patients-years of follow-up during the first two years).⁵⁰ Moreover, current evidence emphasizes that the risk of CRC remains after liver transplantation.^{51,52}

PSC can also be associated with Crohn's colitis or IC, and it may also carry an increased risk of CRC, especially if colitis is extensive.⁵³ Two retrospective case-control studies have reported conflicting results, with a Swedish study reporting an increased risk of CRC in CD-PSC (odds ratio [OR] 6.78, 95% CI 1.65–27.9),⁵⁴ and a British study reporting no increase in risk (OR 1.64, 95% CI 0.14–18.7).⁵⁵

3.4. Other risk factors

Soderlund et al.,⁵⁶ in the previously described cohort, found that males had a 60% higher risk of CRC (RR 1.6, 1.2–2.2), and a greater cumulative incidence after 40 years of disease (8.3% vs. 3.5%); the effect of gender was limited to patients with more than 10 years' of follow-up and in those aged <45 years at diagnosis. No differences have been found in CRC incidence between Caucasian and African-American IBD patients.⁵⁷

4. The pathophysiology of colitis-associated cancer

4.1. Regulatory signals in colitis-associated cancer (CAC)

Numerous positive and negative regulators have been implicated in the development of CRC (see Table 2).

4.1.1. Toll-like receptors (TLRs)

A deficiency in the TLR adaptor, MyD88, has been found to significantly reduce tumour number and size in the *Apcmin/+* mouse model of intestinal tumorigenesis^{58,59} and the absence of bacteria in mouse models of inflammation-associated cancer has been shown to prevent dysplasia or cancer.⁶⁰

4.1.2. Cytokines

4.1.2.1. Tumour necrosis factor- α . The absence of tumour necrosis factor- α (TNF α) signalling in mice has been found to be associated with decreases in mucosal damage, inflammatory cell infiltrates, and cytokine expression, in response to azoxymethane (AOM) challenge and dextran sodium sulphate (DSS), as well as a significant reduction in tumour formation.⁶¹

4.1.2.2. Interleukin-6. High levels of interleukin-6 (IL-6) and a soluble IL-6 receptor in IBD patients promote the accumulation of T cells in the lamina propria of the colon by upregulating anti-apoptotic factors.^{62,63} Moreover, increased IL-6 levels have been observed in the serum of cancer patients and in tumour biopsies.⁶⁴ Finally, expression of IL-6 and STAT3 has been shown to be greater in active UC and after progression to colitis-associated cancer (CAC) than in controls or inactive disease.⁶⁵

4.1.2.3. Interleukin-21. Interleukin-21 (IL-21) was recently found to be elevated in the gut of patients with CAC, and in mice with CAC induced by AOM and DSS.⁶⁶ IL-21-deficient mice also developed fewer and smaller tumours than wild-type mice.⁶⁷

4.1.3. Chemokines

The atypical chemokine receptor D6 selectively recognizes and efficiently depletes most inflammatory CC chemokines from the extracellular milieu.⁶⁸ Furthermore, expression of D6 is increased in the lymphatic vascular bed of colonic sections with IBD-associated colon cancer.⁶⁹

TNF α induces the expression of chemokines from a variety of colonic cells.⁷⁰ Indeed, the mucosa of IBD patients

Table 2 Regulatory signals in colitis-associated cancer.

	Effect	Example	Reference
<i>Positive regulators</i>			
Toll like receptors	Promote carcinogenesis	MyD88 deficiency reduces tumour number and size in the APC ^{min} mice	58–60
Procarcinogenic cytokines	Promote carcinogenesis	Tumour necrosis factor IL6	61–65
Chemokines	Promote carcinogenesis Direct neoplasia suppression-decreased COX-2, attenuated neovascularization	Atypical chemokine receptor D6 CCL2	68,69 62,72,73
<i>Negative regulators</i>			
IL-10	Prevent IBD related colonic neoplasia, through direct inflammation suppression and through IL-6		73
Transforming growth factor- β (TGF β) family	Neoplasia inhibition directly and through inflammation suppression	Transforming growth factor- β (TGF β) family, SMAD-7	74,75
Toll-IL-1R8	Inhibits IL-1 and TLR complexes signals, lack of NF- κ B activation		76–78
Nuclear factor- κ B (NF- κ B)	Promotes cell proliferation and survival.		79–83
<i>Molecular pathways</i>			
Chromosomal instability (CIN) pathway	Prominent neoplastic pathway in IBD-related neoplasia, cause unclear	P53 mutations—common,DPC4 and KRAS—less common	89,92,93
Microsatellite instability pathway	Variable frequency, found in inflamed, non-neoplastic mucosa	MLH1-promoter methylation, TGFBR2 and ACVR2 mutation—lower than in sporadic CRC	94–99
Methylation pathway (CpG island methylation)	Questionable role in IBD related neoplasia	p16 INK4 and p14 ARF methylation.	95,100–102
Serrated neoplasia pathway	Present in some IBD-related neoplasia including inflammatory mucosa—contribution unclear	V600E BRAF mutation, serrated lesions in histology	103
Field effect	Multifocal risk for neoplastic transformation as a result of chronic inflammation and dysbiosis that affects mutation pressure	Markers: aneuploidy, p53 mutations,CpG island hypermethylation	89,104,106,107

has been shown to have enhanced mRNA and protein expression of MCP-1/CCL2,⁷¹ a CC chemokine with potent chemotactic and activating activities for monocytes/macrophages. CCL2 mRNA expression has been shown to be enhanced in the colon of the AOM-challenged mouse model, and blocking the TNF α /TNFR axis reduced colorectal carcinogenesis, intracolonic macrophage infiltration, and CCL2 mRNA expression.⁷²

4.1.4. Negative regulators

4.1.4.1. Interleukin-10. The demonstration that interleukin-10 (IL-10) suppresses IL-6 and controls CAC development in a model of microbial CRC indicates that the control of CAC development by IL-10 may be at least partially mediated through IL-6. Moreover, it was recently demonstrated that patients with IL-10 receptor mutations that abrogate IL-10 signalling develop early and aggressive disease.⁷³

4.1.4.2. Transforming growth factor- β . Transforming growth factor- β (TGF β) receptor-deficient transgenic mice were found to develop significantly more tumours than wild-type mice, while the opposite occurred in TGF β transgenic mice.⁷⁴ Additionally, mechanistic studies showed that TGF β inhibited IL-6-dependent tumour formation, reinforcing the concept

that immunoregulatory molecules such as IL-10 and TGF β have a protective effect on intestinal inflammation and in CAC formation.⁷⁴ Supporting these data, mice transgenic for the TGF β inhibitor smad7 with over-expression specifically on T cells, developed more severe colitis in response to AOM/DSS challenge, but only a few small tumors.⁷⁵

4.1.4.3. Toll-IL-1R8. A recently identified orphan member of the IL-1 receptor family, Toll-IL-1R8 (TIR8), inhibits signalling from IL-1R/TLR complexes,^{76,77} and gene transfer experiments have revealed that it reduces the activation of NF- κ B by the IL-1R complex⁷⁷ and by members of the TLR family.⁷⁶ Moreover, TIR8-deficient mice administered DSS show dramatic weight loss, intestinal bleeding and increased mortality, while susceptibility to CAC in the AOM challenge model was increased.⁷⁸

4.1.4.4. Nuclear factor- κ B. Activation of nuclear factor- κ B (NF- κ B) plays a key role in inflammation and cell proliferation and survival.⁷⁹ Regulation of NF- κ B involves sequestration of NF- κ B in the cytoplasm by the inhibitor I κ B, which is regulated through phosphorylation by the I κ B kinase (IKK β) complex, targeting it for ubiquitin-dependent degradation.⁸⁰

Greten et al.⁸¹ demonstrated that IKK β links inflammation and tumorigenesis in a mouse model of CAC in which IKK β was

deleted either in intestinal epithelial cells or in myeloid cells. NF- κ B has also been shown to induce vascular endothelial growth factor and COX-2, which promote angiogenesis.⁸⁰ In addition, a subgroup of NLR proteins, including NLRP12, inhibits transcription factor NF- κ B, and it was recently reported that NLRP12 ($-/-$) mice were highly susceptible to colitis and CAC, suggesting that NLRP12 is a checkpoint of noncanonical NF- κ B, inflammation and tumorigenesis.^{82,83}

4.1.4.5. Intestinal microbiota. Recent studies point to alterations in the luminal microbiota in patients with IBD. Adherent-invasive *Escherichia coli* is associated both with IBD and CRC.^{84,85} A more recent approach demonstrated microbial changes in tumour-developing IL-10 $^{-/-}$ mice with the prevalence of *E. coli*.⁸⁶ Deletion of the bacterial polyketide synthase (pks) genotoxic island from *E. coli* NC101 reduces the development of dysplasia and cancer, independent of inflammation.

4.2. Molecular pathways of CAC

CRC has been considered to develop through well defined pathways, including chromosomal instability (CIN), microsatellite instability (MSI) and, most recently, the CIMP pathway.⁸⁷ There have been previous attempts to distinguish IBD related to sporadic colonic neoplasia⁸⁸ with some success. A few studies looked at different markers and genes related to CIN, such as aneuploidy, loss of heterozygosity (LOH) and genomic gains or losses of various genes^{89–91} but observations are not consistent.

Notably, p53 mutations occur earlier in IBD-related neoplasia, probably a key initiating event,⁹² than APC mutations, which occur later than commonly observed in sporadic colorectal neoplasia.^{93,94} Allelic deletions of p53 were observed in approximately 50% to 85% of colitis-associated tumours.⁹⁵ p53 LOH has been shown to correlate with malignant potential in the colon. Mutations in some other commonly mutated CIN pathway genes of sporadic CRC such as DPC4 and KRAS were uncommon in CAC.⁹⁶ The CIN pathway can contribute to up to 85% of CAC, while the role of the MSI pathway is less clear. However, MSI has been found in up to 45% of cases.^{97–99} MSI has been demonstrated in histologically normal colonic mucosa of patients with UC, perhaps as a risk indicator for neoplastic progression.¹⁰⁰ A paper looking at MSI and methylation status of 124 IBD-related colonic neoplasias from 78 patients,¹⁰¹ showed 19/124 (9.6%) neoplastic lesions to be MSI-high (MSI-H). Methylation profile, tested by HPP1 and RAB32 methylation, was found to be independent of MSI status, with such methylation found in 28/61 (46%) and 20/60 (33.3%) carcinomas, while HPP1 methylation was frequently (~50%) observed in dysplastic mucosa and even in nondysplastic mucosa (7%). In another study,¹⁰² mutational events secondary to MSI were found to be similar in IBD- and non IBD-related colonic neoplasia. MSI-H IBD neoplasias were not more frequently right sided and there was no age or sex association, whereas sporadic MSI-H CRC was more prevalent in older women. IBD MSI-H tumours were more advanced and less often poorly differentiated than sporadic MSI-H tumours.

Data regarding the methylation pathway are contradictory. Some evidence suggests that the involvement of this pathway is more prevalent in IBD-related neoplasia.¹⁰¹ Methylation of CPG islands in several genes can precede dysplasia and can be

detected throughout the colon in UC patients.¹⁰³ p16^{INK4} has frequently been shown to be hypermethylated in patients with UC even in non-dysplastic mucosa, and is considered a marker for the CIMP pathway.⁹⁸ An additional methylated gene is p14^{RF}, an indirect regulator of p53.¹⁰⁴ Recent evidence does not support the co-occurrence of methylation with IBD CRC neoplasia.¹⁰⁵

The serrated neoplasia pathway, characterized by V600E BRAF mutations, has been recently shown to explain some of the IBD-related neoplastic colonic lesions.¹⁰⁶ Clonality of colitis-associated neoplasia has been shown in several crypts from the same colitis-associated neoplastic lesions.^{92,107} As in sporadic CRC, genomic transformation probably takes place in stem cells at the base of the crypt, as mutations have been shown to spread. These cells apparently sense signals from the microenvironment and the epithelial mesenchymal interaction.¹⁰⁸

4.3. The field effect

One of the main differences between sporadic and IBD-related colonic neoplasia is that in IBD the entire colonic mucosa carries risk for neoplastic transformation that can be multifocal, as opposed to one or few premalignant adenomas or cancers in sporadic cases.¹⁰⁹

Aneuploidy is a marker for genomic instability and has been shown in morphologically normal mucosal cells in IBD. Studies in UC patients with consecutive colonoscopies indicate that cell populations with aneuploidy expand in location over time.¹¹⁰ Aneuploidy does not correlate morphologically with dysplasia and is probably more widespread. Many genomic alterations, including methylation, have been shown in morphologically normal colonic mucosa in IBD patients.^{92,107} p53 mutations that are an initial event in IBD-associated neoplasia can be found in nondysplastic mucosa.⁹⁵ Two recent studies^{92,107} looked into individual crypts and found the same mutations, mainly in TP53 KRAS and CDKN2A, present both in neoplasia and in nondysplastic mucosa, even long before tumour development, strongly supporting the concept of a precancerous field effect.

The cause of the field effect can be explained by the constant re-epithelialization of ulcerated and chronically inflamed colonic mucosa by abnormal healing clones that expand,⁹⁷ as well as local environmental changes such as dysbiosis of the microflora, which can cause the appearance of these mutations in several spots.¹¹¹

5. Outcome of colorectal cancer in IBD

During the last 40 years there have been a few small reports looking at the outcome of CRC in UC^{9,112–115} and CD.^{116,117} The overall 5-year survival rates in these reports range from 33.5% to 55.1% for UC and 18% to 46% for CD, with no significant differences compared with non-IBD cases of CRC.

Over the last 16 years, a number of larger multi-centre cohort studies and population-based reports have been published. Bansal and Sonnenberg¹¹⁸ showed a trend ($p = 0.0681$) towards decreased risk of cancer-related death in CRC among patients with IBD ($n = 371$) compared with sporadic CRC ($n = 52\,243$), but no difference between CD ($n = 134$) and UC ($n = 237$). A 20-year follow-up study from the Mayo Clinic compared IBD-related (UC = 241 and CD = 49) CRC with age-

and sex-matched sporadic CRC.¹¹⁹ IBD-related cancers were more proximal, with only 55% distal to the splenic flexure, compared with 78% among sporadic CRC. Cases of IBD-related CRC were more often grade 3 or 4 tumours (40%) compared with sporadic tumours (33%), and were also more often mucinous (UC 20%, CD 24% and sporadic 13%). Despite this, no differences were found in overall survival between sporadic and IBD-related CRC. Two Danish population-based studies looked specifically at UC-¹²⁰ and CD-¹²¹ related CRC compared with sporadic CRC. In the study on UC by Jensen et al.¹²⁰ 71 259 sporadic CRC cases during 1977–1999 were compared with 279 UC-related cancers. Cancer stage, and rates of lymph node and distant metastasis were similar between the two groups. The overall mortality rates at 1 and 5 years after cancer diagnosis were increased in UC-related CRC compared with sporadic cancers (1.24 (95% CI 1.02–1.51) and 1.17 (95% CI 1.01–1.36), respectively). In a similar study by Larsen et al.,¹²¹ 71 438 sporadic CRC cases were compared with 100 CD-related CRC cases. The hazard ratios for death at 1 and 5 years after cancer diagnosis were 1.82 (95% CI 1.36–2.43) and 1.57 (95% CI 1.24–1.99), respectively. Both in UC and CD, CRC diagnosed <60 years of age was shown to have a worse outcome, especially within the first year with a hazard ratio of 1.92 (95% CI 1.29–2.87). Brackmann et al.¹²² published a population-based study from 1962 to 2005 on 81 780 sporadic cases of CRC and 60 patients with IBD-related dysplasia or CRC (CD colitis *n* = 6, UC *n* = 54). No differences were found regarding cancer stage or distant metastasis at the time of CRC diagnosis. The mortality rate ratio (MRR) was 3.71 (95% CI 2.54–5.42) for IBD-related CRC, compared with sporadic cancers mainly attributable to multifocal dysplasia or cancers, for which the MRR was 4.27 (95% CI 2.83–6). Survival after localised IBD-related CRC was similar to that of sporadic CRC.

In another study, the outcome in UC patients (*n* = 20) was compared with that in CD patients (*n* = 14) diagnosed with CRC between 1990 and 2007.¹²³ CRC in CD was found to be associated with worse outcome in terms of 5-year disease-free survival, 43% and 31% vs. 70% and 63% in UC (*p* = 0.01 respectively). In a Swedish population-based study,¹²⁴ sporadic CRC cases (*n* = 112 948) were compared with tumours occurring in IBD patients (*n* = 251), all with similar tumour stage at CRC diagnosis. Apart from rectal cancer in CD, the CRC-related mortality was increased among IBD patients, with hazard ratios for colon cancer of 2.08 (95% CI 1.46–2.96) in CD, 1.32 (95% CI 1.02–1.72) in UC and 1.77 (95% CI 1.22–2.56) for rectal cancer in UC. Furthermore, this study revealed that IBD patients <60 years at time of diagnosis and males had a worse outcome.

Some studies have found no difference between IBD and sporadic CRC in terms of survival. Peyrin-Biroulet et al. analysed data from the Burgundy digestive cancer registry in France during 1976–2008 (sporadic *n* = 19 413 and IBD related *n* = 38).¹²⁵ As shown previously, IBD patients were younger at the time of CRC diagnosis, 56.9 years vs. 70.9 years (*p* < 0.001). A trend towards an excess death rate was found among IBD patients with CRC (hazard ratio 1.46 95% CI 0.94–2.27). Another more recent US study¹²⁶ compared 29 CRC patients with CD and 53 with UC with 13 938 sporadic CRC patients. Mortality from CRC was 2.3 (95% CI 1.6–3.0) times higher in CD and 2.0 (95% CI 1.3–2.7) times higher in UC.

In summary, the available literature shows an impaired outcome of CRC in IBD, with a 2-fold increase in mortality compared with sporadic cancers, but without major differences between UC and CD except for a more advanced tumour stage at the time of diagnosis in CD patients. It is worth noting that for IBD, the presence of CRC is often not known prior to surgery. Male sex and young age are factors associated with poor outcome.

6. Chemoprevention of colorectal cancer in IBD

A number of agents are potential chemopreventives in IBD, such as 5-aminosalicylates (5-ASAs), ursodeoxycholic acid (UDCA), folic acid and azathioprine.

6.1. 5-Aminosalicylates

5-ASA compounds, used extensively in the management of IBD, particularly UC, have been identified as candidate chemopreventive agents.¹²⁷ The mechanism of action of mesalazine in chemoprevention is diverse, interfering with many pathways in CRC cell biology, rather than simply controlling inflammation. Actions include activation of the PPAR gamma pathway, inhibition of lipoxygenase and cyclooxygenase mediators, free radical scavenger and antioxidant functions, and ability to inhibit interleukins.¹²⁸

Early evidence comes from a community-based study cohort of 175 patients with UC by Moody et al.¹²⁹ who demonstrated a 10-fold reduction in the incidence of CRC in patients on long-term sulphasalazine. In a case-controlled study, Pinchowski et al.¹³⁰ studied this aspect in IBD patients grouped by the duration of 5-ASA use. Patients who had used sulphasalazine for ≥3 months had a 62% risk reduction compared with patients with a shorter duration of use. In a further case-controlled study, Eaden et al.¹³¹ compared 102 patients with UC-related CRC with matched controls, finding a 75% risk reduction for CRC among patients with regular 5-ASA use (OR 0.25; 95% CI 0.13–0.48; *p* < 0.00001). The benefit persisted after adjusting for variables such as disease extent and disease duration (OR 0.19, 95% CI 0.06–0.61; *p* = 0.006). A more recent large population-based general practise database study from the UK¹³² found that regular users of 5-ASAs (defined as >6 prescriptions in the preceding 12 months) had a 40% reduced risk of CRC (adjusted OR 0.60 (0.38–0.96)).

However, not all studies support a protective effect. In a large UC surveillance cohort at St Mark's Hospital, London, primarily designed to examine risk factors for colorectal neoplasia, use of mesalazine for ≥10 years was associated with a more modest and non-significant decrease in CRC risk (OR 0.65; 95% CI 0.26–1.62).¹³³ In addition, this study reported an unexpected increase in risk of CRC in patients taking long-term sulphasalazine. Similarly, Ullman et al.¹³⁴ found no evidence of an effect of mesalazine exposure on subsequent development of advanced neoplasia. Terdiman¹³⁵ compared CRC patients registered on a large administrative claims database in California with controls, finding that 5-ASA therapy of any dose for a duration of 1 year before diagnosis of CRC was not associated with a reduction in CRC risk (OR 0.97, 95% CI 0.77–1.22). Bernstein et al. found a trend towards increased cancer risk associated with the use of 5-ASAs within 2 years of a cancer diagnosis, indicating perhaps that the

benefit only exists if given early in the course of disease in UC.¹³⁶

The most compelling data supporting a beneficial effect of 5-ASAs come from a meta-analysis of 9 observational studies involving a total of 1932 patients, finding a 49% reduction in the risk of CRC or CRC/dysplasia with regular use of 5-ASAs.¹³⁷ The pooled analysis showed a protective association between use of 5-ASA and CRC (OR 0.51, 95% CI 0.37–0.69) and for the combined end point of CRC/dysplasia (OR 0.51, 95% CI 0.38–0.69). A recent systematic review¹³⁸ including 4 observational studies focused on non-referral studies indicated a chemopreventive benefit with 5-ASAs with an adjusted OR of 0.95 (95% CI 0.66–1.38), but there was moderate heterogeneity ($I^2 = 58.2\%$; $p = 0.07$).

Four studies looked at the dose needed to provide a chemoprevention benefit. A study by van Staa et al.¹³² found a doubled risk reduction among those on daily doses above 1.2 g/day compared with those taking <1.2 g/day, but a protective effect was observed in both groups (adjusted OR 0.28 and 0.56, respectively). Similar results were shown by Eaden et al.¹³¹ where mesalazine at doses of 1.2 g or more reduced CRC risk by 91% compared with no treatment (OR 0.09, 95% CI 0.03–0.28) but a protective effect was still found with lower doses (OR 0.08; 95% CI 0.08–0.85). In a more recent case-controlled study involving 1594 IBD patients that included 18 CRC patients, cumulative aminosalicylate dose of over 4500 g reduced the risk of CRC by 97% after adjusting for disease extent and duration.¹³⁹

Overall, despite the lack of good quality prospective randomised trials, there is sufficient evidence now to recommend the use of 5-ASAs for the chemoprevention of CAC. However, a number of unanswered questions remain. There is little data from which to establish recommendations for when to initiate therapy, the duration of therapy needed to achieve the benefit of chemoprevention and whether the effect lasts after cessation of therapy. Future studies should also identify those groups at highest risk among whom the benefit may be higher, and whether 5-ASA can inhibit progression of dysplasia to cancer. The anti-mutational effect of 5-ASA may also be useful in individuals with Lynch syndrome.^{140,141}

6.2. Ursodeoxycholic acid

The first indication of a potential chemoprotective effect of UDCA in UC came from a retrospective study of 59 patients with PSC and UC, of whom 41 were treated with UDCA¹⁴² in which a lower prevalence of dysplasia was reported in the UDCA-treated group (OR 0.18; $p = 0.005$) with a 40% difference in risk between the two groups. The protective effect appeared to persist after adjusting for age, duration of colitis and concomitant sulfasalazine use (adjusted OR 0.14, 95% CI 0.03–0.64). In a placebo-controlled trial in PSC patients with UC (median disease duration 13 years, follow-up time 6.5 years), designed to detect improvement in liver function, UDCA was associated with a 74% risk reduction for CRC and dysplasia (RR 0.26, 95% CI 0.07–0.99).¹⁴³ Due to the limited number of colonic biopsies performed in the study compared to the current surveillance guidelines, the result may be biased.

Further evidence of the benefit of UDCA came from a proof of concept study in IBD patients with pre-existing low-grade

dysplasia and DNA aneuploidy randomised to receive UDCA or placebo.¹⁴⁴ The primary outcome, defined as the need for colectomy for the progression of dysplasia, was reached in 2 patients in the placebo group and none in the UDCA-treated group. This study was limited by very small patient numbers, short treatment duration and the low doses of UDCA used, but it does indicate that UDCA may halt the progression of dysplasia in high-risk patients with extensive colitis and pre-existing premalignant lesions.

The early positive findings for UDCA as a chemopreventive agent have not been replicated in more recent studies, including studies with more patients. In a cohort study that included 120 patients, Wolf et al.¹⁴⁵ found no difference in the risk of developing dysplasia or carcinoma, although they did show a reduction in mortality in the UDCA group. More recent studies suggest that the duration of treatment and the dose of UDCA are important factors in chemoprevention.^{146–150} Eaton and et al.¹⁴⁸ examined the effects of high-dose (28–30 mg/kg/day) UDCA on the development of colorectal neoplasia in 56 patients with UC and PSC with a follow-up duration of 235 patient years. Patients who received high-dose UDCA had a significantly higher risk of developing colorectal neoplasia during the study than those who received placebo (HR: 4.44, 95% CI 1.30–20.10; $p = 0.02$). In another study, also using high-dose UDCA (17–23 mg/kg), there was a similar frequency of dysplasia or cancer after 5 years in patients originally assigned to UDCA or placebo (13% vs. 16%), and no difference was detected in dysplasia/cancer-free survival after 5 years of therapy (13% and 16%) or 10 years of follow-up (30% and 27%).¹⁴⁹ In a study in post-transplant patients with PSC, use of 5-ASA or UDCA was shown on multivariate analysis to be associated with an increased risk of CRC.¹⁵⁰ A large randomised study estimating the survival advantage in PSC patients of high-dose 5-ASA failed to demonstrate any reduction in CRC risk.¹⁵¹ In contrast, a prospective study from Germany in 120 patients with IBD and PSC found that the annual incidence of CRC increased up to 6 years after the start of UDCA but subsequently decreased.¹⁵² A recent meta-analysis of 4 studies, including 281 patients, reported that UDCA was not associated with any improvement in the risk of CRC (OR 0.53; 95% CI: 0.19–1.48; $p = 0.23$) compared with no UDCA or placebo.¹⁵³

Overall, studies so far suggest that the role of UDCA as a chemopreventive agent remains controversial, particularly at high doses. Larger randomised prospective studies including those comparing high doses versus low doses are needed to reach a conclusion. However, this is unlikely given the rarity of treatment-naïve PSC patients with colitis, and the long follow-up necessary.

6.3. Immunomodulators

There is increasing evidence that chronic inflammation favours the development of cancer in IBD, and hence mucosal healing with immunomodulators or biologics may prevent CRC in IBD.¹⁵⁴ However, there is insufficient data on the role of immunomodulators in chemoprevention. In a single-centre study of UC patients treated with azathioprine and followed for 17 years, no patient developed cancer.¹⁵⁵ In a further cohort study from New York of 315 patients followed for an average of 8 years, azathioprine (AZA)/6-mercaptopurine

(6MP) use did not have any effect on the development of CRC.¹⁵⁶ Three further studies examined chemoprevention by 6MP and AZA as chemoprophylaxis, although not as the primary endpoint.^{133,142,157} Rutter et al.¹³³ showed a non-significant trend towards a protective effect of azathioprine use for 1–5 years, or greater than 5 years, with ORs of 0.34 (95% CI 0.09–1.25) and 0.73 (95% CI 0.30–1.78), respectively. Lashner et al.¹⁵⁷ reported that the use of azathioprine for ≥ 6 months had no protective effect (RR 1.12, 95% CI 0.26–4.17). A similar result was reported by Tung et al.¹⁴² These studies were retrospective cohort studies with no data available on the dose used or duration of treatment. In a nested case–control study using a UK primary care database,¹³² among 5-ASA users who had also used oral glucocorticoids or immunosuppressants in the previous 6 months there were no differences in the risk of CRC between regular and irregular 5-ASA use (adjusted OR 1.21 (95% CI 0.43–3.45)). In contrast, regular 5-ASA use was associated with a reduced risk of CRC in non-users of oral glucocorticoids or immunosuppressants (adjusted OR 0.53 (95% CI 0.30–0.92)).¹²⁹ In a more recent nested case–control study in North America, using immunosuppressive therapy (OR 0.3; 95% CI 0.16–0.56; $p < 0.001$) or anti-TNF α (OR 0.09; 95% CI 0.01–0.68; $p < 0.02$) was protective in terms of risk of IBD-related CRC.²⁰ Interestingly, the overall risk of CRC in this cohort was much lower than previously reported, with an overall risk of 0.04%.²⁰

The association between thiopurine and/or 5-ASA use and the risk of advanced neoplasia (AN), including high-grade dysplasia and colorectal cancer, was recently investigated in a large cohort of patients with IBD in the Netherlands.¹⁵⁸ Thiopurine use was associated with a significantly decreased risk of AN (adjusted HR 0.10, 95% CI 0.01 to 0.75). 5-ASA therapy also had a protective effect against AN, but this was not statistically significant (adjusted HR 0.56, 95% CI 0.22 to 1.40).

Recent experimental data identifying TNF α as a mediator involved in initiation and progression of CAC has not yet been reflected in clinical trials, and other risks will preclude their use for the sole aim of chemoprevention. The results from the GETAID CESAME cohort have not yet been fully reported, but preliminary data do not support a chemopreventive role of AZA.¹⁵⁹

6.4. Folic acid

Lashner et al.¹⁵⁷ reviewed data from 99 patients with pancolitis enrolled in a 7-year surveillance programme and studied the effect of folate supplementation on the rate of neoplasia development using case–control methodology. Folate supplementation was associated with a 62% lower incidence of neoplasia but this was not statistically significant (OR, 0.38; 95% CI, 0.12–1.20). The same group also reported a similar non-significant trend towards a protective effect and dose–response effect of folate supplementation of ≥ 6 months in a subsequent cohort study.¹⁶⁰ In this study, the adjusted RR of neoplasia for patients taking folate was 0.72 (95% CI 0.28–1.83). The risk of neoplasia varied with folate dose (RR, 0.54 for 1.0 mg folate; RR, 0.76 for 0.4 mg folate in a multi-vitamin, compared with no folate). The degree of dysplasia also varied with folate use (RR for cancer, 0.45; RR for high-grade dysplasia, 0.52; RR for

low-grade dysplasia, 0.75, compared with patients without dysplasia) ($p = 0.08$). An experimental study on cell kinetics using immunohistochemistry following 3 months of folic acid or placebo supplementation in 12 colitis patients revealed a reduction in cell proliferation.¹⁶¹ However, the role of folic acid has not yet been tested in large randomised controlled trials.

Overall, while there is no clear data to support routine the use of folic acid, there appears to be a trend towards benefit. Since folic acid is inexpensive and safe it may be considered in selected patients at the clinician's discretion.

6.5. Statins

Experimental data from mice indicate that statins may have a protective effect against colitis-associated carcinogenesis, primarily by the induction of selective apoptosis of colon cancer cells.¹⁶² In a population-based case–control study,¹⁶³ statin therapy was associated with a modest reduction in CRC in the non-IBD population, but a substantial 94% risk reduction in patients with IBD was observed in a subset analysis of 55 IBD patients. Furthermore, a recent meta-analysis of 18 trials on statin use in hypercholesterolemia showed only limited benefit in the setting of general population.¹⁶⁴ Overall, further studies are needed in high-risk IBD patients to test the results of initial experimental studies.

6.6. Chemoprevention in CD

The role of chemoprevention in CD has not been well studied, and available data comes from studies with small numbers of CD patients and do not give any firm conclusions. However, a non-significant trend for a protective effect has been reported with regular 5-ASA use (OR 0.30; 95% CI 0.05–1.17; $p = 0.10$) in at least one study.¹⁶⁵

7. Conclusions and areas for future research

Recent data show that CRC risk is decreasing but it still remains a major clinical problem in IBD patients. Better control of intestinal inflammation is likely to be the major reason. Future areas of investigation should focus on understanding better the key pathogenic pathways. In addition, the impact of all the novel medications including anti-TNF in decreasing CRC risk needs to be better assessed, even though interventional studies appear to be unethical and not feasible when it is a matter of choosing between treating or not inflammation. Large prospective cohorts can help us to further investigate and identify the risk factors to develop CRC and to assess the efficacy of IBD related drugs in reducing the risk. Finally, from a surgical point of view the collection of data from cohort studies and managing CRC in similar pathways of care can provide better data on survival and disease related outcomes.

Conflict of interest

Silvio Danese has served as a speaker, a consultant and an advisory board member for Schering-Plough, Abbott

Laboratories, Merck & Co, UCB Pharma, Ferring, Cellcris, Millenium Takeda, Nycomed, Pharmacosmos, Actelion, Alphawasserman, Genentech, Grunenthal, Pfizer, Astra Zeneca, Novo Nordisk, Cosmo Pharmaceuticals, Vifor, and Johnson and Johnson.

References

- Rubin DC, Shaker A, Levin MS. Chronic intestinal inflammation: inflammatory bowel disease and colitis-associated colon cancer. *Front Immunol* 2012;3:107.
- Lakatos PL, Lakatos L. Challenges in calculating the risk for colorectal cancer in patients with ulcerative colitis. *Clin Gastroenterol Hepatol* 2012;10:1179–80.
- Katsanos K, Tsianos E. Inflammatory bowel disease related cancer. *Ann Gastroenterol* 2002;15:134–42.
- Eaden JA, Abrams KR, Mayberry JF. The risk of colorectal cancer in ulcerative colitis: a meta-analysis. *Gut* 2001;48:526–35.
- Katsanos KH, Vermeire S, Christodoulou DK, Riis L, Wolters F, Odes S, et al. Dysplasia and cancer in inflammatory bowel disease 10 years after diagnosis: results of a population-based European collaborative follow-up study. *Digestion* 2007;75:113–21.
- Katsanos KH, Tatsioni A, Pedersen N, Shuhaibar M, Ramirez VH, et al. Cancer in inflammatory bowel disease 15 years after diagnosis in a population-based European collaborative follow-up study. *J Crohns Colitis* 2011;5:430–42.
- Gong W, Lv N, Wang B, Chen Y, Huang Y, Pan W, et al. Risk of ulcerative colitis-associated colorectal cancer in China: a multi-center retrospective study. *Dig Dis Sci* 2012;57:503–7.
- Venkataraman S, Mohan V, Ramakrishna BS, Peter S, Chacko A, Chandy G, et al. Risk of colorectal cancer in ulcerative colitis in India. *J Gastroenterol Hepatol* 2005;20:705–9.
- Kim BJ, Yang SK, Kim JS, Jeon YT, Choi H, Han DS, et al. Trends of ulcerative colitis-associated colorectal cancer in Korea: a KASID study. *J Gastroenterol Hepatol* 2009;24:667–71.
- Hata K, Watanabe T, Kazama S, Suzuki K, Shinozaki M, Yokayama T, et al. Earlier surveillance colonoscopy programme improves survival in patients with ulcerative colitis associated colorectal cancer: results of a 23-year surveillance programme in the Japanese population. *Br J Cancer* 2003;89:1232–6.
- Palli D, Trallori G, Bagnoli S, Saieva C, Tarantino O, Ceroti M, et al. Hodgkin's disease risk is increased in patients with ulcerative colitis. *Gastroenterology* 2000;119:647–53.
- Bernstein CN, Blanchard JF, Kliever E, Wajda A. Cancer risk in patients with inflammatory bowel disease: a population-based study. *Cancer* 2001;91:854–62.
- Winther KV, Jess T, Langholz E, Munkholm P, Binder V. Long-term risk of cancer in ulcerative colitis: a population-based cohort study from Copenhagen County. *Clin Gastroenterol Hepatol* 2004;2:1088–95.
- Jess T, Loftus EV, Velayos FS, Winther KW, Tremaine WJ, Zinsmeister AR, et al. Risk of intestinal cancer in inflammatory bowel disease: a population-based study from Olmsted county, Minnesota. *Gastroenterology* 2006;130:1039–46.
- Lakatos L, Mester G, Erdelyi Z, David G, Pandur T, Balogh M, et al. Risk factors for ulcerative colitis-associated colorectal cancer in a Hungarian cohort of patients with ulcerative colitis: results of a population-based study. *Inflamm Bowel Dis* 2006;12:205–11.
- Jess T, Riis L, Vind I, Winther KV, Borg S, Binder V, et al. Changes in clinical characteristics, course, and prognosis of inflammatory bowel disease during the last 5 decades: a population-based study from Copenhagen, Denmark. *Inflamm Bowel Dis* 2007;13:481–9.
- Söderlund S, Brandt L, Lapidus A, Karlen P, Brostrom O, Lofberg R, et al. Decreasing time-trends of colorectal cancer in a large cohort of patients with inflammatory bowel disease. *Gastroenterology* 2009;136:1561–7.
- Shaukat A, Virnig DJ, Salfiti NI, Howard DH, Sitaraman SV, Liff JM. Is inflammatory bowel disease an important risk factor among older persons with colorectal cancer in the United States? A population-based case-control study. *Dig Dis Sci* 2011;56(8):2378–83.
- Jess T, Simonsen J, Jørgensen KT, Pederson BV, Nielson NM, Frisch M. Decreasing risk of colorectal cancer in patients with inflammatory bowel disease over 30 years. *Gastroenterology* 2012;143 [375–381.e1].
- Baars JE, Looman CWN, Steyerberg EW, Beukers R, Tan AC, Weusten BL, et al. The risk of inflammatory bowel disease-related colorectal carcinoma is limited: results from a nationwide nested case-control study. *Am J Gastroenterol* 2011;106:319–28.
- Castano-Milla C, Chaparro M, Gisbert JP. Has the risk of developing colorectal cancer in patients with ulcerative colitis been overstated? A meta-analysis. *Gastroenterology* 2012;142:S-251.
- Jess T, Rungoe C, Peyrin-Biroulet L. Risk of colorectal cancer in patients with ulcerative colitis: a meta-analysis of population-based cohort studies. *Clin Gastroenterol Hepatol* 2012;10:639–45.
- Jess T, Gomborg M, Matzen P, Munkholm P, Sørensen TI, et al. Increased risk of intestinal cancer in Crohn's disease: a meta-analysis of population-based cohort studies. *Am J Gastroenterol* 2005;100:2724–9.
- Canavan C, Abrams KR, Mayberry J. Meta-analysis: colorectal and small bowel cancer risk in patients with Crohn's disease. *Aliment Pharmacol Ther* 2006;23:1097–104.
- von Roon AC, Reese G, Teare J, Constantinides V, Darzi AW, Tekkis PP, et al. The risk of cancer in patients with Crohn's disease. *Dis Colon Rectum* 2007;50:839–55.
- Laukoetter MG, Mennigen R, Hannig CM, Osada N, Rijcken E, Vowinkel T, et al. Intestinal cancer risk in Crohn's disease: a meta-analysis. *J Gastrointest Surg* 2011;15(4):576–83.
- Friedman S, Rubin PH, Bodian C, Harpaz N, Present DH. Screening and surveillance colonoscopy in chronic Crohn's colitis: results of a surveillance program spanning 25 years. *Clin Gastroenterol Hepatol* 2008;6(9):993–8.
- Basseri RJ, Basseri B, Vassilaki ME, Melmed GY, Ippoliti A, Vasilias EA, et al. Colorectal cancer screening and surveillance in Crohn's colitis. *J Crohns Colitis* 2012;6(8):824–9.
- Stewénus J, Adnerhill I, Anderson H, Ekelund GR, Florén CH, Fork FT, et al. Incidence of colorectal cancer and all cause mortality in non-selected patients with ulcerative colitis and indeterminate colitis in Malmö, Sweden. *Int J Colorectal Dis* 1995;10(2):117–22.
- Ekbom A, Helmick C, Zack M, Adami HO. Ulcerative colitis and colorectal cancer. A population-based study. *N Eng J Med* 1990;323(18):1228–33.
- Ekbom A, Helmick C, Zack M, Adami HO. Increased risk of large-bowel cancer in Crohn's disease with colonic involvement. *Lancet* 1990;336(8711):357–9.
- Askling J, Dickman PW, Karlén P, Broström O, Lapidus A, Löfberg R, et al. Family history as a risk factor for colorectal cancer in inflammatory bowel disease. *Gastroenterology* 2001;120(6):1356–62.
- Soetikno RM, Lin OS, Heidenreich PA, Young HS, Blackstone MO. Increased risk of colorectal neoplasia in patients with primary sclerosing cholangitis and ulcerative colitis: a meta-analysis. *Gastrointest Endosc* 2002;56(1):48–54.
- Rutter MD, Saunders BP, Wilkinson KH, Rumbles S, Schofield G, Kamm MA, et al. Cancer surveillance in longstanding ulcerative

- colitis: endoscopic appearances help predict cancer risk. *Gut* 2004;**53**(12):1813–6.
35. Breynaert C, Vermeire S, Rutgeerts P, Van Assche G. Dysplasia and colorectal cancer in inflammatory bowel disease: a result of inflammation or an intrinsic risk? *Acta Gastroenterol Belg* 2008;**71**(4):367–72.
 36. Gupta RB, Harpaz N, Itzkowitz S, Hossain S, Matula S, Kornbluth A, et al. Histologic inflammation is a risk factor for progression to colorectal neoplasia in ulcerative colitis: a cohort study. *Gastroenterology* 2007;**133**(4):1099–105.
 37. Cairns SR, Scholefield JH, Steele RJ, Dunlop MG, Thomas HJ, Evans GD, et al. Guidelines for colorectal cancer screening and surveillance in moderate and high risk groups (update from 2002). *Gut* 2010;**59**(5):666–89.
 38. Farraye FA, Odze RD, Eaden J, Itzkowitz SH. AGA medical position statement on the diagnosis and management of colorectal neoplasia in inflammatory bowel disease. *Gastroenterology* 2010;**138**(2):738–45.
 39. Van Assche G, Dignass A, Bokemeyer B, Danese S, Gionchetti P, Moser G, et al. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 3: special situations. *J Crohns Colitis* 2013;**7**(1):1–33.
 40. Lutgens MW, Vleggaar FP, Schipper MEI, Stokkers PC, van der Woude CJ, Hommes DW, et al. High frequency of early colorectal cancer in inflammatory bowel disease. *Gut* 2008;**57**(9):1246–51.
 41. Rutter MD, Saunders BP, Wilkinson KH, Rumbles S, Schofield G, Kamm MA, et al. Thirty-year analysis of a colonoscopic surveillance program for neoplasia in ulcerative colitis. *Gastroenterology* 2006;**130**(4):1030–8.
 42. Söderlund S, Brandt L, Lapidus A, Karlén P, Broström O, Löfberg R, et al. Reply. *Gastroenterology* 2009;**137**(5):1858.
 43. Gillen CD, Walmsley RS, Prior P, et al. Ulcerative colitis and Crohn's disease: a comparison of the colorectal cancer risk in extensive colitis. *Gut* 1994;**35**(11):1590–2.
 44. Choi PM, Zelig MP. Similarity of colorectal cancer in Crohn's disease and ulcerative colitis: implications for carcinogenesis and prevention. *Gut* 1994;**35**(7):950–4.
 45. Averboukh F, Ziv Y, Kariv Y, Zmora O, Dotan I, Klausner JM, et al. Colorectal carcinoma in inflammatory bowel disease: a comparison between Crohn's and ulcerative colitis. *Colorectal Dis* 2011;**13**(11):1230–5.
 46. Kiran RP, Khoury W, Church JM, Lavery IC, Fazio VW, Remzi FH. Colorectal cancer complicating inflammatory bowel disease: similarities and differences between Crohn's and ulcerative colitis based on three decades of experience. *Ann Surg* 2010;**252**(2):330–5.
 47. Brackmann S, Andersen SN, Aamodt G, et al. Relationship between clinical parameters and the colitis-colorectal cancer interval in a cohort of patients with colorectal cancer in inflammatory bowel disease. *Scand J Gastroenterol* 2009;**44**(1):46–55.
 48. Karvellas CJ, Fedorak RN, Hanson J, Wong CK. Increased risk of colorectal cancer in ulcerative colitis patients diagnosed after 40 years of age. *Can J Gastroenterol* 2007;**21**(7):443–6.
 49. Baars JE, Kuipers EJ, van Haastert M, Nicolaï JJ, Poen AC, van der Woude CJ. Age at diagnosis of inflammatory bowel disease influences early development of colorectal cancer in inflammatory bowel disease patients: a nationwide, long-term survey. *J Gastroenterol* 2012;**47**(12):1308–22.
 50. Thackeray EW, Charatcharoenwitthaya P, Elfaki D, Sinakos E, Lindor KD. Colon neoplasms develop early in the course of inflammatory bowel disease and primary sclerosing cholangitis. *Clin Gastroenterol Hepatol* 2011;**9**(1):52–6.
 51. Hanounieh IA, Macaron C, Lopez R, Zein NN, Lashner BA. Risk of colonic neoplasia after liver transplantation for primary sclerosing cholangitis. *Inflamm Bowel Dis* 2012;**18**(2):269–74.
 52. Vera A, Gunson BK, Ussatoff V, Nightingale P, Candinas D, Radley S, et al. Colorectal cancer in patients with inflammatory bowel disease after liver transplantation for primary sclerosing cholangitis. *Transplantation* 2003;**75**(12):1983–8 [27].
 53. Torres J, Pineton de Chambrun G, Itzkowitz S, Sachar DB, Colombel JF. Review article: colorectal neoplasia in patients with primary sclerosing cholangitis and inflammatory bowel disease. *Aliment Pharmacol Ther* 2011;**34**(5):497–508.
 54. Lindström L, Lapidus A, Ost Å, Bergquist A. Increased risk of colorectal cancer and dysplasia in patients with Crohn's colitis and primary sclerosing cholangitis. *Dis Colon Rectum* 2011;**54**(11):1392–7.
 55. Braden B, Halliday J, Aryasingha S, Sharifi Y, Checchin D, Warren BF, et al. Risk for colorectal neoplasia in patients with colonic Crohn's disease and concomitant primary sclerosing cholangitis. *Clin Gastroenterol Hepatol* 2012;**10**(3):303–8.
 56. Söderlund S, Granath F, Broström O, Karlén P, Löfberg R, Ekblom A, et al. Inflammatory bowel disease confers a lower risk of colorectal cancer to females than to males. *Gastroenterology* 2010;**138**(5):1697–703.
 57. Hou JK, Kramer JR, Richardson P, Mei M, El-Serag HB. Risk of colorectal cancer among Caucasian and African American veterans with ulcerative colitis. *Inflamm Bowel Dis* 2012;**18**(6):1011–7.
 58. Rakoff-Nahoum S, Medzhitov R. Regulation of spontaneous intestinal tumorigenesis through the adaptor protein MyD88. *Science* 2007;**317**:124–7.
 59. Fukata M, Abreu MT. Role of Toll-like receptors in gastrointestinal malignancies. *Oncogene* 2008;**27**:234–43.
 60. Salcedo R. MyD88-mediated signaling prevents development of adenocarcinomas of the colon: role of interleukin 18. *J Exp Med* 2010;**207**(8):1625–36.
 61. Popivanova BK, Kostadinova FI, Furuichi K, Shamekh MM, Kondo T, Wada T, et al. Blockade of a chemokine, CCL2, reduces chronic colitis-associated carcinogenesis in mice. *Cancer Res* 2009;**69**:7884–92.
 62. Atreya R, Mudter J, Finotto S, Müllberg J, Jostock T, Wirtz S, et al. Blockade of interleukin 6 trans signaling suppresses T-cell resistance against apoptosis in chronic intestinal inflammation: evidence in Crohn disease and experimental colitis in vivo. *Nat Med* 2000;**6**:583–8.
 63. McLoughlin RM, Witowski J, Robson RL, Wilkinson TS, Hurst SM, Williams AS, et al. Interplay between IFN-gamma and IL-6 signaling governs neutrophil trafficking and apoptosis during acute inflammation. *J Clin Invest* 2003;**112**:598–607.
 64. Kai H, Kitadai Y, Kodama M, Cho S, Kuroda T, Ito M, et al. Involvement of proinflammatory cytokines IL-1beta and IL-6 in progression of human gastric carcinoma. *Anticancer Res* 2005;**25**:709–13.
 65. Li Y, de HC, Chen M, Deuring J, Gerrits MM, Smits R, et al. Disease-related expression of the IL6/STAT3/SOCS3 signalling pathway in ulcerative colitis and ulcerative colitis-related carcinogenesis. *Gut* 2010;**59**(2):227–32.
 66. Stolfi C. Involvement of interleukin-21 in the regulation of colitis-associated colon cancer. *J Exp Med* 2011;**208**(11):2279–90.
 67. Jauch D. Interleukin 21 controls tumour growth and tumour immunosurveillance in colitis-associated tumorigenesis in mice. *Gut* 2011;**60**(12):1678–86.
 68. Nibbs RJ, Gilchrist DS, King V, Ferra A, Forrow S, Hunter KD, et al. The atypical chemokine receptor D6 suppresses the development of chemically induced skin tumors. *J Clin Invest* 2007;**117**:1884–92.
 69. Vetrano S, Borroni EM, Sarukhan A, Savino B, Bonecchi R, Correale C, et al. The lymphatic system controls intestinal inflammation and inflammation-associated colon cancer through the chemokine decoy receptor D6. *Gut* 2010;**59**(2):197–206.

70. Wang H, Czura C, Tracey K. The Cytokine Handbook. In: Thomson A, Lotze M, editors. London: Elsevier Science Ltd.; 2003. p. 837–60.
71. Uguccioni M, Gionchetti P, Robbiani DF, Rizzello F, Peruzzo S, Campieri M, et al. Increased expression of IP-10, IL-8, MCP-1, and MCP-3 in ulcerative colitis. *Am J Pathol* 1999;155:331–6.
72. Popivanova BK, Kitamura K, Wu Y, Kondo T, Kagaya T, Kaneko S, et al. Blocking TNF- α in mice reduces colorectal carcinogenesis associated with chronic colitis. *J Clin Invest* 2008;118:560–70.
73. Glocker EO, Kotlarz D, Boztug K, Gertz EM, Schäffer AA, Noyan F, et al. Inflammatory bowel disease and mutations affecting the interleukin-10 receptor. *N Engl J Med* 2009;361:2033–45.
74. Becker C, Fantini MC, Neurath MF. TGF- β as a T cell regulator in colitis and colon cancer. *Cytokine Growth Factor Rev* 2006;17:97–106.
75. Rizzo A, Waldner MJ, Stolfi C, Sarra M, Fina D, Becker C, et al. Smad7 expression in T cells prevents colitis-associated cancer. *Cancer Res* 2011;71(24):7423–32.
76. Wald D, Qin J, Zhao Z, Qian Y, Naramura M, Tian L, et al. SIGIRR, a negative regulator of Toll-like receptor-interleukin 1 receptor signaling. *Nat Immunol* 2003;4:920–7.
77. Polentarutti N, Rol GP, Muzio M, Bosisio D, Camnasio M, Riva F, et al. Unique pattern of expression and inhibition of IL-1 signaling by the IL-1 receptor family member TIR8/SIGIRR. *Eur Cytokine Netw* 2003;14:211–8.
78. Garlanda C, Riva F, Polentarutti N, Buracchi C, Sironi M, De Bortoli M, et al. Intestinal inflammation in mice deficient in Tir8, an inhibitory member of the IL-1 receptor family. *Proc Natl Acad Sci* 2004;101:3522–6.
79. Naugler WE, Karin M. NF- κ B and cancer-identifying targets and mechanisms. *Curr Opin Genet Dev* 2008;8:19–26.
80. Greten FR, Karin M. The IKK/NF- κ B activation pathway—a target for prevention and treatment of cancer. *Cancer Lett* 2004;206:193–9.
81. Greten FR, Eckmann L, Greten TF, Park JM, Li ZW, Egan LJ, et al. IKK β links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 2004;118:285–96.
82. Allen IC, Wilson JE, Schneider M, Lich JD, Roberts RA, Arthur JC, et al. NLRP12 suppresses colon inflammation and tumorigenesis through the negative regulation of noncanonical NF- κ B signaling. *Immunity* 2012;25:36(5):742–54.
83. Pasparakis M. IKK/NF- κ B signaling in intestinal epithelial cells controls immune homeostasis in the gut. *Mucosal Immunol* 2008;1(Suppl 1):S54–7.
84. Swidsinski A, Khilkin M, Kerjaschki D, Schreiber S, Ortner M, Weber J, et al. Association between intraepithelial *Escherichia coli* and colorectal cancer. *Gastroenterology* 1998;115(2):281–6.
85. Darfeuille-Michaud A, Neut C, Lederman E, Di Martino P, Desreumaux P, et al. Presence of adherent *Escherichia coli* strains in ileal mucosa of patients with Crohn's disease. *Gastroenterology* 1998;115(6):1405–13.
86. Arthur JC, Perez-Chanona E, Mühlbauer M, Tomkovich S, Uronis JM, Fan TJ, et al. Intestinal inflammation targets cancer-inducing activity of the microbiota. *Science* 2012;338(6103):120–3.
87. Markowitz SD, Bertagnolli MM. Molecular origins of cancer: molecular basis of colorectal cancer. *N Engl J Med* 2009;361(25):2449–60 [17].
88. Selaru FM, Xu Y, Yin J, Zou T, Liu TC, Mori Y, et al. Artificial neural networks distinguish among subtypes of neoplastic colorectal lesions. *Gastroenterology* 2002;122(3):606–87.
89. Greenwald BD, Harpaz N, Yin J, Huang Y, Tong Y, Brown VL, et al. Loss of heterozygosity affecting the p53, Rb, and mcc/apc tumor suppressor gene loci in dysplastic and cancerous ulcerative colitis. *Cancer Res* 1992;52(3):741–5 [1].
90. Willenbacher RF, Zelman SJ, Ferrell LD, Moore 2nd DH, Waldman FM. Chromosomal alterations in ulcerative colitis-related neoplastic progression. *Gastroenterology* 1997;113(3):791–801.
91. Fogt F, Zhuang Z, Poremba C, Dockhorn-Dworniczak B, Vortmeyer A. Comparison of p53 immunoreactivity with allelic loss of p53 in ulcerative colitis-associated dysplasia and carcinoma. *Oncol Rep* 1998;5(2):477–80.
92. Leedham SJ, Graham TA, Oukrif D, McDonald SA, Rodriguez-Justo M, Harrison RF, et al. Clonality, founder mutations, and field cancerization in human ulcerative colitis-associated neoplasia. *Gastroenterology* 2009;136(2):542–50.
93. Fogt F, Vortmeyer AO, Goldman H, Giordano TJ, Merino MJ, Zhuang Z. Comparison of genetic alterations in colonic adenoma and ulcerative colitis-associated dysplasia and carcinoma. *Hum Pathol* 1998;29(2):131–6.
94. Hussain SP, Amstad P, Raja K, Ambis S, Nagashima M, Bennett WP, et al. Increased p53 mutation load in noncancerous colon tissue from ulcerative colitis: a cancer-prone chronic inflammatory disease. *Cancer Res* 2000;60(13):3333–7.
95. Burner GC, Rabinovitch PS, Haggitt RC. Neoplastic progression in ulcerative colitis: histology, DNA content, and loss of a p53 allele. *Gastroenterology* 1992;103(5):1602–10.
96. Hoque AT, Hahn SA, Schutte M. DPC4 gene mutation in colitis associated neoplasia. *Gut* 1997;40(1):120–2.
97. Lyda MH, Noffsinger A, Belli J, et al. Microsatellite instability and K-ras mutations in patients with ulcerative colitis. *Hum Pathol* 2000;31(6):665–71.
98. Fleisher AS, Esteller M, Harpaz N, Leytin A, Rashid A, Xu Y, et al. Microsatellite instability in inflammatory bowel disease-associated neoplastic lesions is associated with hypermethylation and diminished expression of the DNA mismatch repair gene, hMLH1. *Cancer Res* 2000;60(17):4864–8.
99. Cawkwell L, Sutherland F, Murgatroyd H. Defective hMSH2/hMLH1 protein expression is seen infrequently in ulcerative colitis associated colorectal cancers. *Gut* 2000;46(3):367–9.
100. Brentnall TA, Crispin DA, Bronner MP. Microsatellite instability in nonneoplastic mucosa from patients with chronic ulcerative colitis. *Cancer Res* 1996;56(6):1237–40 [99].
101. Schulmann K, Mori Y, Croog V, Yin J, Olaru A, Sterian A, et al. Molecular phenotype of inflammatory bowel disease-associated neoplasms with microsatellite instability. *Gastroenterology* 2005;129(1):74–85.
102. Svrcek M, El-Bchiri J, Chalastanis A, Capel E, Dumont S, Buhard O, et al. Specific clinical and biological features characterize inflammatory bowel disease associated colorectal cancers showing microsatellite instability. *J Clin Oncol* 2007;25(27):4231–8.
103. Issa JP, Ahuja N, Toyota M, Bronner MP, Brentnall TA. Accelerated age-related CpG island methylation in ulcerative colitis. *Cancer Res* 2001;61(9):3573–7.
104. Sato F, Harpaz N, Shibata D, Xu Y, Yin J, Mori Y, et al. Hypermethylation of the p14(ARF) gene in ulcerative colitis-associated colorectal carcinogenesis. *Cancer Res* 2002;62(4):1148–51.
105. Konishi K, Shen L, Wang S. Rare CpG island methylator phenotype in ulcerative colitis-associated neoplasias. *Gastroenterology* 2007;132(4):1254–60.
106. Bossard C, Denis MG, Béziau S. Involvement of the serrated neoplasia pathway in inflammatory bowel disease-related colorectal oncogenesis. *Oncol Rep* 2007;18(5):1093–7.
107. Galandiuk S, Rodriguez-Justo M, Jeffery R. Field cancerization in the intestinal epithelium of patients with Crohn's ileocolitis. *Gastroenterology* 2012;142(4):855–864.e8.
108. Shaker A, Rubin DC. Intestinal stem cells and epithelial–mesenchymal interactions in the crypt and stem cell niche. *Transl Res* 2010;156(3):180–7.
109. Ullman TA, Itzkowitz SH. Intestinal inflammation and cancer. *Gastroenterology* 2011;140:1807–16.
110. Rubin CE, Haggitt RC, Burner GC, Brentnall TA, Stevens AC, Levine DS, et al. DNA aneuploidy in colonic biopsies predicts

- future development of dysplasia in ulcerative colitis. *Gastroenterology* 1992;103(5):1611–20.
111. Itzkowitz SH, Yio X. Inflammation and cancer IV. Colorectal cancer in inflammatory bowel disease: the role of inflammation. *Am J Physiol Gastrointest Liver Physiol* 2004;287(1):G7–G17.
 112. Hughes RG, Hall TJ, Block GE, Levin B, Moossa AR. The prognosis of carcinoma of the colon and rectum complicating ulcerative colitis. *Surg Gynecol Obstet* 1978;146(1):46–8.
 113. Lavery IC, Chiulli RA, Jagelman DG, Fazio VW, Weakley FL. Survival with carcinoma arising in mucosal ulcerative colitis. *Ann Surg* 1982;195(4):508–12.
 114. Ritchie JK, Hawley PR, Lennard-Jones JE. Prognosis of carcinoma in ulcerative colitis. *Gut* 1981;22(9):752–5.
 115. Sugita A, Greenstein AJ, Ribeiro MB, Sachar DB, Bodian C, Panday AK, et al. Survival with colorectal cancer in ulcerative colitis. A study of 102 cases. *Ann Surg* 1993;218(2):189–95.
 116. Ribeiro MB, Greenstein AJ, Sachar DB, Barth J, Balasubramanian S, Harpaz N, et al. Colorectal adenocarcinoma in Crohn's disease. *Ann Surg* 1996;223(2):186–93.
 117. Richards ME, Rickert RR, Nance FC. Crohn's disease-associated carcinoma. A poorly recognized complication of inflammatory bowel disease. *Ann Surg* 1989;209(6):764–73.
 118. Bansal P, Sonnenberg A. Risk factors of colorectal cancer in inflammatory bowel disease. *Am J Gastroenterol* 1996;91(1):44–8.
 119. Delaunoy T, Limburg PJ, Goldberg RM, Lymp JF, Loftus Jr EV. Colorectal cancer prognosis among patients with inflammatory bowel disease. *Clin Gastroenterol Hepatol* 2006;4(3):335–42.
 120. Jensen AB, Larsen M, Gislum M, Skriver MV, Jepsen P, Norgaard B, et al. Survival after colorectal cancer in patients with ulcerative colitis: a nationwide population-based Danish study. *Am J Gastroenterol* 2006;101(6):1283–7.
 121. Larsen M, Mose H, Gislum M, Skriver MV, Jepsen P, Norgaard B, et al. Survival after colorectal cancer in patients with Crohn's disease: a nationwide population-based Danish follow-up study. *Am J Gastroenterol* 2007;102(1):163–7.
 122. Brackmann S, Aamodt G, Andersen SN, Roald B, Langmark F, Clausen OP, et al. Widespread but not localized neoplasia in inflammatory bowel disease worsens the prognosis of colorectal cancer. *Inflamm Bowel Dis* 2010;16(3):474–81.
 123. Ouaiissi M, Maggiori L, Alves A, Giger U, Sielezneff I, Valleur P, et al. Colorectal cancer complicating inflammatory bowel disease: a comparative study of Crohn's disease vs ulcerative colitis in 34 patients. *Colorectal Dis* 2011;13(6):684–8.
 124. Shu X, Ji J, Sundquist J, Sundquist K, Hemminki K. Survival in cancer patients hospitalized for inflammatory bowel disease in Sweden. *Inflamm Bowel Dis* 2011;17(3):816–22.
 125. Peyrin-Biroulet L, Lepage C, Jooste V, Gueant JL, Faivre J, et al. Colorectal cancer in inflammatory bowel diseases: a population-based study (1976–2008). *Inflamm Bowel Dis* 2012;18(12):2247–51.
 126. Herrinton LJ, Liu L, Levin TR, Allison JE, Lewis JD, Velayos F. Incidence and mortality of colorectal adenocarcinoma in persons with inflammatory bowel disease from 1998 to 2010. *Gastroenterology* 2012;143(2):382–9.
 127. Cheng Y, Desreumaux P. 5-Aminosalicylic acid is an attractive candidate agent for chemo prevention of colon cancer in patients with inflammatory bowel disease. *World J Gastroenterol* 2005;11(3):309–14.
 128. Lyakhovich A, Gasche C. Systematic review: molecular chemopreventive properties of 5-ASA. *Aliment Pharmacol Ther* 2010;31(2):202–9.
 129. Moody GA, Jayanthi V, Probert CS, Mac Kay H, Mayberry JF. Long term therapy with sulphasalazine protects against colorectal cancer in ulcerative colitis: a retrospective study of colorectal cancer and compliance with treatment in Leicestershire. *Eur J Gastroenterol Hepatol* 1996;8:1179–83.
 130. Pinczowski D, Ekblom A, Baron J, Yuen J, Adami HO. Risk factors for colorectal cancer in patients with ulcerative colitis: a case-control study. *Gastroenterology* 1994;107:117–20.
 131. Eaden J, Abrams K, Ekblom A, Jackson E, Mayberry J. Colorectal cancer prevention in ulcerative colitis: a case-control study. *Aliment Pharmacol Ther* 2000;14:145–53.
 132. Van Staa TP, Card T, Logan RF, Leufkens HG. Aminosalicilate use and colorectal cancer risk in inflammatory bowel disease: a large epidemiological study. *Gut* 2005;54:1573–8.
 133. Rutter M, Saunders B, Wilkinson K, Rumbles S, Schofield G, Kamm MA, et al. Severity of inflammation is a risk factor for colorectal neoplasia in ulcerative colitis. *Gastroenterology* 2004;126:451–9.
 134. Ullman T, Croog V, Harpaz N, Hossain S, Kombluth A, Bodian C, et al. Progression to colorectal neoplasia in ulcerative colitis. Effect of mesalamine. *Clin Gastroenterol Hepatol* 2008;6(11):1225–30.
 135. Terdiman JP, Steinbuch M, Blumentals WA, Ullman TA, Rubin DT. 5-Aminosalicylic acid therapy and the risk of colorectal cancer among patients with inflammatory bowel disease. *Inflamm Bowel Dis* 2007;13(4):367–71.
 136. Bernstein CN, Blanchard JF, Metge C, Yogendran M. Does the use of 5-aminosalicylates in inflammatory bowel disease prevent the development of colorectal cancer? *Am J Gastroenterol* 2003;98(12):2784–8.
 137. Velayos FS, Terdiman JP, Walsh JM. Effect of 5-aminosalicylate use on colorectal cancer and dysplasia risk: a systematic review and metaanalysis of observational studies. *Am J Gastroenterol* 2005;100:1345–53.
 138. Nguyen GC, Gulamhusein A, Bernstein CN. 5-aminosalicylic acid is not protective against colorectal cancer in inflammatory bowel disease: a meta-analysis of non-referral populations. *Am J Gastroenterol* 2012;107:1298–304.
 139. Tang J, Sharif O, Pai C, Silverman AL. Mesalamine protects against colorectal cancer in inflammatory bowel disease. *Dig Dis Sci* 2010;55(6):1696–703.
 140. Gasche C, Goel A, Natarajan L, Boland CR. Mesalazine improves replication fidelity in cultured colorectal cells. *Cancer Res* 2005;65:3993–7.
 141. Campregheer C, Honeder C, Chung H, Carethers JM, Gasche C, et al. Mesalazine reduces microsatellite instability in TGFBR2 and ACVR2 by improvement of replication fidelity in mononucleotide repeats. *Clin Cancer Res* 2010;16(6):1950–6.
 142. Tung BY, Emond MJ, Haggitt RC, Bronner MP, Kimmey MB, Kowdley RV, et al. Ursodiol use is associated with lower prevalence of colonic neoplasia in patients with UC and primary sclerosing cholangitis. *Ann Intern Med* 2001;134:89–95.
 143. Pardi DS, Loftus Jr EV, Kremers WK, Keach J, Lindor KD. Ursodeoxycholic acid as chemo preventive agent in patients with ulcerative colitis and primary sclerosing cholangitis. *Gastroenterology* 2003;124:889–93.
 144. Sjöqvist UU, Tribukait B, Ost Å, Einarson C, Oxelmark L, Lofberg R. Ursodeoxycholic acid treatment in IBD patients with colorectal dysplasia and/or DNA-aneuploidy: a prospective, double blind randomised controlled pilot study. *Anticancer Res* 2004;24:3121–7.
 145. Wolf JM, Rybicki LA, Lashner BA. The impact of ursodeoxycholic acid on cancer, dysplasia and mortality in ulcerative colitis patients with primary sclerosing cholangitis. *Aliment Pharmacol Ther* 2012;35:846.
 146. Santander C, Moreno-Otero R. Commentary: ursodeoxycholic acid as chemoprevention in inflammatory bowel disease and primary sclerosing cholangitis. *Aliment Pharmacol Ther* 2005;22:783–8148.
 147. Wolf JM, Rybicki LA, Lashner BA. The impact of ursodeoxycholic acid on cancer, dysplasia and mortality in ulcerative colitis

- patients with primary sclerosing cholangitis. *Aliment Pharmacol Ther* 2005;**22**(9):783–8.
148. Eaton JE, Silveira MG, Pardi DS, Sinakos E, Kowdley KV, Luketic AV, et al. High-dose ursodeoxycholic acid is associated with the development of colorectal neoplasia in patients with ulcerative colitis and primary sclerosing cholangitis. *Am J Gastroenterol* 2011;**106**:1638–45.
 149. Lindström L, Boberg KM, Wikman O, Friis-Liby I, Hultcrantz R, Prytz H, et al. High dose ursodeoxycholic acid in primary sclerosing cholangitis does not prevent colorectal neoplasia. *Aliment Pharmacol Ther* 2012;**35**:451–7.
 150. Jørgensen KK, Lindström L, Cvancarova M, Castedal M, Friman S, Schrupf E, et al. Colorectal neoplasia in patients with primary sclerosing cholangitis undergoing liver transplantation: a Nordic multicenter study. *Scand J Gastroenterol* 2012;**8–9**(47):1021–9.
 151. Olsson R, Boberg KM, de Muckadell OS, Lindgren S, Hultcrantz R, Folvik G, et al. High-dose ursodeoxycholic acid in primary sclerosing cholangitis: a 5-year multicenter, randomized, controlled study. *Gastroenterology* 2005;**129**:1464–72.
 152. Rudolph G, Gotthardt DN, Kloeters-Plachky P, Kulaksiz H, Schirmacher P, Stiehl A. In PSC with colitis treated with UDCA, most colonic carcinomas develop in the first years after the start of treatment. *Dig Dis Sci* 2011;**56**(12):3624–30.
 153. Ashraf I, Choudhary A, Arif M, Matteson ML, Hammad HT, Puli SR, et al. Ursodeoxycholic acid in patients with ulcerative colitis and primary sclerosing cholangitis for prevention of colon cancer; a metaanalysis. *Indian J Gastroenterol* 2012;**31**(2):69–74.
 154. Actis GC, Pellicano R, David E, Sapino A. Azathioprine, mucosal healing in ulcerative colitis and the chemoprevention of colitis cancer: a practice based forecast. *Inflamm Allergy Drug Targets* 2010;**9**(1):6–9.
 155. Actis GC, Fadda M, Pellicano R, David E, Rizzetto M, Sapino A. The 17 year single centre experience with the use of azathioprine to maintain remission in ulcerative colitis. *Biomed Pharmacother* 2009;**63**:362–5.
 156. Matuala S, Coorg V, Itzkowitz S, Harpaz N, Bodian C, Hossain S, et al. Chemoprevention of colorectal neoplasia in ulcerative colitis; the effect of 6-mercaptopurine. *Clin Gastroenterol Hepatol* 2005;**3**(10):1015–21.
 157. Lashner BA, Heidenreich PA, Su GL, Kane SV, Hanauer SB. Effect of folate supplementation on the incidence of dysplasia and cancer in chronic ulcerative colitis. A case–control study. *Gastroenterology* 1989;**97**(2):255–9.
 158. Bvan Schaik FD, van Oijen MG, Smeets HM, van der Heijden GJ, Siersema PD, Oldenburg B. Thiopurines prevent advanced colorectal neoplasia in patients with inflammatory bowel disease. *Gut* 2012;**61**(2):235–40.
 159. Carrat F, Seksik P, Bouvier AM, et al. Case–control study nested in the CESAME cohort. *Gastroenterology* 2010:A255.
 160. Lashner BA, Provencher KS, Seidner DL, Kneesebeck A, Brzezinski A. The effect of folic acid supplementation on the risk for cancer or dysplasia in ulcerative colitis. *Gastroenterology* 1997;**112**(1):29–32.
 161. Biasco G, Zannoni U, Paganelli GM, Santucci R, Gionchetti P, Rivolta G, et al. Folic acid supplementation and cell kinetics of rectal mucosa in patients with ulcerative colitis. *Cancer Epidemiol Biomarkers Prev* 1997;**6**:469–71.
 162. Cho SJ, Kim JS, Kim JM, Lee JY, Jung HC, Song IS. Simvastatin induces apoptosis in human colon cancer cells and in tumor xenografts, and attenuates colitis associated colon cancer in mice. *Int J Cancer* 2008;**123**:952–7.
 163. Poynter JN, Gruber SB, Higgins PD, Almog R, Bonner JD, Rennert HS, et al. Statins and the risk of colorectal cancer. *N Engl J Med* 2005;**352**:2184–92.
 164. Bonovas S, Filioussi K, Floedellis CR, Sitaras NM. Statins and the risk of colorectal cancer: a meta-analysis of 18 studies involving more than 1.5 million patients. *J Clin Oncol* 2007;**25**:342–3468.
 165. Siegel CA, Sands BE. Risk factors for colorectal cancer in Crohn's colitis: a case–control study. *Inflamm Bowel Dis* 2006;**12**(6):491–6.