

Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2012 April ; 21(4): 645–657. doi:10.1158/1055-9965.EPI-11-0888.

Common variants at the 19p13.1 and *ZNF365* loci are associated with ER subtypes of breast cancer and ovarian cancer risk in *BRCA1* and *BRCA2* mutation carriers

Fergus J. Couch^{1, #}, Mia M. Gaudet^{2, #}, Antonis C. Antoniou³, Susan J. Ramus⁴, Karoline B. Kuchenbaecker³, Penny Soucy⁵, Jonathan Beesley⁶, Xiaoqing Chen⁶, Xianshu Wang¹, Tomas Kirchhoff⁷, Lesley McGuffog³, Daniel Barrowdale³, Andrew Lee³, Sue Healey⁶, Olga M. Sinilnikova^{8, 9}, Irene L. Andrulis^{10, 11, 12}, Hilmi Ozelik¹¹, Anna Marie Mulligan^{11, 13}, for OCGN, Mads Thomassen¹⁴, Anne-Marie Gerdes¹⁵, Uffe Birk Jensen¹⁶, Anne-Bine Skytte¹⁷, Torben A. Kruse¹⁴, Maria A. Caligo¹⁸, Anna von Wachenfeldt¹⁹, Gisela Barbany-Bustinza²⁰, Niklas Loman²¹, Maria Soller²², Hans Ehrencrona²³, Per Karlsson²⁴, for SWE-BRCA, Katherine L. Nathanson²⁵, Timothy R. Rebbeck²⁶, Susan M. Domchek²⁵, Ania Jakubowska²⁷, Jan Lubinski²⁷, Katarzyna Jaworska^{27, 28}, Katarzyna Durda²⁷, Elżbieta Złowocka²⁷, Tomasz Huzarski²⁷, Tomasz Byrski²⁷, Jacek Gronwald²⁷, Cezary Cybulski²⁷, Bohdan Górski²⁷, Ana Osorio²⁹, Mercedes Durán³⁰, María Isabel Tejada³¹, Javier Benitez³², Ute Hamann³³, Frans B.L. Hogervorst³⁴, Theo A. van Os³⁵, Flora E. van Leeuwen³⁶, Hanne E.J. Meijers-Heijboer³⁷, Juul Wijnen³⁸, Marinus J. Blok³⁹, Marleen Kets⁴⁰, Maartje J. Hoening⁴¹, Rogier A. Oldenburg⁴², Margreet G.E.M. Ausems⁴³, for HEBON, Susan Peock³, Debra Frost³, Steve D. Ellis³, Radka Platte³, Elena Fineberg³, D. Gareth Evans⁴⁴, Chris Jacobs⁴⁵, Rosalind A. Eeles⁴⁶, Julian Adlard⁴⁷, Rosemarie Davidson⁴⁸, Diana M. Eccles⁴⁹, Trevor Cole⁵⁰, Jackie Cook⁵¹, Joan Paterson⁵², Carole Brewer⁵³, Fiona Douglas⁵⁴, Shirley V. Hodgson⁵⁵, Patrick J. Morrison⁵⁶, Lisa Walker⁵⁷, Mary E. Porteous⁵⁸, M. John Kennedy⁵⁹, Lucy E. Side⁶⁰, for EMBRACE, Betsy Bove⁶¹, Andrew K. Godwin⁶², Dominique Stoppa-Lyonnet^{63, 64, 65, 66}, Marion Fassy-Colcombet⁶⁶, Laurent Castera⁶⁶, François Cornelis^{67, 68, 69}, Sylvie Mazoyer⁹, Mélanie Léoné⁸, Nadia Boutry-Kryza⁸, Brigitte Bressac-de Paillerets⁷⁰, Olivier Caron⁷¹, Pascal Pujol⁷², Isabelle Coupier⁷², Capucine Delnatte⁷³, Linda Akloul⁷³, Henry T. Lynch⁷⁴, Carrie L. Snyder⁷⁴, for GEMO Study Collaborators, Sandra S. Buys⁷⁵, Mary B. Daly⁷⁶, MaryBeth Terry⁷⁷, Wendy K. Chung⁷⁷, Esther M. John⁷⁸, Alexander Miron⁷⁹, Melissa C. Southey⁸⁰, John L. Hopper⁸¹, David E. Goldgar⁸², Christian F. Singer⁸³, Christine Rappaport⁸³, Muy-Kheng M. Tea⁸³, Anneliese Fink-Retter⁸³, Thomas V. O. Hansen⁸⁴, Finn C. Nielsen⁸⁴, Aðalgeir Arason^{85, 86}, Joseph Vijai⁷, Sohela Shah⁷, Kara Sarrel⁷, Mark E. Robson⁷, Marion Piedmonte⁸⁷, Kelly Phillips⁸⁸, Jack Basil⁸⁹, Wendy S. Rubinstein^{90, 91}, John Boggess⁹², Katie Wakeley⁹³, Amanda Ewart-Toland⁹⁴, Marco Montagna⁹⁵, Simona Agata⁹⁵, Evgeny N. Imyanitov⁹⁶, Claudine Isaacs⁹⁷, Ramunas Janavicius⁹⁸, Conxi Lazaro⁹⁹, Ignacio Blanco¹⁰⁰, Lidia Feliubadalo⁹⁹, Joan Brunet¹⁰⁰, Simon A Gayther¹⁰¹, Paul PD Pharoah¹⁰², Kunle O. Odunsi¹⁰³, Beth Y. Karlan¹⁰⁴, Christine S. Walsh¹⁰⁴, Edith Olah¹⁰⁵, Soo Hwang Teo¹⁰⁵, Patricia A. Ganz¹⁰⁷, Mary S. Beattie¹⁰⁸, Elizabeth J. van Rensburg¹⁰⁹, Cecelia M. Dorfling¹⁰⁹, Orland Diez¹¹⁰, Ava Kwong¹¹¹, Rita K. Schmutzler¹¹², Barbara Wappenschmidt¹¹², Christoph Engel¹¹³, Alfons Meindl¹¹⁴, Nina

Corresponding Author: Fergus Couch, Mayo Clinic, Stabile 2-42, 200 First Street SW, Rochester, MN 55905, USA; Tel: 507-284-3623; Fax: 507-538-1937; couch.fergus@mayo.edu.

[#]F.J. Couch and M.M. Gaudet contributed equally as primary authors.

*K. Offit and J. Simard contributed equally as senior authors

Disclosure of Potential Conflicts of Interest

The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the BCFR, nor does mention of trade names, commercial products, or organizations imply endorsement by the US Government or the BCFR. No potential conflicts of interest are disclosed.

Ditsch¹¹⁵, Norbert Arnold¹¹⁶, Simone Heidemann¹¹⁷, Dieter Niederacher¹¹⁸, Sabine Preisler-Adams¹¹⁹, Dorothea Gadzicki¹²⁰, Raymonda Varon-Mateeva¹²¹, Helmut Deissler¹²², Andrea Gehrig¹²³, Christian Sutter¹²⁴, Karin Kast¹²⁵, Britta Fiebig¹²⁶, Wolfram Heinritz¹²⁷, Trinidad Caldes¹²⁸, Miguel de la Hoya¹²⁸, Taru A. Muranen¹²⁹, Heli Nevanlinna¹²⁹, Marc D. Tischkowitz^{130,131}, Amanda B. Spurdle⁶, Susan L. Neuhausen¹³², Yuan Chun Ding¹³², Noralane M. Lindor¹³³, Zachary Fredericksen¹³⁴, V. Shane Pankratz¹³⁴, Paolo Peterlongo¹³⁵, Siranoush Manoukian¹³⁶, Bernard Peissel¹³⁶, Daniela Zaffaroni¹³⁶, Monica Barile¹³⁷, Loris Bernard¹³⁸, Alessandra Viel¹³⁹, Giuseppe Giannini¹⁴⁰, Liliana Varesco¹⁴¹, Paolo Radice¹³⁵, Mark H. Greene¹⁴², Phuong L. Mai¹⁴², Douglas F. Easton³, Georgia Chenevix-Trench⁶, for kConFab investigators, Kenneth Offit^{7,*}, and Jacques Simard^{5,*} on behalf of the Consortium of Investigators of Modifiers of *BRCA1/2*

¹Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA

²Epidemiology Research Program, American Cancer Society, Atlanta, GA, USA ³Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, UK ⁴Department of Preventive Medicine, Keck School of Medicine, University of Southern California, CA, USA ⁵Canada Research Chair in Oncogenetics, Cancer Genomics Laboratory, Centre Hospitalier Universitaire de Québec and Laval University, Quebec City (Quebec), Canada ⁶Department of Genetics, Queensland Institute of Medical Research, Brisbane, Australia ⁷Clinical Cancer Genetics Laboratory, Memorial Sloane Kettering Cancer Center, New York, NY, USA ⁸Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Centre Hospitalier Universitaire de Lyon / Centre Léon Bérard, Lyon, France ⁹INSERM U1052, CNRS UMR5286, Université Lyon 1, Centre de Recherche en Cancérologie de Lyon, Lyon, France ¹⁰Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, Ontario M5G 1X5, Cancer Care Ontario, Department of Molecular Genetics, University of Toronto, Ontario, Canada ¹¹Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, Ontario M5G 1X5, Cancer Care Ontario, Department of Laboratory Medicine and Pathobiology, University of Toronto, Ontario, Canada ¹²Ontario Cancer Genetics Network: Cancer Care Ontario, Canada ¹³Department of Laboratory Medicine, and the Keenan Research Centre of the Li Ka Shing Knowledge Institute, St Michael's Hospital, Toronto, ON, Canada ¹⁴Department of Clinical Genetics, Odense University Hospital, Odense, Denmark ¹⁵Department of Clinical Genetics, Rigshospitalet, København, Denmark ¹⁶Department of Clinical Genetics, Skejby Hospital, Århus, Denmark ¹⁷Department of Clinical Genetics, Vejle Hospital, Vejle, Denmark ¹⁸Section of Genetic Oncology, University Hospital of Pisa, Pisa, Italy ¹⁹Department of Oncology, Karolinska University Hospital, Stockholm, Sweden ²⁰Department of Clinical Genetics, Karolinska University Hospital, Stockholm, Sweden ²¹Department of Oncology, Lund University Hospital, Lund, Sweden ²²Department of Clinical Genetics, Lund University Hospital, Lund, Sweden ²³Department of Genetics and Pathology, Rudbeck Laboratory, Uppsala University, Uppsala, Sweden ²⁴Department of Oncology, Sahlgrenska University Hospital, Gothenburg, Sweden ²⁵Department of Medicine, Abramson Cancer Center, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA ²⁶Department of Biostatistics and Epidemiology, Abramson Cancer Center, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA ²⁷Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland ²⁸Postgraduate School of Molecular Medicine, Warsaw Medical University, Warsaw, Poland ²⁹Human Genetics Group, Human Cancer Genetics Programme, Spanish National Cancer Research Centre, and Spanish Network on Rare Diseases (CIBERER), Madrid, Spain ³⁰Institute of Biology and Molecular Genetics, Universidad de Valladolid (IBGM-UVA), Valladolid, Spain ³¹Molecular Genetics Laboratory (Department of Biochemistry), Cruces Hospital Barakaldo, Bizkaia, Spain ³²Human Genetics Group and Genotyping Unit, Human Cancer Genetics Programme, Spanish National Cancer Research Centre, and Spanish Network on Rare Diseases (CIBERER), Madrid, Spain ³³Molecular Genetics of Breast Cancer, Deutsches Krebsforschungszentrum (DKFZ), Heidelberg, Germany ³⁴Family Cancer Clinic, Netherlands

Cancer Institute, Amsterdam, The Netherlands ³⁵Department of Clinical Genetics, Academic Medical Center, Amsterdam, The Netherlands ³⁶Department of Epidemiology, Head, Division of Psychosocial Research & Epidemiology, Netherlands Cancer Institute, Amsterdam, The Netherlands ³⁷Department of Clinical Genetics, VU Medical Center, Amsterdam, The Netherlands ³⁸Department of Human Genetics & Department of Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands ³⁹Department of Genetics and Cell Biology, University Medical Center, Maastricht, The Netherlands ⁴⁰Hereditary Cancer Clinic, Radboud University Nijmegen Medical Center, The Netherlands ⁴¹Department of Medical Oncology, Family Cancer Clinic, Erasmus University Medical Center, Rotterdam, The Netherlands ⁴²Department of Clinical Genetics, Family Cancer Clinic, Erasmus University Medical Center, Rotterdam, The Netherlands ⁴³Department of Medical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands ⁴⁴Genetic Medicine, Manchester Academic Health Sciences Centre, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK ⁴⁵Clinical Genetics, Guy's and St. Thomas' NHS Foundation Trust, London, UK ⁴⁶Oncogenetics Team, The Institute of Cancer Research and Royal Marsden NHS Foundation Trust, UK ⁴⁷Yorkshire Regional Genetics Service, Leeds, UK ⁴⁸Ferguson-Smith Centre for Clinical Genetics, Yorkhill Hospitals, Glasgow, UK ⁴⁹University of Southampton Faculty of Medicine, Southampton University Hospitals NHS Trust, Southampton UK ⁵⁰West Midlands Regional Genetics Service, Birmingham Women's Hospital Healthcare NHS Trust, Edgbaston, Birmingham, UK ⁵¹Sheffield Clinical Genetics Service, Sheffield Children's Hospital, Sheffield, UK ⁵²Department of Clinical Genetics, East Anglian Regional Genetics Service, Addenbrookes Hospital, Cambridge, UK ⁵³Department of Clinical Genetics, Royal Devon & Exeter Hospital, Exeter, UK ⁵⁴Institute of Genetic Medicine, Centre for Life, Newcastle Upon Tyne Hospitals NHS Trust, Newcastle upon Tyne, UK ⁵⁵Medical Genetics Unit, St George's, University of London, UK ⁵⁶Northern Ireland Regional Genetics Centre, Belfast Health and Social Care Trust, and Department of Medical Genetics, Queens University Belfast, Belfast UK ⁵⁷Oxford Regional Genetics Service, Churchill Hospital, Oxford, UK ⁵⁸South East of Scotland Regional Genetics Service, Western General Hospital, Edinburgh, UK ⁵⁹Academic Unit of Clinical and Molecular Oncology, Trinity College Dublin and St James's Hospital, Dublin, Ireland ⁶⁰North East Thames Regional Genetics Service, Great Ormond Street Hospital for Children NHS Trust, London, UK ⁶¹Clinical Molecular Genetics Laboratory, Fox Chase Cancer Center, Philadelphia, PA, USA ⁶²Department of Pathology and Laboratory Medicine, University of Kansas Medical Center, Kansas City, KS, USA ⁶³GEMO Study Collaborators: Cancer Genetics Network "Groupe Génétique et Cancer", Fédération Nationale des Centres de Lutte Contre le Cancer, France ⁶⁴Service de Génétique Oncologique, Institut Curie, Paris, France ⁶⁵Unité INSERM U830, Institut Curie, Paris, France ⁶⁶Université Paris Descartes, Faculté de Médecine, Paris, France ⁶⁷Genetic Unit, Avicenne Hospital, Assistance Publique-Hôpitaux de Paris, Paris, France ⁶⁸Sud-Francilien Hospital, Evry-Corbeil, France ⁶⁹University Hospital, Clermont-Ferrand, France ⁷⁰Genetics Department, Institut de Cancérologie Gustave Roussy, Villejuif, France ⁷¹Consultation de Génétique, Département de Médecine, Institut de Cancérologie Gustave Roussy, Villejuif, France ⁷²Unité d'Oncogénétique, CHU Arnaud de Villeneuve, Montpellier, France ⁷³Service Oncologie Médicale, Centre René Gauducheau, Nantes, France ⁷⁴Department of Preventive Medicine, Creighton University, Omaha, NE, USA ⁷⁵Department of Oncological Sciences, Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, USA ⁷⁶Fox Chase Cancer Center, Philadelphia, PA, USA ⁷⁷Department of Epidemiology, Columbia University, New York, NY, USA ⁷⁸Department of Epidemiology, Cancer Prevention Institute of California, Fremont, CA, USA ⁷⁹Department of Cancer Biology, Dana Farber Cancer Institute, Boston, MA, USA ⁸⁰Genetic Epidemiology Laboratory, Department of Pathology, The University of Melbourne, Parkville, Victoria, Australia ⁸¹Centre for Molecular, Environmental, Genetic and Analytic Epidemiology, University of Melbourne, Melbourne, Victoria, Australia ⁸²Department of Dermatology, University of Utah School of Medicine, Salt Lake City, UT, USA ⁸³Department of OB/GYN and Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria

⁸⁴Center for Genomic Medicine, Rigshospitalet, Copenhagen University Hospital, Blegdamsvej 9, Copenhagen, Denmark ⁸⁵Department of Pathology, Landspítali University Hospital, Reykjavik, Iceland ⁸⁶Faculty of Medicine, University of Iceland, Reykjavik, Iceland ⁸⁷Gynecologic Oncology Group Statistical and Data Center, Roswell Park Cancer Institute, 666 Elm St, Buffalo, NY, USA ⁸⁸Division of Cancer Medicine, Peter MacCallum Cancer Centre, Melbourne, Victoria, Australia and Department of Medicine, St Vincent's Hospital, The University of Melbourne, Victoria, Australia ⁸⁹Good Samaritan Hospital, Department of Obstetrics and Gynecology, GOG affiliate ⁹⁰Center for Medical Genetics, NorthShore University Health System, Evanston, IL, USA ⁹¹Department of Medicine, University of Chicago Pritzker School of Medicine, Chicago, IL, USA ⁹²University of North Carolina, Chapel Hill, NC, USA ⁹³Dana-Farber/Brigham and Women's Cancer Center in clinical affiliation with South Shore Hospital, South Weymouth, MA, USA ⁹⁴Departments of Molecular Virology, Immunology & Medical Genetics and Internal Medicine, Comprehensive Cancer Center, The Ohio State University, Columbus, OH, USA ⁹⁵Istituto Oncologico Veneto IOV - IRCCS, Immunology and Molecular Oncology Unit, Padua, Italy ⁹⁶Laboratory of Molecular Oncology, N.N. Petrov Institute of Oncology, St.-Petersburg, Russia ⁹⁷Fisher Center for Familial Cancer Research, Lombardi Comprehensive Cancer Center, Georgetown University, Washington D.C., USA ⁹⁸Dept. of Molecular and Regenerative medicine, Hematology, Oncology and Transfusion Medicine Center, Vilnius University Hospital Santariskiu Clinics, Santariskiu st. 2, Vilnius, Lithuania ⁹⁹Molecular Diagnostic Unit, Hereditari Cancer Program, IDIBELL-Catalan Institute of Oncology, Barcelona, Spain ¹⁰⁰Genetic Counseling Unit, Hereditari Cancer Program, IDIBELL-Catalan Institute of Oncology, Barcelona, Spain ¹⁰¹Department of Preventive Medicine, Keck School of Medicine, University of Southern California, CA, USA ¹⁰²Department of Oncology, University of Cambridge, Cambridge, UK ¹⁰³Department of Gynecologic Oncology, Roswell Park Cancer Institute, Buffalo, NY, USA ¹⁰⁴Women's Cancer Program, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA ¹⁰⁵Department of Molecular Genetics, National Institute of Oncology, Budapest, Hungary ¹⁰⁶Breast Cancer Research Unit, University Malaya Cancer Research Institute, University Malaya, Malaysia and Cancer Research Initiatives Foundation, Sime Darby Medical Centre, Subang Jaya, Malaysia ¹⁰⁷UCLA School of Public Health, Department of Health Services, Los Angeles, CA, USA ¹⁰⁸Department of Medicine, Epidemiology and Biostatistics, University of California, San Francisco, CA, USA ¹⁰⁹Cancer Genetics Laboratory, Department of Genetics, University of Pretoria, South Africa ¹¹⁰Molecular Medicine and Genetics Program. Vall d'Hebron University Hospital. Oncogenetics Laboratory. Vall d'Hebron Institute of Oncology (VHIO). Barcelona, Spain ¹¹¹Hong Kong Hereditary Breast Cancer Family Registry and Cancer Genetics Center, Hong Kong Sanatorium and Hospital, Hong Kong ¹¹²Centre of Familial Breast and Ovarian Cancer, Department of Gynaecology and Obstetrics and Centre for Integrated Oncology (CIO), University hospital of Cologne, Germany ¹¹³Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Germany ¹¹⁴Department of Gynaecology and Obstetrics, Division of Tumor Genetics, Klinikum rechts der Isar, Technical University Munich, Germany ¹¹⁵Department of Gynaecology and Obstetrics, Ludwig-Maximilian University Munich, Germany ¹¹⁶Department of Gynaecology and Obstetrics, University Hospital of Schleswig-Holstein, Campus Kiel, Christian-Albrechts University Kiel, Germany ¹¹⁷Institute of Human Genetics, University Hospital of Schleswig-Holstein, Campus Kiel, Christian-Albrechts University Kiel, Germany ¹¹⁸Department of Gynaecology and Obstetrics, University Hospital Düsseldorf, Heinrich-Heine University, Düsseldorf, Germany ¹¹⁹Institute of Human Genetics, University of Münster, Münster, Germany ¹²⁰Institute of Cell and Molecular Pathology, Hannover Medical School, Hannover, Germany ¹²¹Institute of Human Genetics, Campus Virchow Klinikum, Berlin, Germany ¹²²Department of Gynaecology and Obstetrics, University Hospital Ulm, Germany ¹²³Centre of Familial Breast and Ovarian Cancer, Department of Medical Genetics, Institute of Human Genetics, University Würzburg, Germany ¹²⁴Institute of Human Genetics, Division of Molecular Genetics, University Heidelberg, Germany ¹²⁵Department of Gynaecology

and Obstetrics, University Hospital Carl Gustav Carus, Technical University Dresden, Germany ¹²⁶Institute of Human Genetics, University Regensburg, Germany ¹²⁷Institute of Human Genetics, University Leipzig, Germany ¹²⁸Molecular Oncology Laboratory, Hospital Clinico San Carlos. Martin Lagos s/n, 28040 Madrid, Spain ¹²⁹Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Central Hospital, Biomedicum Helsinki, Helsinki, Finland ¹³⁰Program in Cancer Genetics, Departments of Human Genetics and Oncology, McGill University, Montreal, Quebec, Canada ¹³¹Lady Davis Institute, Segal Cancer Centre, Jewish General Hospital, Montreal, Quebec, Canada ¹³²Department of Population Sciences, Beckman Research Institute of City of Hope, Duarte, CA USA ¹³³Department of Medical Genetics, Mayo Clinic Rochester, MN, USA ¹³⁴Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA ¹³⁵Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Preventive and Predicted Medicine, Fondazione IRCCS Istituto Nazionale Tumori (INT), and IFOM, Fondazione Istituto FIRC di Oncologia Molecolare, Milan, Italy ¹³⁶Unit of Medical Genetics, Department of Preventive and Predicted Medicine, Fondazione IRCCS Istituto Nazionale Tumori (INT), Milan, Italy ¹³⁷Division of Cancer Prevention and Genetics, Istituto Europeo di Oncologia (IEO), Milan, Italy ¹³⁸Department of Experimental Oncology, Istituto Europeo di Oncologia, and Consortium for Genomics Technology (Cogentech), Milan, Italy ¹³⁹Unit of Experimental Oncology 1, Centro di Riferimento Oncologico, IRCCS, Aviano (PN), Italy ¹⁴⁰Department of Experimental Medicine, University La Sapienza, Rome, Italy ¹⁴¹Unit of Hereditary Cancer, Department of Epidemiology, Prevention and Special Functions, Istituto Nazionale per la Ricerca sul Cancro (IST), Genoa, Italy ¹⁴²Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, NCI, Rockville, MD, USA

Abstract

Background—Genome-wide association studies (GWAS) identified variants at 19p13.1 and *ZNF365* (10q21.2) as risk factors for breast cancer among *BRCA1* and *BRCA2* mutation carriers, respectively. We explored associations with ovarian cancer and with breast cancer by tumor histopathology for these variants in mutation carriers from the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA).

Methods—Genotyping data for 12,599 *BRCA1* and 7,132 *BRCA2* mutation carriers from 40 studies were combined.

Results—We confirmed associations between rs8170 at 19p13.1 and breast cancer risk for *BRCA1* mutation carriers (hazard ratio (HR)=1.17; 95%CI 1.07–1.27; $p=7.42 \times 10^{-4}$) and between rs16917302 at *ZNF365* (HR=0.84; 95%CI 0.73–0.97; $p=0.017$) but not rs311499 at 20q13.3 (HR=1.11; 95%CI 0.94–1.31; $p=0.22$) and breast cancer risk for *BRCA2* mutation carriers. Analyses based on tumor histopathology showed that 19p13 variants were predominantly associated with estrogen receptor (ER)-negative breast cancer for both *BRCA1* and *BRCA2* mutation carriers, whereas rs16917302 at *ZNF365* was mainly associated with ER-positive breast cancer for both *BRCA1* and *BRCA2* mutation carriers. We also found for the first time that rs67397200 at 19p13.1 was associated with an increased risk of ovarian cancer for *BRCA1* (HR=1.16; 95%CI 1.05–1.29; $p=3.8 \times 10^{-4}$) and *BRCA2* mutation carriers (HR=1.30; 95%CI 1.10–1.52; $p=1.8 \times 10^{-3}$).

Conclusions—19p13.1 and *ZNF365* are susceptibility loci for ovarian cancer and ER subtypes of breast cancer among *BRCA1* and *BRCA2* mutation carriers.

Impact—These findings can lead to an improved understanding of tumor development and may prove useful for breast and ovarian cancer risk prediction for *BRCA1* and *BRCA2* mutation carriers.

Keywords

BRCA1; BRCA2; breast cancer risk; ovarian cancer risk; 19p13.1; ZNF365

Introduction

Genome-wide association studies (GWAS) have been used to identify several loci containing common variants that are associated ($p < 1.0 \times 10^{-7}$) with breast cancer risk in the general population. Variants from twelve of these loci have also been investigated as modifiers of cancer risk in *BRCA1* and *BRCA2* mutation carriers (1–3). While only variants in *CASP8*, *TOX3*, 2q35, and 6q25.1 have been associated with breast cancer risk in *BRCA1* mutation carriers, variants in *FGFR2*, *TNRC9/TOX3*, *MAP3K1*, *LSP1*, 2q35, *SLC4A7/NEK10*, 5p12 and 1p11.2 loci have been associated with breast cancer in *BRCA2* mutation carriers (1–3). This is consistent with the known associations between these SNPs and estrogen receptor (ER) status of breast cancers in the general population (4).

Most recently, a GWAS of *BRCA1* mutation carriers conducted through CIMBA identified five SNPs on 19p13 that were associated with breast cancer risk for *BRCA1* mutation carriers (5). Two of these showed independent associations (rs8170 hazard ratio (HR)=1.26; 95% CI 1.17–1.35; $P_{\text{trend}} = 2.3 \times 10^{-9}$ and rs2363956 HR=0.84; 95% CI 0.80–0.89; $P_{\text{trend}} = 5.5 \times 10^{-9}$). Imputation analysis of the 19p13 region, using 1000 Genomes Project data, identified several correlated SNPs with more significant associations than rs8170 and rs2363956. The 19p13.1 locus was also found to be associated with ER negative breast cancer (rs8170 OR=1.21, $p=0.003$) and triple negative breast cancer (tumors lacking expression of ER, progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2)) (rs8170 OR=1.28, $p=1.2 \times 10^{-6}$) in the general population (5). In addition, the 19p13.1 locus has been associated with ovarian cancer in the general population (rs8170 OR=1.12; $p=3.6 \times 10^{-6}$) (6), but was not found to be associated with ovarian cancer in *BRCA1* mutation carriers (rs8170 HR=1.07; $p=0.33$) (5). A separate GWAS in *BRCA2* mutation carriers identified two breast cancer susceptibility alleles (rs16917302 at *ZNF365* (10q21.2), HR=0.75; 95% CI 0.66–0.86; $p=3.8 \times 10^{-5}$ and rs311499 at 20q13.33, HR=0.72; 95% CI 0.61–0.85; $p=6.6 \times 10^{-5}$) (7). A weakly correlated SNP at the *ZNF365* locus (rs10995190) has also been associated with breast cancer overall (OR=0.83; $p=5.1 \times 10^{-15}$) and ER positive ($p=4.1 \times 10^{-6}$) but not ER negative breast cancer in the general population (8).

Here, we genotyped more than 12,000 *BRCA1* and 7,000 *BRCA2* mutation carriers from the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA), for the previously genotyped variant at 19p13.1, rs8170, and one of the imputed SNPs that was found to have a stronger association with breast cancer risk for *BRCA1* mutation carriers (rs67397200). We also genotyped SNPs at *ZNF365* (rs16917302), and 20q13.3 (rs311499) in an effort to verify these loci as risk factors for ovarian cancer and to further validate these loci as risk factors for breast cancer in *BRCA1* and *BRCA2* mutation carriers.

Materials and Methods

Subjects

All mutation carriers participated in clinical or research studies at the host institutions under ethically approved protocols and provided written informed consent. Subjects were *BRCA1* and *BRCA2* mutation carriers recruited by 40 study centers in 22 countries and assembled through the CIMBA initiative (Supplementary Table 1). The majority were recruited through cancer genetics clinics and enrolled into national or regional studies. Others were

identified in research studies of high-risk families, by population-based sampling of cases and some by community recruitment. Eligibility to participate in CIMBA is restricted to female carriers of pathogenic *BRCA1* or *BRCA2* mutations, defined by generally recognized criteria (Breast Cancer Information Core), who were 18 years old or older at recruitment. Information collected included the year of birth; mutation description (including nucleotide position and base change); age at last follow-up; ages at breast and ovarian cancer diagnoses; and age or date at bilateral prophylactic mastectomy. Information was also available on the country of residence. Related individuals were identified through a unique family identifier. Women with pathogenic mutations in both *BRCA1* and *BRCA2* were excluded from the current analysis. The primary analysis was restricted to women self-reported as “white European”. Overlap of carriers between studies was evaluated by comparing the year of birth, exact mutation description, the reported ages, and previous SNP genotype data available within the CIMBA database. Duplicated mutation carriers were included only once in the analysis.

Genotyping

Rs311499 at 20q13.3, rs16917302 at *ZNF365* and both rs8170 and rs67397200 at 19p13.1 were genotyped using the iPLEX Mass Array platform at four genotyping centers as part of a larger study of 24 candidate SNPs. All centers included at least 2% duplicate samples and a random mixture of affected and unaffected carriers on each plate. Samples that failed for five or more of the SNPs genotyped were excluded from the analysis. Studies with a SNP call rate of <95% were excluded from the analysis of the SNP. The concordance between duplicates had to be at least 98%. To assess the accuracy of genotyping across genotyping centers, all centers genotyped 95 DNA samples from a standard test plate (Coriell Institute) for all SNPs. Genotyping centers with more than one concordance failure on the test plate for a SNP were excluded for analyses of that SNP. Deviation from Hardy-Weinberg equilibrium (HWE) was assessed for unrelated subjects separately for each SNP and study. The observed genotype frequencies were not significantly different from the expected under HWE for any of the SNPs and studies. After the above exclusions, a total of 19,731 unique mutation carriers (12,599 *BRCA1* and 7,132 *BRCA2*) from 40 studies had an observed genotype for at least one SNP (Supplementary Table 1).

Tumor pathology data collection

Tumor pathology data were collected from patient pathology reports, medical records, pathology review data, tumor registry records and results from tissue microarrays. ER status was identified as negative or positive, with immunohistochemistry scoring data and methodology provided when available. Most studies applied a cut-off of >10% tumor cells stained positive for ER positive status. For a small number of cases, where other scoring methods based on the proportion and intensity of staining were applied (Allred score, Remmele score and H-score), widely-accepted cut-offs were used. Consistency checks were performed to validate receptor data against supplementary scoring information if provided.

Statistical analysis

The aim of the primary analysis was to evaluate the association between each genotype and breast cancer risk. We conducted the analysis by modelling the retrospective likelihood of the observed genotypes conditional on the disease phenotypes as previously described (9). The phenotype of each individual was defined by age at diagnosis of breast cancer or age at last follow-up. Individuals were censored at the earliest of age of first breast cancer diagnosis, ovarian cancer diagnosis, bilateral prophylactic mastectomy or age at last observation. Mutation carriers censored at ovarian cancer diagnosis were considered unaffected in the analysis of breast cancer. The effect of each SNP was modelled either as a per-allele HR (multiplicative model) or as separate HRs for heterozygotes and homozygotes.

We used a Cox proportional-hazards model and tested the assumption of proportional hazards by adding a “genotype×age” interaction term in order to fit models in which the HR changed with age. We examined heterogeneity across studies by comparing models that allowed for study-specific log-hazard ratios against models in which the same log-hazard ratio was assumed to apply to all studies. All analyses were stratified by country of residence and applied cohort specific breast cancer incidence rates for *BRCA1* and *BRCA2* (10). A robust variance-estimation approach was used to adjust for the non-independence among related carriers.

To evaluate the evidence of replication for each of the SNPs, analyses were restricted to mutation carriers who had not been used in any of the previous *BRCA1* and *BRCA2* studies. The number of new samples used in each of the SNP analyses are shown in Supplementary Table 2. In addition, analysis were performed using all available *BRCA1* and *BRCA2* carriers. The combined effects of the SNPs on breast cancer risk, were evaluated by fitting retrospective likelihood models while allowing for linkage disequilibrium between the loci. To test for potential effects of survival bias, prevalent cases, defined as mutation carriers diagnosed more than five years prior to the age at recruitment, were excluded. Associations with specific functional class of mutation were also assessed. Class 1 mutations are predicted to undergo nonsense mediated RNA decay resulting in reduced levels of mutant transcript while Class 2 mutations are predicted to generate stable mutant proteins (11). The associations with breast cancer subtypes defined by the estrogen receptor (ER) status of the tumors in *BRCA1* and *BRCA2* mutation carriers were assessed by an extension of the retrospective likelihood approach that models the simultaneous effect of each SNP on more than one tumor subtype (12). Associations with ovarian cancer risk were evaluated within a competing risk analysis framework (13) by estimating HRs simultaneously for breast and ovarian cancers. Since each mutation carrier was at risk of breast and ovarian cancer we assumed that the probabilities of developing each disease were independent conditional on the underlying genotype. In this analysis, individuals were followed to the age of the first breast or ovarian cancer diagnosis and were considered to have developed the corresponding disease. Individuals were censored for breast cancer at the age of bilateral prophylactic mastectomy and for ovarian cancer at the age of bilateral oophorectomy and were assumed to be unaffected for the corresponding disease. The remaining individuals were censored at the age at last observation and were assumed to be unaffected for both diseases.

Results

After quality control exclusions, genotype data from 12,599 *BRCA1* and 7,132 *BRCA2* mutation carriers including 5,408 *BRCA1* and 3,780 *BRCA2* mutation carriers not studied in the original GWAS were available for analysis. Of the *BRCA1* mutation carriers, 6,390 were affected with breast cancer and 6,209 were considered unaffected in the breast cancer analysis (censored at bilateral prophylactic mastectomy, ovarian cancer, or age at last follow up). Similarly, among the *BRCA2* mutation carriers, 3,810 were affected with breast cancer and 3,322 were unaffected. The characteristics of these mutation carriers are shown in Table 1 and the origins of the samples are summarized in Supplementary Table 1.

The associations between breast cancer risk in *BRCA1* and *BRCA2* mutation carriers and the minor alleles of rs8170 and rs67397200 (19p13.1), rs16917302 (*ZNF365*), and rs311499 (20q13.33) are summarized in Table 2. The minor allele of rs8170 at 19p13.1 was strongly associated with risk of breast cancer in *BRCA1* mutation carriers (HR=1.20; 95%CI 1.13–1.28; $p=8.7\times 10^{-9}$) but not *BRCA2* mutation carriers. This result for 12,599 *BRCA1* mutation carriers was consistent with the original finding in the *BRCA1* GWAS using 8,363 *BRCA1* mutation carriers (HR=1.26; 95%CI 1.17–1.35; $p=2.3\times 10^{-9}$). A separate analysis restricted to carriers not used in the *BRCA1* GWAS also confirmed the association

(HR=1.17; 95%CI 1.07–1.27; $p=7.42\times 10^{-4}$) (Supplementary Table 2). Similarly, rs67397200 at 19p13.1, which was imputed in the *BRCA1* GWAS, was strongly associated with breast cancer risk in *BRCA1* carriers (HR=1.17; 95%CI 1.11–1.23; $p=2.4\times 10^{-8}$) (Table 2). There was no evidence of heterogeneity in the HRs across studies for *BRCA1* mutation carriers (Figure 1). However, there was evidence that the per-allele HRs in *BRCA1* mutation carriers for rs8170 ($p=0.015$) and rs67397200 ($p=0.007$) at 19p13.1 decreased with increasing age of diagnosis of breast cancer. Since rs8170 and rs67397200 are located in the same region of 19p13.1 ($r^2=0.58$), we conducted an analysis for the joint effects of these SNPs on breast cancer risk in *BRCA1* mutation carriers ($n=10,173$). When accounting for haplotype structure, rs67397200 remained significant (P for inclusion= 2.75×10^{-3}) and was retained in the model, whereas rs8170 was excluded (P for inclusion=0.18). Rs8170 and rs67397200 were not associated with breast cancer risk for *BRCA2* mutation carriers (Table 2).

Among SNPs identified from the original *BRCA2* GWAS, an analysis of genotype data from 7,132 *BRCA2* mutation carriers confirmed that rs16917302 at the *ZNF365* locus was associated with a decreased risk of breast cancer (HR=0.83; 95%CI 0.75–0.93; $p=7.0\times 10^{-4}$). The association also replicated in the additional carriers, not previously included in the *BRCA2* GWAS (HR=0.84; 95%CI 0.73–0.97; $p=0.017$) (Supplementary Table 2). In contrast, rs311499 from 20q13.3, which was associated with breast cancer risk in the *BRCA2* GWAS (HR=0.72; 95%CI 0.61–0.85; $p=6.6\times 10^{-5}$) (7), was not associated with risk of breast cancer in *BRCA2* carriers in the overall analysis (HR=0.95; 95%CI 0.84–1.07; $p=0.36$) (Table 2) nor the replication study (HR=1.11; 95%CI 0.94–1.31; $p=0.22$) (Supplementary Table 2). There was no evidence for heterogeneity in the HRs across studies for *BRCA2* mutation carriers (Figure 1). HRs for rs16917302 and rs311499 did not vary by age at diagnosis.

To determine whether the inclusion of long-term survivors influenced the results, we repeated our analyses of the four SNPs, excluding *BRCA1* and *BRCA2* mutation carriers diagnosed with breast cancer more than five years before recruitment (prevalent cases). The strength of the associations for rs16917302 at *ZNF365* (per-allele HR=0.85) for *BRCA2* mutation carriers, and for rs8170 (per-allele HR=1.19) and rs67397200 at 19p13.1 (per-allele HR=1.16) for *BRCA1* mutation carriers were essentially unchanged (Supplementary Table 3). There was no influence of mutation type for *BRCA1* mutation carriers on breast cancer risk in the associations between mutations conferring susceptibility to nonsense mediated RNA decay (NMD) (Class 1) and missense or truncating mutations not triggering NMD (Class 2) for any of the SNPs (Supplementary Table 4).

Breast tumors in *BRCA1* mutation carriers are predominantly ER-negative (14) and rs8170 from 19p13.1 is strongly associated with ER-negative but not ER-positive breast cancer in the general population (5). Because of these previous findings, we evaluated whether rs8170 and rs67397200 at 19p13.1, as well as rs311499 at 20q13.3 and rs16917302 at *ZNF365*, were differentially associated with ER-positive and/or ER-negative tumor status in *BRCA1* and *BRCA2* mutation carriers. Although the stratified results suggested a slightly stronger association for the 19p13.1 rs67397200 SNP with ER-negative disease than with ER-positive disease in *BRCA1* mutation carriers (per allele ER-negative HR=1.22; 95%CI 1.14–1.30; $p=4.4\times 10^{-9}$; per allele ER-positive HR=1.14; 95%CI 1.01–1.30; $p=0.040$), the difference was not significant ($p=0.41$) (Table 3). Rs67397200, however, was associated with ER-negative disease (per allele HR=1.29; 95%CI 1.11–1.49; $p=8.7\times 10^{-4}$) but not ER-positive disease (per allele HR=0.92; 95%CI 0.85–1.01; $p=0.074$) in *BRCA2* mutation carriers (p -heterogeneity= 1.5×10^{-4}) (Table 3). The lack of association with rs311499 at 20q13.3 did not vary by ER-status in *BRCA1* or *BRCA2* mutation carriers. For *BRCA2* mutation carriers, the minor allele of rs16917302 at *ZNF365* was inversely associated with

both ER-positive (per allele HR=0.86; 95% CI 0.75–0.97; p=0.016) and ER-negative tumors (per allele HR=0.79; 95% CI 0.62–1.00; p=0.048) (p-heterogeneity=0.56) (Table 3). However, in *BRCA1* mutation carriers rs16917302 was associated with ER-positive (per allele ER-positive HR=0.77; 95% CI 0.62–0.95; p=0.016), but not ER-negative status (p-heterogeneity=0.028) (Table 3).

BRCA1 and *BRCA2* mutations are associated with elevated risk of ovarian cancer. In this CIMBA study 1,465 *BRCA1* mutation carriers and 453 *BRCA2* mutation carriers who developed ovarian cancer were also genotyped for the four SNPs under study. To assess the influence of these SNPs on ovarian cancer risk in *BRCA1* and *BRCA2* mutation carriers we used a competing risk analysis that evaluated the associations with breast and ovarian cancer risk simultaneously. While previous studies did not detect an association between rs8170 at 19p13.1 and ovarian cancer in *BRCA1* or *BRCA2* mutation carriers (5), in this competing risk analysis with larger numbers of *BRCA1* and *BRCA2* mutation carriers, rs8170 was significantly associated with ovarian cancer risk in both *BRCA1* (HR=1.15; 95% CI 1.03–1.29; p=0.015) and *BRCA2* (HR=1.34; 95% CI 1.12–1.62; p=1.9×10⁻³) mutation carriers (Table 4). Similarly rs67397200 at 19p13.1 was associated with ovarian cancer risk in both *BRCA1* (HR=1.16; 95% CI 1.05–1.29; p=3.8×10⁻⁴) and *BRCA2* (HR=1.30; 95% CI 1.10–1.52; p=1.8×10⁻³) mutation carriers (Table 4). Rs311499 at 20q13.3 and rs16917302 at *ZNF365* were not associated with ovarian cancer risk for either *BRCA1* or *BRCA2* mutation carriers (Table 4).

Discussion

GWAS of *BRCA1* and *BRCA2* mutation carriers previously identified variants at 19p13.1, *ZNF365* and 20q13.3 as candidate breast cancer risk modifiers (5, 7). In this study, we further evaluated associations between variants at these loci and both breast and ovarian cancer in *BRCA1* and *BRCA2* mutation carriers. For the first time, we found that both rs8170 and the previously imputed rs67397200 at 19p13.1 were strongly associated with ovarian cancer in both *BRCA1* and *BRCA2* mutation carriers. In addition, we found that rs8170 and rs67397200 at 19p13.1 were associated with breast cancer risk for *BRCA1* and rs16917302 at *ZNF365* was associated with breast cancer in *BRCA2* mutation carriers in this replication study using an independent set of mutation carriers and in the combined analyses of data from the original study and the replication study. In contrast, rs311499 at 20q13.3 showed no association with breast cancer in the replication study. We also report for the first time that the *BRCA1* GWAS SNP rs67397200 is associated with ER-negative breast cancer in *BRCA2* mutation carriers and that the *BRCA2* GWAS SNP rs16917302 is associated with ER-positive disease in *BRCA1* mutation carriers.

The GWAS for breast cancer in *BRCA1* mutation carriers originally identified significant associations between variants at the 19p13.1 locus and risk of breast cancer. Five SNPs including rs8170 from a 39 kb region were associated with risk of disease. In an analysis of joint effects of these SNPs on breast cancer risk, the best model included rs8170 or rs4808611 and rs8100241 or rs2363956 (P for inclusion=7.7×10⁻⁵ and P=6.7×10⁻⁵ for rs8170 and rs8100241, respectively) (5), suggesting that the associations were driven by a single causative variant partially correlated with all five SNPs. Imputation of additional SNPs in the region from the 1000 Genome Project identified eight perfectly correlated SNPs within a 13-kb region that were more significantly associated with breast cancer risk. Of these, we chose rs67397200, which has an $r^2=0.58$ with rs8170 and $r^2=0.37$ with rs8100241/rs2363956, for further genotyping in an effort to determine whether this SNP (or one of the seven other highly correlated SNPs) exhibited stronger associations with breast cancer. In an analysis of rs8170 in 11,669 and rs67397200 in 10,312 *BRCA1* mutation carriers, we observed similarly strong associations with breast cancer for *BRCA1* mutation carriers. In a

joint analysis of rs8170 and rs67397200, allowing for haplotype structure, only rs67397200 remained significant. We were unable to genotype some of the original GWAS SNPs (rs2363956/rs8100241) in the present study and, as a consequence, could not evaluate the joint associations with rs67397200. It is therefore still unclear whether rs67397200 accounts solely for the association signal. The 35kb region containing rs8170 and rs67397200 includes the *ABHD8* (abhydrolase domain containing 8), *ANKLE1* (ankyrin repeat and LEM domain containing 1) and *C19orf62* genes. *C19orf62*, encodes MERIT40 (Mediator of Rap80 Interactions and Targeting 40 kD), a BRCA1 interacting protein that forms a complex with BRCA1-BARD1, Abraxas1, RAP80, BRCC36 and BRCC45 and is required for recruitment and retention of the BRCA1-BARD1 ubiquitin ligase at sites of DNA damage (15). Because alterations in MERIT40 expression or function may modify *BRCA1* activity, variants in the *C19orf62* locus are attractive candidate breast cancer risk modifiers. However, since rs67397200 and the seven other imputed SNPs, that showed the most significant associations with breast cancer risk in *BRCA1* mutation carriers, are located at the 3' end of *ANKLE1* near *ABHD8* it is also possible that one of these genes rather than *C19orf62* is influenced by the underlying causative variants in this region. Further comprehensive genotyping of other common variants and/or rare SNPs from this locus and detailed functional studies will be required to resolve this issue.

Our GWAS for breast cancer in *BRCA2* mutation carriers previously identified strong associations between rs16917302 in the *ZNF365* (dbGENE id: 22891) locus and breast cancer (7). We have now replicated this association for *BRCA2* mutation carriers. Rs16917302 is located within intron 4 of *ZNF365* and is unique to isoform C, the longest of the four isoforms created by alternative splicing sites (16). In independent studies, rs10995195 in *ZNF365*, which is 27kb upstream from and only weakly correlated ($r^2=0.1$) with rs16917302, has been associated with breast cancer risk (8) and with mammographic density (17) in the general population. In addition, a cluster of SNPs located 154kb from rs16917302 in isoform D of *ZNF365* has been associated with Crohn's disease (18–20), and the region has also been implicated in family-based linkage analyses with uric acid nephrolithiasis (21) and hypotrichosis (22). It is unclear whether there is genetic or biologic linkage between these seemingly disparate phenotypes. Further fine-mapping of the *ZNF365* region and functional analyses will be needed to identify the causative variants for each phenotype and to understand the downstream biological effects.

Likewise, rs311499 at 20q13.3 was associated with breast cancer risk in *BRCA2* mutation carriers in the *BRCA2* GWAS (per allele HR=0.72; 95% CI 0.61–0.85; $p=6.6\times 10^{-5}$). However, this association was not confirmed in the replication study described above (HR=1.11; 95% CI 0.94–1.31; $p=0.22$) or in the combined analysis of the discovery and replication stages (HR=0.95; 95% CI 0.84–1.07; $p=0.36$). This result was not unexpected because the association between rs311499 and breast cancer did not reach significance ($p<0.05$) in Stage 2 of the *BRCA2* GWAS (HR=0.86; 95% CI 0.67–1.06; $p=0.13$) (7).

To further characterize the influence of the 19p13.1 and *ZNF365* loci on breast cancer risk, we assessed the strength of association with ER-negative and ER-positive breast cancer in *BRCA1* and *BRCA2* mutation carriers. As reported above, rs67397200 at 19p13.1 was associated with both ER-negative and ER-positive breast cancer in *BRCA1* mutation carriers, whereas rs8170 at 19p13.1 was only associated with ER-negative disease. Interestingly, rs67397200 and rs8170 were also associated with ER-negative breast cancer but not ER-positive breast cancer in *BRCA2* mutation carriers. This is consistent with our previous finding that rs8170 at 19p13.1 is more strongly associated with ER-negative than ER-positive breast cancer in the general population (5). Given that the majority of *BRCA1* breast tumors exhibit a basal breast cancer phenotype (14), it remains to be determined

whether ER-positive basal cases account for the mild association with ER-positive disease in *BRCA1* mutation carriers.

In contrast, we found that rs16917302 in the *ZNF365* locus was associated with both ER-positive and ER-negative disease in *BRCA2* mutation carriers. This was consistent with associations for both ER-positive and ER-negative breast cancer in a recent GWAS of breast cancer cases with a family history of the disease (8). In contrast, among *BRCA1* mutation carriers the association with breast cancer risk was restricted to ER-positive cases. This suggests that refinement of phenotype, perhaps in specific subpopulations, may result in detection of previously hidden associations.

Ovarian cancer is an important component of the cancer phenotype in both *BRCA1* and *BRCA2* mutation carriers. Because breast and ovarian cancer can occur in the same mutation carriers, it has been suggested that susceptibility SNPs common to breast and ovarian cancer may exist in these populations. However, to date none of the SNPs associated with breast cancer risk in *BRCA1* or *BRCA2* mutation carriers have been associated with ovarian cancer risk. Similarly, SNPs in the *BCN2* locus that are associated with ovarian cancer risk in *BRCA1* and *BRCA2* mutation carriers are not associated with breast cancer risk (13). Furthermore, in the general population, only SNPs in the 8q24 locus are known to influence both breast and ovarian cancer, and these appear to be independent disease-specific effects (23). Thus, the recent finding that SNPs at 19p13.1 were associated with breast cancer in *BRCA1* mutation carriers (5) and also with ovarian cancer in the general population (6) raised the possibility that a locus with common influences on breast and ovarian cancer does exist. However, the *BRCA1* GWAS failed to detect any association for 19p13.1 SNPs with ovarian cancer among *BRCA1* mutation carriers (843 ovarian cases) (HR=1.07; 95%CI 0.93–1.24; p=0.33) (5). Here we re-evaluated associations between 19p13.1 SNPs and ovarian cancer using larger numbers of *BRCA1* (n=1312) and *BRCA2* (n=429) carriers diagnosed with ovarian cancer. Rs67397200 at 19p13.1 was associated with ovarian cancer risk in both *BRCA1* (HR=1.16; 95%CI 1.05–1.29; p=3.8×10⁻⁴) and *BRCA2* (HR=1.30; 95%CI 1.10–1.52; p=1.8×10⁻³) mutation carriers. The magnitude of the effect on ovarian cancer risk in *BRCA1* carriers (HR=1.16) was similar to that observed for breast cancer. This is the first locus found to influence both breast and ovarian cancer risk in either *BRCA1* or *BRCA2* mutation carriers.

Including the SNPs from the present study, six loci are now known to modify the risk of breast cancer for *BRCA1* mutation carriers (*CASP8*, *TOX3*, 2q35, 6q25.1, 19p13, and *ZNF365* (ER-positive disease only)) (1–3, 5, 10, 24) and ten loci are known to modify the risk of breast cancer for *BRCA2* mutation carriers (*FGFR2*, *TOX3*, *MAP3K1*, *LSP1*, 2q35, *SLC4A7*, 5p12, *ZNF365*, 1p11.2 and 19p13.1 (ER-negative only)) (5, 7, 10, 24). Taken together, these SNPs result in large variation in the absolute risk of breast cancer for *BRCA1* and *BRCA2* mutation carriers and may further improve our ability to provide individualized risks of breast cancer for *BRCA1* and *BRCA2* mutation carriers.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank all the individuals who took part in these studies and all the researchers, clinicians, technicians and administrative staff who have enabled this work to be carried out. In particular, we thank (BFBOCC) Laima Tikhomirova, (CBCS) Bent Ejlersen, (CNIO) R.M. Alonso, G Pita and R.M. Milne, the Molecular Genetics Laboratory from Hospital de Cruces, (CONSIT) Marco A. Pierotti, Gaia Roversi, Elisa Cattaneo Carla B. Ripamonti and Marilena Morganti (Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy), Bernardo

Bonanni (Istituto Europeo di Oncologia, Milan, Italy); (EMBRACE) Cambridge: Susan Peock, Debra Frost, Steve D. Ellis, Elena Fineberg, Radka Platte. North of Scotland Regional Genetics Service, Aberdeen: Zosia Miedzbrodzka, Helen Gregory. Northern Ireland Regional Genetics Service, Belfast: Patrick Morrison, Lisa Jeffers. West Midlands Regional Clinical Genetics Service, Birmingham: Trevor Cole, Kai-ren Ong, Jonathan Hoffman. South West Regional Genetics Service, Bristol: Alan Donaldson, Margaret James. East Anglian Regional Genetics Service, Cambridge: Joan Paterson, Sarah Downing, Amy Taylor. Medical Genetics Services for Wales, Cardiff: Alexandra Murray, Mark T. Rogers, Emma McCann. St James's Hospital, Dublin & National Centre for Medical Genetics, Dublin: M. John Kennedy, David Barton. South East of Scotland Regional Genetics Service, Edinburgh: Mary Porteous, Sarah Drummond. Peninsula Clinical Genetics Service, Exeter: Carole Brewer, Emma Kivuva, Anne Searle, Selina Goodman, Kathryn Hill. West of Scotland Regional Genetics Service, Glasgow: Rosemarie Davidson, Victoria Murday, Nicola Bradshaw, Lesley Snadden, Mark Longmuir, Catherine Watt, Sarah Gibson, Eshika Haque, Ed Tobias, Alexis Duncan. South East Thames Regional Genetics Service, Guy's Hospital London: Louise Izatt, Chris Jacobs, Caroline Langman, Anna Whaite. North West Thames Regional Genetics Service, Harrow: Huw Dorkins. Leicestershire Clinical Genetics Service, Leicester: Julian Barwell. Yorkshire Regional Genetics Service, Leeds: Julian Adlard, Gemma Serra-Feliu. Cheshire & Merseyside Clinical Genetics Service, Liverpool: Ian Ellis, Catherine Houghton. Manchester Regional Genetics Service, Manchester: D Gareth Evans, Fiona Laloo, Jane Taylor. North East Thames Regional Genetics Service, NE Thames, London: Lucy Side, Alison Male, Cheryl Berlin. Nottingham Centre for Medical Genetics, Nottingham: Jacqueline Eason, Rebecca Collier. Northern Clinical Genetics Service, Newcastle: Fiona Douglas, Oonagh Claber, Irene Jobson. Oxford Regional Genetics Service, Oxford: Lisa Walker, Diane McLeod, Dorothy Halliday, Sarah Durell, Barbara Stayner. The Institute of Cancer Research and Royal Marsden NHS Foundation Trust: Ros Eeles, Susan Shanley, Nazneen Rahman, Richard Houlston, Elizabeth Bancroft, Lucia D'Mello, Elizabeth Page, Audrey Arden-Jones, Kelly Kohut, Jennifer Wiggins, Elena Castro, Anita Mitra, Lisa Robertson. North Trent Clinical Genetics Service, Sheffield: Jackie Cook, Oliver Quarrell, Cathryn Bardsley. South West Thames Regional Genetics Service, London: Shirley Hodgson, Sheila Goff, Glen Brice, Lizzie Winchester, Charlotte Eddy, Vishakha Tripathi, Virginia Attard. Wessex Clinical Genetics Service, Princess Anne Hospital, Southampton: Diana Eccles, Anneke Lucassen, Gillian Crawford, Donna McBride, Sarah Smalley (EMBRACE); (GOG) the investigators of the Australia New Zealand Gynaecological Oncology Group (ANZGOG); (GEMO) Coordinating Centres, Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Centre Hospitalier Universitaire de Lyon / Centre Léon Bérard, & Equipe «Génétique du cancer du sein», Centre de Recherche en Cancérologie de Lyon: Olga Sinilnikova, Sylvie Mazoyer, Laure Barjhoux, Carole Verny-Pierre, Sophie Giraud, Mélanie Léone; and Service de Génétique Oncologique, Institut Curie, Paris: Dominique Stoppa-Lyonnet, Marion Gauthier-Villars, Bruno Buecher, Claude Houdayer, Virginie Moncoutier, Muriel Belotti, Carole Tirapo, Antoine de Pauw. Institut Gustave Roussy, Villejuif: Brigitte Bressac-de-Paillerets, Audrey Remenieras, Véronique Byrde, Olivier Caron, Gilbert Lenoir. Centre Jean Perrin, Clermont-Ferrand: Yves-Jean Bignon, Nancy Uhrhammer. Centre Léon Bérard, Lyon: Christine Lasset, Valérie Bonadona. Centre François Baclesse, Caen: Agnès Hardouin, Pascaline Berthet. Institut Paoli Calmettes, Marseille: Hagay Sobol, Violaine Bourdon, Tetsuro Noguchi, François Eisinger. Groupe Hospitalier Pitié-Salpêtrière, Paris: Florence Coulet, Chrystelle Colas, Florent Soubrier. CHU de Arnaud-de-Villeneuve, Montpellier: Isabelle Couplier, Pascal Pujol. Centre Oscar Lambret, Lille: Jean-Philippe Peyrat, Joëlle Fournier, Françoise Révillon, Philippe Vennin, Claude Adenis. Hôpital René Huguenin/Institut Curie, St Cloud: Etienne Rouleau, Rosette Lidereau, Liliane Demange, Catherine Nagues. Centre Paul Strauss, Strasbourg: Danièle Muller, Jean-Pierre Fricker. Institut Bergonié, Bordeaux: Emmanuelle Barouk-Simonet, Françoise Bonnet, Virginie Buben, Nicolas Sevenet, Michel Longy. Institut Claudius Regaud, Toulouse: Christine Toulas, Rosine Guimbaud, Laurence Gladieff, Viviane Feillel. CHU de Grenoble: Dominique Leroux, Hélène Dreyfus, Christine Rebuschung, Magalie Peysselon. CHU de Dijon: Fanny Coron, Laurence Faivre. CHU de St-Etienne: Fabienne Prieur, Marine Lebrun, Caroline Kientz. Hôtel Dieu Centre Hospitalier, Chambéry: Sandra Fert Ferrer. Centre Antoine Lacassagne, Nice: Marc Frénay. CHU de Limoges: Laurence Vénat-Bouvet. CHU de Nantes: Capucine Delnatte. CHU Bretonneau, Tours: Isabelle Mortemousque. Creighton University, Omaha, USA: Henry T.Lynch, Carrie L.Snyder; (HCSC) Red Tematica de Investigacion Corporativa en Cancer (ISCIII); (HEBCS) Kristiina Aittomäki, Kirsimari Aaltonen and Carl Blomqvist and Tuomas Heikkinen and research nurse Irja Erkkilä; (HEBON) Netherlands Cancer Institute, Amsterdam, NL: F.B.L. Hogervorst, S. Verhoef, M. Verheus, L.J. van 't Veer, F.E. van Leeuwen, M.A. Rookus; Erasmus Medical Center, Rotterdam, NL: M. Collée, A.M.W. van den Ouweland, A. Jager, M.J. Hoening, M.M.A. Tilanus-Linthorst, C. Seynaeve; Leiden University Medical Center, NL, Leiden: C.J. van Asperen, J.T. Wijnen, M.P. Vreeswijk, R.A. Tollenaar, P. Devilee; Radboud University Nijmegen Medical Center, Nijmegen, NL: M.J. Ligtenberg, N. Hoogerbrugge; University Medical Center Utrecht, Utrecht, NL: M.G. Ausems, R.B. van der Luijt; Amsterdam Medical Center, NL: C.M. Aalfs, T.A. van Os; VU University Medical Center, Amsterdam, NL: J.J.P. Gille, Q. Waisfisz, H.E.J. Meijers-Heijboer; University Hospital Maastricht, Maastricht, NL: E.B. Gomez-Garcia, C.E. van Roozendaal, Marinus J. Blok, B. Caanen; University Medical Center Groningen University, NL: J.C. Oosterwijk, A.H. van der Hout, M.J. Mourits; The Netherlands Foundation for the detection of hereditary tumours, Leiden, NL: H.F. Vasen; (INHERIT) Stéphane Dubois, Martine Dumont, Martine Tranchant (Cancer Genomics Laboratory, CRCHUQ) and Sylvie Desjardins and Marc-André Rodrigue (Plateforme de séquençage et de génotypage des génome du CRCHUL/CHUQ); (kConFab)

Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study; (KUMC) JoEllen Weaver; (OCGN) Teresa Selander, Nayana Weerasooriya and members of the Ontario Cancer Genetics Network; (OSU-CCG) Leigha Senter and Kevin Sweet

and the Ohio State University Human Genetics Sample Bank; (SEABASS) Cancer Research Initiatives Foundation (Malaysia), University Malaya (Malaysia), National University Hospital (Singapore), University Kebangsaan Malaysia (Malaysia), Hospital Kuala Lumpur (Malaysia) and Putrajaya Hospital (Malaysia); (SWE-BRCA) Per Karlsson, Margareta Nordling, Annika Bergman and Zakaria Einbeigi, Gothenburg, Sahlgrenska University Hospital, Marie Stenmark-Askmal and Sigrun Liedgren, Linköping University Hospital, Åke Borg, Niklas Loman, Håkan Olsson, Maria Soller, Helena Jernström, Katja Harbst and Karin Henriksson, Lund University Hospital, Annika Lindblom, Brita Arver, Anna von Wachenfeldt, Annelie Liljegren, Gisela Barbany-Bustinza and Johanna Rantala, Stockholm, Karolinska University Hospital, Beatrice Melin, Henrik Grönberg, Eva-Lena Stattin and Monica Emanuelsson, Umeå University Hospital, Hans Ehrencrona, Richard Rosenquist and Niklas Dahl, Uppsala University Hospital; (UKGRFOCR) Roswell Park Alliance Foundation for their continued support of the Gilda Radner Ovarian Family Cancer Registry, Carole Pye, Patricia Harrington, Eva Wozniak, Kirsten Moysich and Lara Sucheston

Grant Support


This research was supported by NIH grant CA128978, an NCI Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA116201), a U.S. Department of Defence Ovarian Cancer Idea award (W81XWH-10-1-0341) and grants from the Breast Cancer Research Foundation and the Komen Foundation for the Cure. This work was also supported by Cancer Research UK grants C12292/A11174 and C1287/A10118. The research leading to these results has received funding from the European Community's Seventh Framework Programme under grant agreement n° 223175 (HEALTH-F2-2009-223175). Support was also provided by the Canadian Institutes of Health Research for the "CIHR Team in Familial Risks of Breast Cancer" program and by the Canadian Breast Cancer Research Alliance-grant #019511. A.C.A. is a CR-UK Senior Cancer Research Fellow. D.F.E. is CR-UK Principal Research Fellow. GCT is a NHMRC Senior Principal Research Fellow. BFBOCC was supported by the Research Council of Lithuania grant LIG-19/2010 to R.J. BMBSA was supported by grants from the Cancer Association of South Africa (CANSA) to Elizabeth J. van Rensburg. BCFR was supported by the National Cancer Institute, National Institutes of Health under RFA-CA-06-503 and through cooperative agreements with members of the Breast Cancer Family Registry (BCFR) and Principal Investigators, including Cancer Care Ontario (U01 CA69467), Columbia University (U01 CA69398), Fox Chase Cancer Center (U01 CA69631), Huntsman Cancer Institute (U01 CA69446), Cancer Prevention Institute of California (formerly the Northern California Cancer Center) (U01 CA69417), University of Melbourne (U01 CA69638), and Research Triangle Institute Informatics Support Center (RFP No. N02PC45022-46). CBCS was supported by The Neye Foundation. CNIO was partially supported by Fundación Mutua Madrileña, Asociación Española Contra el Cáncer, the Spanish Ministry of Science and Innovation (FIS PI08 1120) and the Basque Foundation for Health Innovation and Research (BIOEF): BIO07/CA/006. CONSIT TEAM was supported by grants from Ministero della Salute (Extraordinary National Cancer Program 2006 "Alleanza contro il Cancro" to LV and PR, and "Progetto Tumori Femminili" to PR), Ministero dell'Università e Ricerca (RBLAO3-BETH to PR), Fondazione Italiana per la Ricerca sul Cancro (Special Project "Hereditary tumors" to PR), Associazione Italiana per la Ricerca sul Cancro (4017 to PP) and by funds from Italian citizens who allocated the 5×1000 share of their tax payment in support of the Fondazione IRCCS Istituto Nazionale Tumori, according to Italian laws (INT-Institutional strategic projects "5×1000"). The DKFZ study was supported by funds from the DKFZ. EMBRACE was supported by Cancer Research UK Grants C1287/A10118 and C1287/A11990. D. Gareth Evans and Fiona Laloo were supported by an NIHR grant to the Biomedical Research Centre, Manchester. The Investigators at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust were supported by an NIHR grant to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. Ros Eeles, Elizabeth Bancroft and Lucia D'Mello were supported by Cancer Research UK Grant C5047/A8385. GC-HBOC was supported by a grant of the German Cancer Aid (grant 109076) and by the Centre of Molecular Medicine Cologne (CMMC). The GEMO study was supported by the Ligue Nationale Contre le Cancer; Association for International Cancer Research Grant (AICR-07-0454); and the Association "Le cancer du sein, parlons-en!" Award. The Georgetown study was supported by the Familial Cancer Registry at Georgetown University (NIH/NCI grant P30-CA051008), the Cancer Genetics Network (HHSN261200744000C), and Swing Fore the Cure. GOG was supported through funding provided by both intramural (Clinical Genetics Branch, DCEG) and extramural (Community Oncology and Prevention Trials Program—COPTRG) NCI programs. KAP is the Cancer Council Victoria, Colebatch Clinical Research Fellow. HEBCS was supported by the Helsinki University Central Hospital Research Fund, Academy of Finland (132473), the Finnish Cancer Society, and the Sigrid Juselius Foundation. The HEBON study was supported by the Dutch Cancer Society grants NKI1998-1854, NKI2004-3088, NKI2007-3756 and the ZonMW grant 91109024. HUNBOCS was supported by the Hungarian Research Grant KTLA-OTKA CK-80745. ICO was supported by Asociación Española Contra el Cáncer, Spanish Health Research Fund; Carlos III Health Institute; Catalan Health Institute and Autonomous Government of Catalonia; contract grant numbers ISCHIRETIC RD06/0020/1051, PI10/01422, PI10/31488 and 2009SGR290. IHCC was supported by a Polish Foundation of Science award to K.J., a fellow of International PhD program, Postgraduate School of Molecular Medicine, Warsaw Medical University. ILUH was supported by the Icelandic Association "Walking for Breast Cancer Research" and by the Landspítali University Hospital Research Fund. INHERIT was supported J.S., Chairholder of the Canada Research Chair in Oncogenetics. IOVHBOCS was supported by Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR) and "Ministero della Salute" ("Progetto Tumori Femminili and grant numbers RFPS 2006-5-341353, ACC2/R6.9"). kConFab was supported by grants from the National Breast Cancer Foundation, the

National Health and Medical Research Council (NHMRC) and by the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. The kConFab Clinical Follow Up Study was funded by the NHMRC [145684, 288704, 454508]. ABS is supported by a NHMRC Senior Research Fellowship. A.K.G. was funded by U01CA69631, 5U01CA113916, and the Eileen Stein Jacoby Fund while at FCCC. The author acknowledges support from The University of Kansas Cancer Center and the Kansas Bioscience Authority Eminent Scholar Program. A.K.G. is the Chancellors Distinguished Chair in Biomedical Sciences endowed Professor. The McGill study was supported by the Jewish General Hospital Weekend to End Breast Cancer. MT holds a Fonds de la Recherche en Santé du Québec clinician-scientist award. The MSKCC study was supported by the Starr Cancer Consortium, the Breast Cancer Research Foundation, the Norman and Carol Stone Cancer Research Initiative, the Kate and Robert Niehaus Clinical Cancer Research Initiative, the Lymphoma Foundation, and the Sabin Family Research Initiative. The NCI study was supported by the Intramural Research Program of the US National Cancer Institute, and by support services contracts NO2-CP-11019-50 and N02-CP-65504 with Westat, Inc, Rockville, MD. NNPIO was supported by the Russian Federation for Basic Research (grants 10-04-92601, 10-04-92110, 11-04-00227) and the Federal Agency for Science and Innovations (contract 16.512.11.2237). OCGN was supported by Cancer Care Ontario and the US National Cancer Institute, National Institutes of Health under RFA # CA-06-503 and through cooperative agreements with members of the Breast Cancer Family Registry (BCFR) and Principal Investigators. OSU-CCG was supported by the Ohio State University Comprehensive Cancer Center. PBCS was supported by an Istituto Toscano Tumori grant to MAC. SEABASS was supported by CARIF and University Malaya. The UCSF study was supported by the Helen Diller Family Comprehensive Cancer Center at UCSF, the Avon Foundation, and the Center for Translational and Policy Research in Personalized Medicine (TRANSPERS), NIH/NCI P01 CA130818-02A1. UKFOCR was supported by a project grant from CRUK to Paul Pharoah. The UPENN study was supported Komen Foundation for the Cure to SMD, the Breast Cancer Research Foundation to KLN, and National Institutes of Health grants R01-CA083855 and R01-CA102776 to TRR. WCRI was supported by the American Cancer Society Clinical Research Professorship #SIOP-06-258-06-COUN.

References


1. Antoniou AC, Beesley J, McGuffog L, Sinilnikova OM, Healey S, Neuhausen SL, et al. Common breast cancer susceptibility alleles and the risk of breast cancer for BRCA1 and BRCA2 mutation carriers: implications for risk prediction. *Cancer Res.* 2010; 70:9742–9754. [PubMed: 21118973]
2. Antoniou AC, Kartsonaki C, Sinilnikova OM, Soucy P, McGuffog L, Healey S, et al. Common alleles at 6q25.1 and 1p11.2 are associated with breast cancer risk for BRCA1 and BRCA2 mutation carriers. *Hum Mol Genet.* 2011; 20:3304–3321. [PubMed: 21593217]
3. Engel C, Versmold B, Wappenschmidt B, Simard J, Easton DF, Peock S, et al. Association of the variants CASP8 D302H and CASP10 V410I with breast and ovarian cancer risk in BRCA1 and BRCA2 mutation carriers. *Cancer Epidemiol Biomarkers Prev.* 2010; 19:2859–2868. [PubMed: 20978178]
4. Broeks A, Schmidt MK, Sherman ME, Couch FJ, Hopper JL, Dite GS, et al. Low penetrance breast cancer susceptibility loci are associated with specific breast tumor subtypes: findings from the Breast Cancer Association Consortium. *Hum Mol Genet.* 2011; 20:3289–3303. [PubMed: 21596841]
5. Antoniou AC, Wang X, Fredericksen ZS, McGuffog L, Tarrell R, Sinilnikova OM, et al. A locus on 19p13 modifies risk of breast cancer in BRCA1 mutation carriers and is associated with hormone receptor-negative breast cancer in the general population. *Nat Genet.* 2010; 42:885–892. [PubMed: 20852631]
6. Bolton KL, Tyrer J, Song H, Ramus SJ, Notaridou M, Jones C, et al. Common variants at 19p13 are associated with susceptibility to ovarian cancer. *Nat Genet.* 2010; 42:880–884. [PubMed: 20852633]
7. Gaudet MM, Kirchhoff T, Green T, Vijai J, Korn JM, Guiducci C, et al. Common genetic variants and modification of penetrance of BRCA2-associated breast cancer. *PLoS Genet.* 2010; 6:e1001183. [PubMed: 21060860]
8. Turnbull C, Ahmed S, Morrison J, Pernet D, Renwick A, Maranian M, et al. Genome-wide association study identifies five new breast cancer susceptibility loci. *Nat Genet.* 2010; 42:504–507. [PubMed: 20453838]
9. Antoniou AC, Sinilnikova OM, Simard J, Leone M, Dumont M, Neuhausen SL, et al. RAD51 135G-->C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies. *Am J Hum Genet.* 2007; 81:1186–1200. [PubMed: 17999359]

10. Antoniou AC, Cunningham AP, Peto J, Evans DG, Lalloo F, Narod SA, et al. The BOADICEA model of genetic susceptibility to breast and ovarian cancers: updates and extensions. *Br J Cancer*. 2008; 98:1457–1466. [PubMed: 18349832]
11. Perrin-Vidoz L, Sinilnikova OM, Stoppa-Lyonnet D, Lenoir GM, Mazoyer S. The nonsense-mediated mRNA decay pathway triggers degradation of most BRCA1 mRNAs bearing premature termination codons. *Hum Mol Genet*. 2002; 11:2805–2814. [PubMed: 12393792]
12. Mulligan AM, Couch FJ, Barrowdale D, Domchek SM, Eccles D, Nevanlinna H, et al. Common breast cancer susceptibility alleles are associated with tumor subtypes in BRCA1 and BRCA2 mutation carriers: results from the Consortium of Investigators of Modifiers of BRCA1/2. *Breast Cancer Research BCR*. 2011; 13:R110. [PubMed: 22053997]
13. Ramus SJ, Kartsonaki C, Gayther SA, Pharoah PD, Sinilnikova OM, Beesley J, et al. Genetic variation at 9p22.2 and ovarian cancer risk for BRCA1 and BRCA2 mutation carriers. *J Natl Cancer Inst*. 2010; 103:105–116. [PubMed: 21169536]
14. Lakhani SR, Van De Vijver MJ, Jacquemier J, Anderson TJ, Osin PP, McGuffog L, et al. The pathology of familial breast cancer: predictive value of immunohistochemical markers estrogen receptor, progesterone receptor, HER-2, and p53 in patients with mutations in BRCA1 and BRCA2. *J Clin Oncol*. 2002; 20:2310–2318. [PubMed: 11981002]
15. Shao G, Patterson-Fortin J, Messick TE, Feng D, Shanbhag N, Wang Y, et al. MERIT40 controls BRCA1-Rap80 complex integrity and recruitment to DNA double-strand breaks. *Genes Dev*. 2009; 23:740–754. [PubMed: 19261746]
16. Gianfrancesco F, Esposito T, Casu G, Maninchedda G, Roberto R, Pirastu M. Emergence of Talanin protein associated with human uric acid nephrolithiasis in the Hominidae lineage. *Gene*. 2004; 339:131–138. [PubMed: 15363853]
17. Lindstrom S, Vachon CM, Li J, Varghese J, Thompson D, Warren R, et al. Common variants in ZNF365 are associated with both mammographic density and breast cancer risk. *Nat Genet*. 2011; 43:185–187. [PubMed: 21278746]
18. Rioux JD, Xavier RJ, Taylor KD, Silverberg MS, Goyette P, Huett A, et al. Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat Genet*. 2007; 39:596–604. [PubMed: 17435756]
19. Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature*. 2007; 447:661–678. [PubMed: 17554300]
20. Barrett JC, Hansoul S, Nicolae DL, Cho JH, Duerr RH, Rioux JD, et al. Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's disease. *Nat Genet*. 2008; 40:955–962. [PubMed: 18587394]
21. Gianfrancesco F, Esposito T, Ombra MN, Forabosco P, Maninchedda G, Fattorini M, et al. Identification of a novel gene and a common variant associated with uric acid nephrolithiasis in a Sardinian genetic isolate. *Am J Hum Genet*. 2003; 72:1479–1491. [PubMed: 12740763]
22. Naz G, Ali G, Naqvi SK, Azeem Z, Ahmad W. Mapping of a novel autosomal recessive hypotrichosis locus on chromosome 10q11.23–22.3. *Hum Genet*. 2010; 127:395–401. [PubMed: 20054564]
23. Ghousaini M, Song H, Koessler T, Al Olama AA, Kote-Jarai Z, Driver KE, et al. Multiple loci with different cancer specificities within the 8q24 gene desert. *J Natl Cancer Inst*. 2008; 100:962–966. [PubMed: 18577746]
24. Antoniou AC, Sinilnikova OM, McGuffog L, Healey S, Nevanlinna H, Heikkinen T, et al. Common variants in LSP1, 2q35 and 8q24 and breast cancer risk for BRCA1 and BRCA2 mutation carriers. *Hum Mol Genet*. 2009; 18:4442–4456. [PubMed: 19656774]



1B

- UK/Eire
- Austria
- Australia
- Canada
- Denmark
- France
- Finland
- Germany
- Hungary
- Italy
- Latvia/Lithuania
- Russia
- SouthAfrica
- Spain/Greece
- Sweden
- USA
- SUMMARY**






Figure 1. Forest plots of the associations by country of residence of *BRCA1* and *BRCA2* mutation carriers with breast cancer risk overall

(A–C) Squares indicate the country specific per-allele HR estimates for SNPs (A) rs8170 for *BRCA1* mutation carriers, (B) rs67397200 for *BRCA1* mutation carriers and (C) rs16917302 for *BRCA2* mutation carriers. The area of the square is proportional to the inverse of the variance of the estimate. Horizontal lines indicate 95% Confidence Intervals (CIs).

Table 1Summary characteristics for the 19,731 eligible *BRCA1* and *BRCA2* mutation carriers* used in the analysis

Characteristic	<i>BRCA1</i>		<i>BRCA2</i>	
	Unaffected	Breast Cancer	Unaffected	Breast Cancer
Number	6209	6390	3322	3810
Person-Years follow-up	264903	263068	147053	168201
Median Age at Censure (IQR ¹)	42 (34–50)	40 (34–47)	43 (34–53)	43 (37–50)
Age at Censure, N (%)				
<30	1189 (19.2)	691 (10.8)	611 (18.4)	306 (8.0)
30–39	1661 (26.8)	2445 (38.3)	834 (25.1)	1141 (30.0)
40–49	1765 (28.4)	2191 (34.3)	865 (26.0)	1394 (36.6)
50–59	1058 (17.0)	812 (12.7)	566 (17.0)	687 (18.0)
60–69	380 (6.1)	198 (3.1)	302 (9.1)	226 (5.9)
70+	156 (2.5)	53 (0.8)	144 (4.3)	56 (1.5)
Year of birth, N (%)				
<1920	28 (0.5)	30 (0.5)	23 (0.7)	44 (1.2)
1920–29	131 (2.1)	196 (3.1)	99 (3.0)	167 (4.4)
1930–39	369 (5.9)	516 (8.1)	232 (7.0)	430 (11.3)
1940–49	832 (13.4)	1341 (21.0)	458 (13.8)	896 (23.5)
1950–59	1409 (22.7)	1989 (31.1)	691 (20.8)	1160 (30.5)
1960–69	1703 (27.4)	1666 (26.1)	902 (27.2)	868 (22.8)
1970+	1737 (28.0)	652 (10.2)	917 (27.6)	245 (6.4)
Mutation Class, N (%)				
Class 1 ²	4063 (65.4)	3878 (60.7)	3114 (93.7)	3520 (92.4)
Class 2 ²	1780 (28.7)	1973 (30.9)	72 (2.2)	100 (2.6)
Other	366 (5.9)	539 (8.4)	136 (4.1)	190 (5.0)

¹IQR: Interquartile range²See methods for definitions

* Carriers of self reported European ancestry only.

Table 2

Evaluation of associations between SNPs and breast cancer risk among *BRCA1* and *BRCA2* mutation carriers of European ancestry

SNP/ Mutation	Genotype	Unaffected N (%)	Affected ^a N (%)	HR	95% CI	p-value
rs8170 – 19p13.1						
<i>BRCA 1</i>	GG	3870 (67.5)	3755 (63.3)	1.00		
	AG	1689 (29.4)	1950 (32.9)	1.22	1.14 – 1.31	
	AA	178 (3.1)	227 (3.8)	1.35	1.13 – 1.62	
	per allele			1.20	1.13 – 1.28	8.7×10 ⁻⁹
<i>BRCA 2</i>						
	GG	2047 (66.3)	2501 (68.2)	1.00		
	AG	931 (30.2)	1026 (28.0)	0.93	0.84 – 1.03	
	AA	108 (3.5)	138 (3.8)	1.16	0.89 – 1.52	
	per allele			0.98	0.90 – 1.07	0.67
rs67397200 – 19p13.1						
<i>BRCA 1</i>	CC	2536 (51.0)	2455 (46.0)	1.00		
	GC	2022 (40.6)	2397 (44.9)	1.24	1.16 – 1.34	
	GG	415 (8.4)	487 (9.1)	1.25	1.10 – 1.43	
	per allele			1.17	1.11 – 1.23	2.4×10 ⁻⁸
<i>BRCA 2</i>						
	CC	1553 (49.8)	1871 (50.7)	1.00		
	GC	1302 (41.7)	1494 (40.5)	0.95	0.87 – 1.04	
	GG	265 (8.5)	323 (8.8)	1.07	0.91 – 1.27	
	per allele			1.00	0.93 – 1.07	0.97
rs311499 – 20q13.3						
<i>BRCA 1</i>	GG	5346 (86.2)	5484 (85.9)	1.00		
	AG	816 (13.2)	873 (13.7)	1.03	0.94 – 1.13	
	AA	41 (0.7)	28 (0.4)	0.67	0.42 – 1.08	
	per allele			1.00	0.91 – 1.09	0.94
<i>BRCA 1</i>	GG	2873 (86.6)	3312 (87.0)	1.00		
	AG	429 (13.0)	475 (12.5)	0.94	0.82 – 1.07	
	AA	16 (0.5)	21 (0.6)	0.97	0.60 – 1.57	
	per allele			0.95	0.84 – 1.07	0.36

SNP/ Mutation	Genotype	Unaffected N (%)	Affected ^a N (%)	HR	95% CI	p-value
rs16917302 - 10q21.2						
<i>BRCA 1</i>	AA	4913 (79.3)	5084 (79.7)	1.00		
	CA	1216 (19.6)	1222 (19.2)	0.96	0.88 – 1.01	
	CC	71 (1.1)	73 (1.1)	0.94	0.69 – 1.27	
	per allele			0.96	0.89 – 1.03	0.27
<i>BRCA 2</i>	AA	2583 (77.9)	3101 (81.5)	1.00		
	CA	691 (20.8)	674 (17.7)	0.82	0.74 – 0.92	
	CC	41 (1.2)	32 (0.8)	0.78	0.49 – 1.23	
	per allele			0.83	0.75 – 0.93	7.0×10 ⁻⁴

^aBreast cancer

Table 3

Associations between SNPs and breast cancer risk by estrogen receptor (ER) status of breast cancer cases among women with *BRCA1* and *BRCA2* mutations

SNP/	Unaff	Affected ^a (N)	ER+		ER-		Case het		
Mutation	N	ER-	ER+	HR	95% CI	HR	95% CI	p-value ^b	
								p-trend	
rs8170 – 19p13.1									
<i>BRCA1</i>	4483	1820	541	1.12	0.96 – 1.29	0.15	1.23	1.14 – 1.33	2.0×10 ⁻⁷
<i>BRCA2</i>	2738	401	1343	0.94	0.85 – 1.05	0.26	1.18	0.99 – 1.40	0.058
rs67397200 – 19p13.1									
<i>BRCA1</i>	4486	1821	542	1.14	1.01 – 1.30	0.040	1.22	1.14 – 1.30	4.4×10 ⁻⁹
<i>BRCA2</i>	2733	401	1349	0.92	0.85 – 1.01	0.074	1.29	1.11 – 1.49	8.7×10 ⁻⁴
rs311499 – 20q13.3									
<i>BRCA1</i>	4898	1890	559	1.07	0.87 – 1.31	0.51	0.95	0.85 – 1.06	0.35
<i>BRCA2</i>	2930	406	1372	0.95	0.82 – 1.09	0.48	0.83	0.63 – 1.10	0.19
rs16917302 - 10q21.2									
<i>BRCA1</i>	4897	1888	558	0.77	0.62 – 0.95	0.016	1.01	0.92 – 1.11	0.85
<i>BRCA2</i>	2927	406	1372	0.86	0.75 – 0.97	0.016	0.79	0.62 – 1.00	0.048

^aBreast cancer

^bp-value for heterogeneity in the associations with ER-positive and ER-negative breast cancer

ER+: ER positive, ER-: ER negative, Unaff: Unaffected

Table 4

Associations with SNPs and breast and ovarian cancer risk using a competing risk analysis model among *BRCA1* and *BRCA2* mutation carriers of European ancestry

SNP/	Unaffected		Breast Cancer		Ovarian Cancer		Breast Cancer		Ovarian Cancer	
Mutation	Genotype	N (%)	N (%)	HR	95% CI	p-value	HR	95% CI	p-value	
rs8170 – 19p13.1										
<i>BRCA1</i>	GG	2972 (67.9)	3730 (63.3)	923 (66.0)	1.00		1.00			
	AG	1269 (29.0)	1936 (32.9)	434 (31.0)	1.26	1.17 – 1.36	1.23	1.08 – 1.42		
	AA	139 (3.2)	224 (3.8)	42 (3.0)	1.34	1.10 – 1.63	1.04	0.72 – 1.50		
	per allele				1.22	1.14 – 1.30	2.1 × 10 ⁻⁹	1.15	1.03 – 1.29	0.015
<i>BRCA2</i>	GG	1788 (67.0)	2494 (68.2)	266 (62.2)	1.00		1.00			
	AG	796 (29.9)	1024 (28.0)	137 (32.0)	0.95	0.85 – 1.05	1.17	0.93 – 1.47		
	AA	83 (3.1)	138 (3.8)	25 (5.8)	1.37	1.05 – 1.80	2.72	1.65 – 4.48		
	per allele				1.02	0.94 – 1.12	0.62	1.12 – 1.62		1.9 × 10 ⁻³
rs67397200 – 19p13.1										
<i>BRCA1</i>	CC	1903 (51.5)	2436 (46.0)	652 (49.7)	1.00		1.00			
	GC	1498 (40.5)	2381 (44.9)	540 (41.2)	1.28	1.18 – 1.38	1.16	1.01 – 1.33		
	GG	298 (8.1)	484 (9.1)	120 (9.2)	1.33	1.16 – 1.53	1.36	1.07 – 1.73		
	per allele				1.20	1.13 – 1.27	4.5 × 10 ⁻¹⁰	1.16	1.05 – 1.29	3.8 × 10 ⁻⁴
<i>BRCA2</i>	CC	1363 (50.5)	1866 (50.7)	194 (45.2)	1.00		1.00			
	GC	1123 (41.6)	1489 (40.5)	184 (42.9)	0.96	0.87 – 1.06	1.15	0.92 – 1.44		
	GG	214 (7.9)	323 (8.8)	51 (11.9)	1.18	0.99 – 1.41	1.95	1.37 – 2.77		
	per allele				1.03	0.96 – 1.11	0.39	1.10 – 1.52		1.8 × 10 ⁻³
rs311499 – 20q13.3										
<i>BRCA1</i>	GG	4115 (86.0)	5442 (85.9)	1273 (86.9)	1.00		1.00			
	AG	637 (13.3)	869 (13.7)	183 (12.5)	1.01	0.92 – 1.12	0.88	0.74 – 1.05		
	AA	32 (0.7)	28 (0.4)	9 (0.6)	0.70	0.42 – 1.17	1.16	0.47 – 2.87		
	per allele				0.99	0.90 – 1.08	0.77	0.91	0.77 – 1.07	0.25
<i>BRCA2</i>	GG	2492 (86.7)	3303 (87.0)	390 (86.1)	1.00		1.00			
	AG	372 (12.9)	474 (12.5)	58 (12.8)	0.93	0.82 – 1.07	0.92	0.68 – 1.26		
	AA	11 (0.4)	21 (0.6)	5 (1.1)	1.09	0.68 – 1.74	2.23	0.80 – 6.22		

SNP/	Unaffected		Breast Cancer		Ovarian Cancer		Breast Cancer		Ovarian Cancer		
Mutation	Genotype	N (%)	N (%)	N (%)	N (%)	HR	95% CI	p-value	HR	95% CI	p-value
	per allele					0.95	0.84 – 1.08	0.44	1.02	0.76 – 1.37	0.88
rs16917302 - 10q21.2											
<i>BRCA1</i>	AA	3784 (79.2)	5044 (79.7)	1169 (79.8)	1.00				1.00		
	CA	937 (19.6)	1216 (19.2)	285 (19.5)	0.96	0.88 – 1.04			0.97	0.84 – 1.13	
	CC	60 (1.3)	73 (1.2)	11 (0.8)	0.86	0.62 – 1.19			0.47	0.25 – 0.92	
	per allele								0.92	0.80 – 1.05	0.20
<i>BRCA2</i>	AA	2237 (77.9)	3094 (81.5)	353 (78.1)	1.00				1.00		
	CA	601 (20.9)	671 (17.7)	93 (20.6)	0.81	0.72 – 0.92			0.93	0.72 – 1.21	
	CC	35 (1.2)	32 (0.8)	6 (1.3)	0.80	0.50 – 1.30			1.21	0.46 – 3.18	
	per allele								0.83	0.74 – 0.92	5.8×10 ⁻⁴
									0.96	0.76 – 1.22	0.76