



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Comorbidity and COVID-19: investigating the relationship between medical and psychological well-being**

Stafford, O., Berry, A., Taylor, L. K., Wearen, S., Prendergast, C., Murphy, E., Shevlin, M., McHugh, L., Carr, A., & Burke, T. (2021). Comorbidity and COVID-19: investigating the relationship between medical and psychological well-being. *Irish Journal of Psychological Medicine*. <https://doi.org/10.1017/ipm.2021.37>

### **Published in:**

Irish Journal of Psychological Medicine

### **Document Version:**

Peer reviewed version

### **Queen's University Belfast - Research Portal:**

[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

© 2021 The Authors.

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

**Title: Comorbidity and COVID-19: Investigating the Relationship between Medical and Psychological Well-being**

Owen Stafford<sup>1</sup> (MSc; Research Assistant, UCD),

Anna Berry<sup>1,2</sup> (MPhil; Trainee Clinical Psychologist, HSE/UCD),

Laura K. Taylor<sup>1, 3</sup> (PhD; Assistant Professor, UCD; Reader, QUB),

Sinead Wearen<sup>1</sup> (MSc; Research Assistant, UCD),

Cian Prendergast<sup>1</sup> (MSc; Research Assistant; UCD)

Eddie Murphy<sup>1,2</sup> (DPsychSc; Principal Psychology Manager, HSE),

Mark Shevlin<sup>4</sup> (PhD; Professor, University of Ulster),

Louise McHugh<sup>1</sup> (PhD; Professor, UCD),

Alan Carr<sup>1</sup> (PhD; Clinical Psychologist and Professor, UCD)

Tom Burke<sup>1</sup> (PhD DPsychSc; Clinical Psychologist and Adjunct Assistant Professor, UCD).

<sup>1</sup> School of Psychology, University College Dublin, Dublin, Ireland

<sup>2</sup> Health Service Executive, CHO 8, Laois and Offaly, Ireland

<sup>3</sup> School of Psychology, Queen's University Belfast, Northern Ireland

<sup>4</sup> University of Ulster (Psychology), Coleraine, Northern Ireland

\*Corresponding author: Dr Tom Burke, School of Psychology, University College Dublin. E:

[tom.m.burke@ucd.ie](mailto:tom.m.burke@ucd.ie); Tel: +353(0) 17168120

## Abstract

**Objective:** The unprecedented occurrence of a global pandemic is accompanied by both physical and psychological burdens that may impair quality of life. Research relating to COVID-19 aims to determine the effects of the pandemic on vulnerable populations who are at high risk of developing negative health or psychosocial outcomes. Having an ongoing medical condition during a pandemic may lead to greater psychological distress. Increased psychological distress may be due to preventative public health measures (e.g. lockdown), having an ongoing medical condition, or a combination of these factors.

**Methods:** This study analyses data from an online cross-sectional national survey of adults in Ireland and investigates the relationship between comorbidity and psychological distress. Those with a medical condition (n=128) were compared to a control group without a medical condition (n=128) and matched according to age, gender, annual income, education, and work status during COVID-19. Participants and data were obtained during the first public lockdown in Ireland (27.03.2020–08.06.2020).

**Results:** Individuals with existing medical conditions reported significantly higher levels of anxiety ( $p < .01$ ) and felt less gratitude ( $p \leq .001$ ). Exploratory analysis indicates that anxiety levels were significantly associated with illness perceptions specific to COVID-19. Post-hoc analysis reveal no significant difference between the number of comorbidities and condition type (e.g. respiratory disorders).

**Conclusion:** This research supports individualised supports for people with ongoing medical conditions through the COVID-19 pandemic, and has implications for the consideration of follow-up care specifically for mental health. Findings may also inform future public health policies and post-vaccine support strategies for vulnerable populations.

**Keywords:** Ireland, Comorbidity, COVID-19, Pandemic, Mental Health

## Introduction

On March 11<sup>th</sup> 2020, a global pandemic was declared by the World Health Organisation following the spread of a novel strain of coronaviruses labelled as COVID-19 (Smith *et al.* 2020; Xiong *et al.* 2020). Preventative public health measures such as national and regional lockdowns were implemented by governments in order to reduce the fatality and spread of the disease. These preventative measures including containment and quarantine, are enforced for public safety and were proven successful in previous epidemics (Reynolds *et al.* 2008). Nonetheless, such measures may result in elevated psychological distress for example anxiety, stress, and depression, through mediating factors such as social disconnectedness and self-isolation (Özdin & Bayrak Özdin, 2020; Brook *et al.* 2020; Santini *et al.* 2020). Previous research indicates that psychological responses to epidemics may remain over time and thus pose an acute threat to mental health (Kelly, 2020). Individual variables such as perceived vulnerability, poor self-rated health, and anxiety proneness may result in psychological vulnerability from pandemic-associated stressors (Asmundson & Taylor, 2020).

Research pertaining to the psychological effects of COVID-19 appears heterogenous with countries revealing varying health outcomes and psychological responses. A population-based cross-sectional study revealed that symptoms of depression, anxiety, and stress were prevalent among a cohort of Spanish individuals in the initial phase of the pandemic, this being most pronounced for anxiety (González-Sanguino *et al.* 2020). These findings were mirrored in a study from China which evaluated the immediate and adverse psychological response of COVID-19 on mental health. The study revealed that the most notable expression was found for anxiety, compared to depression and stress, among the general population in China (Wang *et al.* 2020; Smith *et al.* 2020). Within Ireland, these findings were also replicated (Burke *et al.* 2020).

It is necessary for the implementation of newly developed health services and public health policies to address the negative burden that COVID-19 may place on individuals and vulnerable populations (Hao *et al.* 2020; Xiong *et al.* 2020). Advanced age and comorbid chronic illness are significant risk factors for developing negative health outcomes and contracting disease, with these cohorts being considered as ‘high risk’ for contracting COVID-19 (Emami *et al.* 2020; Yu *et al.* 2011; Özdin & Bayrak Özdin, 2020). Not only do patients with comorbidity yield poorer clinical outcomes and prognosis, but they are also more susceptible to greater psychological burden (Guan *et al.* 2020; Wang *et al.* 2020). This burden may be due to compromised immunity, and/or worries about physical health (Xiong *et al.* 2020; Hao *et al.* 2020).

In response to COVID-19, the current study investigated the relationship between comorbid health conditions and psychological well-being during COVID-19 among a cohort of Irish adults. It was hypothesised that the threat of COVID-19 to one’s health would lead individuals with ongoing medical conditions to have lower self-reported subjective well-being, experience increased psychological distress, have higher levels of personal distress, report lower levels of gratitude, and have elevated scores of illness perception when compared to those without medical diagnoses. Variables such as age, gender, annual income, and education have been found to affect symptoms of anxiety and depression and thus were used to select the matched groups (Albert, 2015; Smith *et al.* 2020; Özdin & Bayrak Özdin, 2020).

## Methods

### Inclusion and Exclusion

Participants were required to be over the age of 18, living in Ireland during the period of quarantine (27.03.2020 - 08.06.2020), and to confirm the presence or absence of a medical health diagnosis. Furthermore, participants were required to read an information sheet and provide consent prior to the questionnaire. Individuals who did not meet the inclusion criteria were excluded from the study.

### Participants

This study is a secondary analysis of data obtained in an online national survey, whereby a public sample was recruited through the use of media outlets in Ireland (See Burke *et al.* 2020). The original study provided a sample of n=847 participants; however, data was screened in order to capture the cohort of individuals needed for this study (N=256).

Participants who reported having a medical condition (n=128) formed one group, whilst the control group (n=128) was created by purposively matching participants on age, gender identity, annual income, educational attainment, and work status during COVID-19. In selecting the control participants, outcomes from each participant in the medical group was blinded from their demographics and then purposively matched with an individual of similar demographic information who reported not having a medical condition. When matching, all participant outcome data was blinded. In this sample the average age for the medical condition group was 39 years ( $\pm 11.41$ ), and 83.6% of participants were female. Similarly, the purposively matched control cohort had a mean age of 39 years ( $\pm 11.3$ ), and 83.6% of participants were female.

## Measures

The Warwick-Edinburgh Mental Wellbeing Scale (WEMWBS; Tennant *et al.* 2007) is a 14-item measure which covers both hedonic and eudaimonic facets of mental health. The positively worded items capture various concepts of well-being including positive affect, psychological functioning, and interpersonal relationships. It is proposed that higher scores are indicative of greater well-being.

The Depression, Anxiety and Stress Scale - 21 (DASS-21; Lovibond & Lovibond, 1995) is composed of three varying subscales, with each scale measuring self-reported depression, anxiety, and stress.

The Effects of COVID-19 Questionnaire (ECQ; Burke *et al.* 2020) is a 34-item tool which is designed to measure individuals' perception of COVID-19 related stresses and associated gratitude. This measure contains four subscales: Personal Stress (items 1-13), Parenting Stress (items 14-21), Older Aging Parent Stress (22-25), and Gratitude (26-34) in which respondents must choose from five response options (see supplementary material 1). Within the ECQ subscales, the ranges for Personal Distress are: Normal 0–12; Mild 13–19; Moderate 20–26; Severe 27–33; Extremely Severe >34.

The Brief Illness Perception Questionnaire (BIPQ; Broadbent *et al.* 2006) is a 7-item scale designed to rapidly assess both the emotional and cognitive representation of illness, and has been adapted for use with COVID-19 (see supplementary information for BIPQ questions).

## Data Analysis Plan

An independent samples t-test was used to analyse scores of the WEMWBS and the ECQ, whilst multivariate analyses of variance (MANOVA) were conducted to analyse scores of the DASS-21 and BIPQ. Multiple linear regressions were conducted as an exploratory analysis to determine whether levels of anxiety were predicted based on responses of the WEMWBS and

the BIPQ. Participants with medical conditions were further stratified based on the number of medical morbidities that were present and the associated clinical features (Guan *et al.* 2020), see supplementary Figure 1. A post-hoc MANOVA analysis was conducted to determine if there was a differential effect present based on condition type with Bonferroni correction. An alpha level of .05 was set for significance testing, with Bonferroni adjustment considered where relevant (adjusted p-value for significance threshold is  $p < .025$ ). Outliers were removed when preliminary analyses indicated violations of the assumptions of linearity, multicollinearity, and homogeneity of variance-covariance matrices. The assumption of homogeneity of variance was satisfied.

## Results

### Effects of COVID-19 Questionnaire (ECQ)

An independent samples t-test was conducted to compare levels of gratitude between groups. There was a significant difference in scores ( $t(243) = 3.46, p \leq .001$ ) with participants in the medical condition group ( $M = 17.21, SD = 6.69$ ) scoring lower than those in the control ( $M = 20, SD = 5.91$ ). The magnitude of the differences in the means (mean difference = 2.79, 95% CI: - 4.38 – -1.2) was small (Cohen's  $d = .43, 95\% CI: .18 - .68$ ).

An independent samples t-test was conducted to compare levels of personal stress between groups. There was no significant difference in scores ( $t(249.61) = 1.13, p = .260$ ).

### Depression, Anxiety, Stress Scale (DASS-21)

A one-way between groups MANOVA was conducted to determine if participants with a medical condition and in the control group differed in scores of the DASS-21. There was a statistically significant difference between groups on the combined dependent variable (Wilks' Lambda = .96,  $F(3, 226) = 3.35, p < .05$ , partial eta squared = .043). As can be seen in



Table 1, individuals with a medical condition reported significantly higher levels of anxiety, compared to the control group. According to DASS-21 classification, mean anxiety scores fall within the mild (8-9) to moderate (10-14) clinical ranges in the control and medical groups, respectively.

**Table 1***MANOVA Results for the DASS-21 and BIPQ*

Variable	Group	N	M	SD	F	p	$\eta^2$
<b>DASS-21</b>							
Depression	Medical	113	11.27	3.43	.51	.48	.002
	Control	116	10.95	3.46			
Anxiety	Medical	113	10.26	2.89	8.48*	.004	.04
	Control	116	9.22	2.52			
Stress	Medical	113	13.12	3.67	1.55	.21	.005
	Control	116	12.59	3.67			
<b>BIPQ</b>							
Item 1 (consequences)	Medical	128	6.88	1.97	.08	.79	.000
	Control	125	6.95	2.06			
Item 2 (timeline)	Medical	128	6.65	1.53	1.38	.24	.005
	Control	125	6.42	1.51			
Item 3 (personal control)	Medical	128	5.01	2.09	1.61	.21	.006
	Control	125	5.36	2.32			
Item 4 (treatment)	Medical	128	4.13	2.08	2.15	.14	.008
	Control	125	4.52	2.21			
Item 5 (concern)	Medical	128	7.77	1.71	6.76**	.01	.026
	Control	125	7.15	2.08			
Item 6 (identity)	Medical	128	8.90	1.68	.43	.51	.002
	Control	125	8.76	1.66			
Item 7 (emotional representation)	Medical	128	7.55	2.11	4.67**	.03	.018
	Control	125	6.96	2.27			

*Note.* \*  $p < .01$ , \*\*  $p < .05$ ,  $\eta^2$  = partial eta squared. Item brackets (e.g., consequences) refers to each dimension of illness perception the BIPQ assesses.

### **Brief Illness Perception Questionnaire (BIPQ)**

A one-way between groups MANOVA was conducted to determine if participants with a medical condition and in the control group differed on measures of self-reported illness perception. There was a statistically significant difference between groups on the combined dependent variable (Wilks' Lambda = .94,  $F(7, 246) = 2.06$ ,  $p < .05$ , partial eta squared = .056). As can be seen in Table 1, individuals with a medical condition scored significantly higher on items 5 (concern) and 7 (emotional representation) of the BIPQ (see supplementary material 2).

### **Multiple Linear Regression**

As significant differences were found between groups on BIPQ subscales, a standard multiple regression analysis was performed as an exploratory analysis to assess whether being concerned about the pandemic (concern), peoples' perception of how COVID-19 has affected them emotionally (emotional representation), and measures of self-reported well-being, were significantly associated with scores of the DASS-21 Anxiety subscale. The ECQ subscales were considered as outcome variables, and therefore not included. In the medical condition group, participants' concern of how COVID-19 was affecting them emotionally and scores of self-reported well-being revealed significance, as shown in Table 2. However, only well-being scores revealed a significant association with the criterion variable in the control group (see Table 2). The model as a whole explained 15.4% of the variance in anxiety scores in the medical condition group ( $F(3,108) = 6.58$ ,  $p < .001$ ) and 10.7% of the variance in the control group ( $F(3,104) = 4.14$ ,  $p < .01$ ).

**Table 2***Multiple regression model predicting DASS-21 Anxiety scores*

	Group	R <sup>2</sup>	Adj R <sup>2</sup>	β	B	SE	p	CI 95% (B)
<b>Model</b>	<b>Medical</b>	.154***	.131***					
	<b>Control</b>	.107**	.081**					
BIPQ Q5	Medical			-.02	-.03	.18	.86	-.39 / .33
	Control			.06	.08	.15	.58	-.22 / .39
BIPQ Q7	Medical			.29**	.42	.16	.008	.11 / .73
	Control			.16	.23	.16	.14	-.08 / .54
WEMWBS	Medical			-.24**	-.09	.03	.009	-.16 / -.02
	Control			-.22*	-.07	.03	.02	-.14 / -.01

*Note.* R<sup>2</sup> = R-squared; Adj. R<sup>2</sup> = Adjusted R-squared; β = standardized beta value; B =

unstandardized beta value; SE = Standard errors of B; CI 95% (B) = 95% confidence interval for B; N = 398; Statistical significance: \*p < .05; \*\*p < .01; \*\*\*p < .001

### **The Warwick-Edinburgh Mental Well-being Scale (WEMWBS)**

An independent samples t-test was conducted to compare total scores of the WEMWBS between groups. There was no significant difference in scores ( $t(254) = 1.14, p = .255$ ).

### **Post hoc Analyses**

Oneway ANOVA analyses were conducted for the WEMWBS and the ECQ subscales to compare whether participants' scores varied based on condition type. A multiple regression was also conducted to examine whether condition type was a significant predictor of DASS-21 anxiety scores. Within the medical group, 16.8% have more than one medical condition. There were no statistically significant findings nor differential effects found based on

condition type. Supplementary Figure 1 shows the distribution of medical conditions within the group. Of note, respiratory disorders were the highest self-reported medical morbidity.

### **Discussion**

Adverse psychological effects are commonly expressed at the beginning of a lockdown and in response to a pandemic (Xiong *et al.* 2020; Xiao *et al.* 2020). This can be due to a number of factors, such as the lockdown itself, or the risk of contagion. Significant group differences in anxiety scores were of small effect size ( $\eta p^2 = .04$ ) with mean scores in the medical group ( $10.26 \pm 2.89$ ) falling at the lower end of the moderate range (10-14) for clinical severity. The medical condition cohort reported lower feelings of gratitude and also had higher levels of concern about COVID-19 and how the pandemic was effecting them emotionally. Of note, when compared to healthy controls, participants with a medical condition did not significantly differ on measures of subjective well-being and personal distress. Participants perception of how the pandemic effects them emotionally was significantly associated with anxiety, over and above stress and depression. This pattern of findings is consistent with previous research indicating that worry of a novel virus is related to psychological distress, however it is important to acknowledge that participants with a medical condition appear relatively psychologically healthy, with the exception of anxiety and gratitude subscales (Xiong *et al.* 2020). Grateful individuals often report greater physical health, however further research is recommended to better interpret the direction of results found in this study (Hill *et al.* 2013). Anxiety and depression are common in a wide range of medical conditions, however post-hoc analyses revealed that there was no significant differential effect between participant scores based on condition type, despite the majority of participants with a medical condition having respiratory-related illnesses, as shown in supplementary figure 1( Lenzo *et al.* 2020; Swartz & Jantz, 2014).

This study contributes to, and supports, existing research showing that those with an ongoing medical condition are more prone to worry and concern (Wheaton *et al.* 2012; Özdin & Bayrak Özdin, 2020; Hao *et al.* 2020). However, the findings of this study have to be seen in light of some limitations. The unequal gender amount in both groups makes it difficult to generalize results, furthermore the cross-sectional design is limited to a single timepoint and thus research is needed to evaluate whether these effects are sustained over time, and/or fluctuate with the pandemic infection and mortality rates. Post hoc analyses may also have been underpowered and therefore unable to capture group effects based on the low sample size available when participants were sub-stratified by condition type.

It is evident that with a pandemic brings uncertainty and fear in peoples' lives (Taylor & Asmundson, 2004; Taylor, 2019). In spite of inflated changes in anxiety levels, those with existing medical conditions appear psychologically healthy when compared to those without medical diagnosis at this time. It is important for research to evaluate perpetuating, protective, and predictive factors in order to consider specific interventions for vulnerable populations and those who require them most.

**Funding:** This research was in part funded and supported by the Health Research Board and Irish Research Council [COV19-2020-044].

**Conflicts of Interest:** The authors declare no conflict of interest.

**Ethical approval:** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committee on human experimentation with the Helsinki Declaration of 1975, as revised in 2008.

## References

**Albert PR** (2015). Why is depression more prevalent in women?. *Journal of psychiatry & neuroscience: JPN* **40**, 219–221.

**Asmundson GJG, Taylor S** (2020). Coronaphobia: Fear and the 2019-nCoV outbreak. *Journal of Anxiety Disorders* **70**, 102196.

**Broadbent E, Petrie KJ, Main J, Weinman J** (2006). The Brief Illness Perception Questionnaire. *Journal of Psychosomatic Research* **60**, 631–637.

**Burke T, Berry A, Taylor LK, Stafford O, Murphy E, Shevlin M, McHugh L, Carr A** (2020). Increased Psychological Distress during COVID-19 and Quarantine in Ireland: A National Survey. *Journal of Clinical Medicine* **9**, 3481.

**Emami A, Javanmardi F, Pirbonyeh N, Akbari A** (2020). Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: A Systematic Review and Meta-Analysis. *Archives of Academic Emergency Medicine* **8**, e35.

**González-Sanguino C, Ausín B, Castellanos MÁ, Saiz J, López-Gómez A, Ugidos C, Muñoz M** (2020). Mental health consequences during the initial stage of the 2020 Coronavirus pandemic (COVID-19) in Spain. *Brain, Behavior, and Immunity* **87**, 172–176.

**Guan W, Liang W, Zhao Y, Liang H, Chen Z, Li Y, Liu X, Chen R, Tang C, Wang T, Ou C, Li L, Chen P, Sang L, Wang W, Li J, Li C, Ou L, Cheng B, He J** (2020). Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *European Respiratory Journal* **55**, 2000547.

**Hao F, Tan W, Jiang L, Zhang L, Zhao X, Zou Y, Hu Y, Luo X, Jiang X, McIntyre RS, Tran B, Sun J, Zhang Z, Ho R, Ho C, Tam W** (2020). Do psychiatric patients experience more psychiatric symptoms during COVID-19 pandemic and lockdown? A case-control study with service and research implications for immunopsychiatry. *Brain, Behavior, and Immunity* **87**, 100–106.

**Hill PL, Allemand M, Roberts BW** (2013). Examining the Pathways between Gratitude and Self-Rated Physical Health across Adulthood . *Personality and Individual Differences* **54**, 92-96.

**Kelly BD** (2020). Plagues, pandemics and epidemics in Irish history prior to COVID-19 (coronavirus): what can we learn?. *Irish Journal of Psychological Medicine* **37**, 269–274.

**Lenzo V, Sardella A, Martino G, Quattropiani MC** (2020). A Systematic Review of Metacognitive Beliefs in Chronic Medical Conditions. *Frontiers in Psychology* **10**, 2875.

**Lovibond SH, Lovibond PF** (1995). *Manual for the Depression Anxiety Stress Scales 2nd ed.* Sydney: Psychology Foundation.

**Özdin S, Bayrak Özdin Ş** (2020). Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *International Journal of Social Psychiatry* **66**, 504–511.

**Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsi E, Katsaounou P** (2020). Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain, Behavior, and Immunity* **88**, 901–907.



**Reynolds DL, Garay JR, Deamond SL, Moran MK, Gold W, Styra R (2008).**

Understanding, compliance and psychological impact of the SARS quarantine experience.

*Epidemiology and Infection* **136**, 997–1007.

**Santini ZI, Jose PE, York Cornwell E, Koyanagi A, Nielsen L, Hinrichsen C, Meilstrup**

**C, Madsen KR, Koushede V (2020).** Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): a longitudinal

mediation analysis. *The Lancet Public Health* **5**, 62–70.

**Smith L, Jacob L, Yakkundi A, McDermott D, Armstrong NC, Barnett Y, López-**

**Sánchez GF, Martin S, Butler L, Tully MA (2020).** Correlates of symptoms of anxiety and depression and mental wellbeing associated with COVID-19: a cross-sectional study of UK-

based respondents. *Psychiatry Research* **291**, 113138.

**Swartz JA, Jantz I (2014).** Association Between Nonspecific Severe Psychological Distress

as an Indicator of Serious Mental Illness and Increasing Levels of Medical Multimorbidity.

*American Journal of Public Health* **104**, 2350–2358.

**Taylor S (2019).** *The psychology of pandemics: Preparing for the next global outbreak of infectious disease*. Cambridge Scholars Publishing.

**Taylor S, Asmundson GJG (2004).** *Treating Health Anxiety: A Cognitive-behavioral*

*Approach*. Guilford Press.

**Tennant R, Hiller L, Fishwick R, Platt S, Joseph S, Weich S, Parkinson J, Secker J,**

**Stewart-Brown S (2007).** The Warwick-Edinburgh Mental Well-being Scale (WEMWBS):

development and UK validation. *Health and Quality of Life Outcomes* **5**, 63.

**Wang B, Li R, Lu Z, Huang Y** (2020). Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis. *Aging* **12**, 6049–6057.

**Wang C, Pan R, Wan X, Tan Y, Xu L, McIntyre RS, Choo FN, Tran B, Ho R, Sharma VK, Ho C** (2020). A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain, behavior, and immunity* **87**, 40–48.

**Wheaton MG, Abramowitz JS, Berman NC, Fabricant LE, Olatunji BO** (2012). Psychological Predictors of Anxiety in Response to the H1N1 (Swine Flu) Pandemic. *Cognitive Therapy and Research* **36**, 210–218.


**Xiao H, Zhang Y, Kong D, Li S, Yang N** (2020). The Effects of Social Support on Sleep Quality of Medical Staff Treating Patients with Coronavirus Disease 2019 (COVID-19) in January and February 2020 in China. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research* **26**, e923549.

**Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, Chen-Li D, Iacobucci M, Ho R, Majeed A, McIntyre RS** (2020). Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *Journal of Affective Disorders* **277**, 55–64.

**Yu H, Feng Z, Uyeki TM, Liao Q, Zhou L, Feng L, Ye M, Xiang N, Huai Y, Yuan Y, Jiang H, Zheng Y, Gargiullo P, Peng Z, Feng Y, Zheng J, Xu C, Zhang Y, Shu Y, Wang Y** (2011). Risk Factors for Severe Illness with 2009 Pandemic Influenza A (H1N1) Virus Infection in China. *Clinical Infectious Diseases* **52**, 457–465.

**Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, Guan L, Wei Y, Li H, Wu X, Xu J, Tu S, Zhang Y, Chen H, Cao B** (2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* **395**, 1054–1062.

Supplementary Figure 1:



Supplementary Figure 1: A breakdown of medical disorders within the cohort who self-reported a medical condition. Note: Categories are based on Guan et al, 2020.

## Supplementary material 1:

### Effects of COVID-19 questionnaire (ECQ)

The following questions ask about the effect that the COVID-19 or Corona virus is having on you.

For each item, click on the answer that applies to you in the **PAST MONTH**.

N/A means the item is not applicable to you.

**In the past month, how much stress have you experienced as a result of the following things**

1	Financial hardship for you or your family arising from the COVID-19 crisis, due to job loss, or loss of earnings	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
2	Having difficulty getting supplies when you need them, including face masks, hand sanitizers, medicines, food, drinks or other essentials	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
3	Not being able to meet with your extended family and friends	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
4	Not being able to go to your church or place of religious worship	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
5	Loss of your own, or your family's daily routine (such as sleeping patterns; meal times; work, school and recreation schedules)	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
6	Family conflict arising from the COVID19 crisis, due to arguing, or fighting with other family members more than usual because you are spending more time together at home	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
7	Getting a lot of conflicting information and misinformation online and in the media about COVID-19	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
8	You, or members of your family becoming ill with COVID-19	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
9	Worrying that you may become infected with COVID-19 and then infect other people	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
10	You, or members of your family being hospitalised for COVID-19 illness	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
11	Death of a family member or very close friend as a result of COVID-19	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
12	Witnessing others in your community suffering because of COVID-19	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
13	Worrying about the effects COVID-19 on you or your family, now or in the future	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
<b>If you have children</b>							
14	Your child's school closing	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
15	Helping your child keep a safe distance from their friends, or preventing them from mixing with their friends	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
16	Helping your child keep a safe distance from members of your extended family, or preventing them from visiting with the extended family (for example grandparents)	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
17	Helping your child avoid crowded places, and activities that they like, such as going	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5

	to sports or musical events, scouts or guides, clubs, the playground, or to church						
18	Helping your child to not shake hands, hug, or touch other people	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
19	Helping your child to wash or sanitise their hands regularly	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
20	Helping your child to remember to cough or sneeze into their elbow	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
21	Being worried that your child will catch COVID-19 because they have an underlying medical condition such as cancer or asthma, that makes them vulnerable to severe illness if they become infected	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
<b>If you have aging parents</b>							
22	Worrying that you aging parents will become lonely during the COVID-19 crisis						
23	Worrying that you aging parents will not get supplies during the COVID-19 crisis	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
24	Worrying that your aging parents will become infected with COVID-19	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
25	Worrying that your aging parents will not receive adequate medical care if they become infected with COVID-19	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
<b>In the past month, how much has your experience of the COVID 19 crisis led you to feel grateful for the following things</b>							
26	Your health, and the health of your family	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
27	Your relationships with your extended family and friends	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
28	Your job	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
29	Attending social, sports, and cultural events	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
30	Your community	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
31	Your child's regular attendance at school	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
32	Your child's relationships with their friends	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
33	Your child's involvement in activities such as sports, music, scouts, guides, clubs etc	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5
34	Your aging parents health and safety	N/A 0	None 1	A little 2	Some 3	Quite a lot 4	A great deal 5

**Note:** Items 1-13 COVID-19 personal stress scale. Items 14-21 COVID-19 parenting stress scale. Items 22-25 COVID-19 aging parents stress scale. Items 26-33 COVID-19 gratitude scale. To get a scale score, sum scores of items in the scale and divide this by the number of items which did not have an N/A response.

Supplementary material 2:

**The Brief Illness Questionnaire (BIPQ; Broadbent *et al.* 2006)**

1. How much has the COVID-19 pandemic effected your life? (BIPQ1)  
0 1 2 3 4 5 6 7 8 9 10  
No effect at all severely effects my life
2. How long do you think the COVID-19 pandemic will continue? (BIPQ2)  
0 1 2 3 4 5 6 7 8 9 10  
A very short time Forever
3. How much control do you feel you have over the COVID-19 situation (e.g. not getting infected or getting over it)? (BIPQ3)  
0 1 2 3 4 5 6 7 8 9 10  
Absolutely no control Extreme amount of control
4. How much do you think existing treatments can help COVID-19 patients? (BIPQ4)  
0 1 2 3 4 5 6 7 8 9 10  
Not at all Extremely helpful
5. How concerned are you about the COVID-19 pandemic? (BIPQ5)  
0 1 2 3 4 5 6 7 8 9 10  
Not at all concerned Extremely concerned
6. How well do you feel you understand the COVID-19 situation? (BIPQ6)  
0 1 2 3 4 5 6 7 8 9 10  
Don't understand at all Understand very clearly
7. How much does the COVID-19 pandemic effect you emotionally (e.g. does it make you angry, scared, upset or depressed)? (BIPQ7)  
0 1 2 3 4 5 6 7 8 9 10  
Not at all effected emotionally Extremely effected emotionally

### SCORING INFORMATION

Each item of the Brief Illness Questionnaire assesses one dimension of illness perceptions:  
The **consequences** score is simply response to item 1

The **timeline** score is the response to item 2

The **personal control** score is the response to item 3

The **treatment control** score is the response to item 4

The illness **concern** score is the response to item 5

The **identity** score is the response to item 6

The **emotional representation** is measured by item 7. This reflects a combination of emotional and cognitive representations

In some circumstances it may be possible to compute an overall score which represents the degree to which the illness is perceived as threatening or benign. The internal consistency of this score will depend on the illness studied and it is recommended that this is checked. To compute the score, reverse score items 3,4, and 7 and add these to items 1,2,5,6, and 8. A higher score reflects more threatening view of the illness. (BIPQTot)