

Comparative Seroprevalence and Risk Factors of Toxoplasmosis among Three Subgroups in Nigeria

Emmanuel C. Uttah* Raymond Ajang Jude Ogbeche¹ Hannah Etta Lawrence Etim
Department of Biological Sciences, Cross River University of Technology, Calabar, Nigeria

* E-mail of the corresponding author: drecuttah@yahoo.com

Abstract

The study was aimed at ascertaining the seroprevalence and some risk factors of toxoplasmosis among those attending the National Hospital, Abuja, Nigeria. Using the ImmunoComb Toxo IgG Kit for the detection of IgG antibodies to *Toxoplasma gondii* in human serum/ plasma, 216 persons were examined for toxoplasmosis. This comprised of equal number (72 persons each) of persons from three subgroups, namely, gravid women, immuno-compromised persons consisting of patients with HIV and hepatitis, and the immuno-competent persons who are the controls. The overall seroprevalence of toxoplasmosis was 31.5%. Seroprevalence was significantly higher among pregnant women than among both the immuno-compromised and the immuno-competent persons (χ^2 -test; $p < 0.05$ for both tests). Seroprevalence was comparable between the immuno-compromised and the immuno-competent (Controls) (χ^2 -test; $p > 0.05$). Seroprevalence increased with age among gravid women, as well as among the immuno-compromised subgroups but not among the controls. Overall, seroprevalence among those living with cats/dogs (45.61%) was significantly higher than that of those living without cats/dogs (15.69%) (χ^2 -test; $p < 0.05$); while the risk of infection with toxoplasmosis was four times higher than among those not living with cats (Odds Ratio 4.51). This was the same among the immune-compromised as well as the immune-competent where the risks of getting infected were five times higher among those cohabiting with cats (Odds Ratio 5.45 and Odds Ratio 5.36 respectively). Among the pregnant women, the risk of getting infected with toxoplasmosis was three times higher among those cohabiting with cats (Odds Ratio 3.00). Seroprevalence was independent of the tribal origin (χ^2 -test; $p < 0.05$ for all the tests).

Keywords: Toxoplasmosis, *Toxoplasma gondii*, serprevalence, pregnant women, immuno-compromised, immuno-competent, Nigeria

1. Introduction

Toxoplasmosis is an important parasitic infection with global distribution and significance. About a third of the world's human population is estimated to harbor the causative coccidian protozoan, *Toxoplasma* (Ryan and Ray, 2004). The distribution of this parasite depends on regions and weather condition where the oocysts survive in environment (Dubey and Jones 2008; Dubey 2004; Fayer 1981). It is estimated that between 30% and 65% of all people worldwide are infected with toxoplasmosis (Tenter, 2000). Prevalence of infection varies between countries. In France, for example, about 88% of the population is carriers, probably due to a high consumption of raw and lightly cooked meat (Adam, 2003). High prevalence rates of between 67% and 80% have been reported in Germany, the Netherlands and Brazil (Heckerroth, 2004), while in Britain and South Korea, and about 22% and 4.3% respectively are carriers (Weiss, 2004). Infection is reportedly highest in countries where undercooked meat is traditionally.

Toxoplasmosis could be severe and life-threatening during pregnancy, and to fetuses, and new born babies (Robert-Gangneux et al., 2009). Vertical transmission occurs causing mental retardation, blindness, epilepsy, and death (Petersen, 2007). One of the late sequelae of congenital toxoplasmosis is chorioretinitis (Al-Azawi et al., 2013). Among the immuno-competent people, toxoplasmosis is usually asymptomatic, subclinical or benign, and can be classified as congenital, acquired or ocular (Oyibo et al., 2009). It may precursor spontaneously resolved symptoms such as fever, malaise, and lymphadenopathy, indicating symptomless latent infection (Montoya and Liesenfield, 2004). Toxoplasmosis can be severe and life-threatening to immune-compromised patients (Robert-Gangneux et al., 2009), causing severe encephalitis through acute infection or reactivation of latent infection (Innes, 2010, Hang et al., 2007).

Cats are the definitive hosts since they are the only animals that excrete resistant oocysts into the environment (Silva et al; 2001). Cohabiting with cats increases the chances of getting infected (Sukthana, 2006). However, direct infection through handling cats is generally believed to be rare (Agrappi, 2006). Some animals including humans serve as intermediate hosts in which the parasite may cause systemic infection that result in the formation of tissue cysts. Transmission may occur through ingestion of raw or partly cooked meat, especially pork, lamb, or venison containing *Toxoplasma* cysts. Oocysts may also be ingested through knives, utensils, or cutting boards contaminated by raw meat (Joss, 2004) or through ingestion of oocysts shed by cats in the environment, transplacentally, and through organ trans-plantation ((Nissapatorn et al., 2011).

Toxoplasma gondii is a major cause of economic losses in endemic communities as they are responsible for

abortions, still birth and neonatal losses among various classes of livestock (Raeghi et al., 2011, Buxton et al., 2007, Masala et al., 2003). It is associated with congenital defects in humans, and the risk of the infection being passed on to the fetus increases to between 60% and 90% in the third trimester (Tenter, 2007). The severity of congenital infections depends on the stage of pregnancy when the acute infection occurred, and spontaneous abortions or neurological disorders (Black and Boothroyd, 2000).

Toxoplasmosis is a neglected parasitic infection although it is extremely important economically, medically and epidemiologically (Uttah et al., 2013). Compared with other parasitic infections such as malaria and filariasis, it is grossly underreported. The paucity of research data on various aspects of toxoplasmosis in Nigeria is palpable. This study is an attempt to bridge this gap, and is aimed at ascertaining the seroprevalence and some risk factors of toxoplasmosis among those attending the National Hospital, Abuja, Federal Capital Territory, Nigeria.

2. Materials and Methods

2.1 Description of study area

The National Hospital, Abuja is located in central District phase 11 Garki, City of Abuja. It is located within latitude 7° 25' N and 9° 20' North of the Equator and longitude 5° 45' and 7° 39'. The hospital is owned by the Government, and therefore being visited by people of diverse backgrounds from different sections of the country, including the both the poor and rich. Abuja is replete with hills, highlands and undulating plains.

2.2 The study design

Equal number of three subgroups was enlisted into the study, namely the pregnant women, the Immuno-compromised patients, and the Immuno-competent persons (Controls). The immuno-compromised persons enlisted consisted of patients with such medical conditions such as HIV and hepatitis. The Immuno-competent persons enlisted were healthy persons who volunteered to participate in the study.

2.3 Administration of structured questionnaire

A well-structured questionnaire capturing information regarding age, sex, tribe, possession of cats or other domestic animals in the house were administered on the individuals enlisted into the study.

2.4 Collection of samples and serological testing

From each of the 216 patients, 2ml of blood sample was collected between February 2011 and July 2011, and processed using The ImmunoComb Toxo IgG Kit following standard methods. Blood samples were centrifuged for 5 minutes; serum was collected and stored at 20°C. The developing plates, cards, reagents and specimens were all brought to room temperature of between (22°C and 26°C).

2.5 Ethical considerations

The work was approved by the Ethics Committee of Cross River University of Technology, Calabar, and the authorities of the National Hospital, Abuja. Safety protocols were strictly followed all through the study. The work table was covered with absorbent tissues that were discarded as bio-hazardous waste at the end of tests. Reagents were mixed by shaking the Developing Plate.

3. Results

3.1 Seroprevalence of toxoplasmosis among the three study groups

The overall seroprevalence of toxoplasmosis was 31.48% (see Table 1). Seroprevalence was significantly higher among pregnant women than among both the immuno-compromised and the immuno-competent persons (χ^2 -test; $p < 0.05$). Seroprevalence was comparable in both the immuno-compromised and the immuno-competent (Controls) (χ^2 -test; $p > 0.05$).

3.2 Seroprevalence in relation to age and sex

Seroprevalence increased with age among pregnant women (See Table 2), and the immunocompromised groups (see Table 3). Among the immunocompromised group, seroprevalence was independent of sex (χ^2 -test: 0.3136; $p < 0.05$).

Seroprevalence was comparable among males and females and there was no definite age-related pattern among the immunocompetent (control) group (see Table 4).

3.3 Seroprevalence among gravid females

The overall prevalence among gravid females was 44.4% (see Table 2). Prevalence increased significantly with age to 66.7% in the last age group (χ^2 -test; $p < 0.05$).

3.4 Seroprevalence among the Immuno-compromised patients

The overall prevalence among the immune-compromised patients was 27.8% (see Table 3). Prevalence was significantly higher among females than males (χ^2 -test; $p < 0.05$); and significantly increased with age in both sexes (χ^2 -test; $p < 0.05$ for all tests).

3.5 Seroprevalence among the immunocompetent persons (the controls)

The overall prevalence among the immunocompetent persons was 22.2% (see Table 4). Prevalence decreased significantly with age, especially among females (χ^2 -test; $p < 0.05$ for all tests).

3.6 Comparison of seroprevalence among the three subgroups

In all the three subgroups, there was no positive case observed among those 20 years old or younger. Prevalence among females were significantly higher than that among males in both the immuno-compromised and the immune-competent subgroups (χ^2 -test; $p < 0.05$ in both tests).

3.7 Seroprevalence of toxoplasmosis among those living with cats

Prevalence among those who have cats living with them in their residence (see Table 5) was significantly higher than among those not living with cats (χ^2 -test; $p < 0.05$ for all tests). The risk of getting infected among the various subgroups is presented in Table 6. Among those living with cats overall, the risk of infection with toxoplasmosis was four times higher than among those not living with cats (OR 4.51). This was the same among the immune-compromised as well as the immune-competent where the risks of getting infected were five times higher among those cohabiting with cats (OR 5.45 and OR 5.36 respectively). Among the pregnant women, the risk of getting infected with toxoplasmosis was three times higher among those cohabiting with cats (OR 3.00).

3.8 Seroprevalence of toxoplasmosis in relation to the tribe of origin of participants in the study

The seroprevalence of toxoplasmosis in relation to the tribe of origin of participants in the study is presented in Table 7. Prevalence was comparable among the tribes (χ^2 -test; $p < 0.05$ for all the tests).

4. Discussion

The results from this study indicate that toxoplasmosis is highly prevalent among those attending the National Hospital Abuja especially among pregnant women. This agrees with the findings of Torda (2001). An even higher seroprevalence rates for pregnant women in a Nigerian population ranged from 72.5% to 88.8% with an overall rate of 75.4% (Onadeko et al., 1996). In Columbia, increase in seroprevalence to *Toxoplasma* antibodies for the past 25 years among pregnant women has been reported (Rosso et al., 2008).

Toxoplasmosis seroprevalence increased with age and was not sex-related. Similar findings were made in Malaysia (Nissapatorn et al., 2011), and in a northern Mexican city of Durango among elderly people (Alvarado-Esquivé et al., 2012). However, living with cats was an important risk factor in the epidemiology of toxoplasmosis. Perhaps it was a major determinant of toxoplasmosis infection in the study area, as those living with cats constituted a highly significant proportion of positive case in all the three study groups. This is in agreement with findings elsewhere (Joss, 2004; Sukthana, 2006).

Toxoplasmosis is the most frequent protozoal opportunistic infection in immunocompromised individuals. Its association with immunosuppression has been known for several decades. ((Nissapatorn et al., 2011).).

T. gondii antibody prevalence rates vary greatly by geographic distribution and population among healthy persons (Pordeus et al, 2008), pregnant women (Nissapatorn et al, 2003), immunocompromised patients, including those with HIV/AIDS (Lindström et al, 2006), cancer (Rai et al, 2003). The seroprevalence of toxoplasmosis obtained in this study among the immunocompromised was lower than (54%) reported for the same group in a study in Uganda (Lindstrom et al., 2006). Similarly seroprevalence obtained in this study among pregnant women was lower than that (70%) reported in Cameroon and 53% in Brazil for the same subgroup (Vaz et al., 2010, Njunda et al., 2011).

In a comparative study, the seroprevalence of toxoplasmosis in pregnant women from the inner area of Ibadan (78%) was significantly higher than that among pregnant women from the Swansea area of the UK (22%); and it was concluded that reinfection or recrudescence was responsible for maintaining high antibody levels in African women, which is also worrisome considering its implications for the immune-compromised subset (Onadeko et al., 1992).

There was no significant difference in prevalence between the various ethnic/ tribal groups in this study. This is not in agreement with findings in Asia where *Toxoplasma* infection was reported to be more common among Malays than other ethnic groups, especially in those with a lower education level (Nissapatorn et al., 2011).

Close contact with cats as domestic pets is common among these people, and this is a well-known risk factor of toxoplasmosis transmission (Alvarado-Esquivé et al., 2012). In another study in neighbouring Cameroon, age, cat ownership, consumption of raw vegetables, source of potable water, meat consumption, and gestational age were identified as risk factors for toxoplasmosis in pregnancy (Njunda et al., 2011). In endemic areas wells located in farms could be contaminated with *Toxoplasma* (Sroka et al., 2006). Health education regarding toxoplasmosis and its consequences need to be given to prevent primary infection among the general population irrespective of their race or socioeconomic status. ((Nissapatorn et al., 2011).).

5. Conclusion

Toxoplasmosis is highly prevalent among those attending the National Hospital, Abuja. These findings buttress

the need for a nation-wide Toxoplasma screening program for a reliable data for proper planning by Health authorities, to be preceded by a comprehensive health education. There should be a definite policy of ensuring that toxoplasmosis screening and monitoring is included in the tests protocols in the first trimester of pregnancy in the Nigeria, considering the high prevalence recorded in this study. This is highly essential as early diagnosis during pregnancy allows for prompt intervention through treatment in order to reduce the probability of foetal infection and consequent substantial damage to the foetus (Njunda et al., 2011). Furthermore, campaign for general improvement in personal hygiene standards among inhabitants should be intensified and sustained. Evolving a strategy for sustainable inspection of meats meant for human consumption for Toxoplasma infection is a challenge that must be surmounted to check the rising profiles of toxoplasmosis in Nigeria.

References

- Al-Azawi, A.K.A., Al-Rawe, I.H.A., and Al-Bayati, R.Y.J. (2013). Seroprevalence of toxoplasmic chorioretinitis in Baghdad Province. *International Journal of Science and Nature* 4(1), 68-71.
- Alvarado-Esquivé, C., Liesenfeld, O., Burciaga-López, B.D., Ramos-Nevárez, A., Estrada-Martínez, S., Cerrillo-Soto, S.M., Carrete-Ramírez, F.A., López-Centeno, M., and Ruiz-Martínez, M.M. (2012). Vector-Borne and Zoonotic Diseases 12(7), 568-574. doi:10.1089/vbz.2011.0875.
- Black, M.W. and Boothroyd, J.C. (2000). Lytic Cycle of *Toxoplasma gondii*. *Microbiol. Mol. Biol. Rev.* 64, 607-623.
- Buxton, D., Maley, S.W., Wright, S.E., Rodger, S., Bartley, P., Innes, E.A. (2007). *Toxoplasma gondii* and ovine toxoplasmosis: new aspects of an old story. *Vet Parasitol.* 149, 25-28.
- Dubey, J.P. (2004). Toxoplasmosis-a waterborne zoonosis. *Vet Parasitol.* 126, 57-72.
- Dubey JP, Jones JL. (2008). *Toxoplasma gondii* infection in humans and animals in the United States. *Int J Parasitol.* 38, 1257-1278.
- Fayer, R. (1981). Toxoplasmosis update and public health implications. *Canadian Veterinary Journal*, 22, 344-352.
- Innes, E.A. (2010). A brief history and overview of *Toxoplasma gondii*. *Zoon Pub Heal.* 57, 1-7.
- Les, J.T., Agudelo, A., Villalobos, C., Chaves, J.A., Tunubala, G.A., Messa, A., Remington, J.S. and Montoya, J.G. (2008). Prevalence of Infection with *Toxoplasma gondii* among Pregnant Women in Cali, Colombia, South America. *Am. J. Trop. Med. Hyg.* 78, 504-508.
- Lindstroma, I., Deogratias, I., Kaddu-Mulindwa, H., Kironde, F., Lindh J. (2006). Prevalence of latent and reactivated *Toxoplasma gondii* parasites in HIV-patients from Uganda. *Acta Tropica* 100, 218-222.
- Masala, G., Porcu, R., Madau, L., Tanda, A., Ibba, B., Satta, G., Tola, S. (2003). Survey of ovine and caprine toxoplasmosis by IFAT and PCR assays in Sardinia, Italy. *Vet Parasitol.* 117, 15-21.
- Montoya, J.G., and Liesenfeld, O. (2004). Toxoplasmosis. *Lancet*, 363, 1965-1976.
- Nissapatorn, V., Leong, T.H., Lee, R., Inthoi, Ibrahim, J., and Yen, T.S. (2011). Seroepidemiology of toxoplasmosis in renal patients. *Southeast Asian J Trop Med Public Health* 42 (2), 237-247.
- Nissapatorn, V., Noor, Azmi, M.A., Cho, S.M., et al. (2003). Toxoplasmosis: prevalence and risk factors. *J Obstet Gynaecol.* 23, 618-624.
- Njunda, A.L., Assob, J.C.N., Nsagha, D.S., Kamga, H.L., Nde, P.F., Yugah, V.C. (2011). Seroprevalence of *Toxoplasma gondii* infection among pregnant women in Cameroon. *Journal of Public Health in Africa*, 2 (e24), 98-101.
- Onadeko, M.O., Joynson, D.H., Payne, R.A. (1992). The prevalence of *Toxoplasma* infection among pregnant women in Ibadan, Nigeria. *J Trop Med Hyg.* 95(2), 143-145.
- Onadeko, M.O., Joynson, D.H., Payne, R.A., Francis, J. (1996). The prevalence of toxoplasma antibodies in pregnant Nigerian women and the occurrence of stillbirth and congenital malformation. *Afr J Med Med Sci.* 25(4), 331-334.
- Oyibo, W.A, Oladosu, O.O., Agomo, C.O., Ojuromi, O.T., Anunobi, C.C., and Soyebi, K. (2009). Congenital Toxoplasmosis: A Review of its Pathology, Immune Response and Current Treatment Options. *Sierra Leone J Biomed Res* 1 (1), 9-20,
- Petersen, E. (2007). Toxoplasmosis. *Semin Fetal Neonatal Med*, 12, 214-223.
- Pordeus, V., Barzilai, O., Sherer, Y., et al. 2008 A latitudinal gradient study of common anti-infectious agent antibody prevalence in Italy and Colombia. *Isr Med Assoc J* 10, 65-68.
- Raeghi, S., Akaberi, A., and Sedeghi, S. (2011). Seroprevalence of *Toxoplasma gondii* in Sheep, Cattle and Horses in Urmia North-West of Iran. *Iranian J Parasitol* 6 (4), 90-94.
- Rai, S.K., Upadhyay, M.P., Shrestha, H.G. (2003). Toxoplasma infection in selected patients in Kathmandu, Nepal. *Nepal Med Coll J* 5, 89-91.
- Robert-Gangneux, F., Year, H., D'Herve, D., Guiguen, C. (2009). Congenital toxoplasmosis after a

preconceptional or periconceptional maternal infection. *Pediatr Infect. Dis. J.* 28, 660-661

Ryan, K.J. and Ray, C.G. (2004). *Sherris Medical Microbiology* (4th ed.) McGraw Hill. New York, pp. 723-727.

Silva, C.H., de Andrade, G.Q., Januário, J.N., Carneiro, A.C.A.V., Carneiro, M., Vasconcelos-Santos, D.V., Vitor, R.W. (2012). Early diagnosis of congenital toxoplasmosis in newborn infants using IgG subclasses against two *Toxoplasma gondii* recombinant proteins. *Mem Inst Oswaldo Cruz, Rio de Janeiro*, 107(3), 342-347.

Sroka, J., Wójcik-Fatla, A. and Dutkiewicz, J. (2006). Occurrence of *Toxoplasma gondii* in water from Wells located on farms. *Ann Agric Environ Med*, 13, 169–175.

Sukthana, Y., Chintana, T., Damrongkitchaiporn, S., Lekkla, A. (2001). Serological study of *Toxoplasma gondii* in kidney recipients. *J Med Assoc Thai* 84, 1137-1141.

Tenter, A.M., Heckeroth, A.R., and Weiss, L.M. (2000). *Toxoplasma gondii*: from animals to humans. *International Journal of Parasitology* 30, 1217–1258

Vaz, R.S., Thomaz-Soccol, V., Sumikawa, E., Guimarães, A.T.B. (2010). Serological prevalence of *Toxoplasma gondii* antibodies in pregnant women from Southern Brazil, *Parasitology Research* 106, 661–665. DOI 10.1007/s00436-009-1716-2

Table 1. Seroprevalence of toxoplasmosis among pregnant women, immunocompromised and immunocompetent patients (control) attending national hospital, Abuja.

Group	No Examined	No. Positive	Prevalence (%)
Pregnant	72	32	44.4
Immunocompromised patients	72	20	27.8
Control	72	16	22.2
Total	216	68	31.5

Table 2: Seroprevalence of toxoplasmosis among the age related group in pregnant women.

Age Group (years)	No Examined	No. Positive	Prevalence (%)
≤ 20	6	0	0
21-40	54	24	44.4
41+	12	8	66.7
Total	72	32	44.4

Table 3: Seroprevalence of toxoplasmosis among immunocompromised patients in relation to age and sex

age group (yrs)	Males			Females			Total		
	No. exam ^a	No. +ve	Prev (%)	No. exam	No. +ve	Prev (%)	No. exam	No. +ve	Prev (%)
≤ 20	2	0	0	4	0	0	6	0	0
21-40	14	2	14.3	14	4	28.6	28	6	21.4
41+	20	6	30.0	18	8	44.4	38	14	36.8
Total	36	8	22.2	36	12	33.3	72	20	27.8

^aNote: No. exam means number examined, while No. +ve means number of positive cases; Prev (%) stands for prevalence of infection.

Table 4: Seroprevalence of toxoplasmosis among immunocompetent patients in relation to sex and age

Age group (yrs)	Males			Females			Total		
	^a No exam	No +ve	Prevalence (%)	No exam	No +ve	Prevalence (%)	No exam	No +ve	Prevalence (%)
≤ 20	4	0	0	6	0	0	10	0	0
21-40	18	4	22.2	22	8	36.4	40	12	30.0
41 +	10	2	20.0	12	2	16.7	22	4	18.2
Total	32	6	18.8	40	10	25.0	72	16	22.2

^a Note: No. exam means number examined, while No. +ve means number of positive cases

Table 5: Seroprevalence of toxoplasmosis among those living with cats/dogs

Group	Response	No. examined	No. positive cases	Prevalence (%)
Pregnant women	YES	44	24	54.6
	NO	28	8	28.6
Immuno-compromised patients	YES	38	16	42.1
	NO	34	4	11.8
Immuno-competent (control) patients	YES	32	12	37.5
	NO	40	4	10.0
Total	YES	114	52	45.6
	NO	102	16	15.7
	Total	216	68	31.5

Table 6: Assessment of the risk of getting infected with toxoplasmosis between those cohabiting with cats and those who are not.

Subgroups	Infection status	Not living with cat	Living with cat	Odds ratio
Pregnant women	Positive for toxoplasmosis	8	24	3.00
	Negative for toxoplasmosis	20	20	
Immuno-compromised	Positive for toxoplasmosis	4	16	5.46
	Negative for toxoplasmosis	30	22	
Immuno-competent	Positive for toxoplasmosis	4	12	5.36
	Negative for toxoplasmosis	36	20	
Overall	Positive for toxoplasmosis	16	52	4.51
	Negative for toxoplasmosis	86	62	

Table 7: Prevalence of toxoplasmosis in relation to tribe among the pregnant women, immunocompromised patients and immunocompetent (control) patients

Tribe of origin	Number examined	Number of positive cases	Prevalence (%)
Hausa	68	24	35.3
Igbo	46	14	30.4
Yoruba	46	16	34.8
Others	14	4	28.6
Total	216	68	31.5

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

