



## **Comparative Study of Chemical Composition of Three Different Eggplant Fruit Species**

**Amadi Benjamin<sup>1\*</sup>, Kanu Winner<sup>2</sup>, Nwadike Constance<sup>3</sup>, Ezekwe Ahamefula<sup>4</sup>  
Eboagwu Ijeoma<sup>5</sup>, Onyeabo Chimaraoke<sup>6</sup> and Onuoha Nchekwube<sup>7</sup>**

<sup>1</sup>Department of Biochemistry, University of Port Harcourt, Choba, Nigeria.

<sup>2</sup>Department of Chemical Sciences (Biochemistry Unit), Rhema University, Aba, Nigeria.

<sup>3</sup>Department of Medical Laboratory Science, Imo State University, Owerri, Nigeria.

<sup>4</sup>Department of Medical Biochemistry, Faculty of Basic Medical Sciences, Rivers State University, Nkpolu Oroworokwo, Port Harcourt, Nigeria.

<sup>5</sup>Department of Food Technology, Federal Institute of Industrial Research Oshodi, Nigeria.

<sup>6</sup>Department of Biochemistry, Michael Okpara University of Agriculture, Umudike, Nigeria.

<sup>7</sup>Federal College of Education Technical Asaba Delta State, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors AB, KW, NC and EA designed the study. Authors AB, NC, EA and EI performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors KW, EI, OC and ON managed the analyses of the study. Authors NW, EA, EI, OC and ON managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/AFSJ/2020/v19i330239

#### Editor(s):

(1) Dr. Chouitah Ourida, Professor, University of Mascara, Algeria.

#### Reviewers:

(1) Iwona Rybakowska, Medical University of Gdansk, Poland.

(2) Ernani Clarete Da Silva, Federal University of Sao João del - Rei, Brazil.

(3) Mohamed Fadel, National Research Center, Egypt.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/54271>

**Review Article**

**Received 25 November 2019**

**Accepted 03 February 2020**

**Published 12 December 2020**

### **ABSTRACT**

Chemical composition of three different eggplant fruit species was studied using standard methods. *Solanum aethiopicum*, *Solanum melongena* and *Solanum macrocarpon* fruit species were used as samples. Results obtained showed high moisture, phytate and oxalate contents in the investigated fruit species. Vitamin concentrations of the investigated fruits ranged from high to low. One or two of the investigated fruits possessed a higher amount of the chemical compounds than others. This study has shown the comparative study of the chemical composition of three different eggplant fruit species.

\*Corresponding author: Email: amadiachor@gmail.com;

**Keywords:** Fruit specie; phytochemicals; proximate composition; vitamins.

## 1. INTRODUCTION

The importance of plants and the roles they play in nutrition especially in humans cannot be overstated [1-10]. Studies have shown that products of plants contain chemicals that offer protection to the body [11-19]. In Africa, some plant products are employed as a laxative and in treating many disease conditions. They also serve as medicines and are used in traditional or complementary medicine industry [10,12,19-27].

Different researchers have identified the chemical compounds responsible for the pharmacological action of different parts of plants [28-30]. It is a known fact that most of these compounds that confer protective and medicinal property to them. These compounds possess the characteristic of being biologically active [31-34]. Among such compounds are phytochemicals, vitamins, minerals, amino acids, etc [35-36]. The roles these compounds play in both plants and animals have long been noted [31-34]. Due to the roles of these compounds, there is a need ascertain their presence in plants that come in varieties.

Eggplant is one of such plants that come in varieties with fruits [16,37-38]. Both the leaves and fruits of eggplant are edible. Aside from the leaves and fruits, all parts of eggplant are employed in traditional medicine against one illness or another [26,32]. Most varieties of the fruits that come from a different variety of eggplants are edible. The edible variety of these fruits especially in African continent covers a large group, whereas the major edible species of the fruits are *Solanum aethiopicum*, *Solanum melongena*, and *Solanum macrocarpon*. In Nigeria, the fruits are called "Añara" or "Afufa" or "Mkpuruofe" in Igbo; "Sokoyokoto or "Igba" in Yoruba; and "Gauta" in Hausa. The consumption of these *Solanum* fruits is spread to all parts of the country. The fruits are used during the ceremony, as medicines by herbalists [30,35-36], and in place of kola nuts at homes [26, 32,39-40]. Due to the importance of this variety of *Solanum* fruits within Nigeria, there is need to investigate their active compounds as to advise the people properly on their consumption.

The present study, therefore, compared some chemical composition of *Solanum aethiopicum*,

*Solanum melongena* and *Solanum macrocarpon* fruits.

## 2. MATERIALS AND METHODS

**Sample collection and preparation:** The *Solanum* specie fruit samples used in the present study were purchased from a farm within Owerri Municipal, Imo State, Nigeria and were properly identified by Dr. F. N. Mbagwu of Plant Science and Biotechnology Department, Imo State University, Owerri, Nigeria. Fresh samples of the identified fruits were ground using a simple electric blender. The ground samples were used for analysis.

### 2.1 Chemical Compound Analysis

**Proximate content:** Moisture, ash, lipid and fibre contents were determined according to the procedures of [41]. Nitrogen was determined using kjeldahl method and crude protein value was determined by multiplying the obtained nitrogen value by a factor of 6.25 [41]. The carbohydrate content was obtained by the difference between 100 and sum total of other proximate content of the present study while energy value was obtained using the Atwater factor of 4:9:4 of carbohydrate, lipid and protein respectively by following the procedures of Onyeike et al. [42].

**Anti-nutrients:** Saponins were determined as described by [43], alkaloids of this study were done using the method of [44], tannins were determined using the method of [45], phytate and oxalate contents were done with the methods of [42]; and [46] respectively.

**Vitamins:** The vitamins A, B complex, C and E were screened in the present study with the methods of [47].

## 3. RESULTS AND DISCUSSION

Proximate composition result of the studied fruits showed the highest moisture content (90.30 g/100 g) in *S. macrocarpon*. Moisture content is related to shelf life and susceptibility of the food material to microbial attack [47]. The observed values of moisture in this study followed the order of *S. macrocarpon* > *S. aethiopicum* > *S. melongena*. In general, the moisture contents of the studied fruit specie are high and could mean

low shelf life for the fruits. The observed moisture content values of the investigated fruit species in the present study were higher than the values reported by Agoreyo et al. [48] on two varieties of *Solanum melongena*. The ash content of food materials is related to mineral contents [1,47]. The ash contents of the studied fruit species were low and followed the order *S. melongena* > *S. aethiopicum* > *S. macrocarpon*. Relatively, the high value of ash in *S. melongena* specie against other fruit species in the present study could mean more minerals in the fruit specie than others. The studied fruit species have protein range 1.73-4.40 g/100 g with *S. melongena* specie producing the highest protein when compared to other species. It could be that it can supply this protein to the biological system on consumption. Studies have shown that fibre in foods has numerous advantages such as making stools softer and bulkier; lowering cholesterol levels, reduction of risk of colon cancers, etc [49]. *S. aethiopicum* may be best adapted to perform these functions since it produced the highest crude fibre value of 2.20 g/100 g when compared to other studied fruit species. The observed crude fibre content of *S. aethiopicum* in this study is in line with the earlier report by Shalom et al. [38] on the fruit. *S. macrocarpon* had the least crude fibre content in the present study. The studied fruit species could be classified as poor in carbohydrate contents and this could as well be the cause of the poor energy values of the fruits as observed in the present study. The observed low carbohydrate and energy values of the studied fruit species could be that they are poor in energy-giving compounds.

Anti-nutrients are mostly planted chemicals that protect them. When ingested in animals (humans inclusive), they could be beneficial or deleterious depending on their levels [2-3,6-10,14-31,50]. Observed saponin levels in the present study followed the order *S. aethiopicum* > *S. macrocarpon* > *S. Melongena* (Table 1). Saponins are non-toxic but can generate adverse physiological responses in animals that consume them. Akindahunsi and Salawu [51] noted that saponins exhibit cytotoxic effect and growth inhibition against a variety of cells, making them have anti-inflammatory and anticancer properties. *S. aethiopicum* may be best suited to produce these effects in the system when consumed due to the observed level of saponins in the fruit when compared to other species considered in this study. Alkaloids are ranked as

the most efficient therapeutically significant plant compound [24,52]. They also exhibit marked physiological activity when administered to animals. Some alkaloid compounds have been reported as convulsant alkaloids and cause central nervous system paralysis in animals [53]. Care should, therefore, be taken in the consumption of *S. aethiopicum* fruit, which produced the highest alkaloid content in the present study. Tannins are noted for their astringency and bitterness [3,7,9,14,19,23,25-27]. The bitter taste of *S. melongena* fruit specie could be attributed to its high value of tannic compound as observed in the present study. *S. melongena* fruit specie had the highest phytate and oxalate contents in the present study. Phytates chelate important elements and bind to proteins hence reducing their absorption in the body [47]. Oxalates remove important minerals such as calcium, etc and precipitate them as kidney stone [47]. The presence of saponins, alkaloids, tannins, phytates and oxalates as observed in this study agree with the earlier work of Shalom et al. [38] on proximate and phytochemical analysis of *Solanum aethiopicum* L and *Solanum macrocarpon* L. fruit.

Table 3 reveals the vitamin composition of the investigated fruit species. The investigated fruit species showed low concentrations of vitamin A, B<sub>2</sub> and E. *S. aethiopicum* and *S. melongena* fruit species had moderate concentrations of vitamin B<sub>1</sub>; *S. aethiopicum* and *S. macrocarpon* were moderate in terms of vitamin B<sub>3</sub>; *S. melongena* and *S. macrocarpon* showed moderate concentrations of vitamin B<sub>6</sub>; while vitamin C was highest in *S. aethiopicum* fruit specie. In a related study, Shalom et al. [38] reported high and moderate concentrations of vitamin C in *Solanum aethiopicum* and *Solanum macrocarpon* respectively. The observed vitamins become important when their functions are considered in the system [1,24,54-57]. From the vitamin result of the present study (Table 3), and sequel to the functions of the observed vitamins, *S. aethiopicum* and *S. melongena* fruit species could be against thiamine deficiency; *S. aethiopicum* and *S. macrocarpon* against pellagra disease, *S. melongena* and *S. macrocarpon* may be against deformation of red cells; whereas *S. aethiopicum* specie could be for anti-scurvy following the consumption of the studied fruit species.

**Table 1. Proximate composition of *S. aethiopicum*, *S. melongena* and *S. macrocarpon* fruits**

Proximate content	<i>S. aethiopicum</i>	<i>S. melongena</i>	<i>S. macrocarpon</i>
Moisture content (g/100 g)	86.57±1.01	81.43±0.85	90.30±0.41
Ash(g/100 g)	1.08±0.02	1.94±0.05	0.64±0.11
Protein ( g/100 g)	2.59±0.13	4.40±0.18	1.73±0.09
Crude fibre (g/100 g)	2.20±0.70	1.80±0.12	1.31±0.43
lipid(g/100 g)	0.92±0.01	1.05±0.05	0.70±0.03
Carbohydrate (g/100 g)	6.64±1.06	9.38±0.90	5.32±0.82
Energy value (Kcal/100 g)	45.20±0.17	64.66±1.02	64.88±0.30

Results are means and standard deviations of quadruplicate determinations

**Table 2. Anti-nutrient composition of *S. aethiopicum*, *S. melongena* and *S. macrocarpon* fruits**

Anti-nutrient (mg/100 g)	<i>S. aethiopicum</i>	<i>S. melongena</i>	<i>S. macrocarpon</i>
Saponins	11.69±0.32	5.34±0.13	9.26±0.29
Alkaloids	4.12±0.10	1.16±0.09	1.09±0.04
Tannins	1.07±0.03	9.05±0.11	1.14±0.07
Phytates	17.01±1.04	28.19±1.15	16.43±1.20
Oxalates	38.37±0.12	41.27±0.10	22.01±0.11

Results are means and standard deviations of quadruplicate determinations

**Table 3. Vitamin composition of *S. aethiopicum*, *S. melongena* and *S. macrocarpon* fruits**

Vitamin	<i>S. aethiopicum</i>	<i>S. melongena</i>	<i>S. macrocarpon</i>
Vitamin A	+	+	+
Vitamin B <sub>1</sub>	++	++	+
Vitamin B <sub>2</sub>	+	+	+
Vitamin B <sub>3</sub>	++	+	++
Vitamin B <sub>6</sub>	+	++	++
Vitamin C	+++	++	++
Vitamin E	+	+	+

+++ (high concentration); ++ (moderate concentration); +(low concentration)

#### 4. CONCLUSION

The high phytate and oxalate contents of the investigated fruit species are alarming if attention is drawn to [58]. This observation calls for a further study on the fruit species. The present study has shown the comparative study of chemical compounds present in three different species of the fruit of an eggplant.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Olusanya JO. Essential of food and nutrition.1<sup>st</sup> edition Apex Book Limited, Lagos. 2008;125-130.
2. Amadi BA,Duru MKC, Ayalogu EO. Proximate and anti-nutritional composition of “ntiti-ikpa”, and “ntubiri”, traditional diets of the ikwerre ethnic national in Nigeria. Elixir Food Science 2014;71:24751-24754.
3. Okoroh PN, Duru MKC, Onuoha SC, Amadi BA. Proximate composition, phytochemical and mineral analysis of the fruits of *Ficus capensis*. International Journal of Innovative Research & Development 2019;8(9):83-88.
4. Onibon VO, Abulude FO, Lawal LO. Nutritional and anti-nutritional composition of some Nigeria fruits. Journal of Food Technology. 5(2):120-122.
5. Nwachukwu MI, Duru MKC, Amadi BA, Nwachukwu IO. Comparative evaluation of phytoconstituents, antibacterial activities and proximate contents of fresh, oven-dried uncooked and cooked samples of *Buchholzia coriacea* seed and their effects on hepatocellular integrity International. Journal of Pharmaceutical Science Invention. 2014;3(6):41-4.
6. Nwachukwu MI, Duru MKC, Nwachukwu IO. Antifungal properties and effect of

- fresh, oven-dried uncooked and cooked seeds of *Buchholzia coriacea* on haematology and kidney. Elixir Food Science. 2013;64:19350-19356.
7. Duru M, Amadi C, Ugbogu A, Eze A, Amadi B. Phytochemical, vitamin and proximate composition of *Dacryodes edulis* fruit at different stages of maturation. Asian Journal of Plant Science and Research. 2012;2(4):437-441.
  8. Amadi BA, Arukwe U, Duru MKC, Adindu EA, Ufornewa EC, Odika PC. The effect of fermentation on anti-nutrients, carbohydrates and vitamin contents of *Pentaclethra macrophylla* Seed. International Science Research Journal 2011;3:74-77.
  9. Amadi BA, Arukwe U, Duru MKC, Amadi CT, Adindu EA, Egejuru L, Odika PC. Phytonutrients and antinutrients screening of *D. edulis* fruits at different maturation stages. Journal of Natural Product Plant Resource. 2012;2(4):530-533.
  10. Duru M, Ugbogu A, Amadi B, Odika P, Chima-Ezika R, Anudike J, Osuocha K. Chemical constituents of *Buchholzia coriacea* seed. Proceedings of the 35<sup>th</sup> Annual International Conference, Workshop & Exhibition of Chemical Society of Nigeria. 2012;2:39-45.
  11. Agomuo EN, Amadi BA, Duru MKC. Some biochemical studies on the leaves and fruits of *Persea americana*. International Journal of Research and Reviews in Applied Sciences. 2011;11(3):565-569.
  12. Amadi BA, Duru MKC, Agomuo EN. The chemical profiles of leaf, stem, and flower of *Ageratum conyzoides*. Asian Journal of Plant Sciences and Research. 2012; 2(4):428-43.
  13. Sofowara A. Medicinal plants and traditional medicine in African. Spectrum Books Ltd, Ibadan, Nigeria. 1993;191-289.
  14. Duru M, Eboagwu I, Kalu W, Odika P. Nutritional, anti-nutritional and biochemical studies on the oyster mushroom, *Pleurotus ostreatus*. EC Nutrition. 2019;14(1):36- 59.
  15. Duru M, Nwadike C, Ezekwe A, Nwaogwugwu C, Eboagwu I, Odika P, Njoku S, Chukwudoruo C. Evaluation of nutritional, anti-nutritional and some biochemical studies on *Pleurotus squarrosulus* (Mont.) singer using rats. African Journal of Biochemistry Research. 2018;12(2):7-27.
  16. Duru M, Amadi B, Agomuo E, Eze A. Chemical profile of an anti-malarial concoction "Udu" used in Umunchi autonomous community in Isiala Mbanu L.G.A of Imo State, Nigeria. Journal of Emerging Trends in Engineering and Applied Science. 2012;3(3):444-447.
  17. Duru MKC, Amadi BA, Amadi CT, Lele KC, Anudike JC, Chima-Ezika OR, Osuocha K. Toxic effect of *carica papaya* bark on body weight, haematology, and some biochemical parameters. Biokemistri. 2012; 24(2):67-71.
  18. Amadi BA, Agomuo EN, Duru MKC. Toxicological studies of *Asmina triloba* leaves on haematology, liver, kidney using rat model. International Science Research Journal. 2013;4(2):11-17.
  19. Duru MKC, Agomuo EN, Amadi BA. Biochemical studies on "Udu" an antimalarial concoction used in Umunchi village, Isiala Mbanu L.G.A of Imo State, Nigeria. Continental J. Pharmacology and Toxicology Research. 2012;5(2):28-34.
  20. Monago-Ighorodje C, Duru M, Adindu E, Nwauche K, Ezekwe A, Nosiri I, Odika P, Onyeabo C, Ogar I, Berezi EP, Ugoh AI, Eboagwu I and Otta E. Effect of ethanolic leaf extract of *Vinca major* L. on biochemical parameters and glucose level of alloxan induced diabetic rats. 2019; 18(32):1054-1068.
  21. Duru M, Agomuo E, Amadi B, Iheanacho A, Nutritional evaluation and some biochemical studies on *Ageratum conyzoides* using its different parts. Proceedings of the 35<sup>th</sup> Annual International Conference, Workshop & Exhibition of Chemical Society of Nigeria. 2012;1:372-379.
  22. Nwachukwu MI, Duru MKC, Nwachukwu IO, Obasi CC, Uzoechi AU, Ezenwa CM, Anumodu CK. In-vitro phytochemical characterization and antibacterial activity of *Newbouldia laevis* (boundary tree) on *Escherichia coli* and *Staphylococcus aureus*. Asian Journal of Microbiology and Biotechnology. 2017;2(1):30-36.
  23. Agomuo EN, Duru MKC, Amadi BA. Some bioactive constituents of *Asmina triloba* (paw paw) leaf variety. International Science Research Journal. 2013;4(2):18-22.
  24. Okwu DE. Phytochemicals and vitamin content of indigenous species of South Eastern Nigeria. J. Sustain. Agric Environ. 2004; 6:30-34.
  25. Duru MKC, Arukwe U, Amadi BA. Bioactive constituents and macronutrients composition of anti-malarial concoction

- used in Umunchi village in Isiala Mbanu L.G.A of Imo State, Nigeria. International Science Research Journal. 2011;3:61-64.
26. Amadi B, Onuoha N, Amadi C, Ugbogu A, Duru M. Elemental, amino acid and phytochemical constituents of fruits of three different species of eggplants. International Journal of Medicinal and Aromatic Plants. 2013;3(2):200-202.
  27. Gill LS. Ethnomedical uses of plants in Nigeria. University of Benin Press, Nigeria. 1992;215.
  28. Kubmarawa D, Ajoku G., Enwerem NM, Okorie DA. Preliminary phytochemical and antimicrobial Screening of 50 Medicinal Plants from Nigeria, Afr. J. Biotechnology 2007;6(14):1690-1696.
  29. Adebajo AO, Adewumi, CO, Essein EE. Antiinfective agents of higher plants. International Symposium of Medicinal Plants, 5<sup>th</sup> edn. University of Ife, Nigeria. 1983;152–158.
  30. Schipper RR. African indigenous vegetables: An overview of the cultivated Species chatham, UK: National Resource Institute /ACPEU Technical Center or Agriculture and Rural Cooperation. 2004; 214.
  31. Ugbogu AE, Okezie E, Uche-Ikonne C, Duru M, Atasie OC. Toxicity evaluation of the aqueous stem extracts of *Senna alata* in wistar rats. American Journal of Biomedical Research, 2016;4(4):80-86.
  32. Duru MKC, Amadi BA, Eze AE, Ugbogu AE, Onuoha N. *In vivo* studies of *Solanum aethiopicum* fruit on some biochemical parameters using rats. Journal of Chemical and Pharmaceutical Research. 2013;5(2): 1-4.
  33. Nwachukwu MI, Duru MKC, Nwachukwu IO, Obasi CC, Uzoechi AU, Ezenwa CM, Anumodu CK. *In-vitro* phytochemical characterization and antibacterial activity of *Newbouldia laevis* (boundary tree) on *Escherichia coli* and *Staphylococcus aureus*. Asian Journal of Microbiology and Biotechnology. 2017;2(1):30-36.
  34. Duru M, Amadi B, Ugbogu A, Eze A. Effect of “udu” an antimalarial herbal preparation on visceral organ weight and blood lipid profiles in wistar rats. Journal of Pharmacy and Clinical Sciences. 2014;8:1-7.
  35. Addae-Mensah I. Towards a rational scientific basis for herbal medicine a phytochemist’s two decades contribution. An inaugural lecture delivered at the University of Ghana, Legon, Ghana Universities Press, Accra. 1992;63.
  36. Okoegwale, EE, Olumese GO. Folk medicine practices in Nigeria: some medicinal plants of Esan people in Edo State Nigeria. Niger. J. Appl. Sci. 2001;4: 2350-2358.
  37. Auguste K, Eric B, Tah Yves-Nathan T, Oulo A, Abou BK, Jean-Claude N, Assanvo SPN and Bakary T. Identification of three distinct eggplant subgroups within the *Solanum aethiopicum* Gilo group from Côte d'Ivoire by morpho-agronomic characterization. Agriculture 2014;4:260-273.
  38. Shalom NC, Abayomi CO, Okwuchukwu KE, Opeyemi CE, Olajumoke KA, Damilola ID. Proximate and phytochemical analyses of *Solanum aethiopicum* L. and *Solanum macrocarpon* L. fruits. Research Journal of Chemical Sciences 2011;1(3):13.
  39. Majubwa RO, Msogoya TJ, Maerere AP. Effects of local storage practices on deterioration of African eggplant (*Solanum aethiopicum* L.) fruits. Tanzania Journal of Agricultural Sciences 2015;14(2):106-111.
  40. Okoli RI, Aigbe O, Ohaju-obodo JO, Mensah JK. Medicinal Herbs used for Managing Some Common Ailments among Esan people of Edo State, Nigeria. Pak. J. Nutr. 2007;6(5):490-696.
  41. AOAC. Official Methods of Analysis (15<sup>th</sup> Edn.) Washington D.C USA Association of Official Analytical Chemists Inch 400-22— Wilson Boalevard Artlinton Virginia USA. 1990;2:910-928.
  42. Onyeike, E. Nand Omubo- Dede TT. Effect of heat treatment on the proximate composition, energy values and levels of some toxicants in African yam bean (*Sphenostylis stenocarpum*) seed varieties. Plant Food Resour. 2002;57:233-231.
  43. Oloyed OI. Chemical profile of unripe pulp of *Carica papaya*. Pak. J. Nutr. 2005;4: 379-381.
  44. AOAC Association of Official Analytical Chemists. Official Methods of Analysis (21<sup>st</sup> Edition) Washington DC USA; 1999.
  45. Trease GE, Evans WC. Pharmacognosy. 13<sup>th</sup>. Brailliere Tindall, London. 1989;290.
  46. Krisna G, Ranjhan SK. Laboratory manual for nutritional research, Vikas Publishing House PVT Ltd. Ghaziabad, UP (India); 1980.
  47. Onwuka GI. Food analysis and instrumentation (Theory and Practice). 1<sup>st</sup>

- Edn, Naphthal Prints, Surulere, Lagos – Nigeria. 2005;140-160.
48. Agoreyo BO, Obansa ES, Obanor EO. Comparative nutritional and phytochemical analysis of two varieties of *Solanum melogena*. Science World Journal. 2012; 7(1):5-8.
49. Anderson J, Perryman S, Young L, Prior S. Dietary fiber, health food and nutrition series. No. 9.333. Colorado State University Extension.12/98; Revised 5/07 (www.ext.colostate.edu); 2007.
50. Thompson LU. Potential health effect benefits and problems associated with Anti-nutrients in Foods. Int J. Foods Human Nutr.1993;57:223-231.
51. Akindahunsi AA, Salawu, SO. Phytochemical screening and nutrient-antinutrient composition of selected tropical green leafy vegetables. African Journal of Biotechnology. 2005;4:497-501.
52. Stephen KS. Medicinal plant alkaloids. The University of Toronto Press Canada; 1970.
53. Okwu DE, Ndu CU. Evaluation of the phytonutrients, mineral and vitamin content of some varieties of yam (*Dioscorea sp*). Int. J. Med. Adv. Sci. 2006;2(2):199-203.
54. Amadi BA, Arukwe U, Duru MKC, Adindu EA, Uforonwa EC, Odika PC. The effect of fermentation on anti-nutrients, carbohydrates and vitamin contents of *Pentaclethra macrophylla* Seed. International Science Research Journal. 2011;3:74-77.
55. Uwakwe AA, Anyalogu EO. Biochemistry (A Tropical Approach), Fius Publishers, Port Harcourt, Nigeria. 1998;1:3-5.
56. Paul G, Pearson S. The vitamin 2<sup>nd</sup> Edn., Academic Press, New York, USA. 2005; 31-34.
57. Okaka JC, Akobundu ENJ, Okaka ANC (1992). Human Nutritional and Integrated Approach. ESUT Publishers, Enugu. 1992; 1:190.
58. NRC National Research Council Recommended Dietary Allowance. National Academy press Washington DC; 1989.

© 2020 Benjamin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/54271>